

Veterinary Research Communications

Tat-MyoD FUSED PROTEINS, TOGETHER WITH C2C12 CONDITIONED MEDIUM, ARE ABLE TO INDUCE EQUINE ADULT MESENCHYMAL STEM CELLS TOWARDS THE MYOGENIC FATE.

--Manuscript Draft--

Manuscript Number:	VERC-D-17-00002	
Full Title:	Tat-MyoD FUSED PROTEINS, TOGETHER WITH C2C12 CONDITIONED MEDIUM, ARE ABLE TO INDUCE EQUINE ADULT MESENCHYMAL STEM CELLS TOWARDS THE MYOGENIC FATE.	
Article Type:	Original Article	
Keywords:	Tat-MyoD; equine PB-MSCs; C2C12; coculture; myogenic induction.	
Corresponding Author:	Marco Patrino, PH.D Faculty of Medicine Veterinary Legnaro, Padova ITALY	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Faculty of Medicine Veterinary	
Corresponding Author's Secondary Institution:		
First Author:	Marco Patrino, PH.D	
First Author Secondary Information:		
Order of Authors:	Marco Patrino, PH.D	
	Chiara Gomiero	
	Roberta Sacchetto	
	Ohad Topel	
	Alessandro Negro	
	Tiziana Martinello	
Order of Authors Secondary Information:		
Funding Information:	University of Padua, PRAT 2013 (CPDA138242)	Prof. Marco Patrino
Abstract:	<p>The Tat protein is able to translocate through the plasma membrane and when it is fused with other peptides may act as a protein transduction system. This ability appears particularly interesting to induce tissue-specific differentiation when the Tat protein is associated to transcription factors. In the present work, the potential of the complex Tat-MyoD in inducing equine peripheral blood mesenchymal stem cells (PB-MSCs) towards the myogenic fate, was evaluated. Results showed that the internalization process of Tat-MyoD happens only in serum free conditions and that the nuclear localization of the fused complex is observed after 15 hours of incubation. However, the supplement of Tat-MyoD only was not sufficient to induce myogenesis and, therefore, in order to achieve the myogenic differentiation of PB-MSCs, conditioned medium was added. The latter was obtained coculturing PB-MSCs with C2C12 without direct contact. These results suggest that TAT- transduction of Tat-MyoD, when supported by conditioned medium, represents a useful methodology to induce myoblasts differentiation.</p>	

[Click here to view linked References](#)

1 **Tat-MyoD FUSED PROTEINS, TOGETHER WITH C2C12 CONDITIONED MEDIUM,**
2 **ARE ABLE TO INDUCE EQUINE ADULT MESENCHIMAL STEM CELLS TOWARDS**
3 **THE MYOGENIC FATE.**

4
5
6
7 Marco Patruno ^{a*}, Chiara Gomiero ^a, Roberta Sacchetto ^a, Ohad Topel ^a, Alessandro Negro ^b,
8 Tiziana Martinello ^{a*}.

9
10
11 ^a *Department of Comparative Biomedicine and Food Science, University of Padova, Italy*

12 ^b *VTH - Koret School of Veterinary Medicine, The Robert H. Smith Faculty of Agriculture, Food*
13 *and Environment, The Hebrew University of Jerusalem, Israel*

14 ^c *Department of Biomedical Sciences University of Padova,, Italy*

15
16
17 *= Corresponding authors

18
19 **Abstract**

20
21
22 The Tat protein is able to translocate through the plasma membrane and when it is fused with other
23
24 peptides may act as a protein transduction system. This ability appears particularly interesting to
25
26 induce tissue-specific differentiation when the Tat protein is associated to transcription factors. In
27
28 the present work, the potential of the complex Tat-MyoD in inducing equine peripheral blood
29
30 mesenchymal stem cells (PB-MSCs) towards the myogenic fate, was evaluated. Results showed that
31
32 the internalization process of Tat-MyoD happens only in serum free conditions and that the nuclear
33
34 localization of the fused complex is observed after 15 hours of incubation. However, the
35
36 supplement of Tat-MyoD only was not sufficient to induce myogenesis and, therefore, in order to
37
38 achieve the myogenic differentiation of PB-MSCs, conditioned medium was added. The latter was
39
40 obtained coculturing PB-MSCs with C2C12 without direct contact. These results suggest that TAT-
41
42 transduction of Tat-MyoD, when supported by conditioned medium, represents a useful
43
44 methodology to induce myoblasts differentiation.
45
46
47
48
49
50

51
52 **KEYWORDS:** Tat-MyoD, equine PB-MSCs, C2C12, coculture, myogenic induction.
53
54
55

56
57
58 **1. Introduction**
59
60
61
62
63
64
65

32 Adult skeletal muscle presents a low cellular turnover in the absence of disease or damages
1
23 (Cheung et al 2013). On the contrary, during regenerative mechanisms the muscle tissue becomes
3
4
54 very dynamic thanks to the involvement of satellite cells. The use of these cells for therapeutic
6
75 purpose appears promising for treatment of diseases and injuries affecting skeletal muscle,
8
9
36 including muscular dystrophy (Partridge 2003). Both skeletal muscle injuries and disorders are
10
11
127 actually quite common among athletic animals such as horses (Freestone and Carlson, 1991; Lee et
13
14
1538 al., 2016). However, the self-renewal potential of adult satellite cells is per se limited, decreases
16
1739 with age, sarcopenia (Chen and Goldhamer 2003) and is depleted by wasting muscular dystrophies
18
1940 (Yusuf and Brand-Saberi 2012). Given the need to use an unlimited cell population, mesenchymal
20
21
2241 stem cell (MSCs) deserves a particular attention to offer an alternative therapeutic solution for
23
2442 muscle diseases (Mizuno 2010). MSCs can be isolated from various anatomical districts such as
25
26
2743 bone marrow, adipose tissue, amniotic fluid, peripheral blood (Kuznetsov et al. 2001; Kern et al.
28
2944 2006; Koerner et al. 2006; Martinello et al. 2010; Martinello et al. 2011) and they share the ability
30
31
3245 to differentiate along several pathways (Chamberlain et al. 2007; Giovannini et al. 2008). Up to
33
3446 now, scarce data are present in literature about the differentiation of MSCs into myoblasts. *In vitro*,
35
3647 it has been shown that MSCs may differentiate into skeletal muscle cells with conditioned medium
37
38
3948 as well as in coculture with a fusion between MSCs and myoblasts (Dezawa et al. 2005; Sung et al.
40
4149 2013; Dugan et al. 2014). Specific signaling molecules, such as dexamethasone together with
42
43
4450 insulin and EGF (epidermal growth factor) (Tehrani et al. 2014), are able to induce the
45
4651 differentiation into skeletal muscle. Furthermore, MSCs isolated from bone marrow and treated
47
48
4952 with FGF (Fibroblast Growth Factors), forskolin, PDGF (Platelet-Derived Growth Factor) and
50
5153 transfected with an NICD plasmid were able to express MyoD (Dezawa et al. 2005), although the
52
53
5454 frequency of spontaneous cell fusion was very low. Recently, Rabiee et al. demonstrated that the
55
5655 overexpression of FND5, using an inducible lentivirus system, increased the transcription level for
57
5856 cardiac progenitors in embryonic stem cells (Rabiee et al. 2014) and Sung et al. induced equine
59
60
6157 MyoD expression in equine adipose-derived mesenchymal stem cell using a MyoD lentiviral vector
62
63
64
65

(Sung et al. 2016). Moreover, embryonic stem cells were induced to differentiate also into smooth muscle cells if *Olfm2* (olfactomedin 2) overexpression was promoted (Shi et al. 2014). In a coculture of stem cells from amniotic fluid and cardiac cells, the physical contact between the two types of cells seems to be necessary but not sufficient to induce the cardiogenic potential (Gao et al. 2014); this fact means that a specific microenvironment is required to induce the maturation of myogenic cells. Therefore, the innovative approach of protein transduction with Tat domain fused with various transcription factors (Lin and Kao 2015; Woo et al. 2015), including MyoD (Sung et al. 2013; Hidema et al. 2014), appears to be a valid technical approach. Even though some data indicate that Tat-MyoD induces myogenic differentiation in naturally predisposed cells only, like the C2C12 cell line (Noda et al. 2009) or the mouse muscle primary cells (Hidema et al. 2014) Sung et al. demonstrated that myogenic differentiation of human adipose-derived stem cells was reached using Tat-MyoD transduction when the cells were fused with C2C12 myoblasts (Sung et al. 2013). In the present study, we described that myogenic differentiation of equine peripheral blood mesenchymal stem cells (PB-MSCs) using the Tat-MyoD transduction can be achieved simply with a coculture C2C12 myoblasts. .

2. Materials and methods

2.1. Generation of Tat-MyoD fused proteins

The nucleotide sequence encoding human MyoD was amplified from a human cDNA library with the following oligonucleotides (CAGCTAGCATGTCCTTCGCCATGCTGCGTTCAG - TGCAAGCTTCTAACTTCGAATCGCCGTCTTTTC) and cloned in plasmid Tat-Prp (Vicario et al. 2014) between *NheI* and *HindIII* restriction site, in order to obtain plasmid pTat-MyoD. The plasmid pTAT-MyoD is able to coding for MyoD sequence fused to peptide containing the translocation of HIV-1 protein TAT with 6x Histidine tag at N-terminus.

84 (NH₂-MRGSHHHHHHGMARGYGRKKGRQRRR-).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

85 The plasmid pTat-MyoD was transformed in Escherichia Coli BL21 bacteria cells. The bacteria were
86 grown at 37°C in Luria Broth (LB) medium containing ampicillin (100 µg/ml) to an OD600 of 600
87 nm. Protein expression was induced by adding IPTG (Isopropil-β-D-1-Thiogalactopyranoside)
88 about 4 hours at 25°C. To collect the Tat-MyoD protein, bacteria were harvested and cell membrane
89 was lysed by sonication under denaturing condition using 6 M guanidinium. The proteins were
90 bound to the resin IMAC and then were eluted with 8 M urea and 300 mM imidazole (pH 6.3). The
91 fractions containing the larger quantity of protein were purified using a gel filtration PD10 column
92 (GE Healthcare) to eliminate urea and imidazole. The purified protein was quantified using a
93 spectrophotometer and then an SDS-PAGE was made to verify the purity of Tat-MyoD (44 KDa).
94 The final protein concentration obtained was 0,5 mg/ml.

2.2. Transduction of Tat-MyoD into peripheral blood derived-mesenchymal stem cells (PB- MSCs)

95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165

100 MSCs were isolated from equine peripheral blood (Martinello et al. 2010) and were cultured in GM
101 (growth medium, DMEM Dulbecco's Modified Eagle's Medium, 10% fetal bovine serum FBS, and
102 antibiotics 100 mg/ml streptomycin, 100 U/ml penicillin, Euroclone) at 37°C. In order to evaluate
103 the internalization of Tat-MyoD, PB-MSCs (when reaching confluence) were incubated in the
104 presence of 0,1 µg/ml Tat-MyoD for 2, 6, 15, 24 and 48 hours in medium without serum. The time
105 course analysis was repeated in quadruplicate.

2.3. Coculture of PB-MSCs and C2C12

106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165

108 PB-MSCs and C2C12 cells were cocultured independently by using transwell insert (BD Falcon)
109 with a 1 µm pore size of membrane to separate each cell type. PB-MSCs were plated at the bottom

110 of 6-well plates at concentration of $1,5 \times 10^5$ cells/well in GM and the day after the cells were treated
1
111 with Tat-MyoD for 15h in medium without serum. Concurrently, C2C12 were seeded at density of
3
112 3×10^5 cells per insert in GM, when the cells reached 80% of confluence the medium was changed to
4
6
113 DM (differentiation medium, DMEM, horse serum 2%, antibiotics 1%, Euroclone). After 3 days the
8
9
114 inserts with C2C12 were transferred into the wells with PB-MSCs in DM. The coculture was
10
11
115 maintained for 7 days in DM and the experiment was repeated in triplicate.
13
14
15
16

117 **2.4.Immunostaining**

18
19
20
21
22
119 To perform immunostaining experiments cells were washed with PBS and fixed in 4%
23
24
120 paraformaldehyde for 10 min; after further washing they were permeabilized with 0,3% Triton X-
26
27
121 100 for 5 min and blocked for 1h using 1% FBS. Anti-His tag antibody (1:100, Sigma) was
28
29
122 employed to evaluate the internalization of Tat-MyoD. To evaluate the differentiation of cells, anti-
31
32
123 MyoD (1:100, Santa Cruz), anti-Myf5 (1:100, Santa Cruz) and anti-Myogenin antibodies (1:500,
33
34
124 Chemicon) were used. All antibodies were maintained overnight at 4°C. Fixed cells were washed
35
36
125 with PBS followed by addition of anti-mouse or anti-rabbit Alexa 568 conjugated antibody
38
39
126 (Molecular Probes) at a 1:500 (v/v) dilution. Finally, staining of nuclei was obtained with DAPI
40
41
127 (Sigma). As controls, PB-MSCs treated with Tat-MyoD without coculture and PB-MSCs in
43
44
128 coculture, but without Tat-MyoD treatment, were used.
45
46
47
48

130 **3. Results**

132 **3.1.Purification of Tat-MyoD protein**

133
134 Tat-MyoD was expressed in *E. Coli* B121 and purification was performed using a Ni-NTA column.
60
135 Tat-MyoD purified to homogeneity shows and apparent molecular weight of 44 KDa on SDS-
61
62
63
64
65

136 PAGE and migrate on gel slower respect its theoretical molecular weight of 37905.1 Da (Fig. 1).
1
137 This common behavior may be explained due to the high number of basic amino acids (17.2%
2
3
4
138 respect to total amino acids).
5
6

139 140 **3.2. Localization of Tat-MyoD into PB-MSCs**

141
142 In order to evaluate the cellular pathway of Tat-MyoD protein construct, an immunofluorescence
143 assay was chosen (Fig. 2). Using confocal microscopy, it was found that after 2 and 6 hours of PB-
144 MSCs treatment with MyoD-Tat, the protein permeated cell membrane and was present in the
145 cytoplasm; only after 15 hours of incubation, the construct was confined in the nucleus and this
146 localization was persistent after 24 and 48 hours of treatment (Fig. 2). Experiments were performed
147 in serum free medium since the latter inhibits this process (data not shown).
148
149

150 151 **3.3. Myogenic differentiation of PB-MSCs**

152 Myogenic differentiation was achieved using Tat-MyoD transduction and the inductive medium of
153 the cellular line C2C12. To study the effect of our set up on myogenic marker expression in PB-
154 MSCs, we performed an indirect coculture using transwell insert (Fig. 3B). The scheme of
155 experiment is illustrated in Figure 3A. The effective differentiation was evaluated observing the
156 localization of Myf5 and Myogenin by immunofluorescence (Fig. 4). Results indicated that to
157 activate the myogenic pathway in mesenchymal stem cells it was necessary the co-action of MyoD
158 transduction and the molecular signals present in the medium of C2C12. Figure 4 (A, B) shows
159 Myf5 and Myogenin expression in PB-MSCs treated for 15 hours with Tat-MyoD in serum free
160 medium and, subsequently grown for 7 days in coculture with C2C12 myotubes in differentiative
161 medium. The myogenic differentiation of PB-MSCs was not achieved using, separately, Tat-MyoD
162
163
164
165

161 (Fig. 4 D, E) or the C2C12 conditioned medium (Fig. 4 G, H). Fig 4C and 4F show the internal
1
162 localization of Tat-MyoD complex by means of His-Tag antibody and fig. 4I confirms the absence
3
4
163 of myogenic differentiation with only C2C12 conditioned medium with the use of MyoD antibody.
5
6
7
164
9

10 **4. Discussion**

1165 The equine model offers a unique opportunity to explore treatment strategies for musculoskeletal
12
13
1466 disorders under conditions similar to the pathophysiology of human patients. Current treatments are
15
16
167 often restricted to the management of symptoms or replacement with inert materials; therefore,
17
18
168 there is a need for alternative biological approaches. MSCs may differentiate into cell types relevant
20
21
169 to amend musculoskeletal diseases (Gupta et al. 2007; Lee et al. 2011; Galli et al. 2014) and are
22
23
170 able to secrete growth factors to promote a repairing environment. However, for cell therapy
24
25
171 purposes is necessary that MSCs are able to participate in the formation of new muscle fibers, a
26
27
28
172 critical process that has not been fully elucidated so far. In vitro, hASCs (Human adipose-derived
29
30
3173 stem cells) treated with 5-azacytidine and fibroblast growth factor-2 (FGF-2) stimulates the early
32
33
174 muscle differentiation steps (Eom et al. 2011); more, the expression of MyoD using high efficient
34
35
3675 lentiviral transduction induces myogenic differentiation while adipogenic differentiation is inhibited
37
38
176 (Goudenege et al. 2009). Moreover, using MyoD lentiviral vector Sung et al. induced the expression
39
40
177 of MyoD but not of Myogenin, (Sung et al. 2016). However, these methods are not appropriated for
41
42
178 clinical use due to their mutagenic potential. In the last decade, several groups have demonstrated
43
44
179 that the Tat protein transduction domain (PTD) is a great transactivator of gene expression (Dietz
45
46
180 and Bähr 2004; Fittipaldi and Giacca 2005); its short amino acid motif, highly enriched in basic
47
48
181 amino acids, binds to the cell surface and internalize in a variety of different cell types. In the recent
49
50
182 past, various cellular proteins were described to interact with Tat and mediate or control its
51
52
183 transcriptional activity (Kashanchi et al. 1996; Benkirane et al. 1998; Marzio et al. 1998; Col et al.
53
54
184 2001). In the present study, the human MyoD protein was engineered with the Tat sequence in order
55
56
57
58
59
60
185
61
62
63
64
65

186 to evaluate a safe method for the induction of mesenchymal stem cells towards the myogenic
1
187 differentiation. This approach was already proposed in cells that naturally follow the myogenic fate,
2
3
4
188 as mouse myogenic primary cells (Noda et al. 2009) and C2C12 cell line (Hidema et al. 2012) but
5
6
189 to our knowledge was never tried on PB-MSCs. Additionally, experiments from Sung et al. (2013)
7
8
190 underlines the importance of the extracellular environment, as they were able to differentiated
9
10
11
191 human adipose-derived stem cells into myogenic cells using a fusion with C2C12 cells.
12
13
192 We were successful in inducing myoblasts differentiation in PB-MSCs. Our experiment indicates
14
15
193 that the development of myogenic phenotypes of mesenchymal stem cells by Tat-MyoD construct
16
17
194 depends on time and culture conditions, highlighting the role of *in vitro* microenvironment in terms
18
19
20
21
195 of secreted factors and cell contacts.
22
23
196 Indeed, an important observation raised from our experiments was the necessity to add Tat-MyoD
24
25
197 in a cell culture with serum free medium. It has been demonstrated that short peptides (Green and
26
27
298 Loewenstein 1988) rich in arginine (Suzuki et al. 2002) are rapidly internalized by cells, in a
28
29
30
31
199 receptor-independent manner and without energy consumption. This does not happen for Tat basic
32
33
200 domain when fused to protein cargos (Fittipaldi and Giacca 2005). It was suggested that the process
34
35
36
201 of Tat internalization occurs through adsorptive endocytosis. Several investigators (Hakansson et al.
37
38
202 2001; Mann and Frankel 1991) state that Tat sequence binds homologue of heparin sulfate (HS)
39
40
203 glycosaminoglycan (GAG), a major constituent of extracellular matrix, suggesting that the bound
41
42
204 HS/Tat might be involved in the internalization process. In accordance with this hypothesis, our
43
44
205 study suggests that the presence of heparin in serum competes with the bound of HS/Tat, decreasing
45
46
47
206 the uptake progression. To stimulate myogenic differentiation, Tat-MyoD has to be localized in the
48
49
50
207 nucleus. Our results demonstrated that after 2 and 6 hours the construct remained in the cytoplasm,
51
52
208 probably in vesicle as hypothesized by (Noda et al. 2009). Only after 15 hrs of incubation, Tat-
53
54
55
209 MyoD was localized in the nucleus where it persisted after 24 and 48 hrs. However, the activation
56
57
210 of myogenic pathway by nuclear MyoD was not sufficient to induce cellular differentiation..
58
59
60
61
62
63
64
65

211 Likewise PB-MSCs cocultured with C2C12 grown in cell insert (prevent the cell direct contact but
1
212 permits the interaction of culture medium) was not enough to induce the myogenic commitment.
2
3
4
213 To our knowledge, this is the first study that shows a myogenic differentiation in equine adult stem
5
6
214 cells using the TAT-mediated protein transduction system; the advantage of our method consists in
7
8
215 obtaining committed myogenic cells derived from an abundant cell source, as PB-MSCs, without
9
10
11
1216 the need of fusion with other cells. It is important to state that our model might easily be reproduced
13
14
217 also in human mesenchymal stem cells too (Martinello et al, unpublished results) although further
15
16
1718 studies will be necessary to develop this methodology for clinical purposes.
18
19
20
21

220 **Acknowledgments**

221 We thank Prof. Anthea Rowleron (King's College London, UK) for manuscript language
22
23
24
2521 revision. This work was supported by a grant from the University of Padova, Italy (PRAT 2013,
26
27
2822 code number CPDA138242).
29
30
31
32
33
34

3525 **Conflict of Interest:** None
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

234
1
2
235
4
5
236
7
8
9
237
11
238
13
14
239
15
16
240
18
19
241
20
21
242
23
24
243
25
26
244
28
29
245
30
31
246
32
33
247
35
36
248
37
38
249
40
41
250
42
43
251
45
46
252
47
48
253
49
50
254
52
53
255
54
55
256
57
58
59
60
61
62
63
64
65

References

1. Benkirane M, Chun RF, Xiao H, Ogryzko VV, Howard BH, Nakatani Y, Jeang K (1998) Activation of Integrated Provirus Requires Histone Acetyltransferase. *J Biol Chem* 273, 24898–24905
2. Bentzinger CF, Wang YX, Rudnicki MA (2012) Building muscle: Molecular regulation of myogenesis. *Cold Spring Harb Perspect Biol* 4, a008342
3. Chamberlain G, Fox J, Ashton B, Middleton J (2007) Concise review: mesenchymal stem cells: their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem Cells* 25, 2739–2749
4. Chen JCJ, Goldhamer DJ (2003) Skeletal muscle stem cells. *Reprod Biol Endocrinol* 13, 1:101
5. Cheung TH, Rando TA (2013) Molecular regulation of stem cell quiescence. *Nat Rev Mol Cell Biol* 14, 329–340
6. Col E, Caron C, Seigneurin-Berny D, Gracia J, Favier A, Khochbin S (2001) The histone acetyltransferase, hGCN5, interacts with and acetylates the HIV transactivator, Tat. *J Biol Chem* 276, 28179–28184
7. Dezawa M, Ishikawa H, Itokazu Y, Yoshihara T, Hoshino M, Takeda S, Chizuka I, Nabeshima Y (2005) Bone marrow stromal cells generate muscle cells and repair muscle degeneration. *Science* 309, 314–317
8. Dietz GPH, Bähr M (2004) Delivery of bioactive molecules into the cell: the Trojan horse approach. *Mol Cell Neurosci* 27, 85–131

- 257 9. Dugan JM, Cartmell SH, Gough JE (2014) Uniaxial cyclic strain of human
1 mesenchymal stem cells and C2C12 myoblasts in coculture. *J Tissue Eng* 5,
258 2041731414530138
3
4
259 2041731414530138
5
6
260 10. Eom YW, Lee JE, Yang MS, Jang IK, Kim HE, Lee DH, Kim YJ, Park WJ, Kong JH, Shim
7
8
261 KY, Lee JI, Kim HS (2011) Effective myotube formation in human adipose tissue-derived
9
10 stem cells expressing dystrophin and myosin heavy chain by cellular fusion with mouse
11
12 stem cells expressing dystrophin and myosin heavy chain by cellular fusion with mouse
13
14 C2C12 myoblasts. *Biochem Biophys Res Commun* 408, 167–173
15
16
264 11. Fittipaldi A, Giacca M (2005) Transcellular protein transduction using the Tat protein of
17
18 HIV-1. *Adv Drug Deliv Rev* 57, 597-608
19
20
265 12. Freestone JF, Carlson GR (1991) Muscle disorders in the horse: a retrospective study.
21
22
266 Equine veterinary journal 23, 86-90.
23
24
267 13. Galli D, Vitale M, Vaccarezza M (2014) Bone marrow-derived mesenchymal cell
25
26
268 differentiation toward myogenic lineages: facts and perspectives. *BioMed Res Int*
27
28
269 ID762695
29
30
31
270 14. Gao Y, Connell JP, Wadhwa L, Ruano R, Jacot JG (2014) Amniotic Fluid-Derived Stem
32
33
271 Cells Demonstrated Cardiogenic Potential in Indirect Co-culture with Human Cardiac Cells.
34
35
272 An Biomed Eng 42, 2490-2500
36
37
38
273 15. Giovannini S, Brehm W, Mainil-Varlet P, Nestic D (2008) Multilineage differentiation
39
40
274 potential of equine blood-derived fibroblast-like cells. *Differentiation* 76, 118–129
41
42
275 16. Goudenege S, Pisani DF, Wdziekonski B, Di Santo JP, Bagnis C, Dani C, Dechesne CA
43
44
276 (2009) Enhancement of myogenic and muscle repair capacities of human adipose-derived
45
46 stem cells with forced expression of MyoD. *Mol Ther* 17, 1064–1072
47
48
277 17. Green M, Loewenstein PM (1988) Autonomous functional domains of chemically
49
50
278 synthesized human immunodeficiency virus Tat trans-activator protein. *Cell* 55, 1179–
51
52
279 1188
53
54
55
280
56
57
281
58
59
60
61
62
63
64
65

- 282 18. Gupta A, Leong DT, Bai HF, Singh SB, Lim TC, Hutmacher DW (2007) Osteo-maturation
1 of adipose-derived stem cells required the combined action of vitamin D3, beta-
283 2 of adipose-derived stem cells required the combined action of vitamin D3, beta-
3 glycerophosphate, and ascorbic acid. *Biochem Biophys Res Commun* 362, 17–24
4
284 3 glycerophosphate, and ascorbic acid. *Biochem Biophys Res Commun* 362, 17–24
5
6
285 19. Hakansson S, Jacobs A, Caffrey M (2001) Heparin binding by the HIV-1 Tat protein
7
8 transduction domain. *Protein Sci* 10, 2138–2139
9
286 10 transduction domain. *Protein Sci* 10, 2138–2139
11
12
287 20. Hidema S, Tonomura Y, Date S, Nishimori K (2012) Effects of protein transduction with
13
14 intact myogenic transcription factors tagged with HIV-1 Tat-PTD (T-PTD) on myogenic
15
288 15 intact myogenic transcription factors tagged with HIV-1 Tat-PTD (T-PTD) on myogenic
16
17 differentiation of mouse primary cells. *J Biosci Bioeng* 113, 5–11
18
19
290 21. Kashanchi F, Khleif SN, Duvall JF, Sadaie MR, Radonovich MF, Cho M, Martin MA, Chen
20
21 SY, Weinmann R, Brady JN (1996) Interaction of human immunodeficiency virus type 1
22
291 21 SY, Weinmann R, Brady JN (1996) Interaction of human immunodeficiency virus type 1
23
24 Tat with a unique site of TFIID inhibits negative cofactor Dr1 and stabilizes the TFIID-
25
292 24 Tat with a unique site of TFIID inhibits negative cofactor Dr1 and stabilizes the TFIID-
26
27 TFIIA complex. *J Virol* 70, 5503–5510
28
29
294 22. Kern S, Eichler H, Stoeve J, Klüter H, Bieback K (2006) Comparative analysis of
30
31 mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem*
32
295 31 mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem*
33
34
296 32 mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue. *Stem*
35
36
297 33 Cells 24, 1294–1301
37
38
298 34 Cells 24, 1294–1301
39
40
299 35 Cells 24, 1294–1301
41
42
300 36 Cells 24, 1294–1301
43
44
301 37 Cells 24, 1294–1301
45
46
302 38 Cells 24, 1294–1301
47
48
303 39 Cells 24, 1294–1301
49
50
304 40 Cells 24, 1294–1301
51
52
305 41 Cells 24, 1294–1301
53
54
55
56
57
58
59
60
61
62
63
64
65

- 306 26. Lee EM, Kim AY, Lee EJ, Park JK, Park SI, Cho SG, Kim HK, Kim SY, Jeong KS (2016)
1
307 Generation of Equine-Induced Pluripotent Stem Cells and Analysis of Their Therapeutic
3
4
308 Potential for Muscle Injuries. *Cell transplantation* 25, 2003-2016.
5
6
309 27. Lin BY, Kao MC (2015) Therapeutic applications of the Tat-mediated protein transduction
8
9
310 system for complex I deficiency and other mitochondrial diseases. *Ann N Y Acad Sci* 1350,
10
11
311 17–28.
13
14
312 28. Mann DA, Frankel AD (1991) Endocytosis and targeting of exogenous HIV-1 Tat protein,
15
16
313 *EMBO J* 10, 1733–1739
18
19
314 29. Martinello T, Bronzini I, Maccatrozzo L, Iacopetti I, Sampaolesi M, Mascarello F, Patruno
20
21
315 M (2010) Cryopreservation does not affect the stem characteristics of multipotent cells
23
24
316 isolated from equine peripheral blood. *Tissue Eng Part C Methods* 16, 771–781
25
26
317 30. Martinello T, Bronzini I, Maccatrozzo L, Mollo A, Sampaolesi M, Mascarello F,
27
28
318 Decaminada M, Patruno M (2011) Canine adipose-derived-mesenchymal stem cells do not
30
31
319 lose stem features after a long-term cryopreservation. *Res Vet Sci* 91, 18–24
32
33
320 31. Marzio G, Tyagi M, Gutierrez MI, Giacca M (1998) HIV-1 Tat transactivator recruits p300
35
36
321 and CREB-binding protein histone acetyltransferases to the viral promoter. *PNAS* 95,
37
38
322 13519–13524
40
41
323 32. Mizuno H (2010) The potential for treatment of skeletal muscle disorders with adipose-
42
43
324 derived stem cells. *Curr Stem Cell Res Ther* 5, 133-136
45
46
325 33. Noda T, Fujino T, Mie M, Kobatake E (2009) *Biochemical and Biophysical Research*
47
48
326 *Communications* Transduction of MyoD protein into myoblasts induces myogenic
49
50
327 differentiation without addition of protein transduction domain. *Biochem Biophys Res*
52
53
328 *Commun* 382, 473–477
54
55
329 34. Partridge TA (2003) Stem cell route to neuromuscular therapies. *Muscle Nerve* 27, 133–141
57
58
330 35. Rabiee F, Forouzanfar M, Ghazvini Zadegan F, Tanhaei S, Ghaedi K, Motovali Bashi M,
59
60
331 Baharvand H, Nasr-Esfahani MH (2014) Induced expression of Fndc5 significantly
62
63
64
65

- 332 increased cardiomyocyte differentiation rate of mouse embryonic stem cells. *Gene* 551,
1 127–137
- 333
3 36. Shi N, Guo X, Chen SY (2014) Olfactomedin 2, a novel regulator for transforming growth
4
334 factor- β -induced smooth muscle differentiation of human embryonic stem cell-derived
6
335 mesenchymal cells. *Mol Biol Cell* 15, 4106-4114
8
336
10
11
1337 37. Sung MS, Mun JY, Kwon O, Kwon KS, Oh DB (2013) Efficient myogenic differentiation
13
14
338 of human adipose-derived stem cells by the transduction of engineered MyoD protein.
15
16
1339 *Biochem Biophys Res Commun* 437, 156–161
18
19
340 38. Sung SE, Hwang M, Kim AY, Lee EM, Lee EJ, Hwang SK, Kim SY, Kim HK, Jeong KS
20
21
341 (2016). MYOD overexpressed equine adipose-derived stem cells enhanced myogenic
23
24
342 differentiation potential. *Cell Transplant* Nov;25(11):2017-2026.
25
26
343 39. Suzuki T, Futaki S, Niwa M, Tanaka S, Ueda K, Sugiura Y (2002) Possible Existence of
27
28
344 Common Internalization Mechanisms among Arginine-rich Peptides. *J Biol Chem* 277,
30
31
345 2437–2443
32
33
346 40. Tehrani HJ, Parivar K, Ai J, Kajbafzadeh A, Rahbarghazi R, Hashemi M, Sadeghizadeh M
35
36
347 (2014) Effect of Dexamethasone, Insulin and EGF on the Myogenic Potential on Human
37
38
348 Endometrial Stem Cell. *Iran J Pharm Res* 13, 659–664
40
41
349 41. Vicario M, Zagari A, Granata V, Munari F, Mammi S, Bubacco L. Skaper SD, Negro A
42
43
350 (2014) A novel prion protein-tyrosine hydroxylase interaction. *CNS Neurol Disord Drug*
45
46
351 *Targets* 13, 896-908
47
48
352 42. Woo S J, Shin MJ, Kim DW, Jo HS, Yong IJ, Ryu EJ, Cha HJ, Kim SJ, Yeo HJ, Cho SB,
49
50
353 Park JH, Lee CH, Yeo EJ, Choi, YJ, Park S, Im SK, Kim DS, Kwon OS, Park J, Eum WS,
52
53
354 Choi SY (2015) Effects of low doses of Tat-PIM2 protein against hippocampal neuronal cell
54
55
355 survival. *J Neurol Sci* 358, 226-235
57
58
356 43. Yusuf F, Brand-Saberi B (2012) Myogenesis and muscle regeneration. *Histochem Cell Biol*
59
60
357 138, 187–199
62
63
64
65

358
1
359
3
4
360
6
7
361
9
362
11
12
363
14
15
364
16
17
365
19
366
21
22
367
24
368
26
369
28
370
31
371
33
372
36
373
38
39
374
41
375
43
376
46
377
48
378
50
379
53
380
55
381
58
59
60
61
62
63
64
65

Figure Legends

Fig. 1. Purification of Tat-MyoD by Ni-NTA column. Lane 1, BL21 cell and Sumo-hyrudin. Lane 2, BL21 and pTat-MyoD before induction. Lane 3, BL21 and pTat-MyoD after induction with 0.5 mM IPTG. Lane 4, Purified Tat-MyoD after Ni-NTA column.

Fig. 2. Immunofluorescence analysis of PB-MSCs treated with Tat-MyoD for 2, 6, 15, 24 and 48 hours using the anti-His Tag antibody (red) and DAPI (blue). From 15 hours of incubation anti-His Tag and DAPI colocalized. Bottom right image shows PB-MSCs after 48 hours of Tat-MyoD incubation (PC = Phase contrast). Scale bars: 58 μ m

Fig. 3. (A) Scheme of coculture between PB-MSCs treated with Tat-MyoD and C2C12, GM indicates growth medium and DM differentiation medium. (B) Scheme of transwell insert used for the coculture.

Fig. 4. Myogenic differentiation of PB-MSCs. Immunofluorescence of PB-MSCs after the Tat-MyoD treatment and the contemporary coculture with differentiated C2C12 (A, B, C). Immunofluorescence of PB-MSCs after 7 days of Tat-MyoD treatment (D, E, F) and after 7 days of coculture with differentiated C2C12 (G, H, I). The images show the merge between nuclear DAPI

382 staining (blue) and anti-Myf5 (A, D, G), anti-Myogenin (B, E, H), anti-His Tag (C, F), and anti
1
2
383 MyoD (I) antibodies (red staining). Scale bars: 58 μ m.
3

4
384
5

6
385
8

9
386
10

11
387
13

14
388
15

16
389
17

18
390
19

20
391
21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65







