

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/328163220>

I.S.Mu.L.T. Achilles Tendon Ruptures Guidelines

Article · October 2018

DOI: 10.11138/mltj/2018.8.3.310

CITATIONS

0

READS

242

48 authors, including:



Francesco Oliva

University of Rome Tor Vergata

153 PUBLICATIONS 2,552 CITATIONS

[SEE PROFILE](#)



Umberto Tarantino

University of Rome Tor Vergata

226 PUBLICATIONS 1,985 CITATIONS

[SEE PROFILE](#)



Angelo de carli

Sapienza University of Rome

93 PUBLICATIONS 1,167 CITATIONS

[SEE PROFILE](#)



Matteo Baldassarri

Istituto Ortopedico Rizzoli

23 PUBLICATIONS 83 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



University of Rome Tor Vergata "Consolidate the Foundation" grant project titled "Consolidate the Vitamin D Foundation of the Bone-Muscle Crosstalk in Elderly" [View project](#)



Central and peripheral fatigue [View project](#)

I.S.Mu.L.T. Achilles tendon ruptures guidelines

Francesco Oliva¹
Gabriele Bernardi¹
Vincenzo De Luna¹
Pasquale Farsetti¹
Monica Gasparini¹
Emanuela Marsilio¹
Eleonora Piccirilli¹
Umberto Tarantino¹
Clelia Rugiero²
Angelo De Carli²
Edoardo Gaj²
Domenico Lupariello²
Antonio Vadalà²
Matteo Baldassarri³
Roberto Buda³
Simone Natali³
Luca Perazzo³
Michela Bossa⁴
Calogero Foti⁴
Asmaa Mahmoud^{4,16}
Leonardo Pellicciari^{4,19}
Carlo Biz⁵
Ilaria Fantoni⁵
Daniela Buonocore⁶
Pietro Ruggieri⁵
Maurizia Dossena⁶
Carlotta Galeone⁶
Manuela Verrì⁶
Vito Chianca⁷
Anna Collina⁸
Imma Di Lanno⁸
Luigi Di Lorenzo⁹
Francesco Di Pietro¹⁰
Bernardo Innocenti¹¹
Milena Fini¹²
Paolo Finotti¹³
Antonio Frizziero¹³
Jacopo Gamberini¹³
Alfonso Maria Forte¹⁴
Alessio Gaià Via¹⁵
Biagio Moretti¹⁷
Johnny Padulo¹⁸
Pietro Picerno²⁰
Francesca Veronesi²¹
Mario Vetrano²²
Maria Chiara Vulpiani²²
Marcello Zappia²³
Nicola Maffulli²⁴

- ¹ Department of Orthopaedics and Traumatology, University of Rome "Tor Vergata", Rome, Italy
- ² Department of Orthopaedics and Traumatology, "Sapienza" University of Rome, Sant'Andrea Hospital, Rome, Italy
- ³ Department of Orthopaedics and Traumatology, Rizzoli Orthopaedic Institute, Bologna, Italy
- ⁴ Department of Physical and Rehabilitation Medicine, University of Rome "Tor Vergata", Rome, Italy
- ⁵ Orthopaedics Unit, Department of Surgical, Oncologic and Gastroenterological Sciences DiSCOG, University of Padua, Padua, Italy
- ⁶ Department of Biology and Biotechnology, University of Pavia, Pavia, Italy
- ⁷ Department of Advanced Biomedical Sciences, University of Naples "Federico II", Naples, Italy
- ⁸ Department of Diagnostic Imaging, Campolongo Hospital, Eboli (SA), Italy
- ⁹ Rehabilitation Unit, G. Rummo Hospital, Benevento, Italy
- ¹⁰ Department of Diagnostic Imaging, AORNA, Cardarelli Hospital, Naples, Italy
- ¹¹ Department BEAMS (Bio Electro and Mechanical Systems), University of Brussels, Brussels, Belgium
- ¹² Laboratory of Preclinical and Surgical Studies, Rizzoli Orthopaedic Institute, Bologna, Italy
- ¹³ Department of Physical and Rehabilitation Medicine, University of Padua, Padua, Italy
- ¹⁴ Center of Rehabilitation and Biomedical Research, Biomedical Research Center Gruppo Forte, Salerno, Italy
- ¹⁵ Department of Orthopaedics and Traumatology, Hip Surgery Center, IRCCS San Donato Hospital, San Donato Milanese, Milan, Italy
- ¹⁶ Department of Physical Medicine, Rheumatology and Rehabilitation, University of Cairo "Ain Shams, Cairo, Egypt
- ¹⁷ Department of Orthopaedics and Traumatology, Bari Hospital, Bari, Italy
- ¹⁸ Sport Sciences, University e-Campus, Novedrate, Italy; Tunisian Laboratory of Research for Sporty Performance Optimization, National Center of Medicine and Sport Sciences, Tunis, Tunisia
- ¹⁹ Department Health Technical, USL Toscana Center, Empoli (FI), Italy
- ²⁰ Telematics University e-Campus, Novedrate, Italy
- ²¹ Rizzoli Orthopaedic Institute, Bologna, Italy
- ²² Department of Physical and Rehabilitation Medicine, "Sapienza" University of Rome, Sant'Andrea Hospital, Rome, Italy

²³ Department of Medicine and Health Science, University of Molise, Campobasso, Italia; Varelli Institute, Naples, Italy

²⁴ Department of Physical and Rehabilitation Medicine, San Giovanni di Dio e Ruggi d'Aragona Hospital, University of Salerno, Italy; University of London Queen Mary, Barts and the London School of Medicine and Dentistry, Sport Medicine Center, Mile End Hospital, London, UK

Corresponding author:

Francesco Oliva
Department of Orthopaedics and Traumatology,
University of Rome "Tor Vergata"
Viale Oxford 81
00133 Rome, Italy
E-mail: olivafrancesco@hotmail.com

Summary

This work provides easily accessible guidelines for the diagnosis, treatment and rehabilitation of Achilles tendon ruptures. These guidelines could be considered as recommendations for good clinical practice developed through a process of systematic review of the literature and expert opinion, to improve the quality of care for the individual patient and rationalize the use of resources. This work is divided into two sessions: 1) questions about hot topics; 2) answers to the questions following Evidence Based Medicine principles. Despite the frequency of the pathology and the high level of satisfaction achieved in treatment of Achilles tendon ruptures, a global consensus is lacking. In fact, there is not a uniform treatment and rehabilitation protocol used for Achilles tendon ruptures.

KEY WORDS: Achilles tendon ruptures, guidelines.

Introduction

Achilles tendon rupture is the most frequent tendon rupture in the human body^{1,2}. In 85% of patients, the rupture is 2-7 cm proximal to its calcaneal insertion³. Acute ruptures of the Achilles tendon are most frequent in men⁴, 30-40 years old, in particular in weekend athletes who play football, basketball, tennis and squash⁵. Chronic ruptures are defined as an untreated tendon rupture persisting more than 4 weeks³. The incidence changes in the different countries. Re-rupture of the Achilles tendon is failure of its treatment⁶, conservative (12%) or surgical (4%)⁷. The etiology of the Achilles tendon rupture is multifactorial, including intrinsic and extrinsic factors, but the specific role and weight of each of these factors remains unclear (Tab. I).

Methodology

These guidelines are recommendations developed through a process of systematic review of the literature and expert opinion. The recommendations are based on the scientific evidence and clinical experience and can be used to improve the quality of care for individual patients.

The Authors were divided into four groups:

- *Coordinator*: conceived and organized the work with the group of experts.
- *Overseeing group*: controlled the development of the work and discussed the recommendations.
- *Group of experts*: individually received a question and developed the topic according to the rules of Evidence Based Medicine (EBM), when it was possible.
- *Group of preparation and evaluation of literature*: drew up the text and assisted the group of experts in evaluating the literature.

Methods and criteria study selection

For the research were consulted the following databases:

- PubMed;
- Embase;
- Web of Science;
- CINAHL;
- Scopus;
- Google Scholar;
- Cochrane Library.

Using the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, randomized controlled trials (RCTs) and systematic reviews were included; to follow if missing the first two, the other levels of evidence. Date of publications: 1987-November 2017.

Level of evidence

De Vries JG, Berlet GC. *Understanding levels of evidence for scientific communication*. Foot and Ankle Spec. 2010;3(4):205-9 (Tab. II).

Question n. 1: Animal models

The study of the animal models is consequent to the necessity of regenerate the tendon, identify optimal surgical techniques and rehabilitative protocol, accelerate return to work and return to sport.

The main animal models for Achilles tendon studies are mouse, rat and rabbit. The choice of animal model should be based on the type of study: rupture, tendinopathy, healing physiopathology.

Key points

- Animal models allow to study molecular and cellu-

Table I. Extrinsic and intrinsic factors involved in the etiology of Achilles tendon rupture.

Theory	Author	Year
Extrinsic factors		
Mechanical factors	Hunt KJ, et al. ⁸ Józsa L, et al. ⁹ Kannus P, et al. ¹⁰	2014 1989 1997
Drugs	Laseter JT, et al. ¹¹ Khaliq Y, et al. ¹² Parmar C, et al. ¹³	1991 2003 2007
Footwear, ground and type of training	Wertz J, et al. ¹⁴	2012
Intrinsic factors		
Age	Magnusson SP, et al. ¹⁵ McCarthy MM, et al. ¹⁶	2002 2014
Gender	Claessen FMAP, et al. ¹⁷ Hunt KJ, et al. ⁸ Smith FB, et al. ¹⁸ Frizziero A, et al. ¹⁹ Lemoine JK, et al. ²⁰ Cook JL, et al. ²¹	2014 2014 2002 2014 2009 2000
Genetic factors (group ABO)	Józsa L, et al. ²² Kujala UM, et al. ²³	1989 1992
Hormonal factors	Oliva F, et al. ²⁴	2016
Obesity	Battery L, et al. ²⁵	2011
Hypercholesterolemia	Hast MW, et al. ²⁶	2014

lar characteristics and healing physiopathology through quantitative and qualitative analysis, not possible on human.

- Because of the heterogeneity of models and of studies, it is not possible to establish the best suture technique, the best suture material and whether adjuvant therapies ameliorate tendon healing after suture.
- Most animal models do not mimic rupture, but are simple transition models, and are therefore not relevant to the matter at hand.

Level of recommendation: D.

KEY WORDS: Achilles tendon, clinical trials, animal models, surgery, surgical sutures, tendon sutures.

Question n. 2: Clinical diagnosis

The clinical diagnosis is based on history (sudden and severe pain, audible snap), clinical exam in action (swelling, ecchymosis, tendon discontinuity) and clinical

tests. The main clinical tests used are: Calf squeeze sign (Simmond-Thompson test), Single leg heel rise test, Matles test, Copeland test, O'Brien test.

Key points

- Signs and clinical tests recommended are:
 - tendon discontinuity;
 - calf squeeze sign;
 - simmond triad (Matles test, Calf squeeze test, palpable gap).

Level of recommendation: C.

KEY WORDS: clinical test, physical examination, diagnosis, Achilles tendon rupture.

Question n. 3: Ultrasound diagnosis

Ultrasound is used to identify or to confirm Achilles tendon ruptures (both partial and total) and to identify Achilles tendon alterations. Ultrasound is able to identify silent mechanical and structural tendon

Table II. Level of evidence and criteria for analysis.

Level of evidence	Criteria for analysis and inclusion
I	Meta-analyzes and systematic reviews of randomized controlled trials (RCTs) of high quality, or RCTs with minimum or low risk of bias. Systematic reviews of high quality relative to cohort studies or case-control.
II	Cohort studies or randomized case-control high quality, with minimal risk of confounding or bias and with high or discrete probability of causation.
III	Case-control studies and retrospective comparison of well-conducted with reasonable probability of causation.
IV	Non-analytic studies as case series or individual cases

Level of recommendation	Criteria for analysis
A	Supported by at least two studies of level Ib or from a review level Ia ("It was shown")
B	Supported by at least two independent studies of level II or extrapolations from studies of level I ("it is possible")
C	Not supported by adequate studies of level I or II ("indications")
D	Indications of experts ("there is no evidence")

changes which led to rupture. Ultrasound is also used to identify complications after rupture (deep venous thrombosis) and to prevent complications after surgery (identifying sural nerve). It is necessary focused on: patient position, probe position, acoustic window utilized.

Key points

- Ultrasound is useful to diagnose Achilles tendon ruptures, but also to study Achilles tendon characteristics (length, biomechanics, degenerative features) and results after surgery.

Level of recommendation: C.

- Ultrasound is useful to guide to the best choice of treatment.

Level of recommendation: C.

- Ultrasound allows dynamic study. Dynamic study is more sensible than static study to recognize Achilles tendon diseases.

Level of recommendation: B.

- Ultrasound is helpful to recognized degenerative changes in Achilles tendon of asymptomatic athletes and to identify athletes with higher risk of Achilles tendon rupture.

Level of recommendation: C.

KEY WORDS: Achilles tendon, tear, injury, rupture, ultrasonography, ultrasound, sonography, sonoelastography.

Question n. 4: Magnetic resonance diagnosis

Preoperative magnetic resonance (MR) imaging is useful to distinguish partial from complete ruptures and to assess the site and the extent of the tear.

In acute ruptures, the tendon gap demonstrates intermediate signal intensity on T1-weighted images and high signal intensity on T2-weighted images. These findings are consistent with oedema and haemorrhage. In chronic ruptures, scar or fat may replace the tendon.

Key points

- MR is a valid alternative or complementary diagnostic technique.
- MR is recommended to identify or confirm Achilles tendon ruptures and to distinguish acute or chronic ruptures and partial or complete ruptures.

Level of recommendation: C.

KEY WORDS: Achilles tendon, rupture, tear, diagnosis, magnetic resonance, imagine.

Question n. 5: Conservative treatment

The aim of both conservative and surgical treatment is restoring tendon length and tension to optimize force and function. In the last 10 years, the use of conservative treatment has increased in Europe. Modern rehabilitative protocols after conservative treatment are based on early weight bearing concession and early mobilization. However, it is not possible to establish which is the better treatment because of lack of high quality clinically applicable randomized studies.

Key points

- The choice between surgery and conservative treatment should be based on individual factors (age, comorbidities, functional necessity, physical activity, patient preference).

Level of recommendation: A.

- Conservative treatment is recommended if adequate functional rehabilitation is permitted (early mobilization and weight-bearing).

Level of recommendation: B.

- PRP infiltrations and rehabilitation after conservative treatment do not add benefits.

Level of recommendation: C.

KEY WORDS: *Achilles tendon, rupture, conservative, non surgical, non operative, rehabilitation.*

Question n. 6: Sutures and materials

The suture must restore tendon continuity and resistance, allowing tendon glide and preventing adhesions. In addition, the aim of suture is to support mechanical load during rehabilitation, preventing complications and recurrences.

There is lack of randomized clinical trials comparing the different types of sutures and the various techniques. Some studies are discordant on the recommendation of the most adequate technique.

Key points

- The use of absorbable sutures (Vycril, Polydioxanone) is safe because of strength and because of low rate of complications (granuloma, infections).

Level of recommendation: B.

- The choice of the suture technique (es. Bunnell, Kessler, Dresden, Krackow) depends on the experience and on the preference of the surgeon, because of lack of adequate studies.

Level of recommendation: A.

KEY WORDS: *suture, material, Achilles tendon, repair, technique, tendon rupture.*

Question n. 7: Use of autologous derived

The use of platelet-rich plasma (PRP) is started to aid tendon healing. PRP is rich of platelets and of their products such as vascular endothelial growth factor (VEGF), insulin-like growth factor (IGF), fibroblast

growth factor (FGF), platelet-derived growth factor (PDGF), transforming growth factor beta (TGF β) and epidermal growth (EGF). These agents aid regeneration and tissue healing. The biological action of PRP is clear but it is unknown the best application protocol. There is no consensus in literature above the use of PRP in the Achilles tendon ruptures. The existing studies use different protocols, different kinds of PRP, different surgical techniques and different rehabilitation protocols.

Key points

- PRP regenerative capacity is demonstrated.
- Which is the best type of PRP? PRP or PRF (platelet-rich fibrin)? Which is the best application protocol? Is it necessary to associate surgery? Which is the best surgery technique to associate? Which is the best rehabilitation protocol?
- High level of evidence studies are necessary.

Level of recommendation: A.

KEY WORDS: *Achilles tendon, Achilles tendon rupture, mesenchymal stem cells, MSC, PRP, platelet rich plasma, platelet gel, platelet derived growth factors, platelet concentrate, PRGF, platelet lysate, platelet rich fibrin, platelet rich membrane.*

Question n. 8: Open surgery

The open surgical technique allows to directly see the tendon stumps but it mostly damages paratenon and tendon vascularization. The open technique requires less days of hospitalization compared with both conservative treatment and mini-open surgery. Different suture configurations can be utilized in open technique; the most frequently used are Bunnell, Kessler and Krackow. There are contrasting results on ROM, tropism, return to work, and to sport.

It is impossible to define the gold standard treatment of Achilles tendon acute ruptures and the better open suture technique because of lack of high level literature.

Key points

- There are no differences in clinical results after open or percutaneous surgery.
- Open surgery reduces the risk of re-ruptures.
- Open tenorrhaphy requires a longer surgery time and leads to a major rate of complications during wound healing.
- Open surgery is associated with a greater rate of complications, especially infections.
- The treatment choice should be individualised.

Level of recommendation: B.

KEY WORDS: *Achilles tendon acute rupture, open tenorrhaphy, recurrence, complications.*

Question n. 9: Minimally invasive surgery

The complications of the open treatment (infections, adhesions, paresthesia, incision delayed healing)

led to development of mini-invasive and percutaneous techniques. The main mini-invasive techniques studied are mini-open techniques, mini-open Dresden technique, mini-open Kakiuchi technique, Achillon device. The results are satisfactory (rate of complications, return to previous activities, objective and subjective questionnaires, imaging).

The literature does not offer high level studies. Adequate studies are necessary.

Key points

- Mini-invasive surgery techniques, used to treat the acute subcutaneous Achilles tendon ruptures, lead to optimal results and clinical recovery rate is at least 85%.
- Absorbable sutures and the post-surgery weight-bearing reduce the risk of complications.
- The use of PRP in the acute ruptures does not significantly ameliorate clinical and functional outcomes.

Level of recommendation: C.

KEY WORDS: Achilles tendon, rupture, mini-open, repair.

Question n. 10: Percutaneous surgery

Percutaneous techniques consist in no exposition of tendon stumps with intact skin. In this way, the two stumps are approached but not sutured. The first percutaneous technique was described by Ma and Griffith (1977). Subsequently, many modifications were introduced and different instruments used.

Key points

- Percutaneous surgery reduces surgery time and wound complications.

Level of recommendation: A.

- There are no statistically significant difference in clinical outcome between percutaneous and open surgery.

Level of recommendation: A.

- Earlier return to daily activities and to sport.

Level of recommendation: C.

- Higher rate of re-ruptures.

Level of recommendation: C.

- Percutaneous technique leads to a higher rate of sural nerve's lesions than open surgery.

Level of recommendation: A

- Lower rate of infective complications.

Level of recommendation: C.

KEY WORDS: Achilles tendon, tendon rupture, Achilles tendon repair, tendon suture, open repair, percutaneous suture.

Question n. 11: Tendon transfers for chronic tears

Surgery treatment is necessary for the chronic Achilles tendon ruptures because of the retraction of tendon stumps. Tendon transfers are used for the treatment of inveterate Achilles tendon ruptures.

There are different tendon transfer techniques: autograft, allograft, xenograft (based on the source of donor) and flexor hallucis longus, peroneus brevis, gastrocnemius-soleus, fascia lata, semitendinosus, gracilis (based on the donor site). The results are good but randomized controlled clinical trials are necessary.

Key points

- Autograft transfer to treat chronic Achilles tendon ruptures with tendon loss > 50%.

Level of recommendation: A.

- Allograft or xenograft transfer to treat inveterate Achilles tendon ruptures.

Level of recommendation: D.

- Lower rate of return to sport at the same level.

Level of recommendation: A.

- Higher post-surgery outcomes (AOFAS score, calf circumference) after tendon autograft.

Level of recommendation: D.

- Re-ruptures incidence after tendon autograft not statistically significant.

Level of recommendation: D.

- Infection (deep and superficial) incidence of the surgical wound not statistically significant.

Level of recommendation: D.

KEY WORDS: Achilles tendon and transfer, neglected Achilles tendon rupture, chronic Achilles tendon rupture, tendon transfer, Achilles tendon and flexor hallucis longus transfer, Achilles tendon and peroneus brevis tendon transfer.

Question n. 12: Imaging post-surgery

Imaging post-surgery allows to study the intrinsic characteristics of tendon fibers. Follow-up of an operated tendon is clinical. Post-surgery examination can include magnetic resonance imaging (MRI) or Ultrasound (US). Imaging examination may give important information regarding general morphology, tendon structure, grade of vascularisation and tissue mobility. In particular, US plays a crucial role in the follow-up of operated tendons because of the dynamic nature of this technique and the contribution of colour-doppler tool and MRI has shown to be a useful method to evaluate the healing process of surgically treated Achilles tendon. In addition, the use of elastosonography and diffusion tensor imaging (DTI) is increased. Elastosonography and DTI represent innovative and effective quantitative tools that might be able to provide microstructural abnormalities not appreciable using conventional radiological techniques. In last years, the use of DTI in musculoskeletal field keeps on growing in clinical practice. After surgical procedures the use of DTI may ascertain the microstructural properties and integrity restoration of the ruptured tendon during the healing process, even if DTI technique needs more studies on musculoskeletal structures. However, imaging post-surgery appearance of Achilles tendon repair is dependent on the surgical technique used.

Key points

- Imaging post-surgery does not offer clinical and functional benefits.
- Use of DTI allows to have quantitative informations on tendon structure.
- Using Elastography, healing tendons are shown to be softer than healthy tendons.

Level of recommendation: D.

KEY WORDS: *imaging, follow-up, post-surgery, Achilles tendon, rupture, magnetic resonance, ultrasonography.*

Question n. 13: Rehabilitation protocol after acute ruptures

Recently, the rehabilitation regimen after Achilles tendon ruptures has become more active. Immobilization and weight bearing prohibition for 6 weeks has been replaced by functional rehabilitation, characterized by partial or full weight bearing in the first 2 weeks after surgery, and active controlled mobilizations in the first few days after surgery. Functional rehabilitation can include early mobilization or early weight bearing, or both early mobilization and early weight bearing.

Key points

- Functional rehabilitation after surgery is safe and more advantageous than conventional immobilization.

Level of recommendation: A.

- There are no scientific evidences among the best rehabilitation protocol.

Level of recommendation: A.

KEY WORDS: *Achilles, ruptur*, surg*, operat*, mobili*, immobili*, cast*, weight bearing, rehab*, comparison.*

Question n. 14: Rehabilitation protocol after chronic ruptures

The rehabilitation protocol after chronic Achilles tendon ruptures proposed by these guidelines is as follows.

WEEKS 1-4

Cast/Boot (30° plantar flexion), weight-bearing after 3 weeks, cautious mobilizations.

WEEKS 4-8

Complete weight-bearing with cast (5-6 weeks), progressive mobilizations.

WEEKS 8-12

Free deambulation, mobilizations against resistance, cyclette and swimming.

MONTHS 3-6

Sport specific exercises (closed chain), muscular strengthening.

6° MONTH

Jogging, running, jumping and eccentric exercises.

8°-9° MONTH

Return to sport if possible.

Key points

- There are no scientific evidences among the best rehabilitation protocol.

Level of recommendation: A.

KEY WORDS: *Achilles tendon, rehabilitation, program, chronic rupture.*

Question n. 15: Nutraceuticals

The word nutraceutical derived from "nutrition + pharmaceutical". Nutraceuticals are food supplements: L-arginine- α -ketoglutarate, methylsulfonylmethane, type I collagen, bromelain, polyphenols, vitamins (C, A, B6, E), minerals (selenium, zinc), essential fatty acids (omega-3, omega-6). Nutraceuticals can help the normal functions of human body. They have different mechanisms of action: antiinflammatory, analgesic, antioxidant, collagen synthesis promotion, immunomodulation, free radicals scavenging.

Key points

- There are only studies on animal models (studies on human are necessary).
- The use of nutraceuticals, in different combinations, can be helpful to tendon healing and to Achilles tendon rupture prevention, with or without the addition of other strategies.

Level of recommendation: D.

KEY WORDS: *supplement*, nutraceutical*, phytochemicals, extract*, plant, herbal, herbals, glucosamine, glycosaminoglycans, mucopolysaccharides, mucopolisaccharides, glycosaminoglycan polysulphate, glycosaminoglycan polysulfate, chondroitin sulphate, chondroitin sulfate, vitamin C, ascorbate, ascorbic acid, type I collagen, arginine, curcumin, boswellic acid, Boswellia, methylsulfonylmethane, bromelain, tendon*, tendinopathy, tendinitis, Achilles, peritendinitis, tendinitis, tendinosis.*

Question n. 16: Return to sport

Achilles tendon rupture is frequent during sport activities, only 50% of patients return to sport after 1 year. Return to sport is on average 6 months after rupture. 4 of 5 patients return to play after Achilles tendon rupture. Different methods to evaluate function are utilized: AOFAS (American Orthopaedic Foot and Ankle Society Ankle-Hindfoot Score), ARPS (Achilles Rupture Performance Score), ATRS (Achilles Tendon Total Rupture Score), FAAM (Foot and Ankle Ability Measure), FAOS (Foot and Ankle Outcome Score—Ankle and Hindfoot), PAS (Physical Activity Scale), PER (Player Efficiency Rating). Therefore, it is not possible to compare the results of scientific researches.

Key points

- 80% of patients return to sport after Achilles tendon rupture.
- The literature is heterogeneous.
- Scientific evidence about return to play is needed to establish recovery time.

Level of recommendation: D.

KEY WORDS: Achilles tendon and injury, Achilles tendon and rupture, recovery of function or performance outcome, athletic performance, return to play, return to sport, treatment outcome.

Question n. 17: Outcome evaluation devices

There are different types of outcome evaluation devices:

- non invasive laboratory techniques to estimate *in vivo* Achilles tendon force during deambulation;
- movement analysis through methodological and technological instruments: planar trajectories measurement of selected anatomic landmarks, constrain force returned by ground, inertial parameters and muscular geometries evaluation to calculate tendon force through reverse dynamic.

Key points

- AT force during terrestrial human locomotion can be estimated non-invasively through inverse dynamics by means of motion analysis techniques and musculoskeletal modeling.
- Such an approach, although clinical-friendly, presents several limitations due to the reliability of the collected experimental data and to the specificity of musculoskeletal models.
- State-of-the-art high-resolution imaging techniques are being used to record subject-specific musculoskeletal geometries to fit to motion data collected into the laboratory to improve the accuracy in estimating muscle force through inverse dynamics.

Level of recommendation: D.

KEY WORDS: joint kinematics, inverse dynamics, gait analysis, Achilles tendon force, musculoskeletal model.

Question n. 18: Acute ruptures in the childhood

Acute Achilles tendon ruptures in the childhood are rare. The rupture can be initially partial and can become total after few weeks because of a new trauma.

Key points

- In patients under 10 years old treatment can be conservative, with good results.

Level of recommendation: C.

- Chronic ruptures usually require open surgical treatment; if there is a wide gap, autografts can be used to bridge such gap.

Level of recommendation: C.

- Acute ruptures in skeletally mature patients can be treated both surgically (percutaneous technique) or conservative.

Level of recommendation: C.

KEY WORDS: pediatric Achilles tendon tear, pediatric Achilles tendon repair, pediatric Achilles tendon injury.

Answer n. 1: Animal models in Table III.

Answer n. 2: Clinical diagnosis in Table IV.

Answer n. 3: Ultrasound as diagnostic tool in Table V. Ultrasound as outcome measurement to establish treatment validity in Table VI.

Answer n. 4: Magnetic resonance diagnosis

Preoperative MR imaging is useful for distinguishing partial from complete rupture and assessing the site and extent of the tear^{93,94}. At MR, partial tendon tears can be defined on MR images in the sagittal and axial planes demonstrating heterogeneous signal intensity and thickening of the tendon without complete interruption⁹⁵. Longitudinal splits in chronic Achilles tendinopathy that are low to intermediate in signal intensity on long-TR/TE images may be seen in association with a superimposed acute partial tear. Linear or focal regions of increased signal and thickening of fibers without a tendinous gap are characteristic⁹⁵.

Differentiation between partial tear and severe chronic Achilles tendinosis may be difficult apart from clinical history. In general, acute partial tears are often associated with subcutaneous edema, haemorrhage within the Kager fat pad and intratendinous haemorrhage at MR imaging, whereas chronic tendinosis does not usually demonstrate increased subcutaneous or intratendinous signal intensity on T2-weighted images^{96,97}.

Complete Achilles tendon rupture manifests as discontinuity with fraying and retraction of the torn edges of the tendon. In acute rupture, the tendon gap demonstrates intermediate signal intensity on T1-weighted images and high signal intensity on T2-weighted images, findings that are consistent with edema and haemorrhage, whereas in chronic ruptures, scar or fat may replace the tendon⁹⁷.

Key MRI findings include: a fluid-filled gap with or without interposed fat at the tear site in complete tendinous disruptions with discontinuity; fraying or corkscrewing of the tendon edges associated with proximal tendon retraction; in the absence of overlapping tendon edges, no tendon fibers can be seen at the tear site on axial images; tendon disruption with discontinuity and a wavy retracted tendon; associated haemorrhage or edema in intratendinous or peritendinous soft tissues on axial or sagittal images; effacement of Kager's triangle⁹⁵.

The main differential features between partial and complete tears include the following: partial tears demonstrate hyperintense signal with incomplete anterior-to-posterior or posterior-to-anterior extension on fat sat FSE PD images; complete tears demonstrate a hyperintense fluid-filled tendinous gap; tendon rupture usually occurs 2 to 6 cm superior to the os calcis; the size of the rupture varies, based on the degree of tendon retraction; ruptures demonstrate dif-

Table III. Answer n. 1: Animal models.

Authors	Year	Animal	Type of lesion	Type of suture +/- additional techniques
Dogan A, et al. ²⁷	2009	36 Sprague-Dawley rats	Z-plasty	Group 1: suture with 5-0 Ethibond; Group 2: no suture
Lusardi DA, Cain J E ²⁸	1994	24 New Zealand rabbits	Longitudinal	Group 1: 4-0 prolene "horizontal mattress" suture; Group 2: fibrin sealant
Jielile J, et al. ²⁹	2016	135 New Zealand rabbits	Unilateral tenotomy 1.6 cm by calcaneal insertion	Yurt-bone suture method Group 1: suture + cast Group 2: suture + mobilization; Group 3: control
Aydin BK, et al. ³⁰	2015	12 Wistar albino rats	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 4/0 polypropylene Group 1: suture + topic hemostatic agent Group 2: suture only
Dabak TK, et al. ³¹	2015	72 Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 5/0 absorbable. Group 1: single phospholipids injection post-surgery; Group 2: multiple phospholipids injections post-surgery; Group 3: hyaluronic acid injection post-surgery Control group: physiological solution injection
Aliodoust M, et al. ³²	2014	88 Wistar rats with and without diabetes - streptozotocin induced	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 4.0 nylon. Group 1: non diabetics, suture + low-level laser therapy; Group 2: non diabetics, suture; Group 3: diabetics+ suture+ low-level laser therapy; Group 4: diabetics + suture
Gereli A, et al. ³³	2014	21 albino Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 5/0 monofilament polypropylene. Group 1: suture + 0.01 ml solution with organic silicone; Group 2: suture + 0.01 ml physiological solution
Liang JJ, et al. ³⁴	2014	120 Sprague-Dawley rats	Cross sectional, in the half tendon	Modified Bunnell technique with 4-0; Nylon. Group 1: suture + 0,2 ml hyaluronic acid + tenocytes; Group 2: suture + 0,2 ml hyaluronic acid; Group 3: suture + physiological solution
Selek O, et al. ³⁵	2014	40 albino Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 3-0 Ethibond. Group 1: suture + mesenchymal cells; Group 2: suture + physiological solution
Zeytin K, et al. ³⁶	2014	16 albino diabetic Sprague-Dawley rats	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 5-0 monofilament polypropylene. Group 1: suture + perichondral autologous graft with suture 6-0 monofilament polypropylene; Group 2: suture
Hapa O, et al. ³⁷	2013	32 samples of bovine Achilles tendon	Cross sectional, 5 mm by calcaneal insertion	Krackow technique. Group 1: 2 sutures with 2 sutures and 2 locked loops; Group 2: 2 sutures with 2 strands and 4 locked loops; Group 3: 2 sutures with 2 strands and 4 locked loops; Group 4: 2-0 suture with 4 strands and 2 loops

To be continued

Continued from Table III

Huri G, et al. ³⁸	2013	27 Merino Wether sheep	Cross sectional, 2 cm by calcaneal insertion	Group 1: Modified Bunnell technique Endobutton-assisted; Group 2: Krackow technique; Group 3: native tendon
Nouruzian M, et al. ³⁹	2013	33 diabetic streptozotocin-induced Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Kessler technique with 4.0 nylon. Group 1: non diabetics + suture + low-level laser therapy 2.9 J/cm; Group 2: non diabetics+ suture + low-level laser therapy 11.5 J/cm; Group 3: diabetics + suture + low-level laser therapy 2.9 J/cm; Group 4: diabetics + suture+ low-level laser therapy a 11.5 J/cm
Leek BT, et al. ⁴⁰	2012	84 New Zealand rabbits	Cross sectional, partial (50%)	Krackow technique. Group 1: 0-ultrabrade suture impregnated with butyric acid; Group 2: non impregnated
Ni T, et al. ⁴¹	2012	64 adult New Zealand white rabbits	Cross sectional, 1-2 cm by calcaneal insertion	Kessler technique. Group 1: 5-0 vicryl coated + epitendinous suture; Gruppo 2: 5-0 vicryl + 1 cm by section electrospun silk (ES) bounded to tendinous surface + lambda 532 nm and 0.3 W/cm ² irradiated for 6 minutes
Ishiyama N, et al. ⁴²	2011	18 Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Kessler technique with 6-0 braided polyestere + cast. Group 1: suture + injected 2- metha cryloyloxyethyl phosphorylcholine (MPC) polymer 2,5%; Group 2: suture + injected 2-metha cryloyloxyethyl phosphorylcholine (MPC) polymer 5.0; Group 3: suture + physiological solution
Ishiyama N, et al. ⁴³	2010	12 Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Kessler technique with 6-0 braided polyestere + cast. Group 1: suture + injected 2-metha cryloyloxyethyl phosphorylcholine (MPC) polymer 2,5%; Group 2: suture + injected 2- metha cryloyloxyethyl phosphorylcholine (MPC) polymer 5.0; Group 3: suture + physiological solution
Lyras DN, et al. ⁴⁴	2011	48 New Zealand white rabbits	Cross sectional, 2 cm by calcaneal insertion	Paratenon with continuous suture 4-0 nylon. Group1: suture + injected 0.5 ml of PRP distal and proximal tendon insertions; Group 2: suture
Saygi B, et al. ⁴⁵	2008	45 Sprague-Dawley rats	Cross sectional, 5 mm by calcaneal insertion	Kessler technique 3/0 Ethibond. Group 1: suture; Group 2: direct exposition to air + irrigation with 3 drops physiological solution each 5 minutes for 60 minutes + suture; Group 3: exposition to air for 60 minutes + suture
Chong AK, et al. ⁴⁶	2007	57 New Zealand white rabbits	Cross sectional, in the half tendon	Modified Kessler technique with prolene 4-0. Group 1: suture + mesenchymal bone marrow cells in a fibrin carrier; Group 2: suture + fibrin carrier
Gilbert TW, et al. ⁴⁷	2007	12 mongrel dogs	Segmental excision, 1.5 cm in the half tendon	Graft marked with carbonio14 2x3 cm extracellular matrix of intestinal submucosa and suture 4-0 prolene

To be continued

Continued from Table III

Duygulu F, et al. ⁴⁸	2006	22 New Zealand rabbits	Cross sectional, in the half tendon	Modified Kessler technique with 4/0 PDS + cast. Group 1: suture + nicotine subcutaneous injection 3 mg/kg/die; Group 2: suture + physiological solution infusion
Strauch B, et al. ⁴⁹	2006	40 Sprague-Dawley rats	Cross sectional	Modified Kessler technique with 6-0 nylon. Active group: suture + PMF (pulsed-magnetic-field) 2 sessions (30 minutes/die) for 3 weeks; Control group: suture
Bolt P, et al. ⁵⁰	2007	90 Sprague-Dawley rats	Cross sectional, in the half tendon	Horizontal mattress with 6-0 Ticon. Group 1: suture + transfection with adenovirus expressing green fluorescent protein gene (AdGFP); Group 2: suture + transfection with adenovirus expressing humane BMP-14 gene and AdBMP-14; Group 3: suture
Zantop T, et al. ⁵¹	2006	40 chimerical rats expressing fluorescent green protein in all mesenchimal cells	Step 1: placing 7-0 prolene suture loops 2 cm apart in the midsubstance of the tendon. Step 2: the tendon was cut within the suture loops to hold the explanted tendon in place. Step 3: the sutures were finally performer to secure the autologous tendon graft	Two 7-0 Vicryl sutures were placed proximal and distal in the Achilles tendon. A single layer of lyophilized porcine small intestinal sub mucosa (SIS) was secured to the cut ends of the tendon with 7-0 prolene suture. Finally, the graft and the graft was hydrated with saline. Group 1: SIS graft; Group 2: autologous tendon repair
Chan BP, et al. ⁵²	2005	48 Sprague-Dawley adult rats	Cross sectional, 6 mm by calcaneal insertion	Modified Kessler technique + cast + injected Rosa bengala (RB) solution (0.1%) at the extremities lesions. Group 1: suture; Group 2: laser Group 3: RB only; Group 4: photochemical tissue bonding (PTB) treatment (RB + laser)
Kashiwagi K, et al. ⁵³	2004	90 Wistar rats	Cross sectional, 5 mm by calcaneal insertion	Tsuge technique with 5/0 nylon. Control group: suture + local injection of physiological solution; Group 1: suture + local injection of TGF-beta1 10 ng; Group 2: suture + local injection of TGF-beta1 100 ng
Orhan Z, et al. ⁵⁴	2004	48 Wistar albino rats	Cross sectional	Modified Kessler technique. Group 1: suture + shock waves (ESWT) post-surgery; Group 2: suture Group 3: suture + 500 15 KV shock waves in 2 nd day post-surgery
Kazimoğlu C, et al. ⁵⁵	2003	75 Sprague-Dawley rats	3 cm lesion	Group 1: only cutaneous incision; Group 2: lesion 1 cm by calcaneal insertion + cast; Group 3: modified Kessler technique; Group 4: plasty with biodegradable film PCL (poly-e-caprolactone); Group 5: lesion 1 cm distal by half tendon

To be continued

Continued from Table III

Palmes D, et al. ⁵⁶	2002	114 Balb-C mice	Cross sectional, 5 mm by calcaneal insertion	Modified Kirchmayr-Kessler technique. Group 1: equine cast; Group 2: passive mobilization; Group 3: contralateral Achilles tendons
Thermann H, et al. ⁵⁷	2002	105 rabbits	5 longitudinal lesion, 1 cm by calcaneal insertion	Group 1: continuous fascia suture; Group 2: suture with 5/0 plantar flexion; Group 3: 1 mm of fibrin glue
Rickert M, et al. ⁵⁸	2001	80 Sprague-Dawley rats	Cross sectional, 5 mm by calcaneal insertion	Suture with 3 points. Group 1: suture impregnated with growth and differentiation factor-5 (GDF-5); Group 2: suture
Pneumaticos SG, et al. ⁵⁹	2000	24 New Zealand rabbits	Cross sectional, 1-1.5 cm by calcaneal insertion	Krackow technique + immobilization at 90° with Kirschner wire Group 1: 35 days of immobilization; Group 2: 14 days + active mobilization
Owoeye I, et al. ⁶⁰	1987	60 Sprague-Dawley rats	Cross sectional	Suture with 5-0 black silk + glue for K wire fixation. Group 1: suture + anodic electrical stimuli (15 minutes for 2 weeks 75 microA and 10/sec frequency); Group 2: suture + catodic electrical stimuli; Control group: no suture, no electricity
Petrou CG, et al. ⁶¹	2009	42 New Zealand white rabbits	Tenotomy, 3 cm by calcaneal insertion	Absorbable epitendon suture. Group 1: calcitonin 21 IU /kg intramuscularly; Group 2: physiological solution
Fukawa T, et al. ⁶²	2015	24 New Zealand white rabbits	Cross sectional, 2 cm by calcaneal insertion	Paratenon suture with standard technique 4-0 nylon. Group 1: 1.0 ml di PRP application; Group 2: 1.0 ml physiological solution application
Adams SB, et al. ⁶³	2014	54 Sprague Dawley rats	2 Cross sectional lesions, 3 mm by muscle-tendon origin musculo tendine with 3mm segmental tendon excision	Suture type 8. Group 1: suture only; Group 2: suture + mesenchymal cells injection
Irkören S, et al. ⁶⁴	2012	8 New Zealand white rabbits	Cross sectional, 5 mm by calcaneal insertion	Modified Kessler technique with 5/0 monofilament polypropylene. Group 1: suture + perichondral autologous graft by right ear and continuous suture with 6-0 monofilament polypropylene; Group 2: suture only
Meimandi-Parizi A, et al. ⁶⁵	2013	75 White New Zealand rabbits	Longitudinal	Kessler technique with monofilament absorbable 4-0 polydioxanon. Group 1: suture + collagen implant; Group 2: suture only
Oryan A, et al. ⁶⁶	2013	40 white New Zealand rabbits	2 Cross sectional lesions, 5 mm by muscle-tendon origin with 5 mm segmental tendon excision	Kessler technique. Group 1: suture + collagen 3-D structure between tendon stumps; Group 2: suture only
Godbout C, et al. ⁶⁷	2009	12 males C57BL/6 mice	Cross sectional	Technique type 8 with VICRYL 6-0. Group 1: suture + suture impregnated with antibodies which induce thrombocytopenia; Group 2: suture + placebo

Table IV. Answer n. 2: Clinical diagnosis.

Sign/Test	Action	Significance	Sensitivity	Specificity
Tendon discontinuity ⁶⁸⁻⁷⁰	Palpation of the tendon in prone position	Positive if palpable gap is felt	0.73	0.89
Calf squeeze sign ⁶⁹⁻⁷⁰ (Thompson's test)	Compression of the triceps muscle in a prone patient	Positive if the manoeuvre cannot elicit foot plantarflexion	0.96	0.93
Matles's test ⁷¹⁻⁷³	Active knee flexion in the prone position	Positive if knee flexion leads to progressive foot dorsiflexion	0.88	0.85
Simmonds triad ^{74,69}	Association of tendon discontinuity, Thompson's test and Matles test	Positive if all three signs are present	1	

fuse convexity of the anterior margin and enlarged tendon ends at the tear site⁹⁷. We point out, however, that even advanced imaging techniques should be interpreted in the light of clinical findings. In case of diagnostic doubts, the fallback position should be more accurate clinical examination, not just this imaging.

Answer n. 5: Conservative treatment in Tables VII-VIII.

Answer n. 6: Sutures and materials in Table IX.

Answer n. 7: Use of autologous derived blood products in Table X.

Answer n. 8: Open surgery in Table XI.

Answer n. 9: Minimally invasive surgery in Table XII.

Answer n. 10 : Percutaneous surgery in Table XIII.

Answer n. 11: Tendon transfers in Table XIV.

Answer n. 12: Imaging post-surgery

Despite follow-up of an operated tendon is primarily clinical, postoperative examination has been improved by the recent technological progress either on

MRI or on ultrasound that allow better representation of tendon structural specimens. Postoperative imaging appearance of Achilles tendon repair is dependent on the surgical technique used. Imaging examination allows to obtain information regarding: general morphology, tendon structure, grade of vascularity, tissue mobility.

Ultrasound

Ultrasound (US) can be used to follow-up operated tendons²¹⁹ because of the dynamic nature of this technique and the contribution of colour-doppler tool²²⁰⁻²²¹.

Both scans are essential for the correct examination of the treated area and for correct measurement of tendon's dimension. The operated tendon is thicker and wider than a normal ones; its mean thickness is about 10 mm (ranged from 7 to 16 mm) whereas the average diameter of a healthy tendon is 5.4 mm (ranged from 4.0 to 7.9 mm)²²². This progressive increase in size occurs during the first 3-6 months after surgery and gradually decrease in thickness 1 year after surgery^{223,224}.

Fluid collections are suggestive of a poor prognosis if greater than 50% of the affected tendon, and extensive intratendinous calcifications should be considered pathological²²⁵. The contours of the tendon may be irregular with hypoechoic peritendinous area, which may persist for up to 3 months²²⁶, and small hypoechoic areas may surround the stitches into 6-24 months after surgical treatment^{220,224}.

The microvasculature assessment with colour-doppler tool shows newer vessels with higher flow rates during the healing process²²⁷⁻²²⁸; the vascular response may indicate tendon healing with initial high flow vas-

Table V. Answer n. 3: Ultrasound as diagnostic tool.

Author	Type of study	Patients	Type of surgery	Outcome assessment	Results	Conclusions	Level of evidence
Lang TR, et al., 2017 ⁵	Systematic revision	26 articles (20 case studies, 5 case series e 1 prospective not controlled study), 61 participants. 53 patients (88%, 53 of 61 cases): calcaneal tendon involved	Different databases (Medline, CINAHL, Biological Abstracts, AMED, Web of Knowledge, SCOPUS, SportDiscus e EMBASE) utilising words MeSH and free text, combined with the boolean operators (AND, OR). Imaging utilised: MRI, ultrasound B-mode and CT	Not applicable	Complete rupture in 25% of subjects. In the article, qualitative description of tendon thickening (25%), partial or incomplete ruptures (11%), signal intensity (10%), tendon thinning (7%), inflammation and hypoechogenicity	Despite the strong clinical indication for fluoroquinolones, data are not sufficient to define specific structural changes that lead to adverse reactions in the tendon	I
Barford KW, et al., 2015 ⁶	Cross sectional study	19 patients (8 men, 11 women, mean age: 43.4 years old, range of age: 26-63 years old) without previous problems of Achilles tendon	Achilles tendons (both 2 sides) of all patients (dominant side: dx) examined with MRI and ultrasound. Two phases of measurement: identification of anatomical references and measurement of the skin distance with a centimeter. Repeated ultrasound measurements compared with MRI measurements	Not applicable	Intra-operator reliability with ultrasound do not have significantly differences between prove days: ICC 0.96, SEM 4 mm and MDC 10 mm. Inter-operator reliability has a systematic difference between ultrasounds: 2-5 mm ($p = 0.001-0.036$); ICC 0.97, SEM 3 mm e MDC 9 mm. MRI measurement is mean 4 mm longer than ultrasound ($p = 0.001$)	Ultrasound has a good reliability and precision. Comparing groups of healthy people it is possible to identify differences of more than 4 mm. With repeated evaluations it is possible identify differences of more than 10 mm	III

To be continued

Continued from Table V.

<p>Pedersen M, et al., 2012⁷⁷</p>	<p>Systematic revision</p>	<p>8 articles about midtendineous elastosonography <i>in vivo</i> (4 AT)</p>	<p>PubMed e EMBASE were utilised with a free text research</p>	<p>Not applicable</p>	<p>Elastosonography (SEL) results correlate with conventional ultrasound results and with MRI clinical exam. In few articles, elastosonography is more sensible than traditional ultrasound. For muscles, it is founded an important correlation between SEL, ultrasound and MRI, but only an article exists. Sonoelastography discerns between healthy muscles and lesioned and is probably more sensible than ultrasound and MRI to identify early dystrophic changes</p>	<p>Elastosonography is utilised to identify tendon alterations, like ultrasound and RMI. Elastosonography can identify subclinical alterations of the tendon, not visible with conventional ultrasound. Elastosonography could be a supplementar imaging technique to evaluate muscle-skeletal alterations, virtually superior to ultrasound and MRI. Currently it must be considered an experimental exam</p>	<p>I</p>
<p>Fredberg U, et al., 2008⁷⁸</p>	<p>Randomized trial</p>	<p>209 danish professional men footballer (Achilles tendon and patellar tendon)</p>	<p>Experimental group (mean age 25 years old; range of age: 18-37): eccentric prevention and stretching of patellar and Achilles tendons Control group (mean age 25 years old; range of age: 18-38)</p>	<p>Follow-up with ultrasound more than 12 months and accidents registration</p>	<p>Eccentric training and stretching do not reduce the risk of lesions and this risk is higher during season in player with abnormal patellar tendon at the start of the study. Training programme reduces ultrasound abnormalities in patellar tendon, but not in Achilles tendon</p>	<p>With ultrasound, changes of footballer tendons could be diagnosed before coming symptomatic. Eccentric prevention and stretching reduce the risk of ultrasound alterations in patellar tendon, but there is not the reduction of risk of lesions. On the contrary, in asymptomatic footballer with patellar tendons altered at ultrasound ultrasonographically, eccentric prevention and stretching increase the risk of lesions</p>	<p>I</p>
<p>Flavin R, et al., 2007⁷⁹</p>	<p>Cross sectional study</p>	<p>10 healthy men (range of age: 25-30)</p>	<p>All patients analysed with ultrasound</p>	<p>Ultrasound evaluation</p>	<p>Average distance between geographical mapping and clinical points is 2,5 mm (range 0-20 mm)</p>	<p>Good correlation between clinical and ultrasound evaluation</p>	<p>III</p>

To be continued

Continued from Table V.

Ofer N, et al., 2004 ⁸⁰	Cross sectional study	Patients with Achilles tendon rupture	Group A (range of age: 31-57): patients with Achilles tendon rupture ; Group B (range of age 31-56): control healthy people	Ultrasound: automatic test for evaluation of symmetrical proprieties of tendon movement	Result better in post-surgery tendons than in healthy contralateral tendon in the same subjects. In case of traumatic rupture, there is not this effect. So, negative asymmetry of tendon movement can be associated to degenerative or pre-degenerative processes	Objective method, low cost, non invasive and maybe more sensible of non invasive technique	III
Bleakney RR, et al., 2002 ⁸¹	Cross sectional study	72 patients (58 men, 14 women; average age 49.3 years old; range of age 30-82 years old) with clinical diagnosis of Achilles tendon rupture	All patients analysed with ultrasound + 70 control healthy people (same age and gender)	Ultrasound (diameter, echogenicity, presence of calcifications)	Average maximum AP diameter of ruptured tendon is 11,7 mm (SD = 2,10); the normal tendons is on average 5,4 mm (SD = 0,9) and it is on average 4,9 mm (SD = 0,5) ($p < 0,0001$) in the controls. No differences in maximum AP diameter of ruptured tendon depending of the treatment method (conservative, open reparation, percutaneous reparation). 17 patients have hypoechoic areas in the ruptured tendon, 2 patients have hypoechoic areas in their healthy contralateral tendon, 10 patients have calcifications in their ruptured tendon	AP diameter of ruptured tendon is significantly greatest of healthy contralateral tendon. However, if compared with control group, contralateral tendons have a significantly maximum AP diameter and a higher prevalence of intratendinous alterations. This difference can signified a subclinical tendinopathy that can lead to rupture	III

To be continued

Continued from Table V.

<p>Cunndne G, et al., 1996⁸²</p>	<p>Cross sectional study</p>	<p>19 patients (10 men, 9 women; average age 42 years old; range of age: 18-72) with talloidinia in associated with chronic inflammatory arthritis</p>	<p>All patients analysed with ultrasound</p>	<p>Ultrasound</p>	<p>8 patients (2 had previous blinded failed injections) had 11 injections of corticosteroids ultrasound-guided to treat retrocalcaneal bursitis (n=6), plantar fasciitis (n=3) and tibial posterior tenosynovitis (n=2). Ultrasound showed Achilles tendon rupture (n=2), Achilles tendinitis (n=8), tibial posterior tenosynovitis (n=6), peroneus longus tenosynovitis (n=2), retrocalcaneal bursitis (n=13) and plantar fasciitis (n=4). Lost of bone profile (n = 13) is related to osseous erosions on radiographs in all patients, except one. 10 of 11 guided injections lead to complete resolution of talloidinia</p>	<p>The different causes of alloidinia were identify and the ultrasound capacity to provide useful informations to clinical management is confirmed. Ultrasound guided injection of corticosteroids is advantageous, mostly after failure of blinded injection</p>	<p>III</p>
---	------------------------------	--	--	-------------------	---	---	------------

Table VI. Answer n. 3: Ultrasound as outcome measurement to establish treatment validity.

Author	Type of study	Patients	Type of surgery	Outcome assessment	Results	Conclusions	Level of evidence
Eliasson P, et al., 2016 ⁸³	Cross sectional study	23 patients (19 men, 4 women; average age \pm SD: 38 \pm 2.1 years old) with Achilles tendon rupture during sport, surgery	Open surgery and cast (6 weeks)	PET, ultrasound with power doppler (PDUS), evaluation questionnaires (ATRS, VISA-A)	Glucose supply is more elevated in repaired tendon than in intact tendons at all follow-up times (6, 3 and 1,6 time more elevated respectively at 3, 6 and 12 months, $p < 0,001$) and it is also more elevated in the central part of the tendon than at extremities at 3 and 6 months ($p \leq 0,02$), but lower at 12 months ($p = 0,06$). Relative glucose absorption is negatively correlated to ATRS at 6 months after repairation ($r = -0,89$, $p < 0,01$). Flow activity at PDUS is more elevated in repaired tendon than in intact tendon at 3 and 6 months (both $p < 0,05$), but it is normalized at 12 months	Healing process based on metabolic activity and on vascularization, continues for 6 months after lesion when heavy loads on the tendon are allowed. In fact, metabolic activity was high for more than 1 year after lesion despite vascularization normalization	III

To be continued

Continued from Table VI.

Jiellie J, et al., 2016 ⁸⁴	RCT	57 patients with misunderstood Achilles tendon rupture	2 groups: 25 patients (21 men, 4 women; mean age: 31-47) early rehabilitation post-surgery (group EPR) and 32 patients (27 men, 5 women; range of age 29-45) immobilization post-surgery with cast (group PCI)	Leppilahti Score (LSS), ultrasound, computed tomography multislice spiral (TCmS), electromyography	Ultrasound and msTC do not revealed presence of tendon elongation or adhesion. Group PCI have higher post-surgery LSS score, but recovery is slower. Post-surgery complications, such as ankle ankylosis and osteoporosis, are present only in PCI group. In both the groups, cross sectional section of ruptured tendon is wider than section of healthy contralateral tendon. However, comparing cross sectional section of ruptured tendon in the different groups, the section in EPR group is significantly wider than in PCI group ($p < 0.01$)	Compared to immobilization with a cast, early post-surgery rehabilitation leads to a better clinical result and a faster global regeneration of tendon with an ignored tendon lesion	II
---------------------------------------	-----	--	--	--	---	--	----

To be continued

Continued from Table VI.

<p>Busilacchi A, et al., 2016⁸⁵</p>	<p>Perspective study</p>	<p>25 patients (22 men, 3 women) spontaneous subcutaneous Achilles tendon rupture</p>	<p>Percutaneous tenorrhaphy using terephthalate polyethylene. Control group: 30 healthy volunteers (25 men, 5 women) compared for ultrasound and elastonography results</p>	<p>Evaluation questionnaire (ATRS) correlated with ultrasound</p>	<p>Strain index (SI) in the treated tendons shows progressive stiffness, mostly at myotendinous junction and a sutured site, with stiffness significantly higher in both the contralateral tendons and in healthy volunteers. Maximum thickness of treated tendons is at 6 months, with a reduction after 1 year, without return to physiological normality. The better remodelling is at lesion site. Contralateral tendon has a significantly thickness at myotendinous and osteotendinous junctions. Strain index of contralateral tendon is more rigid than physiological values in the control group. ATRS score is better between 6 months and 1 year, negatively related to SI ($p < 0,001$)</p>	<p>Elastosonography demonstrated that Achilles tendon become progressively thicker after surgery during follow-up, while ATRS score is better. Basing on biomechanical evaluation, 1 year after surgery Achilles tendons do not have a "restitutio ad integrum". Elastosonography provides to major qualitative and quantitative information for diagnosis and follow-up in Achilles tendon pathologies and evaluating post-surgery evolution of repaired tissue</p>	<p>II</p>
--	--------------------------	---	---	---	---	--	-----------

To be continued

Continued from Table VI.

Chiu CH, et al., 2013 ⁸⁶	Retrospective study	19 patients (18 men, 1 woman; average age 38.7 years old, range of age: 20-50) with acute Achilles tendon lesion related to sport	Diagnosis: anamnesis, objective exam, ultrasound. Percutaneous repairation endoscopic assisted, post-surgery rehabilitation	Physical exam, ultrasound and magnetic resonance (MRI)	Tendon healing in all patients. All patients were evacuate with ultrasound and 16 patients with MRI to evaluate the level of healing. Final dorsiflexion was 16° and plantar flexion 26°. 95% of patients (18/19) returned to sport at previous level	Percutaneous Achilles tendon repairation, endoscopy assisted, allowed tendon treatment and return to sport after 6 months	III
Jielle J, et al., 2012 ⁸⁷	Retrospective study	107 patients (84 women, 23 women; average age 36.2 years old) with acute Achilles tendon rupture	Surgery: new technique "Pa-bone". Early rehabilitation post-surgery	Achilles tendon rupture score (ATRS), bilateral ultrasound	At ultrasound, cross sectional areas of ruptured tendon are significantly major than in the controlateral tendon	Early post-surgery kinesietherapy after "Pa-bone" surgery technique leads to excellent clinical results and it is useful to Achilles tendon reconstruction	III
Gigante A, et al., 2008 ⁸⁸	RCT	40 patients (36 men, 4 women; average age 40.7 years old; range of age: 20-60) with acute Achilles tendon rupture related to indirect trauma	Open repairation (group A) or percutaneous repairation (group B) (randomization with Casio Scientific Calculator fix-88). Same rehabilitation protocol with minimal differences in immobilization time	Evaluation questionnaire (SF-121), bilateral ultrasound, isokinetic test	Not significantly differences in clinical evaluation, except ankle circumference, that significantly wider in group B. Not significantly differences between the groups in SF-121 questionnaire, ultrasound and isokinetic test	Open and percutaneous techniques are safe and effective for repair of calcaneal tendon ruptures. Both the techniques lead to the same clinical, ultrasonography and isokinetic results	II

To be continued

Continued from Table VI.

<p>Maffulli N, et al., 2003⁸⁹</p>	<p>RCT</p>	<p>45 patients with subcutaneous Achilles tendon rupture diagnosed with clinical evaluation and confirmed with surgery</p>	<p>Group 1 (21 men, 4 women; average age 44 years old; range of age: 31-69): immobilization with ankle in physiological position (equine) for 2 weeks and in neutral position for 4 weeks. Weight bearing if comfortable and progressive increase; Group 2 (24 men, 4 women; average age 43.8 years old; range of age: 30-67): immobilization with ankle in equine for 2 weeks and in neutral position for 2 weeks. Plantar flexion between 4 and 6 weeks after surgery. Weight bearing when ankle is immobilizer in neutral position</p>	<p>Anthropometric evaluation, sural triceps isometric force, evaluation questionnaire, ultrasound</p>	<p>Group 1: few out patients visits, crutches for 2.5 weeks after surgery (group 2: on average 5,7 weeks after surgery) more patients satisfied of surgery. On ultrasound average repaired tendon thickness is 12,1 mm (SD=2), without differences in ruptured tendon thickness, regardless of post-surgery protocol. Not significantly differences between the two groups in isometric resistance</p>	<p>Early weight bearing with plantigrade load is not dangerous to result of reparation after Achilles tendon rupture</p>	<p>II</p>
--	------------	--	---	---	--	--	-----------

To be continued

Continued from Table VI.

Costa ML, et al., 2003 ⁹⁰	RCT	28 patients (24 men, 4 women; average age: 41 years old) unilateral Achilles tendon rupture diagnosed with clinical evaluation	Group A: immediate weight bearing with cast; Group B: weight bearing with traditional plaster	Return to sport, flexion deficit; force deficit, ultrasound	Ultrasound evaluation of tenodesis: not negative effects of early weight bearing. Not significantly wider tendon diameter in group with cast. In group with immediate weight bearing: clinical anthropometric and functional improvements	Ultrasound evaluation confirms absence of deleterious effects on tenodesis	II
Maffulli N, et al., 2003 ⁹¹	RCT	53 patients subcutaneous Achilles tendon rupture diagnoses with clinical evaluation and confirmed with surgery	Group 1: post-surgery immobilization with ankle in equine, early weight bearing cast changed after 2 weeks with ankle in plantar flexion; Group 2: immobilization with ankle in equine, cast changed after 2 weeks, ankle in intermediate position after 4 weeks with weight bearing	Anthropometric evaluation; isometric force of sural triceps, ultrasound evaluation with high temporal resolution and at real time, evaluation questionnaire	Group 1: few outpatients visits, crutches for 2.5 weeks, satisfied of surgery. On ultrasound, average repaired tendon thickness is 12,1 mm, no differences in thickness of ruptured tendon regardless of post-surgery protocol. Not significantly differences between the two groups in isometric resistance	Early weight bearing with plantar flexion do not influence the results of repair after Achilles tendon acute rupture and reduces time necessary to rehabilitation. However, force deficit and muscular atrophy are not prevented	II

To be continued

Continued from Table VI.

Möller M, et al., 2002 ²⁹²	RCT	65 patients (55 men, 10 women; average age 38.6 ± 8.3 years old) with Achilles tendon ruptured	Group A (35 patients): surgery; Group B (30 patients) no surgery	Ultrasound and magnetic resonance	Peritendinous reactions, oedema and deficit only in few patients. Not significantly differences between the two groups, except tendon elongation function, that significantly lower in no surgery group. No correlation between radiological and clinical results, such as muscular force, resistance and range of movement	The role of ultrasound and MRI during healing process after Achilles tendon ruptures is limited, because of a weak correlation with clinical results
---------------------------------------	-----	--	---	-----------------------------------	---	--

cularity within and around repaired tendons and the total blood flow amount consistently and predictably decrease with time²²⁹. The increased vascularity showed by Power Doppler indicated a possible healing progress of repaired Achilles tendon and it persisted until avascular scar formation.

In the last years ultrasound elastosonography increased its diagnostic utility with the introduction of shear wave method (SWE), a non-invasive ultrasonographic imaging technique introduced in 2002 which has the advantage of being operator-independent, reproducible, and quantitative²³⁰.

Healthy Achilles tendons have a hard elastographic pattern, whereas pathologic ones show a reduction in stiffness. After surgical treatment of a complete tear, tendon stiffness pattern gradually increases at 12, 24, and 48 weeks as the wound-healing process continues^{230,231}.

If an Achilles tendon re-rupture is suspected, sonographic diagnosis is more difficult due to the structural characteristics of the tendon, particularly if large fluid collections are present; a dynamic evaluation during ankle flexion and extension is helpful in revealing the gap of tendon discontinuity²²⁴.

Magnetic resonance imaging

MR imaging can be useful to evaluate the healing process of a surgically treated Achilles tendon.

In almost all surgically repaired Achilles tendons, high signal intensity areas (on fluid sensitive sequences) at the rejoined tendon ends was identified. This finding was clearly seen between 6 weeks and 3 months postoperatively; 6 months after, this area had reduced greatly in size. The high-signal intensity findings on MR images seems to be correlate with the healing response and with the actual tendon tissue composition with respect to morphology and biochemistry²³².

Fujikawa, et al. explored the MRI features of normal healing of the expected residual gap in the Achilles tendon after surgical repair. MRI images showed visible gap on MR imaging on 4 weeks after surgery on T1-WI and T2-WI images, both after percutaneous repair and after open surgery. At 8 weeks a gap was visible on T1-weighted MR images in 80% after percutaneous repair and in 10% after open surgical repair; T2-weighted MR images showed a tendon gap in 63% but in none of the tendons in the open surgical repair group. After 12 weeks, neither T1-weighted nor T2-weighted images showed a tendon gap in both the two tendon's group²³³.

Karjalainen, et al. analysed 21 surgically repaired Achilles tendon ruptures with imaging at 3 and 6 weeks, and at 3 and 6 months after surgery and found intratendinous area of high-intensity signal in almost all surgically repaired Achilles tendons (19/21) at 3 months after surgery on PD (proton density) and T2-WI²³⁴.

Hahn, et al. demonstrated the postoperative MR course after flexor hallucis longus tendon transfer and described that full tendon integration can be expected only in half the patients and fatty muscle degeneration in

Table VII. Answer n. 5: Conservative treatment.

Author	Type of study	Protocol	Follow-up (months)	Outcome assessment	Results	Level of evidence
Neumayer F, et al. ⁹⁸ 2010	Prospective not randomized	Dynamic cast and early mobilization	60	Leppilahti ankle score, isokinetic strenght	Good functional results	III
Metz R, et al. ⁹⁹ 2008	RCT	Surgery vs conservative treatment	6	Isokinetic strenght, ROM	Not significant differences between the two groups	II
Willits K, et al. ¹⁰⁰ 2010	RCT	Surgery vs conservative treatment	24	Re-ruptures, isokinetic strenght, ROM, Leppilahti score, calf circumference	Less complications with conservative treatment, similar functional results	I
Nillson-Helander K, et al. ⁷ 2010	RCT	Surgery vs conservative treatment	12	ATRS, functional tests	Not significant differences between the two groups	I
Soroceanu A, et al. ¹⁰¹ 2012	Meta-analysis of RCT	Surgery vs conservative treatment	-	Complications, strenght, calf circumference, functional tests	Less complications and similar functional results with early functional rehabilitation	I
Wilkins R, et al. ¹⁰² 2012	Meta-analysis of RCT	Open surgery vs conservative treatment	-	Re-ruptures and other complications	Less re-ruptures but major complications with surgery	I
Olsson N, et al. ¹⁰³ 2013	RCT	Surgery + early rehabilitation vs conservative treatment	12	ATRS, functional tests, quality of life	Not significant differences between the two groups	I
Kaniki N, et al. ¹⁰⁴ 2014	Comparative retrospective	Functional rehabilitation + PRP vs functional rehabilitation	24	Isokinetic strenght, ROM, calf circumference, Leppilahti score	Not significant differences between the two groups	III
Mark-Christensen T, et al. ¹⁰⁵ 2014	Meta-analysis of RCT	Functional rehabilitation vs immobilization	-	Complications, strenght, ROM, return to work and to sport	Better results with the functional rehabilitation	II
Young SW, et al. ¹⁰⁶ 2014	RCT	Early weight bearing vs not weight bearing for 8 weeks	24	Re-ruptures, return to work and to sport, pain, stiffness	Not significant differences between the two groups	I
Zhang H, et al. ¹⁰⁷ 2015	Review of meta-analysis	Surgery vs conservative treatment	-	Complications, ROM, calf circumference, functional tests	Different complications for major re-ruptures with surgery, not other significant differences between the two groups	II
Lantto I, et al. ¹⁰⁸ 2015	RCT	Surgery vs conservative treatment	18	Leppilahti score, isokinetic strenght	Similar functional results, but force, ROM and quality of life better with surgery	I

Table VIII. Answer n. 5: Conservative treatment.

Author	Type of study	N° of studies/patients	Topic	Results	Level of evidence
Khan RJ, et al. ¹⁰⁹ 2010	Meta-analysis (RCTs)	12	<ul style="list-style-type: none"> Conservative treatment vs surgery Different techniques of tenorrhaphy 	Surgery: less risk of recurrence and major risk of complications, in particular with open technique	I
Gigante A, et al. ⁸⁸ 2008	RCT	40	Open vs percutaneous technique	Less complications and recovery time with percutaneous technique	II
Aviña Valencia JA, et al. ¹¹⁰ 2009	RCT	56	Open vs mini-invasive technique	Less complications and recovery time with mini-invasive technique	II
Kou J, ¹¹¹ 2010	Guidelines	8	Open surgery - all outcomes	Attention at diabetic patients, smokers, >65 years old, sedentary, obese (BMI >30), neuropathic and with local or systemic dermatologic pathologies	IV
Wilkins R, et al. ¹⁰² 2012	Review of randomized studies	7	Conservative treatment vs surgery	Less incidence of recurrence with surgery	I
Jiang N, et al. ¹¹² 2012	Review of randomized studies	10	Conservative treatment vs surgery	Surgery: major complications risk but early functional recovery and less risk of recurrence	I
Jones MP, et al. ¹¹³ 2012	Review of randomized studies or almost randomized	8 4	<ul style="list-style-type: none"> Conservative treatment vs surgery Open vs percutaneous technique 	Less complications risk. Not differences in recurrence. Major infection risk with open technique. Not differences in sural nerve lesions, TVP and hematomas.	I
Wu Y, et al. ¹¹⁴ 2016	Review of meta-analysis	9	Conservative treatment vs surgery	Less risk of recurrence and major risk of complications with surgery	I

To be continued

Continued from Table VIII.

Miyamoto W, et al. ¹¹⁵ 2017	Retrospective	44	Double locked suture	Correct tendon tension, good functional results, early recovery	IV
Yang B, et al. ¹¹⁶ 2017	Meta-analysis of RCT and retrospective studies	12	Open vs percutaneous technique	<ul style="list-style-type: none"> • Open technique: major risk of deep infections • Percutaneous technique: major risk of sural nerve lesions, less surgery time, better AOFAS score • No significantly differences in recurrence incidence, in thrombotic risk, in ankle ROM, in sural triceps tropism 	II
Del Buono A, et al. ¹¹⁷ 2014	Meta-analysis of RCT and retrospective studies	12	Open vs mini-invasive technique	Less complications and major ROM with mini-invasive technique	I
Li CG, et al. ¹¹⁸ 2017	Retrospective	24	Single bundle termino-terminal suture	After 1 year: mean AOFAS score: 92.4 ± 5.9. Not differences in dorsiflexion, plantar flexion and muscular tropism with contralateral limb	IV
Lewis N, et al. ¹¹⁹ 2003	Controlled on cadaver	/	Reparation with Teno Fix anchor	Good stumps approach, less risk of gap formation	III
Manent A, et al. ¹²⁰ 2017	Controlled on cadaver	/	Differents techniques of tenorrhaphy	Bunnel technique: less risk of lengthening	III
Aktas S, et al. ¹²¹ 2007	Perspective	30	Termino-terminal suture vs augmentation	Less complications with termino-terminal suture	III
Oze Mr, et al. ¹²² 2016	Retrospective	23	Gastrocnemius rotation flap, associated with crural fascia incision	Mean AOFAS score: 98.2 ± 2.3 (range 93-100)	IV

Table IX. Answer n. 6: Sutures and materials.

Author	Type of study	Protocol	Follow-up (months)	Outcome assessment	Results	Level of evidence
Kocaoglu B, et al. ¹²³ 2015	Perspective not randomized	Absorbable vs not absorbable suture	-	AOFAS hindfoot clinical outcome scores, return to work, complications	Less risk of complications with absorbable suture	II
Kara A, et al. ¹²⁴ 2014	Case report	-	12	Post-surgery complications	Granuloma formation with non absorbable suture	V
Olliviere BJ, et al. ¹²⁵ 2014	Case report	-	8	Post -surgery complications	Granuloma formation with FiberWire suture (silicone and polyethylene)	V
Baig MN, et al. ¹²⁶ 2017	Perspective not randomized	Absorbable vs not absorbable suture	6	Complications (infections), Boyden score	Major risk of complications and worse Boyden score with absorbable suture	II
Sadoghi P, et al. ¹²⁷ 2012	Systematic review	Different suture techniques evaluation (Kessler, Bunnell, Krackow, Achillon, Ma-Griffith, giftbox)	-	Resistance to rupture	Impossible to define better technique	II
Manent A, et al. ¹²⁰ 2017	Perspective not randomized	Different suture techniques evaluation (double Kessler, double Bunnell, Krackow, Ma-Griffith)	-	Resistance to rupture*	• Double Bunnel: major resistance, less risk of tendon lengthening • Krackow technique: same resistance, major lengthening	III
Herbort M, et al. ¹²⁸ 2008	Perspective not randomized	Bunnell vs Kessler on cavader	-	Resistance to cyclic loads	Similar biomechanical properties	II
McCoy BW, et al. ¹²⁹ 2010	Perspective not randomized	Different suture techniques evaluation (double Kessler, double Bunnell, double Krackow)	-	Resistance to rupture	No differences in resistance	III

the gastrocnemius muscle and soleus muscle is commonly seen after this technique.²³⁵

The analysis of gadolinium contrast agent enhancement (Gd-CME) images shows larger high signal intensity alterations than on T1-WI before CME or on T2-WI; this finding slowly decreased with time and, at the 2-year MR follow-up, there was no significant intratendinous signal enhancement. This supports the hypothesis that the Gd-contrast agent interacts with the pathological intratendinous tendon healing process²³².

One year after surgery, adhesions between the tendon and the skin may be reported in as many as 40% of the patients²³⁶. The surgical wound scar may be clearly detected on MR images; there was no high

signal intensity subcutaneous fat tissue on images and the tendon seemed to be attached to the skin at the site of the scar, thereby preventing the correct range of motion of the tendon²³⁷.

Advanced MRI application

The use of diffusion tensor imaging (DTI) in musculoskeletal field keeps on growing not only in experimental settings but also in clinical practice, reflecting the information about the architectural organization of tissue. After surgical procedures the use of DTI may ascertain the microstructural properties and integrity restoration of the ruptured tendon during the healing process²³⁸.

Table X. Answer n. 7: Use of autologous derived blood products.

Author	Year	Type of study	Level of evidence	N. of patients	Follow-up (months)	Technical notes
Sánchez M, et al. ¹³⁰	2007	Retrospective (S vs S+PRP)	III	12 (6 vs 6)	-	Intraoperative injection
Shepull T, et al. ¹³¹	2011	RCT (S vs S+PRP)	II	30 (14 S vs 16 S+PRP)	12	Intraoperative injection
Kaniki N, et al. ¹⁰⁴	2014	Retrospective (S vs PRP)	III	145 (72 vs 73 PRP)	24	No surgery
De Carli A, et al. ¹³²	2016	Comparative (S vs S+PRP)	IV	30 (15 S vs 15 S+PRP)	6	Intraoperative injection and after 14 days
Alvitti F, et al. ¹³³	2017	Retrospective (S vs S+PRF vs control group)	IV	28 (9 S vs 11 S+PRF vs 8 control group)	6	PRF application
Zou J, et al. ¹³⁴	2017	RCT (S vs S+PRP)	II	36 (20 S vs 16 S+PRP)	24	Intraoperative injection

S, Surgery (tenorrhaphy); PRP, platelet-rich plasma; PRF, platelet-rich fibrin.

Sarman, et al. analysed pre and postoperative DTI imaging of the Achilles tendon of 16 patients with median duration of follow-up of 21 (range 6 to 80) months; the tendon fractional anisotropy values of the ruptured Achilles tendon were statistically significantly lower than those of the normal side ($p=.001$)²³⁸.

Answer n. 13: Rehabilitation protocol after acute ruptures (Tabs. XV, XVI)

Answer n. 14: Rehabilitation protocol after chronic ruptures

Regardless of treatment, timing does not change, depending on biological healing²⁴⁹⁻²⁶⁴.

Rate of recurrence

The American Academy of Orthopaedic Surgeons (AAOS) guidelines^{265,111} published in 2010, underline the necessity of a cast in the first phases after accident. A meta-analysis of 2012²⁶⁶ reports a significantly rate of post-surgery re-rupture after plaster (3.5%) and after utilised of functional cast (5%). In other studies²⁶⁷⁻²⁶⁹, the rate of recurrence is 3.3% after an accelerated rehabilitative protocol with functional cast and 11.4% with post-surgery plaster.

Rehabilitation protocol

An evidence based optimal protocol does not exist. In

2008, the Swansea Morrision Achilles Rupture Treatment (SMART) Programme was proposed²⁷⁰.

Usually, it is recommended a cast at 30° of plantar flexion for 2 weeks with progressive weight bearing until 8°-9° weeks²⁴⁰⁻²⁷¹. Other Authors recommend the use of a cast at 20° of equinism for the first weeks after tenorrhaphy until start of rehabilitative programme²⁷². Full ankle and limb motion is recommended after 8-9 weeks and return to sport is allowed after 6-9 months²⁴⁰⁻²⁷¹. There is no standard protocol but only some guidance according to biological healing time considering the better synthesis of collagene and the improvement of tendon viscoelastic properties after the first weeks. Physical therapy is a part of protocol reducing inflammatory processes and pain during physiotherapy²⁷³.

Instrumental physiotherapy

Instrumental physiotherapy has therapeutic effects: analgesia, activation of local metabolism, relaxing or muscle tonification. Therefore, instrumental physiotherapy can be utilised in most of therapeutic and rehabilitative programmes in association with other methods²⁷³.

Answer n. 15: Nutraceuticals (Tabs. XVII, XVIII)

Answer n. 16: Return to sport in Table XIX

Table XI. Answer n. 8: Open surgery.

Author	Year	Type of study	Level of evidence	N. of patients (P vs O vs C)	Follow-up	Surgery technique
Nilsson-Helander K, et al. ⁷	2010	RCT	I	97 (49 vs 48)	1 y	O vs C
Keating JF, et al. ¹³⁵	2011	CT	II	80 (41 vs 39)	1 y	O vs C
Nistor L ¹³⁶	1981	RCT	II	105 (45 vs 60)	2.5 y	O vs C
Cetti R, et al. ¹³⁷	1993	RCT	II	111 (65 vs 55)	1 y	O vs C
Möller M, et al. ¹³⁸	2001	RCT	II	112 (59 vs 53)	2 y	Modified Kessler vs C
Twaddle BC, et al. ¹³⁹	2007	RCT	II	50 (25 vs 25)	1 y	O vs C
Willits K, et al. ¹⁰⁰	2010	RCT	II	144 (72 vs 72)	2 y	O vs C
Kołodziej L, et al. ¹⁴⁰	2013	RCT	II	47 (22 vs 25)	3-24 m	Achillon vs Krackow
Gigante A, et al. ⁸⁸	2008	RCT	II	40 (20 vs 20)	1 y	Tenolig vs Kessler
Cretnik A, et al. ¹⁴¹	2005	CT	II	237 (132 vs 105)	2 y	P vs O
Aktas S, et al. ¹⁴²	2009	RCT	II	40 (20 vs 20)	10-48 m	Achillon vs Krakow
Karabinas PK ¹⁴³	2014	RCT	II	34 (19 vs 15)	9-24 m	Ma and Griffit vs Krackow
Lim J, et al. ¹⁴⁴	2001	RCT	II	66 (33 vs 33)	NA	Ma-Griffit vs Krackow
Aviña Valencia JA, et al. ¹¹⁰	2009	RCT	II	56 (28 vs 28)	4 m	Achillon vs Linn
Henriquez H, et al. ¹⁴⁵	2012	Retrospective	III	32 (17 vs 15)	6-48 m	Dresden vs Kessler
Carmont MR, et al. ¹⁴⁶	2013	Retrospective	III	84 (49 vs 35)	18-70 m	P vs Kessler
Miller D, et al. ¹⁴⁷	2005	Retrospective	III	140 (54 vs 86)	3-12 m	Ma-Griffit vs Kessler
Chan AP, et al. ¹⁴⁸	2011	Retrospective	III	19 (10 vs 9)	2-12 m	Achillon vs Krackow
Goren D, et al. ¹⁴⁹	2005	Retrospective	III	20 (10 vs 10)	6-39 m	P (Ma-Griffit) vs O (Krackow)
Daghino W, et al. ¹⁵⁰	2016	Retrospective	III	140	6 m	M (Achillon) vs O
Haji A, et al. ¹⁵¹	2004	Retrospective	III	108 (38 vs 70)	NA	Ma and Griffith vs Bunnell
Lewis N, et al. ¹¹⁹	2003	Comparative on cadaver	III	10	NA	Teno Fix vs two-strand modified Kessler repair
Zhao HM, et al. ¹⁵²	2011	Case series	IV	6	2 y	Bundle to bundle suture
Li CG, et al. ¹¹⁸	2017	Case series	IV	24	1 y	Tendon-bundle technique
Ozer H, et al. ¹²²	2016	Case series	IV	23	1 y	Tenorrhaphy + gastrocnemius flap
Miyamoto W, et al. ¹¹⁵	2017	Case series	IV	44	2 y	Double side-locking loop suture

P, percutaneous tenorrhaphy; M, mini-invasive tenorrhaphy; O, open surgery; C, conservative treatment; NA, no application.

Table XII. Answer n. 9: Minimally invasive surgery.

Author	N. of patients	Follow-up (months)	Variable evaluated	Results	Complications	Level of evidence	Return to sport	Type of surgery
Rebeccato A, et al. ¹⁵³ (2001)	22	21	Objective and subjective evaluation, RMN	Objective and subjective improvement, RMN improvement	1 re-rupture; 1 incision healing delayed	III	Not evaluated	Open vs mini-open vs percutaneous
De Carl A, et al. ¹⁵⁴ (2009)	20	52	Objective and subjective evaluation, functional tests (Ergo-jump Bosco System)	Objective and subjective improvement, dynamic scores improvement	4 incision adhesions	III	85%	Mini-open
Ng ES, et al. ¹⁵⁵ (2006)	25	65,5	Surgery complications	Less complications in mini-open group, similar clinical results	3 minor complications (1 hypertrophic scar, 2 superficial infections)	III	96%	Open vs mini-open (double-ended needle)
Bhattacharyya M, et al. ¹⁵⁶ (2009)	25	14	Objective and subjective evaluation	Objective and subjective improvement, cost reduction	No complications	III	Not evaluated	Mini-open (Achillon system) vs open
Mukundan C, et al. ¹⁵⁷ (2010)	21	12	Functional scores (Leppilatti score, AOFAS)	Functional scores improvement (Leppilatti score, AOFAS)	No complications	III	95%	Mini-open (Achillon system)
Aktas S, et al. ¹⁴² (2009)	20	22,4	Objective and subjective evaluation, functional scores (AOFAS) and complications	No significantly difference in AOFAS, less complications rate	1 insertional tendinopathy	I	89%	Mini-open (Achillon system) vs open
Vadalà A, et al. ¹⁵⁸ (2012)	80	58	Functional scores (Hannover score, VISA-A), ultrasound	Functional scores improvement (Hannover score, VISA-A), ultrasound improvement	12 minor complications (1 hypertrophic scar, 9 incision adhesions, 2 incision healing delayed)	III	84%	Combined mini-open and percutaneous
Vadalà A, et al. ¹⁵⁹ (2014)	36	28	Functional scores (Hannover score, VISA-A), ultrasound	Functional score improvement (Hannover score, VISA-A), ultrasound improvement	6 minor complications (2 incision adhesions, 1 hypertrophic scar, 3 superficial infections).	III	91%	Combined mini-open and percutaneous

To be continued

Continued from Table XII.

Keller A, et al. ¹⁶⁰ (2014)	100	42.1	Objective and subjective evaluation, AOFAS and complications Isokinetic test (21 patients)	Objective and subjective improvement, isokinetic evaluation: full recovery of gastrocnemius and soleus function	2 re-ruptures; 5 TVP;	IV	85%	Dresden mini-open
Klein EE, et al. ¹⁶¹ (2012)	18	12-108	Objective and subjective evaluation, VISA-A score and complications	Objective and subjective improvement	1 re-rupture; 1 complications incision	III	Not evaluated	Mini-open (Achillon system) vs open
Barte AF, et al. ¹⁶² (2014)	253	19,2	Complications	Incidence of complications acceptable, in relation to the other surgery techniques	Re-ruptures 8; incision complications: 5; sural nerve lesions: 3; infections: 2; suture irritation: 3	Systematic review	Not evaluated	-
De Carli A, et al. ¹³² (2016)	30	28	Functional scores (VAS, FAOS, VISA-A), ultrasound and RMN	Functional scores improvement (VAS, FAOS, VISA-A), ultrasound and RMN improvement	5 minor complications (3 incision healing delayed, 2 incision adhesences)	IV	100%	Mini-open
Daghino W, et al. ¹⁵⁰ (2016)	68	6-53	Objective evaluation and complications	Objective improvement, quality of life improvement	2 major complications (2 re-ruptures); 2 minor complications (2 incision adhesences)	III	87,50%	Mini-open (Achillon system) vs Open
Taştan E, et al. ¹⁶³ (2016)	20	58,5	Functional scores (AOFAS)	Functional scores improvements (AOFAS)	No complications	III	100%	Mini-open (Achillon system)

Table XIII. Answer n. 10: Percutaneous surgery.

Author	Year	Type of study	Level of evidence	N. of patients (P vs O)	Follow-up (months)	Type of surgery
Karabinas PK, et al. ¹⁴³	2014	RCT (P vs O)	I	34 (19 vs 15)	22	Ma and Griffith
Gigante A, et al. ⁸⁸	2008	RCT (P vs O)	I	40 (20 vs 20)	24	Tenolig®
Lim J, et al. ¹⁴⁴	2001	RCT (P vs O)	I	66 (33 vs 33)	6	Ma and Griffith
Jallageas R, et al. ¹⁶⁴	2013	Comparative (P vs O)	II	31 (16 vs 15)	15	Tenolig®
Cretnik A, et al. ¹⁴¹	2005	Comparative (P vs O)	II	237 (132 vs 105)	24	Ma and Griffith
Zayni R, et al. ¹⁶⁵	2017	Retrospective (P vs O)	III	29 (16 vs 13)	46	Tenolig®
Henriquez H, et al. ¹⁴⁵	2012	Retrospective (P vs O)	III	32 (17 vs 15)	18	Tenolig®
Tagliavoro G, et al. ¹⁶⁶	2011	Retrospective (P vs P)	III	60 (30 vs 30)	24	Ma and Griffith vs Tenolig®
Haji A, et al. ¹⁵¹	2004	Retrospective (P vs O)	III	108 (38 vs 70)	Not reported	Ma and Griffith
Bradley JP, et al. ¹⁶⁷	1990	Comparative (P vs O)	III	27 (12 vs 15)	Not reported	Ma and Griffith
Tenenbaum S, et al. ¹⁶⁸	2010	Case series	IV	29	32	Ma and Griffith
Maes R, et al. ¹⁶⁹	2006	Case series	IV	124	23	Tenolig®
Lacoste S, et al. ¹⁷⁰	2014	Case series	IV	75	21	Tenolig®

P, percutaneous tenorrhaphy; O, open surgery.

Answer n. 17: Outcome evaluation devices (Indirect determination of Achilles tendon force during locomotion by motion analysis techniques)

The position of selected anatomical landmarks of the lower limb and the foot-to-ground reaction force, as collected during terrestrial locomotion, represent the experimental data that are sufficient to solve the inverse dynamic problem and estimate the so-called "intersegmental couple" (IC) at the ankle³⁵⁹. IC can be considered as a muscle-equivalent representation of the angular actuator responsible for the motion of the foot about the ankle joint center in the sagittal plane during the ground-contact phase. IC results from the contributions of the moments due to: the ground reaction force acting on the foot; the segment's weight; the acceleration force of the segment's center of mass; the segment's angular acceleration³⁶⁰. All these quantities can be easily gathered in a motion analysis laboratory. When the sign of IC is negative³⁶¹. The tensile force of the Achilles tendon (AT) can be computed as the ratio between IC and the AT lever arm with respect to the ankle joint center³⁶². In fact, as the main plantar-flexor muscles of the ankle converge in the AT and no optimization

may be needed as no plantar-flexor muscles redundancy occurs³⁶³. The AT lever arm is typically estimated from scaled generic musculoskeletal models³⁶⁴. A high level of association and a low bias were found between the AT force estimated through inverse dynamics and that measured *in vivo* with an implanted force transducer³⁶⁵.

Several are, however, the limitations of such approach. First, the assumption that IC can be uniquely addressed to the plantar-flexors muscles (hence, excluding co-contraction of antagonist muscles³⁶² and neglecting the contribution of passive forces exerted by ligaments³⁶⁶). Second, the accuracy of the estimated AT force strongly depends on the reliability of the collected experimental data (anatomical landmarks identification and skin artefact in the first place³⁶⁷⁻³⁶⁹) and on the chosen musculoskeletal model (inertial parameters and musculoskeletal geometries are based on generic models scaled on the subject's proportions)³⁷⁰. For this latter reason, the scientific community has been recently focusing on the availability of imaging techniques to assess subject-specific musculoskeletal geometries simultaneously to motion data collection to estimate ankle dynamics³⁷¹⁻³⁷³.

Table XIV. Answer n. 11: Tendon transfer.

Author	Year	Type of study	Level of evidence	N. of patients	Follow-up (months)	Type of surgery
Maffulli N, et al. ¹⁷¹	2005	Cohort study	III	21	24	Free autologous gracilis tendon graft
El Shewy MT, et al. ¹⁷²	2009	Case series	IV	11	90	Intratendinosus flaps from gastrocnemius-soleus complex
Maffulli N, et al. ¹⁷³	2010	Case series	IV	32	72	Peroneus brevis tendon transfer
Us AK, et al. ¹⁷⁴	1997	Case series	IV	6	16	V-Y gastrocnemius recession, end to end anastomosis and gastrocnemius aponeurotic flap
Kissel CG, et al. ¹⁷⁵	1994	Case series	IV	4	38	V-Y gastrocnemius recession, end to end anastomosis and plantaris tendon weaving
Esenyel CZ, et al. ¹⁷⁶	2014	Case series	IV	10	43,2	Turndown gastrocnemius-soleus fascial flap
Guclu B, et al. ¹⁷⁷	2016	Retrospective comparative study	III	17	195	V-Y tendon plasty with fascia turndown
Rush JH, et al. ¹⁷⁸	1980	Case series	IV	5	18-24	Gastrocnemius-soleus aponeurotic flap turndown
Wapner KL, et al. ¹⁷⁹	1993	Case series	IV	7	17	Flexor hallucis longus tendon transfer
Pintore E, et al. ¹⁸⁰	2001	Comparative (A vs C)	II	59	53	Peroneus brevis tendon transfer
Ademoglu Y, et al. ¹⁸¹	2001	Case series	IV	4	39,2	Peroneus brevis tendon transfer
Wong MW, et al. ¹⁸²	2005	Case series	IV	5	28,8	Flexor hallucis longus tendon transfer
Elias I, et al. ¹⁸³	2007	Case series	IV	15	26,5	V-Y leghtening and flexor hallucis longus tendon transfer
Mahajan RH, et al. ¹⁸⁴	2009	Case series	IV	36	12	Flexor hallucis longus tendon transfer
Maffulli N, et al. ¹⁸⁵	2012	Case series	IV	16	185	Peroneus brevis tendon transfer
Rahm S, et al. ¹⁸⁶	2013	Retrospective comparative series (tt vs to)	III	40	73-35	Flexor hallucis longus tendon transfer
Dumbre Patil SSD, et al. ¹⁸⁷	2014	Case series	IV	35	30,7	Semitendinosus tendon autograft
Singh A, et al. ¹⁸⁸	2014	Case series	IV	22	12	Peroneus brevis tendon augmentation
Khiami F, et al. ¹⁸⁹	2013	Retrospective	IV	23	24,5	Free sural triceps aponeurosis transfer
Maffulli N, et al. ¹⁹⁰	2015	Case series	IV	17	54	Peroneus brevis tendon transfer
Ahmad J, et al. ¹⁹¹	2016	Case series	IV	32	62,3	Flexor hallucis longus tendon transfer
Gedam PN, et al. ¹⁹²	2016	Retrospective comparative	III	14	30,1	Central turndown flap with free semitendinosus tendon graft
Maffulli N, et al. ¹⁹³	2013	Case series	IV	26	31,4	Free semitendinosus tendon graft

To be continued

Continued from Table XIV.

Author	Year	Type of study	Level of evidence	N. of patients	Follow-up (months)	Type of surgery
Mann RA, et al. ¹⁹⁴	1991	Case series	IV	7	39	Flexor digitorum longus tendon graft
Elgohary HEA, et al. ¹⁹⁵	2016	Case series	IV	19	29	Flexor hallucis longus tendon transfer
Miao X, et al. ¹⁹⁶	2016	Case series	IV	32	32,2	Flexor hallucis longus tendon transfer
Maffulli N, et al. ¹⁹⁷	2015	Cohort study	III	21	54	Peroneus brevis tendon transfer
Yeoman TF, et al. ¹⁹⁸	2012	Case series	IV	11	6	Flexor hallucis longus tendon transfer
Park YS, et al. ¹⁹⁹	2012	Retrospective (VY vs G vs FHL)	III	12	36,2	V-Y advancement, gastrocnemius fascial turndown flap, FHL tendon transfer
Sarzaeem MM, et al. ²⁰⁰	2012	Case series	IV	11	25	Free semitendinosus tendon graft
Zheng L, et al. ²⁰¹	2011	Case series	IV	10	8-48	Peroneus brevis tendon transfer
Wegrzyn J, et al. ²⁰²	2010	Case series	IV	11	79	Flexor hallucis longus tendon transfer
Lee KB, et al. ²⁰³	2009	Case series	IV	3	18-24	Flexor hallucis longus tendon transfer
Fotiadis E, et al. ²⁰⁴	2008	Case series	IV	9	44	Plantaris tendon transfer and Duthie's biological repair
Lui TH, et al. ²⁰⁵	2007	Case series	IV	3	15	Flexor hallucis longus tendon transfer
Miskulin M, et al. ²⁰⁶	2005	Case series	IV	5	12	Peroneus brevis tendon transfer and plantaris tendon Augumentation
Dalal RB, et al. ²⁰⁷	2003	Case series	IV	2	Not reported	Flexor hallucis longus tendon transfer
Seker A, et al. ²⁰⁸	2016	Case series	IV	21	145,3	Gastrocnemius fascial flap
Lapidus LJ, et al. ²⁰⁹	2012	Case series	IV	9	60	Achilles tendon island flap
Takao M, et al. ²¹⁰	2003	Case series	IV	10	26-192	Gastrocnemius fascial flap
Ozan F, et al. ²¹¹	2017	Comparative (V vs L)	II	15	19.6	Lindholm and Vulpius tecique
Sanada T, et al. ²¹²	2017	Case series	IV	56	6	Free gastrocnemius aponeurotic flap
Maffulli N, et al. ²¹³	2014	Case series	IV	28	24	Semitendinosus tendon autograft
El Shazly O, et al. ²¹⁴	2014	Case series	IV	15	27	Free hamstring tendon autograft
Tay D, et al. ²¹⁵	2010	Case series	IV	6	24	Turndown tendon flaps
Nilsson-Helander K, et al. ²¹⁶	2008	Case series	IV	28	29	Free gastrocnemius aponeurotic flap
Tawari AA, et al. ²¹⁷	2013	Case series	IV	20	18	Peroneus brevis tendon transfer
Oksanen MM, et al. ²¹⁸	2014	Case series	IV	7	27	Flexor hallucis longus tendon transfer

A, acute rupture; C, chronic rupture; tt, transtendineous technique; to, transosseus technique; VY, V-Y plasty; G, gastrocnemius fascial flap; "FHL", flexor hallucis longus tendon transfer; V, Vulpius tecique; L, Lindholm tecique.

Table XV. Answer n. 13: Rehabilitation protocol after acute ruptures. Open Surgery.

Author	Year	Type of study	Level of evidence	N. of patients (P vs O)	Follow-up (months)	Treatment groups
Valkering KP, et al. ²³⁹	2017	RCT	II	56 (27 vs 29)	12	<ul style="list-style-type: none"> • Mobilized and FWB group • Immobilized and NBW group
Lantto I, et al. ²⁴⁰	2015	RCT	I	50 (25 vs 25)	132	<ul style="list-style-type: none"> • Early mobilization group • Immobilization in tension group
Suchak AA, et al. ²⁴¹	2008	RCT	I	110 (55 vs 55)	6	<ul style="list-style-type: none"> • Weight-Bearing as tolerated Group • NBW group
Costa ML, et al. ²⁴²	2006	RCT	II	48 (23 vs 25)	12	<ul style="list-style-type: none"> • Treatment Group • Control Group
Maffulli N, et al. ⁹¹	2003	Case-control study	III	53 (26 vs 27)	4.5	<ul style="list-style-type: none"> • Group 1 • Group 2
Kangas J, et al. ²⁴³	2003	RCT	II	50 (25 vs 25)	15	<ul style="list-style-type: none"> • Group I • Group II
Kerkhoffs GM, et al. ²⁴⁴	2002	RCT	II	39 (23 vs 16)	80	<ul style="list-style-type: none"> • Cast group • Wrap group
Mortensen HM, et al. ²⁴⁵	1999	RCT	II	61 (31 vs 30)	24	<ul style="list-style-type: none"> • Early Motion group • Cast group

FBW, complete weight bearing; NBW, no weight bearing.

Table XVI. Answer n. 13: Rehabilitation protocol after acute ruptures. Minimally invasive or percutaneous surgery.

Author	Year	Type of study	Level of evidence	N. of patients (P vs O)	Follow-up (months)	Treatment Groups
De la Fuente C, et al. ²⁴⁶	2016	RCT	II	38 (19 vs 19)	3	<ul style="list-style-type: none"> • Conventional group • Aggressive group
Groetelaers RP, et al. ²⁴⁷	2014	RCT	II	60 (32 vs 28)	12	<ul style="list-style-type: none"> • Functional group • Immobilization group
Majewski M, et al. ²⁴⁸	2008	Case-control study	III	28 (14 vs 14)	12	<ul style="list-style-type: none"> • Cast group • Shoe group

Prediction of AT force during terrestrial locomotion: difference with respect to methods, to the computational approach and to the adopted musculoskeletal model in Table XX.

Answer n. 18: Acute ruptures in the childhood in Table XXI.

Project management

I.S.Mu.L.T. - Italian Society of Muscles Ligaments & Tendons.

Coordinator

Francesco Oliva
Department of Orthopaedics and Traumatology, University of Rome "Tor Vergata", Italy.

Table XVII Answer n. 15: Nutraceuticals. Clinical studies about the characteristics in the use of nutraceuticals for therapy of tendinopathies.

Author/Year	Pathology	Type of nutraceutical and composition	Type of study/N. of patients	Groups compared
Notarnicola A, et al. 2012 ²⁷⁴	Insertional Achilles tendinopathy	Tenosan® (L-arginine- α -ketoglutarate, methylsulfonylmethane, type I hydrolyzate collagen, Vinitrox™, bromelain, vitamin C)	RCT (placebo) g-t: 32 g-c: 32>26	g-t: ESWT + Tenosan® g-c: EWTS + placebo Dosage: 2 bags/day for 60 days before main meal
Balius R, et al. 2016 ²⁷⁵	Non-insertional painful Achilles tendinopathy	Tendoactive® (mucopolysaccharids, type I collagen, vitamin C)	RCT (no placebo) g-t 1: 19>17 g-t 2: 20 g-c: 19>18	-t 1: EC + Tendoactive® g-t 2: PS + Tendoactive® g-c: EC Dosage: 3 capsules/day for 12 weeks
Hai-Binh B, et al. 2014 ²⁷⁶	Various tendinopathies (Achilles tendon, supraspinatus, lateral epicondyle, plantar fasciitis)	Tendoactive® (mucopolysaccharids, type I collagen, vitamin C)	RCT (placebo) g-t: 30 g-c: 30	g-t: Tendoactive® g-c: placebo Dosage: 2 capsules/day for 90 days
Nadal F, et al. 2009 ²⁷⁷	Various tendinopathies (Achilles tendon, supraspinatus, lateral epicondyle, plantar fasciitis)	Tendoactive® (mucopolysaccharids, type I collagen, vitamin C)	RCT (no placebo) g-t: 10 g-c: 10	g-t: rehabilitation + Tendoactive® g-c: rehabilitation Dosage: 2.16 g/day for 3 months
Arquer A, et al. 2014 ²⁷⁸	Various tendinopathies (Achilles tendon n=32, patellar tendon n=32, lateral epicondyle n=34)	Tendoactive® (mucopolysaccharids, type I collagen, vitamin C)	Perspective not controlled explorative study of phase IV n=98->70	Dosage: 3 capsules/day for 90 days
Mavrogenis S, et al. 2004 ²⁷⁹	Chronic tendon disorders*	Bio-Sport® Essential fatty acids (EPA, DHA, GLA) + antioxidants (selenium, zinc, vitamin A, vitamin B6, vitamin C, vitamin E)	RCT (placebo, double blinded) on athletes g-t: 20->17 g-c: 20->14	g-t: ultrasounds + supplements g-c: ultrasounds + placebo Dosage: 8 capsules/day essential fatty acids + 1 antioxidants for 32 days

EC, eccentric exercise; PS, passive stretching; g-t, treated group; g-c, control group; ESWT, Extracorporeal shock wave therapy; EPA, eicosapentaenoic acid; DHA, docosahexaenoic acid; GLA, gamma-linolenic acid; *Chronic tendon disorders. NB: Balius - Hai-Bin - Arquer - Nadal: same supplement (Tendoactive®).

Overseeing group

Nicola Maffulli, Pasquale Farsetti, Calogero Foti, Milena Fini, Biagio Moretti, Pietro Ruggieri, Umberto Tarantino, Maria Chiara Vulpiani.

Group of experts

Carlo Biz, Roberto Buda, Daniela Buonocore, Vincenzo De Luna, Luigi Di Lorenzo, Bernardo Innocenti, Alessio Giai Via, Antonio Frizziero, Alfonso Maria Forte, Asmaa Mahmoud, Angelo De Carli, Johnny

Padulo, Pietro Picerno, Francesca Veronesi, Mario Vetrano, Marcello Zappia.

Group of preparation and evaluation of the literature

Matteo Baldassarri, Gabriele Bernardi, Michela Bossa, Vito Chianca, Anna Collina, Imma Di Lanno, Francesco Di Pietto, Maurizia Dossena, Ilaria Fantoni, Paolo Finotti, Edoardo Gaj, Carlotta Galeone, Jacopo Gamberini, Monica Gasparini, Domenico Lupariello,

Table XVIII. Answer n. 15: Nutraceuticals. Clinical studies about the use of nutraceuticals for therapy of tendinopathies.

Author/Year	Outcome assessments	Follow-up	Results
Notarnicola A, et al. 2012 ²⁷⁴	<p>Tenosan® efficacy combined with shock waves in insertional Achilles tendinopathy management</p> <p>Primary endpoints (clinical and functional effects) VAS score^a Ankle-Hindfoot Scale^b (pain, function, alignment) Roles and Maudsley score (subjective improvement perception)^c</p> <p>Secondary endpoint (neoangiogenesis) Tissue oximetry</p>	2 and 6 months	<p>VAS score significantly lower in both groups during the study. At 6 months, VAS score significantly lower in the group with combined treatment (average score: 2.0 vs 2.9, $p=0.04$), although difference <2 points (threshold clinically significantly)</p> <p>Ankle-Hindfoot Scale significantly improved scores only in the group with combined treatment during the study. At 2 and 6 months, improved scores in the group with combined treatment (average at 6 months: 92.4 vs 76.5, $p=0.0002$)</p> <p>At 2 and 6 months, improved scores (lower) in Roles and Maudsley score in the group with combined treatment (average at 6 months: 1.5 vs 2.3, $p<0.0001$)</p> <p>Significantly lower scores at oximetry in both groups due during the study; only at 6 months significantly difference between the two groups in favor of the group with combined treatment (average 60.2 vs 66.0, $p=0.007$)</p>
Balius R, et al. 2016 ²⁷⁵	<p>Tendoactive® efficacy combined with eccentric physical exercise to improve non-insertional painful Achilles tendinopathy symptoms</p> <p>Primary endpoint VISA-A questionnaire score^d (function and pain)</p> <p>Secondary endpoints VAS score for pain^a at rest and during activity Tendon thickness (ultrasound)</p>	6 and 12 weeks	<p>At 12 weeks, VISA-A score significantly improved (higher) in the 3 groups. No significantly difference between the groups at VISA-A score</p> <p>At 12 weeks, VAS score at rest and during activity significantly reduced in the 3 groups. Significantly difference in reduction of VAS score at rest in the Tendoactive® + PS group compared with EC (-3.7 vs -2.7, $p<0.005$); borderline difference at VAS during activity (-4.4 Tendoactive® + PS vs -3.5 EC, $p=0.074$).</p> <p>At 12 weeks, no significantly difference in tendon thickness between the 3 groups; significantly reduction from baseline to 12 weeks only in Tendoactive® +PS group (-0.63 mm).</p> <p>In analysis stratified on pathology stage (reactive/degenerative tendinopathy): no significantly differences between the treated groups in both stages; VAS score at rest significantly lower in Tendoactive® + PS group than in EC (-3.82 vs -2.80, $p<0.005$) in patients with reactive tendinopathy; VAS score at rest and during activity similar between the groups in patients with degenerative tendinopathy; significantly reduction of tendon thickness from baseline only in Tendoactive® + PS group in patients with degenerative tendinopathy</p>
Hai-Binh B, et al. 2014 ²⁷⁶	<p>Tendoactive® efficacy and safety in management of different tendinopathies</p> <p>Swelling, heat, redness (clinical evaluation) VAS score for pain^a Tendinopathy (ultrasound)</p>	Monthly during the study (90 days)	<p>Progressively reduction of presence of swelling, heat, redness in both groups; lower in the experimental group at every monthly control</p> <p>VAS score significantly reduced in both groups during the study. At 90 days, VAS score significantly lower in the experimental group (average: 2.5 vs 3.2, $p<0.05$)</p> <p>At 90 days, no patient in the experimental group has diagnosis of tendinopathy (% placebo group not reported by Authors)</p>

To be continued

Continued from Table XVIII.

Author/Year	Outcome assessments	Follow-up	Results
Nadal F, et al. 2009 ²⁷⁷	Tendoactive® efficacy in treatment of different tendinopathies Pain SF36 (Quality of life) Functional evaluation by physiotherapist	1, 2 and 3 months	Significantly reduction of pain in the experimental group for every pathology, except for epicondylitis Improved of SF36 in every group of pathology At 3 months significantly improvement of function for every tendinopathies. (Results of placebo group not reported by Authors)
Arquer A, et al. 2014 ²⁷⁸	Tendoactive® efficacy and safety in treatment of different tendinopathies VAS score for pain ^a at rest and during activity Function (VISA-A score for Achilles tendon, VISA-P for patellar tendon, PRTEE for elbow) Ultrasound structural parameters (tendon thickness, effacement of the paratenon, heteroechoogenicity and hypoechoogenicity levels, neovascularization)	30, 60, 90 days	3 groups based on pathology: Achilles tendinopathy (AQ), patellar tendinopathy (RO), lateral epicondylitis (EPI) Significantly reduction of VAS score at rest and during activity in the 3 groups at 30, 60 and 90 days. At 90 days, compared to baseline, the pain at rest is reduced of 80% in AQ, of 71% in RO and of 91% in EPI; pain during activity reduced of 82% in AQ, 73% in RO and 81% in EPI Significantly improvement of VISA-A, VISA-P and PRTEE at 30, 60 and 90 days. At 90 days, compared to baseline, improvement of 38%, 46% and 77% in AQ, RO, and EPI Significantly reduction in tendon thickness in the 3 groups (at 90 days: 12% in AQ, 10% in RO and 20% in EPI). In EPI group reduction during all period; in AQ and RO groups reduction at 60 days, after stable at 90 Improved of all structural parameters in the 3 groups. Paratenon blurred and levels of heteroechoogenicity and hypoechoogenicity significantly improved in AQ and EPI; level of hypoechoogenicity not significantly improve in RO group ($p=0.07$); neovascularization significantly improve only in EPI group
Mavrogenis S, et al. 2004 ²⁷⁹	Efficacy of suppluement combined with physiotherapy in treatment of chronic tendinopathies in athletes Primary endpoints VAS score for pain ^a VAS score for pain ^a after isometric test Secondary endpoints Physical activity	8, 16, 24 and 32 days	VAS score lower duraing the study in both groups. At 32 days, statistically significantly difference between the groups in favor of experimental group ($p<0.001$) (VAS score reduced 99% in experimental group and 31% in control group). Similar results of VAS score after isometric test: at 32 days, score significantly lower ($p<0.001$) in experimental group (VAS score reduced 99% in experimental group and 37% control group) At 32 days, improved sport activity compared to basal (53% in experimental group and 11% control group) No adverse events in both groups

EC, eccentric exercise; PRTEE, Patient-Rated Tennis Elbow Evaluation; PS, passive stretching; SF, short-form; VAS, visual analog scale; VISA-A, Victorian Institute of Sports Assessment-Achilles; VISA-P, Victorian Institute of Sports Assessment-Patella.

^a VAS: range 0-10 (10=severe pain; 0=no pain).

^b Ankle-Hindfoot Scale: range 0-100 (100=no pain, no limitations, good alignment; 0=severe pain, severe limitations, severe misalignment).

^c Roles and Maudsley score: range 1-4 (4=no satisfaction or low satisfaction of the treatment, 1=good satisfaction of the treatment).

^d VISA-A questionnaire: range 0-100 (higher scores for better functionality and lower pain).

Table XIX. Answer n. 16: Return to sport.

Author	N. of patients	Groups	% return to sport	Variables analyzed
Ahmad J, et al. ²⁸⁰	30	1	NR	FAAM Sports Subscale
Aktas S, et al. ¹⁴²	40	1	87	AOFAS
Aktas S, et al. ¹²¹	30	1	86.9	AOFAS
Al-Mouazzen L, et al. ²⁸¹	30	1	NR	ATRS
Amin NH, et al. ²⁸²	18	1	61	NBA Player Efficiency Rating
Amlang MH, et al. ²⁸³	39	1	51	AOFAS
Ateschrang A, et al. ²⁸⁴	104	1	64.4	Thermann Score
Barfod KW, et al. ²⁷¹	56	1	18.6	ATRS
Bassi JL, et al. ²⁸⁵	11	2	100	
Bevoni R, et al. ²⁸⁶	66	2	98.5	AOFAS, Leppilahti
Bostick GP, et al. ²⁸⁷	84	2	84	
Boyden EM, et al. ²⁸⁸	10	2	80	Boyden Scale
Carmont MR, et al. ²⁸⁹	26	1	61	Tegner Score
Ceccarelli F, et al. ²⁹⁰	24	1	91.7	AOFAS
Chandrakant V, et al. ²⁹¹	52	1	90	AOFAS
Chen Z, et al. ²⁹²	76	1	100	
Chiu CH, et al. ⁸⁶	19	1	94.7	Tegner Score, AOFAS
Coutts A, et al. ²⁹³	25	1	80	
Cretnik A, et al. ¹⁴¹	237	1	72.1	AOFAS
Cretnik A, et al. ²⁹⁴	116	1	96	AOFAS
Cretnik A, et al. ²⁹⁵	13	2	100	AOFAS
De Carli A, et al. ¹⁵⁴	20	1	70.5	
Demirel M, et al. ²⁹⁶	78	1	77.1	
Doral MN, ²⁹⁷	32	1	100	FAOS, ATRS
Eames MHA, et al. ²⁹⁸	32	1	63	
Feldbrin Z, et al. ²⁹⁹	14	1	100	AOFAS
Fernández-Fairén M, et al. ³⁰⁰	29	2	96.6	AOFAS
Fortis AP, et al. ³⁰¹	20	1	100	
Garabito A, et al. ³⁰²	49	1	89.8	AOFAS

To be continued

Continued from Table XIX.

Garrido IM, et al. ³⁰³	18	2	72.2	AOFAS
Goren D, et al. ¹⁴⁹	20	1	55	
Gorschewsky O, et al. ³⁰⁴	20	2	100	
Gorschewsky O, et al. ³⁰⁵	66	2	100	
Groetelaers RP, et al. ²⁴⁷	55	1	39	ARPS
Guillo S, et al. ³⁰⁶	23	1	80	ATRS, Boyden Scale
Halasi T, et al. ³⁰⁷	144	1	60.7	
Hohendorff B, et al. ³⁰⁸	42	1	88.6	Thermann score
Hufner TM, et al. ³⁰⁹	125	2	75.2	
Jaakkola JI, et al. ³¹⁰	55	2	90.9	AOFAS
Jacob KM, et al. ³¹¹	46	1	88.9	
Jallageas R, et al. ¹⁶⁴	31	1	77.5	AOFAS
Jennings AG, et al. ³¹²	30	1	63.6	Tennier
Josey RA, et al. ³¹³	39	1	66.7	AOFAS, Thermann score
Jung HG, et al. ³¹⁴	30	2	90	
Kakiuchi M, et al. ³¹⁵	22	1	45.5	
Karabinas PR, et al. ¹⁴³	34	2	NR	AOFAS
Karkhanis S, et al. ³¹⁶	107	2	77	ATRS
Keating JF, et al. ¹³⁵	80	1	66.9	
Kelle A, et al. ¹⁶⁰	100	1	80	
Klein EE, et al. ¹⁶¹	34	2	100	VISA-A
Knobe M, et al. ³¹⁷	64	1	36.6	
Kolodziej L, et al. ¹⁴⁰	47	1	46	
Korkmaz M, et al. ³¹⁸	47	1	NR	PASS
Kraus R, et al. ³¹⁹	36	1	53	
Labib SA, et al. ³²⁰	44	1	65.71	
Lacoste S, et al. ¹⁷⁰	75	1	63.6	ATRS, AOFAS
Lansdaal JR, et al. ³²¹	163	1	59.5	Leppilahti Score
Lee DK, ³²²	11	2	NR	
Leppilahti J, et al. ³²³	101	1	85.7	Boyden Scale
Macquet AJ, et al. ³²⁴	87	1	68.1	

To be continued

Continued from Table XIX.

Maffulli N, et al. ⁹¹	53	1	92.5	Modified VISA-A
Maffulli N, et al. ³²⁵	17	2	94	ATRS
Maffulli N, et al. ³²⁶	27	2	50	ATRS
Majewski M, et al. ³²⁷	84	1	100	Hannover Achilles tendon score
Majewski M, et al. ²⁴⁸	28	1	65.2	Hannover Achilles tendon score
Mandelbaum BR, et al. ³²⁸	29	1	100	
Maniscalco P, et al. ³²⁹	7	1	100	Mandelbaum and Pavanini evaluation
Martinelli B, et al. ³³⁰	30	1	100	
McComis GP, et al. ³³¹	15	1	66	
Metz R, et al. ⁹⁹	83	1	72.8	Leppilahti score
Metz R, et al. ³³²	210	1	50	ATRS
Miller D, et al. ¹⁴⁷	111	1	88	
Möller M, et al. ¹³⁸	112	1	54	Functional index of lower limbs
Mortensen HN, et al. ³³³	57	1	70	
Mortensen HN, et al. ²⁴⁵	61	1	54.1	
Motta P, et al. ³³⁴	71	1	28	
Mukundan C, et al. ¹⁵⁷	21	1	95.2	AOFAS, Leppilahti
Nestorson J, et al. ³³⁵	25	1	36	
Nilsson-Helander R, et al. ⁷	97	1	NR	PAS, ATRS
Olsson N, et al. ¹⁰³	100	1	NR	PAS, ATRS, FAOS
Orr J, et al. ³³⁶	15	2	100	AOFAS
Ozsoy M, et al. ³³⁷	13	1	92	AOFAS
Pajala A, et al. ³³⁸	60	1	100	Leppilahti score
Parekh SG, et al. ³³⁹	31	1	64.3	Power rating (pre-surgery and during match)
Park HG, et al. ³⁴⁰	14	2	NR	
Rajasekar K, et al. ³⁴¹	35	1	50	Accidents questionnaire
Rebeccato A, et al. ¹⁵³	59	1	98.4	
Rettig AC, et al. ³⁴²	89	1	100	
Richardson LC, et al. ³⁴³	30	1	77	AOFAS
Sánchez M, et al. ¹³⁰	12	1	58	Functional Cincinnati Scale (modified)
Schepull T, et al. ³⁴⁴	10	1	40	Thermann score

To be continued

Continued from Table XIX.

Silbernagel KG, et al. ³⁴⁵	8	1	NR	ATRS, FAOS
Soldatis J, et al. ³⁴⁶	30	1	61	
Solveborn S, et al. ³⁴⁷	17	1	94	Amer-Lindon Scale
Sorrenti S, et al. ³⁴⁸	52	2	100	
Speck M, et al. ³⁴⁹	20	1	100	
Stein BE, et al. ³⁵⁰	27	1	92	
Strauss E, et al. ³⁵¹	54	1	74	Boyden Score, AOFAS
Suchak AA, et al. ²⁴¹	98	2	65	
Talbot J, et al. ³⁵²	15	1	66.7	AOFAS
Tenenbaum S, et al. ¹⁶⁸	29	1	90	AOFAS, Boyden score (modified)
Troop RL, et al. ³⁵³	13	1	94	
Uchiyama E, et al. ³⁵⁴	100	1	100	
Valente M, et al. ³⁵⁵	35	2	100	AOFAS
Wagnon R, et al. ³⁵⁶	57	1	40	
Wallace RGH, et al. ³⁵⁷	945	1	100	
Wallace RGH, et al. ³⁵⁸	140	1	37	
Young SW, et al. ¹⁰⁶	84	1	NR	Leppilahti score, halasi score

NR, not reported; AOFAS, American Orthopaedic Foot and Ankle Society Ankle-Hindfoot Score; ARPS, Achilles Rupture Performance Score; ATRS, Achilles Tendon Total Rupture Score; FAAM, Foot and Ankle Ability Measure; FAOS, Foot and Ankle Outcome Score-Ankle and Hindfoot; PAS, Physical Activity Scale; PER, Player Efficiency Rating.

Table XX. Prediction of AT force during terrestrial locomotion: difference with respect to methods, to the computational approach and to the adopted musculoskeletal model.

Authors	Protocol	Task	Results
Fukashiro S, et al. ³⁶⁵ 1993	Inverse dynamics vs direct measure	Hopping	diff = 8% r = 0.99
Kernozek T, et al. ³⁶² . 2017	Conventional vs optimized inverse dynamics	Running	diff = 4.7% (p = 0.054)
Gerus P, et al. ³⁷² . 2012	Subject-specific vs generic musculoskeletal models	Hopping/running	diff = 17%

Table XXI. Answer n. 18: Acute ruptures in the childhood.

Author	Year	Type of study	Level of evidence	N. of patients	Follow-up (months)	Type of treatment
Ralston EL, et al. ³⁷⁴	1971	Case series	IV	1	12	Surgery
Eidelman M, et al. ³⁷⁵	2004	Case series	IV	1	12	Conservative
Tudisco C t al. ³⁷⁶	2012	Case series	IV	1	36	Surgery - Bunnell open
Vasileff WK, et al. ³⁷⁷	2014	Case series	IV	1	8	Surgery -Bunnell open

Emanuela Marsilio, Simone Natali, Leonardo Pellicciari, Luca Perazzo, Eleonora Piccirilli, Clelia Rugiero, Antonio Vadalà, Manuela Verri.

Ethics

The Authors declare that this research was conducted following basic ethical aspects and international standards as required by the journal and recently update in³⁷⁸.

References

1. Maffulli N, Waterston SW, Squair J, Reaper J, Douglas AS. Changing incidence of Achilles tendon rupture in Scotland: a 15-year study. *Clin J Sport Med*. 1999 Jul;9(3):157-160.
2. Longo UG, Petrillo S, Maffulli N, Denaro V. Acute Achilles tendon rupture in athletes. *Foot Ankle Clin*. 2013 Jun;18(2):319-338.
3. Egger AC, Berkowitz MJ. Achilles tendon injuries. *Curr Rev Musculoskelet Med*. 2017;10:72-80.
4. Vosseller JT, Ellis SJ, Levine DS, Kennedy JG, Elliott AJ, DeLand JT, Roberts MM, O'Malley MJ. Achilles tendon rupture in women. *Foot Ankle Int*. 2013 Jan;34(1):49-53.
5. Maffulli N, Giai Via A, Oliva F. Chronic Achilles Tendon Rupture. *Open Orthop J*. 2017 Jul 31;11:660-669.
6. Jean-Luc Besse. IFFAS Symposium 3, September. 2014.
7. Nilsson-Helander K, Silbernagel KG, Thomee R, Faxen E, Olsson N, Eriksson BL. Acute Achilles tendon rupture. A randomized, controlled study comparing surgical and nonsurgical treatments using validated outcome measures. *Am J Sports Med*. 2010;38:2186-2193.
8. Hunt KJ, Bundy AM, Maffulli N, Schuberth JM. Achilles tendon ruptures. *Foot&Ankle Specialist*. 2014 June;7(3).
9. Józsa L, Kvist M, Balint BJ, et al. The role of recreational sport activity in Achilles tendon rupture. A clinical, pathoanatomical, and sociological study of 292 cases. *Am J Sports Med*. 1989;17(3):338-343.
10. Kannus P, Natri A. Aetiology and pathophysiology of tendon ruptures in sports. *Scand J Med Sci Sports*. 1997;7(2):107-112.
11. Laseter JT, Russell JA. Anabolic steroid-induced tendon pathology: a review of the literature. *Med Sci Sports Exerc*. 1991;23:1-3.
12. Khaliq Y, Zhanel GG. Fluoroquinolone-associated tendinopathy: a critical review of the literature. *Clin Infect Dis*. 2003;36(11):1404-1410.
13. Parmar C, Meda KP. Achilles tendon rupture associated with combination therapy of levofloxacin and steroid in four patients and a review of the literature. *Foot Ankle Int*. 2007;28:1287-1289.
14. Wertz J, Galli M, Borchers JR. Achilles Tendon Rupture: Risk Assessment for Aerial and Ground Athletes. *Sports Health*. 2012;5(5).
15. Magnusson SP, Qvortup K, Larsen JO, et al. Collagen fibril size and crimp morphology in ruptured and intact Achilles tendons. *Matrix Biol*. 2002;21(4):369-377.
16. McCarthy MM, Hannafin JA. The Mature Athlete: Aging Tendon and Ligament. *SPORTS HEALTH*. 2014 Jan-Feb;6(1).
17. Claessen FMAP, de Vos RG, Reijman M, Meuffels DE. Predictors of Primary Achilles Tendon Ruptures. *Sports Med*. 2014;44:1241-1259.
18. Smith FB, Smith BA. Musculoskeletal differences between males and female. *Sports Med Arth Rev*. 2002;10:98-100.
19. Frizziero A, Vittadini F, Gasparre G, Masiero S. Impact of oestrogen deficiency and aging on tendon: concise review. *MLTJ* 2014;4(3):324-328.
20. Lemoine JK, Lee JT, Trappe TA. Impact of sex and chronic resistance training on human patellar tendon dry mass, collagen content, and collagen cross-linking. *Am J Physiol Regul Integr Comp Physiol*. 2009;296:119-124.
21. Cook JL, Khan KM, Kiss ZS, Griffiths L. Patellar tendinosis in junior basketball players: a controlled clinical and ultrasonographic study of 268 tendons in players aged 14-18 years. *Scand J Med Sci Sports*. 2000;10(4):216-230.
22. Józsa L, Balint JB, Kannus P, et al. Distribution of blood groups in patients with tendon rupture. An analysis of 832 cases. *J Bone Joint Surg Br*. 1989;71(2):272-274.
23. Kujala UM, Järvinen M, Natri A, et al. ABO blood groups and musculoskeletal injuries. *Injury*. 1992;23(2):131-133.
24. Oliva F, Piccirilli E, Berardi AC, et al. Hormones and tendinopathies: the current evidence. *Br Med Bull*. 2016:1-20.
25. Battery L, Maffulli N. Inflammation in Overuse Tendon Injuries. *Sports Med Arthrosc Rev*. 2011;19:13-217.
26. Hast MW, Abboud JA, Soslowsky LJ. Exploring the role of hypercholesterolemia in tendon health and repair. *MLTJ*. 2014; 4:275-279.
27. Dogan A, Korkmaz M, Cengiz N, Kalender AM, Gokalp MA. Biomechanical comparison of Achilles tenotomy and achilloplasty techniques in young rats: an experimental study. *J Am Podiatr Med Assoc*. 2009 May-Jun;99(3):216-222.
28. Lusardi DA, Cain JE Jr. The effect of fibrin sealant on the strength of tendon repair of full thickness tendon lacerations in the rabbit Achilles tendon. *J Foot Ankle Surg*. 1994 Sep-Oct;33(5):443-447.
29. Jielile J, Asilehan B, Wupuer A, et al. Early Ankle Mobilization Promotes Healing in a Rabbit Model of Achilles Tendon Rupture. *Orthopedics*. 2016 Jan-Feb;39(1):e117-126.
30. Aydın BK, Altan E, Acar MA, Erkoçak ÖF, Ugraş S. Effect of Ankaferd blood stopper® on tendon healing: an experimental study in a rat model of Achilles tendon injury. *Eklemler Hastalıkları Cerrahisi*. 2015;26(1):31-37.
31. Dabak TK, Sertkaya O, Acar N, Donmez BO, Ustunel I. The Effect of Phospholipids (Surfactant) on Adhesion and Biomechanical Properties of Tendon: A Rat Achilles Tendon Repair Model. *Biomed Res Int*. 2015;2015:689314.
32. Aliodoust M, Bayat M, Jalili MR, et al. Evaluating the effect of low-level laser therapy on healing of tenotomized Achilles tendon in streptozotocin-induced diabetic rats by light microscopical and gene expression examinations. *Lasers Med Sci*. 2014 Jul;29(4):1495-1503.
33. Gereli A, Akgün U, Uslu S, Ağır I, Ateş F, Nalbantoğlu U. The effect of organic silicon injection on Achilles tendon healing in rats. *Acta Orthop Traumatol Turc*. 2014;48(3):346-354.
34. Liang JI, Lin PC, Chen MY, Hsieh TH, Chen JJ, Yeh ML. The effect of tenocyte/hyaluronic acid therapy on the early recovery of healing Achilles tendon in rats. *J Mater Sci Mater Med*. 2014 Jan;25(1):217-227.
35. Selek O, Buluç L, Muezzinoğlu 3, Ergün RE, Ayhan S, Karaöz E. Mesenchymal stem cell application improves tendon healing via anti-apoptotic effect (Animal study). *Acta Orthop Traumatol Turc*. 2014;48(2):187-195.
36. Zeytin K, Ciloğlu NS, Ateş F, Vardar Aker F, Ercan F. The effects of resveratrol on tendon healing of diabetic rats. *Acta Orthop Traumatol Turc*. 2014;48(3):355-362.
37. Hapa O, Erduran M, Havitçioğlu H, Çeçen B, Akşahin E, Güler

- S, Atalay K. Strength of different Krackow stitch configurations using high-strength suture. *J Foot Ankle Surg.* 2013 Jul-Aug;52(4):448-450.
38. Huri G, Biçer ÖS, Ozgözen L, Uçar Y, Garbis NG, Hyun YS. A novel repair method for the treatment of acute Achilles tendon rupture with minimally invasive approach using button implant: a biomechanical study. *Foot Ankle Surg.* 2013 Dec;19(4):261-266.
 39. Nouruzian M, Alidoust M, Bayat M, Bayat M, Akbari M. Effect of low-level laser therapy on healing of tenotomized Achilles tendon in streptozotocin-induced diabetic rats. *Lasers Med Sci.* 2013 Feb;28(2):399-405.
 40. Leek BT, Tasto JP, Tibor LM, Healey RM, Freemont A, Linn MS, et al. Augmentation of tendon healing with butyric acid-impregnated sutures: biomechanical evaluation in a rabbit model. *Am J Sports Med.* 2012 Aug;40(8):1762-1771.
 41. Ni T, Senthil-Kumar P, Dubbin K, et al. A photoactivated nanofiber graft material for augmented Achilles tendon repair. *Lasers Surg Med.* 2012 Oct;44(8):645-652.
 42. Ishiyama N, Moro T, Ohe T. Reduction of Peritendinous adhesions by hydrogel containing biocompatible phospholipid polymer MPC for tendon repair. *J Bone Joint Surg Am.* 2011 Jan 19;93(2):142-149.
 43. Ishiyama N, Moro T, Ishihara K, et al. The prevention of peritendinous adhesions by a phospholipid polymer hydrogel formed in situ by spontaneous intermolecular interactions. *Biomaterials.* 2010 May;31(14):4009-4016.
 44. Lyras DN, Kazakos K, Georgiadis G, et al. Does a single application of PRP alter the expression of IGF-I in the early phase of tendon healing? *J Foot Ankle Surg.* 2011 May-Jun;50(3):276-282.
 45. Saygi B, Yildirim Y, Cabukoğlu C, Kara H, Ramadan SS, Esemenli T. The effect of dehydration and irrigation on the healing of Achilles tendon: an experimental study. *Ulus Travma Acil Cerrahi Derg.* 2008 Apr;14(2):103-109.
 46. Chong AK, Ang AD, Goh JC, Hui JH, Lim AY, Lee EH, Lim BH. Bone marrow-derived mesenchymal stem cells influence early tendon-healing in a rabbit Achilles tendon model. *J Bone Joint Surg Am.* 2007 Jan;89(1):74-81.
 47. Gilbert TW, Stewart-Akers AM, Simmons-Byrd A, Badylak SF. Degradation and remodeling of small intestinal submucosa in canine Achilles tendon repair. *J Bone Joint Surg Am.* 2007 Mar;89(3):621-630.
 48. Duygulu F, Karaoğlu S, Zeybek ND, Kaymaz FF, Güneş T. The effect of subcutaneously injected nicotine on Achilles tendon healing in rabbits. *Knee Surg Sports Traumatol Arthrosc.* 2006 Aug;14(8):756-761.
 49. Strauch B, Patel MK, Rosen DJ, Mahadevia S, Brindzei N, Pilla AA. Pulsed magnetic field therapy increases tensile strength in a rat Achilles' tendon repair model. *J Hand Surg Am.* 2006 Sep;31(7):1131-1135.
 50. Bolt P, Clerk AN, Luu HH, et al. BMP-14 gene therapy increases tendon tensile strength in a rat model of Achilles tendon injury. *J Bone Joint Surg Am.* 2007 Jun;89(6):1315-1320.
 51. Zantop T, Gilbert TW, Yoder MC, Badylak SF. Extracellular matrix scaffolds are repopulated by bone marrow derived cells in a mouse model of Achilles tendon reconstruction. *J Orthop Res.* 2006 Jun;24(6):1299-1309.
 52. Chan BP, Amann C, Yaroslavsky AN, et al. Photochemical repair of Achilles tendon rupture in a rat model. *J Surg Res.* 2005 Apr;124(2):274-279.
 53. Kashiwagi K, Mochizuki Y, Yasunaga Y, Ishida O, Deie M, Ochi M. Effects of transforming growth factor-beta 1 on the early stages of healing of the Achilles tendon in a rat model. *Scand J Plast Reconstr Surg Hand Surg.* 2004;38(4):193-197.
 54. Orhan Z, Ozturan K, Guven A, Cam K. The effect of extracorporeal shock waves on a rat model of injury to tendo Achillis. A histological and biomechanical study. *J Bone Joint Surg Br.* 2004 May;86(4):613-618.
 55. Kazımoğlu C, Bölükbaşı S, Kanatlı U, Senköylü A, Altun NS, Babaç C, et al. A novel biodegradable PCL film for tendon reconstruction: Achilles tendon defect model in rats. *Int J Artif Organs.* 2003 Sep;26(9):804-812.
 56. Palmes D, Spiegel HU, Schneider TO, Langer M, Stratmann U, Budny T, Probst A. Achilles tendon healing: longterm biomechanical effects of postoperative mobilization and immobilization in a new mouse model. *J Orthop Res.* 2002 Sep;20(5):939-946.
 57. Thermann H, Frerichs O, Holch M, Biewener A. Healing of Achilles tendon, an experimental study: part 2- Histological, immunohistological and ultrasonographic analysis. *Foot Ankle Int.* 2002 Jul;23(7):606-613.
 58. Rickert M, Jung M, Adiyaman M, Richter W, Simank HG. A growth and differentiation factor-5 (GDF-5)-coated suture stimulates tendon healing in an Achilles tendon model in rats. *Growth Factors.* 2001;19(2):115-126.
 59. Pneumatics SG, Phd PCN, McGarvey WC, Mody DR, Trevino SG. The effects of early mobilization in the healing of Achilles tendon repair. *Foot Ankle Int.* 2000 Jul;21(7):551-557.
 60. Owoeye I, Spielholz NI, Fetto J, Nelson AJ. Low-intensity pulsed galvanic current and the healing of tenotomized rat Achilles tendons: preliminary report using load-to-breaking measurements. *Arch Phys Med Rehabil.* 1987 Jul;68(7):415-418.
 61. Petrou CG, Karachalios TS, Khaldi L, Karantanas AH, Lyritis GP. Calcitonin effect on Achilles tendon healing. An experimental study on rabbits. *J Musculoskelet Neuronal Interact.* 2009 Jul-Sep;9(3):147-154.
 62. Fukawa T, Yamaguchi S, Watanabe A, et al. Quantitative Assessment of Tendon Healing by Using MR T2 Mapping in a Rabbit Achilles Tendon Transection Model Treated with Platelet-rich Plasma. *Radiology.* 2015 Sep;276(3):748-755.
 63. Adams SB Jr, Thorpe MA, Parks BG, Aghazarian G, Allen E, Schon LC. Stem cell-bearing suture improves Achilles tendon healing in a rat model. *Foot Ankle Int.* 2014 Mar;35(3):293-299.
 64. İrkören S, Demirdöver C, Akad BZ, Aytuğ Z, Yılmaz E, Oztan Y. Use of a perichondrial autograft on the peritendinous adhesion: an experimental study in rabbits. *Acta Orthop Traumatol Turc.* 2012;46(3):208-214.
 65. Meimandi-Parizi A, Oryan A, Moshiri A. Role of tissue engineered collagen based tridimensional implant on the healing response of the experimentally induced large Achilles tendon defect model in rabbits: a long term study with high clinical relevance. *J Biomed Sci.* 2013 May 14;20:28.
 66. Oryan A, Moshiri A, Parizi Meimandi A, Silver IA. A long-term in vivo investigation on the effects of xenogenous based, electrospun, collagen implants on the healing of experimentally-induced large tendon defects. *J Musculoskelet Neuronal Interact.* 2013 Sep;13(3):353-367.
 67. Godbout C, Bilodeau R, Bouchard P, Frenette J. Thrombocytopenia alters early but not late repair in a mouse model of Achilles tendon injury. *Wound Repair Regen.* 2009 Mar-Apr;17(2):260-267.
 68. Boyd RPR, Dimock R, Solan MC, Porter E. Achilles tendon rupture: how to avoid missing the diagnosis *Br J Gen Pract.* 2015 Dec;65(641):668-669.
 69. Maffulli N. The clinical diagnosis of subcutaneous tear of the Achilles tendon. A prospective study in 174 patients. *Am J Sports Med.* 1998;26(2):266-270.

70. Van Dijk CN, Karlsson J, Maffulli N, Thermann H. Achilles tendon rupture. Current concepts DJO Publications. 2008;4:27-32.
71. Garras DN, Raikin SM, Bhat SB, Taweel N, Karanjia H. MRI is unnecessary for diagnosing acute Achilles tendon ruptures: clinical diagnostic criteria. *Clin Orthop Relat Res*. 2012 Aug;470(8):2268-2273.
72. Maffulli N. Clinical tests in sports medicine: more on Achilles tendon. *Br J Sports Med*. 1996;30:250.
73. Matles AL. Rupture of the tendo Achilles: another diagnostic sign. *Bull Hosp Joint Dis*. 1975;36:48-51.
74. Singh D. Acute Achilles tendon rupture. *Br J Sports Med*. 2017 Aug;51(15):1158-1160.
75. Lang TR, Cook J, Rio E, Gaida JE. What tendon pathology is seen on imaging in people who have taken fluoroquinolones? A systematic review. *Fundam Clin Pharmacol*. 2017;31(1):4-16.
76. Barfod KW, Riecke AF, Boesen A, Hansen P, Maier JF, Døssing S, Troelsen A. Validation of a novel ultrasound measurement of Achilles tendon length. *Knee Surg Sports Traumatol Arth*. 2015;23(11):3398-3406.
77. Pedersen M, Fredberg U, Langberg H. Sonoelastography as a diagnostic tool in the assessment of musculoskeletal alterations: a systematic review. *Ultraschall in der Medizin-European Journal of Ultrasound*. 2012;33(05):441-446.
78. Fredberg U, Bolvig L, Andersen NT. Prophylactic Training in Asymptomatic Soccer Players With Ultrasonographic Abnormalities in Achilles and Patellar. *Tendons Am J Sports Med*. 2008 Mar;36(3):451-460.
79. Flavin R, Gibney R, O'Rourke SK. A Clinical Test To Avoid Sural Nerve Injuries In Percutaneous Achilles Tendon Repairs. In *Orthopaedic Proceedings. Orthopaedic Proceedings*. 2008 Aug;90(SUPP III):493-493.
80. Ofer N, Akselrod S, Nyska M, Werner M, Glaser E, Shabat S. Motion-based tendon diagnosis using sequence processing of ultrasound images. *J Orthop Res*. 2004;22(6):1296-1302.
81. Bleakney RR, Tallon C, Wong JK, Lim KP, Maffulli N. Long-term ultrasonographic features of the Achilles tendon after rupture. *Clin J Sport Med*. 2002;12(5):273-278.
82. Cunndne G, Brophy DP, Gibney RG, FitzGerald O. Diagnosis and treatment of heel pain in chronic inflammatory arthritis using ultrasound. In *Seminars in arthritis and rheumatism*. 1996; 25(6):383-389.
83. Eliasson P, Couppe C, Lonsdale M, Svensson RB, Neergaard C, Kjær M, et al. Ruptured human Achilles tendon has elevated metabolic activity up to 1 year after repair. *Eur J Nucl Med M Imaging*. 2016;43(10): 1868-1877.
84. Jielile J, Badalihan A, Qianman B, Satewalede T, Wuerliebieke J, Kelamu M, Jialihasi A. Clinical outcome of exercise therapy and early post-operative rehabilitation for treatment of neglected Achilles tendon rupture: a randomized study. *Knee Surg Sports Traumatol Arth*. 2016;24(7): 2148-2155.
85. Busilacchi A, Olivieri M, Ulisse S, Gesuita R, Skrami E, Lording T, et al. Real-time sonoelastography as novel follow-up method in Achilles tendon surgery. *Knee Surg Sports Traumatol Arth*. 2016;24(7):2124-2132.
86. Chiu CH, Yeh WL, Tsai MC, Chang SS, Hsu KY, Chan YS. Endoscopy-assisted percutaneous repair of acute Achilles tendon tears. *Foot & Ankle International*. 2013;34(8):1168-1176.
87. Jielile J, Sabirhazi G, Hu G, Chen J, Aldyarhan K, Zheyiken J, et al. Novel surgical technique and early kinesiotherapy for acute Achilles tendon rupture. *Foot & Ankle International*. 2012;33(12):1119-1127.
88. Gigante A, Moschini A, Verdenelli A, Del Torto M, Ulisse S, De Palma L. Open versus percutaneous repair in the treatment of acute Achilles tendon rupture: a randomized prospective study. *Knee Surgery, Sports Traumatol Arthr*. 2008;16(2):204-209.
89. Maffulli N, Tallon C, Wong J, Lim KP, Bleakney R. No adverse effect of early weight bearing following open repair of acute tears of the Achilles tendon. *J Sports Med Physical Fitness*. 2003a;43(3):367.
90. Costa ML, Shepstone L, Darrah C, Marshall T, Donell ST. Immediate full-weight-bearing mobilisation for repaired Achilles tendon ruptures: a pilot study. *Injury*. 2003;34(11):874-876.
91. Maffulli N, Tallon C, Wong J, Lim KP, Bleakney R. Early weightbearing and ankle mobilization after open repair of acute midsubstance tears of the Achilles tendon. *Am J Sports Med*. 2003;31(5):692-700.
92. Möller M, Kålebo P, Tidebrant G, Movin T, Karlsson J. The ultrasonographic appearance of the ruptured Achilles tendon during healing: a longitudinal evaluation of surgical and non-surgical treatment, with comparisons to MRI appearance. *Knee Surg Sports Traumatol Arthr*. 2002;10(1):49-56.
93. Bullock MJ, Mourelatos J, Mar A. Achilles Impingement Tendinopathy on Magnetic Resonance Imaging. *J Foot Ankle Surg*. 2017;56(3):555-563.
94. Aguila Maldonado R, Ruta S, Valuntas ML, García M. Ultrasonography assessment of heel entheses in patients with spondyloarthritis: a comparative study with magnetic resonance imaging and conventional radiography. *Clin Rheumatol*. 2017;36(8):1811-1817.
95. Stoller DW. *Magnetic Resonance Imaging in Orthopaedics and Sports Medicine*. 2007:2336.
96. Berquist. *MRI of the Musculoskeletal System*. 2012.
97. Rosenberg ZS, Beltran J, Bencardino JT. From the RSNA Refresher Courses. Radiological Society of North America. MR imaging of the ankle and foot. *Radiographics*. 2000;20 Spec No:S153-179.
98. Neumayer F, Mouhsine E, Arletta Y, Gremion G, Wettstein M, Crevoisier X. A new conservative-dynamic treatment for the acute ruptured Achilles tendon. *Arch Orthop Trauma Surg*. 2010;130(3):363-368.
99. Metz R, Verleisdonk EJ, van der Heijden GJ, et al. Acute Achilles tendon rupture: minimally invasive surgery versus nonoperative treatment with immediate full weightbearing. A randomized controlled trial. *Am J Sports Med*. 2008;36(9): 1688-1694.
100. Willits K, Amendola A, Bryant D, et al. Operative versus non-operative treatment of acute Achilles tendon ruptures: a multicenter randomized trial using accelerated functional rehabilitation. *J Bone Joint Surg Am*. 2010;92(17):2767-2775.
101. Soroceanu A, Sidhwa F, Aarabi S, Kaufman A, Glazebrook M. Surgical versus nonsurgical treatment of acute Achilles tendon rupture: a meta-analysis of randomized trials. *J Bone Joint Surg Am*. 2012;94(23):2136-2143.
102. Wilkins R, Bisson LJ. Operative versus nonoperative management of acute Achilles tendon ruptures: a quantitative systematic review of randomized controlled trials. *Am J Sports Med*. 2012;40:2154-2160.
103. Olsson N, Silbernagel KG, Eriksson BI, et al. Stable surgical repair with accelerated rehabilitation versus nonsurgical treatment for acute Achilles tendon ruptures: a randomized controlled study. *Am J Sports Med*. 2013;41:2867-2876.
104. Kaniki N, Willits K, Mohtadi N, et al. A Retrospective Comparative Study With Historical Control to Determine the Effectiveness of Platelet-Rich Plasma as Part of Nonoperative Treatment of Acute Achilles Tendon Rupture. *Arthroscopy*. 2014; 30(9):1139-1145.

105. Mark-Christensen T, Troelsen A, Kalleose T, Barfod KW. Functional rehabilitation of patients with acute Achilles tendon rupture: a meta-analysis of current evidence. *Knee Surg Sports Traumatol Arthrosc.* 2014.
106. Young SW, Patel A, Zhu M, et al. Weight-bearing in the non-operative treatment of acute Achilles tendon ruptures: a randomized controlled trial. *J Bone Joint Surg Am.* 2014;96:1073-1079.
107. Zhang H, Tang H, He Q, et al. Surgical versus conservative intervention for acute Achilles tendon rupture: a PRISMA-Compliant systematic review of overlapping meta-analyses. *Medicine.* 2015;94(45):1-7.
108. Lantto I, Heikkinen J, Flinkkila T, et al. Epidemiology of Achilles tendon ruptures: increasing incidence over a 33-year period. *Scand J Med Sci Sports.* 2015;25:133-138.
109. Khan RJ, Carey Smith RL. Surgical interventions for treating acute Achilles tendon ruptures. *Cochrane Database Syst Rev.* 2010 Sep 8;(9):CD003674.
110. Aviña Valencia JA, Guillen Alcalá MA. Repair of acute Achilles tendon rupture. Comparative study of two surgical techniques. *Acta Ortop Mex.* 2009;23(3):125-129.
111. Kou J. AAOS Clinical Practice Guideline: acute Achilles tendon rupture. *J Am Acad Orthop Surg.* 2010 Aug;18(8):511-513.
112. Jiang N, Wang B, Chen A, Dong F, Yu B. Operative versus nonoperative treatment for acute Achilles tendon rupture: a meta-analysis based on current evidence. *Int Orthop.* 2012 Apr;36(4):765-773.
113. Jones MP, Khan RJ, Carey Smith RL. Surgical interventions for treating acute Achilles tendon rupture: key findings from a recent cochrane review. *J Bone Joint Surg Am.* 2012 Jun. 20;94(12):e88.
114. Wu Y, Lin L, Li H, Zhao Y, Liu L, Jia Z, et al. Is surgical intervention more effective than non-surgical treatment for acute Achilles tendon rupture? A systematic review of overlapping meta-analyses. *Int J Surg.* 2016 Dec;36(Pt A):305-311.
115. Miyamoto W, Imade S, Innami K, Kawano H, Takao M. Acute Achilles Tendon Rupture Treated by Double Side-Locking Loop Suture Technique With Early Rehabilitation. *Foot Ankle Int.* 2017 Feb;38(2):167-173.
116. Yang B, Liu Y, Kan S, Zhang D, Xu H, Liu F, Ning G, Feng S. Outcomes and complications of percutaneous versus open repair of acute Achilles tendon rupture: A meta-analysis. *Int J Surg.* 2017 Apr;40:178-186.
117. Del Buono A, Volpin A, Maffulli N. Minimally invasive versus open surgery for acute Achilles tendon rupture: a systematic review. *Br Med Bull.* 2014;109:45-54.
118. Li CG, Li B, Yang YF. Management of acute Achilles tendon rupture with tendon-bundle technique. *J Int Med Res.* 2017 Feb;45(1):310-319.
119. Lewis N, Quitkin HM. Strength analysis and comparison of the Teno Fix Tendon Repair System with the two-strand modified Kessler repair in the Achilles tendon. *Foot Ankle Int.* 2003 Nov;24(11):857-860.
120. Manent A, Lopez L, Vilanova J, Mota T, Alvarez J, Santamaría A, Oliva XM. Assessment of the Resistance of Several Suture Techniques in Human Cadaver Achilles Tendons. *J Foot Ankle Surg.* 2017 Sep-Oct;56(5):954-959.
121. Aktas S, Kocaoglu B, Nalbantoglu U, Seyhan M, Guven O. End-to-end versus augmented repair in the treatment of acute Achilles tendon ruptures. *J Foot Ankle Surg.* 2007 Sep-Oct;46(5):336-340.
122. Ozer H, Selek HY, Harput G, Oznur A, Baltaci G. Achilles Tendon Open Repair Augmented With Distal Turndown Tendon Flap and Posterior Crural Fasciotomy. *J Foot Ankle Surg.* 2016 Nov-Dec;55(6):1180-1184.
123. Kocaoglu B, Ulku TK, Gereli A, et al. Evaluation of absorbable and non absorbable sutures for repair of Achilles tendon rupture with a suture-guiding device. *Foot Ankle Int.* 2015 Jun;36(6):691-695.
124. Kara A, Celik H, Seker A, et al. Granuloma formation secondary to Achilles tendon repair with non absorbable suture. *Int J Surg Case Rep.* 2014;5:720-722.
125. Ollivere BJ, Bosman HA, Bearcroft PW, Robinson AH. Foreign body granulomatous reaction associated with polyethylene "Fiberwire®" suture material used in Achilles tendon repair. *Foot Ankle Surg.* 2014 Jun;20(2):e27-29.
126. Baig MN, Yousaf I, Galbraith JG, Din R. Absorbable Polydioxanone (PDS) suture provides fewer wound complications than polyester (ethibond) suture in acute Tendo-Achilles rupture repair. *Ir Med J.* 2017 May 10;110(5):566.
127. Sadoghi P, Rosso C, Valderrabano V, et al. Initial Achilles tendon repair strength-synthesized biomechanical data from 196 cadaver repairs. *Int Orthop (SICOT).* 2012;36:1947-1951.
128. Herbolt M, Haber A, Zantop T, et al. Biomechanical comparison of the primary stability of suturing Achilles tendon rupture: a cadaver study of Bunnell and Kessler techniques under cyclic loading conditions. *Arch Orthop Trauma Surg.* 2008 Nov;128(11):1273-1277.
129. McCoy BW, Haddad SL. The strength of Achilles tendon repair: a comparison of three suture techniques in human cadaver tendons. *Foot Ankle Int.* 2010 Aug;31(8):701-705.
130. Sánchez M, Anitua E, Azofra J, et al. Comparison of surgically repaired Achilles tendon tears using platelet-rich fibrin matrices. *Am J Sports Med.* 2007;35(2):245-251.
131. Schepull T, Kvist J, Norrman H, et al. Autologous platelets have no effect on the healing of human Achilles tendon ruptures: a randomized single-blind study. *Am J Sports Med.* 2011;39(1):38-47.
132. De Carli A, Lanzetti RM, Ciompi A, et al. Can platelet-rich plasma have a role in Achilles tendon surgical repair? *Knee Surg Sports Traumatol Arthrosc.* 2016;24(7):2231-2237.
133. Alvitti F, Gurzi M, Santilli V, Paoloni M, Padua R, Bernetti A, et al. Achilles Tendon Open Surgical Treatment With Platelet-Rich Fibrin Matrix Augmentation: Biomechanical Evaluation. *J Foot Ankle Surg.* 2017;56:581-585.
134. Zou J, Mo X, Shi Z, Li T, Xue J, Mei G, Li X. A Prospective Study of Platelet-Rich Plasma as Biological Augmentation for Acute Achilles Tendon Rupture Repair. *Biomed Res Int.* 2017;2016:9364170.
135. Keating JF, Will EM. Operative versus non-operative treatment of acute rupture of tendo Achillis. *J Bone Joint Surg Br.* 2011 Aug;93(8):1071-1078.
136. Nistor L. Surgical and non-surgical treatment of Achilles Tendon rupture. A prospective randomized study. *J Bone Joint Surg Am.* 1981 Mar;63(3):394-399.
137. Cetti R, Christensen SE, Ejsted R, Jensen NM, Jorgensen U. Operative versus nonoperative treatment of Achilles tendon rupture. A prospective randomized study and review of the literature. *Am J Sports Med.* 1993 Nov-Dec;21(6):791-799.
138. Möller M, Movin T, Granhed H, Lind K, Faxén E, Karlsson J. Acute rupture of tendon Achillis. A prospective randomised study of comparison between surgical and non-surgical treatment. *J Bone Joint Surg Br.* 2001;Aug;83(6):843-848.
139. Twaddle BC, Poon P. Early motion for Achilles tendon ruptures: is surgery important? A randomized, prospective study. *Am J Sports Med.* 2007 Dec;35(12):2033-2038.
140. Kolodziej L, Bohatyrewicz A, Kromuszczyska J, et al. Efficacy and Complications of Open and Minimally Invasive Surgery in Acute Achilles Tendon Rupture: a Prospective Randomised

- Clinical Study and preliminary Report. *Int Orthop*. 2013 Apr;37(4):625-629.
141. Cretnik A, Kosanovic M, Smrkolj V. Percutaneous versus open repair of the ruptured Achilles tendon: a comparative study. *Am J Sports Med*. 2005;33(9):1369-1379.
 142. Aktas S, Kocaoglu B. Open versus minimal invasive repair with Achillon device. *Foot Ankle Int*. 2009 May;30(5):391-397.
 143. Karabinas PK, Benetos IS, Lampropoulou-Adamidou K, et al. Percutaneous versus Open Repair of Acute Achilles Tendon Ruptures. *Eur J Orthop Surg Traumatol*. 2014 May;24(4):607-613.
 144. Lim J, Dalal R, Waseem M. Percutaneous vs Open Repair of the Ruptured Achilles Tendon: A Prospective Randomized Controlled Study. *Foot Ankle Int*. 2001 Jul;22(7):559-568.
 145. Henriquez H, Munoz R, Carcuro G, et al. Is percutaneous repair better than open repair in acute Achilles tendon rupture? *Clin Orthop Relat Res* 470. 2012. 998e1003.
 146. Carmont MR, Heaver C, Pradhan A, Mei-Dan O, Gravare Silbernagel K. Surgical repair of the ruptured Achilles tendon: the cost-effectiveness of open versus percutaneous repair. *Knee Surg Sports Traumatol Arthrosc*. 2013 Jun;21(6):1361-1368.
 147. Miller D, Waterston S, Reaper J, et al. Conservative management, percutaneous or open repair of acute Achilles tendon rupture: a retrospective study. *Scott Med J*. 2005 Nov;50(4):160-165.
 148. Chan AP, Chan YY, Fong DT, Wong PY, Lam HY, Lo CK, et al. Clinical and biomechanical outcome of minimal invasive and open repair of the Achilles tendon. *Sports Med Arthrosc Rehabil Ther Technol*. 2011 Dec. 20;3(1):32.
 149. Goren D, Ayalon M, Nyska M. Isokinetic strength and endurance after percutaneous and open surgical repair of Achilles tendon ruptures. *Foot Ankle Int*. 2005 Apr;26(4):286-290.
 150. Daghino W, Enrietti E, Sprio AE, di Prun NB, Berta GN, Massè A. Subcutaneous Achilles tendon rupture: A comparison between open technique and mini-invasive tenorrhaphy with Achillon® suture system. *Injury*. 2016 Nov;47(11):2591-2595.
 151. Haji A, Sahai A, Symes A, Vyas JK. Percutaneous versus open tendon Achillis repair. *Foot Ankle Int*. 2004 Apr;25(4):215-218.
 152. Zhao HM, Yu GR, Yang YF, Zhou JQ, Aubeeluck A. Outcomes and complications of operative versus non-operative treatment of acute Achilles tendon rupture: a meta-analysis. *Chin Med J (Engl)*. 2011 Dec;124(23):4050-4055.
 153. Rebeccato A, Santini S, Salmaso G, Nogarini L. Repair of the Achilles tendon rupture: a functional comparison of three surgical techniques. *J Foot Ankle Surg*. 2001 Jul-Aug;40(4):188-194.
 154. De Carli A, Vadalà A, Ciardini R, Iorio R, Ferretti A. Spontaneous Achilles tendon ruptures treated with a mini-open technique: clinical and functional evaluation. *J Sports Med Phys Fitness*. 2009 Sep;49(3):292-296.
 155. Ng ES, Ng YO, Gupta R, Lim F, Mah E. Repair of acute Achilles tendon rupture using a double-ended needle. *J Orthop Surg (Hong Kong)*. 2006 Aug;14(2):142-146.
 156. Bhattacharyya M, Gerber B. Mini-invasive surgical repair of the Achilles tendon—does it reduce post-operative morbidity? *Int Orthop*. 2009 Feb;33(1):151-156.
 157. Mukundan C, El Husseiny M, Rayan F, Salim J, Budgen A. “Mini-open” repair of acute tendo Achilles ruptures—the solution? *Foot Ankle Surg*. 2010 Sep;16(3):122-125.
 158. Vadalà A, De Carli A, Vulpiani MC, Iorio R, Vetrano M, Scapellato S, Suarez T, Di Salvo F, Ferretti A. Clinical, functional and radiological results of Achilles tenorrhaphy surgically treated with mini-open technique. *J Sports Med Phys Fitness*. 2012 Dec.
 159. Vadalà A, Lanzetti RM, Ciompi A, Rossi C, Lupariello D, Ferretti A. Functional evaluation of professional athletes treated with a mini-open technique for Achilles tendon rupture. *MLTJ*. 2014 Jul 14;4(2):177-181.
 160. Keller A, Ortiz C, Wagner E, Wagner P, Mococain P. Mini-open tenorrhaphy of acute Achilles tendon ruptures: medium-term follow-up of 100 cases. *Am J Sports Med*. 2014 Mar;42(3):731-736.
 161. Klein EE, Weil L Jr, Baker JR, Weil LS Sr, Sung W, Knight J. Retrospective analysis of mini-open repair versus open repair for acute Achilles tendon ruptures *Foot Ankle Spec*. 2013 Feb;6(1):15-20.
 162. Bartel AF, Elliott AD, Roukis TS. Incidence of complications after Achillon® mini-open suture system for repair of acute mid-substance Achilles tendon ruptures: a systematic review. *J Foot Ankle Surg*. 2014 Nov-Dec;53(6):744-746.
 163. Taştan E, Emre TY, Demircioğlu DT, Demiralp B, Kirdemir V. Long Term Results of Mini-Open Repair Technique in the Treatment of Acute Achilles Tendon Rupture: A Prospective Study. *J Foot Ankle Surg*. 2016 Sep-Oct;55(5):971-975.
 164. Jallageas R, Bordes J, Daviet JC, Mabit C, Costec C. Evaluation of surgical treatment for ruptured Achilles tendon in 31 athletes. *Orthop Traumatol: Sur Res*. 2013;99:577-584.
 165. Zayni R, Coursier R, Zakaria M, Desrousseaux JF, Cordonnier D, Polveche G. Activity level recovery after acute Achilles tendon rupture surgically repaired: a series of 29 patients with a mean follow-up of 46 months. *MLTJ*. 2017;7:69-77.
 166. Tagliavoro G, Biz C, Mastrangelo G, Aldegheri R. The repair of the Achilles tendon rupture: comparison of two percutaneous techniques. *Strat Traum Limb Recon*. 2011;6:147-154.
 167. Bradley JP, Tibone JE. Percutaneous and open surgical repairs of Achilles tendon ruptures. A comparative study. *Am J Sports Med*. 1990;18:188-195.
 168. Tenenbaum S, Dreierangel N, Segal A, Herman A, Israeli A, Chechik A. The percutaneous surgical approach for repairing acute Achilles tendon rupture: a comprehensive outcome assessment. *J Am Podiatr Med Assoc*. 2010;100:270-275.
 169. Maes R, Copin G, Averous C. Is percutaneous repair of the Achilles tendon a safe technique? A study of 124 cases. *Acta Orthop Belg*. 2006;72:179-183.
 170. Lacoste S, Féron JM, Cherriera B. Percutaneous Tenolig® repair under intra-operative ultrasonography guidance in acute Achilles tendon rupture. *Orthop Traumatol Sur Res*. 2014;100:925-930.
 171. Maffulli N and Leadbetter WB. Free gracilis tendon graft in neglected tears of the Achilles tendon. *Clin J Sport Med*. 2005 Mar;15(2):56-61.
 172. El Shewy MT, El Barbary HM, Abdel-Ghani H. Repair of chronic rupture of the Achilles tendon using 2 intratendinous flaps from the proximal gastrocnemius-soleus complex. *Am J Sports Med*. 2009 Aug;37(8):1570-1577.
 173. Maffulli N, Spiezia F, Longo UG, Denaro V. Less-invasive reconstruction of chronic Achilles tendon ruptures using a peroneus brevis tendon transfer. *Am J Sports Med*. 2010 Nov;38(11):2304-2312.
 174. Us AK, Bilgin SS, Aydin T, Mergen E. Repair of neglected Achilles tendon ruptures: procedures and functional results. *Arch Orthop Trauma Surg*. 1997;116(6-7):408-411.
 175. Kissel CG, Blackledge DK, Crowley DL. Repair of neglected Achilles tendon ruptures: procedure and functional results. *J Foot Ankle Surg*. 1994 Jan-Feb;33(1):46-52.
 176. Esenyel CZ, Tekin C, Cakar M, et al. Surgical treatment of the neglected Achilles tendon rupture with Hyalofect. *J Am Podiatr Med Assoc*. 2014 Sep-Oct;104(5):434-443.

177. Guclu B, Basat HC, Yildirim T, Bozduman O, Us AK. Long-term Results of Chronic Achilles Tendon Ruptures Repaired With V-Y Tendon Plasty and Fascia Turndown. *Foot Ankle Int.* 2016 Jul;37(7):737-742.
178. Rush JH. Operative repair of neglected rupture of the tendo Achillis. *Aust N Z J Surg.* 1980 Aug;50(4):420-422.
179. Wapner KL, Pavlock GS, Hecht PJ, Naselli F, Walther R. Repair of chronic Achilles tendon rupture with flexor hallucis longus tendon transfer. *Foot Ankle.* 1993 Oct;14(8):443-449.
180. Pintore E, Barra V, Pintore R, Maffulli N. Peroneus brevis tendon transfer in neglected tears of the Achilles tendon. *J Trauma.* 2001 Jan;50(1):71-78.
181. Ademoğlu Y, Ozerkan F, Ada S, Bora A, Kaplan I, Kayalar M, Kul F. Reconstruction of skin and tendon defects from wound complications after Achilles tendon rupture. *J Foot Ankle Surg.* 2001 May-Jun;40(3):158-165.
182. Wong MW, Ng VW. Modified flexor hallucis longus transfer for Achilles insertional rupture in elderly patients. *Clin Orthop Relat Res.* 2005 Feb;(431):201-206.
183. Elias I, Besser M, Nazarian LN, Raikin SM. Reconstruction for missed or neglected Achilles tendon rupture with V-Y lengthening and flexor hallucis longus tendon transfer through one incision. *Foot Ankle Int.* 2007 Dec;28(12):1238-1248.
184. Mahajan RH, Dalal RB. Flexor hallucis longus tendon transfer for reconstruction of chronically ruptured Achilles tendons. *J Orthop Surg (Hong Kong).* 2009 Aug;17(2):194-198.
185. Maffulli N, Spiezia F, Pintore E, Longo UG, Testa V, Capasso G, Denaro V. Peroneus brevis tendon transfer for reconstruction of chronic tears of the Achilles tendon: a long-term follow-up study. *J Bone Joint Surg Am.* 2012 May 16;94(10):901-905.
186. Rahm S, Spross C, Gerber F, Farshad M, Buck FM, Espinosa N. Operative treatment of chronic irreparable Achilles tendon ruptures with large flexor hallucis longus tendon transfers. *Foot Ankle Int.* 2013 Aug;34(8):1100-1110.
187. Dumbre Patil SS, Dumbre Patil VS, Basa VR, Dombale AB. Semitendinosus Tendon Autograft for Reconstruction of Large Defects in Chronic Achilles Tendon Ruptures. *Foot Ankle Int.* 2014 Jul;35(7):699-705.
188. Singh A, Nag K, Roy SP, Gupta RC, Gulati V, Agrawal N. Repair of Achilles tendon ruptures with peroneus brevis tendon augmentation. *J Orthop Surg (Hong Kong).* 2014 Apr;22(1):52-55.
189. Khiami F, Di Schino M, Sariali E, Cao D, Rolland E, Catonné Y. Treatment of chronic Achilles tendon rupture by shortening suture and free sural triceps aponeurosis graft. *Orthop Traumatol Surg Res.* 2013 Sep;99(5):585-591.
190. Maffulli N, Oliva F, Costa V, Del Buono A. The management of chronic rupture of the Achilles tendon: minimally invasive peroneus brevis tendon transfer. *Bone Joint J.* 2015 Mar;97-B(3):353-357.
191. Ahmad J, Jones K, Raikin SM. Treatment of Chronic Achilles Tendon Ruptures With Large Defects. *Foot Ankle Spec.* 2016 Oct;9(5):400-408.
192. Gedam PN, Rushnawala FM. Endoscopy-Assisted Achilles Tendon Reconstruction With a Central Turndown Flap and Semitendinosus Augmentation. *Foot Ankle Int.* 2016 Dec;37(12):1333-1342. Epub. 2016 Sep. 20.
193. Maffulli N, Loppini M, Longo UG, Maffulli GD, Denaro V. Minimally invasive reconstruction of chronic Achilles tendon ruptures using the ipsilateral free semitendinosus tendon graft and interference screw fixation. *Am J Sports Med.* 2013 May;41(5):1100-1107.
194. Mann RA, Holmes GB Jr, Seale KS, Collins DN. Chronic rupture of the Achilles tendon: a new technique of repair. *J Bone Joint Surg Am.* 1991 Feb;73(2):214-219.
195. Elgohary HEA, Elmoghazy NA, Abd Ellatif MS. Combined flexor hallucis longus tendon transfer and gastrocnemius recession for reconstruction of gapped chronic Achilles tendon ruptures. *Injury.* 2016 Dec;47(12):2833-2837. Doi: 10.1016/j.injury.2016.10.029. Epub. 2016 Nov 3.
196. Miao X, Wu Y, Tao H, Yang D, Huang L. Reconstruction of Kuwada grade IV chronic Achilles tendon rupture by minimally invasive technique. *Indian J Orthop.* 2016 Sep;50(5):523-528.
197. Maffulli N, Oliva F, Del Buono A, Florio A, Maffulli G. Surgical management of Achilles tendon re-ruptures: a prospective cohort study. *Int Orthop.* 2015 Apr;39(4):707-714.
198. Yeoman TF, Brown MJ, Pillai A. Early post-operative results of neglected tendo-Achilles rupture reconstruction using short flexor hallucis longus tendon transfer: a prospective review. *Foot (Edinb).* 2012 Sep;22(3):219-223.
199. Park YS, Sung KS. Surgical reconstruction of chronic Achilles tendon ruptures using various methods. *Orthopedics.* 2012 Feb 17;35(2):e213-218.200. Sarzaeem MM, Lemraski MM, Safdari F. Chronic Achilles tendon rupture reconstruction using a free semitendinosus tendon graft transfer. *Knee Surg Sports Traumatol Arthrosc.* 2012 Jul;20(7):1386-1391.
201. Zheng L, Zhang XS, Dong ZG, Liu LH, Wei JW. One-staged reconstruction of Achilles tendon and overlying skin defects with suppuration: using peroneus brevis tendon transfer and reversed sural neurofasciocutaneous flap. *Arch Orthop Trauma Surg.* 2011 Sep;131(9):1267-1272.
202. Wegrzyn J, Luciani JF, Philippot R, Brunet-Guedj E, Moyen B, Besse JL. Chronic Achilles tendon rupture reconstruction using a modified flexor hallucis longus transfer. *Int Orthop.* 2010 Dec;34(8):1187-1192.
203. Lee KB, Park YH, Yoon TR, Chung JY. Reconstruction of neglected Achilles tendon rupture using the flexor hallucis tendon. *Knee Surg Sports Traumatol Arthrosc.* 2009 Mar;17(3):316-320.
204. Fotiadis E, Chatzissimeon A, Samoladas E, Antonarakos P, Akritopoulos P, Akritopoulou K. A Combined Repair Technique for Early Neglected Achilles Tendon Ruptures. *Eur J Trauma Emerg Surg.* 2008 Feb;34(1):37-42.
205. Lui TH. Endoscopic assisted flexor hallucis tendon transfer in the management of chronic rupture of Achilles tendon. *Knee Surg Sports Traumatol Arthrosc.* 2007 Sep;15(9):1163-1166. Epub. 2007 May 30.
206. Miskulin M, Miskulin A, Klobucar H, Kuvajla S. Neglected rupture of the Achilles tendon treated with peroneus brevis transfer: a functional assessment of 5 cases. *J Foot Ankle Surg.* 2005 Jan-Feb;44(1):49-56.
207. Dalal RB, Zenios M. The flexor hallucis longus tendon transfer for chronic tendo-Achilles ruptures revisited. *Ann R Coll Surg Engl.* 2003 Jul;85(4):283.
208. Seker A, Kara A, Armagan R, Oc Y, Varol A, Sezer HB. Reconstruction of neglected Achilles tendon ruptures with gastrocnemius flaps: excellent results in long-term follow-up. *Arch Orthop Trauma Surg.* 2016 Oct;136(10):1417-1423.
209. Lapidus LJ, Ray BA, Hamberg P. Medial Achilles tendon island flap: a novel technique to treat ruptures and neglected ruptures of the Achilles tendon. *Int Orthop.* 2012 Aug;36(8):1629-1634.
210. Takao M, Ochi M, Naito K, Uchio Y, Matsusaki M, Oae K. Repair of neglected Achilles tendon rupture using gastrocnemius fascial flaps. *Arch Orthop Trauma Surg.* 2003 Nov;123(9):471-474. Epub. 2002 Oct 25.
211. Ozan F, Dogar F, Gurbuz K, Ekinci Y, Koyuncu S, Sekban H. Chronic Achilles Tendon Rupture Reconstruction Using the Lindholm Method and the Vulpius Method. *J Clin Med Res.* 2017 Jul;9(7):573-578.

212. Sanada T, Uchiyama E. Gravity Equinus Position to Control the Tendon Length of Reversed Free Tendon Flap Reconstruction for Chronic Achilles Tendon Rupture. *J Foot Ankle Surg.* 2017 Jan-Feb;56(1):37-41
213. Maffulli N, Del Buono A, Loppini M, Denaro V. Ipsilateral free semitendinosus tendon graft with interference screw fixation for minimally invasive reconstruction of chronic tears of the Achilles tendon. *Oper Orthop Traumatol.* 2014 Oct;26(5):513-519.
214. El Shazly O, Abou El Soud MM, El Mikkawy DM, El Ganzoury I, Ibrahim AM. Endoscopic-assisted Achilles tendon reconstruction with free hamstring tendon autograft for chronic rupture of Achilles tendon: clinical and isokinetic evaluation. *Arthroscopy.* 2014 May;30(5):622-628.
215. Tay D, Lin HA, Tan BS, Chong KW, Rikhraj IS. Chronic Achilles tendon rupture treated with two turnover flaps and flexor hallucis longus augmentation: two-year clinical outcome. *Ann Acad Med Singapore.* 2010 Jan;39(1):58-60.
216. Nilsson-Helander K, Swärd L, Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. A new surgical method to treat chronic ruptures and reruptures of the Achilles tendon. *Knee Surg Sports Traumatol Arthrosc.* 2008 Jun;16(6):614-620.
217. Tawari AA, Dhamaogankar AA, Goregaonkar AB, Chhapan JB. Augmented repair of degenerative tears of tendo Achilles using peroneus brevis tendon: early results. *Malays Orthop J.* 2013 Mar;7(1):19-24.
218. Oksanen MM, Haapasalo HH, Elo PP, Laine HJ. Hypertrophy of the flexor hallucis longus muscle after tendon transfer in patients with chronic Achilles tendon rupture. *Foot Ankle Surg.* 2014 Dec;20(4):253-257.
219. Zappia M, Berritto D, Oliva F, et al. High resolution real time ultrasonography of the sural nerve after percutaneous repair of the Achilles tendon. *Foot Ankle Surg.* 2017;37:636-643.
220. Cohen M. US imaging in operated tendons. *J Ultrasound.* 2012;15:69-75. Doi: 10.1016/j.jus.2011.11.001.
221. Zappia M, Cuomo G, Martino MT, et al. The effect of foot position on Power Doppler Ultrasound grading of Achilles enthesitis. *Rheumatol Int.* 2016;36:871-874.
222. Fornage BD. Achilles tendon: US examination. *Radiology.* 1986;159:759-764.
223. Blei CL, Nirschl RP, Grant EG. Achilles tendon: US diagnosis of pathologic conditions. Work in progress. *Radiology.* 1986;159:765-767.
224. Rupp S, Tempelhof S, Fritsch E. Ultrasound of the Achilles tendon after surgical repair: morphology and function. *Br J Radiol.* 1995;68:454-458.
225. Möller M, Kälebo P, Tidebrant G, et al. The ultrasonographic appearance of the ruptured Achilles tendon during healing: a longitudinal evaluation of surgical and nonsurgical treatment, with comparisons to MRI appearance. *Knee Surg Sports Traumatol Arthrosc.* 2002;10:49-56.
226. Gitto S, Draghi AG, Bortolotto C, Draghi F. Sonography of the Achilles Tendon After Complete Rupture Repair: What the Radiologist Should Know. *J Ultrasound Med.* 2016. 35:2529-2536. D
227. Diao Z-B, Chu H-K, Li N, et al. [Short-term clinical effects of Achillon in repair of acute Achilles tendon rupture]. *Zhongguo Gu Shang.* 2012;25:959-961.
228. Klein EE, Weil L, Baker JR, et al. Retrospective Analysis of Mini-Open Repair Versus Open Repair for Acute Achilles Tendon Ruptures. *Foot Ankle Spec.* 2013;6:15-20. .
229. Chun KA, Cho K-H. Postoperative ultrasonography of the musculoskeletal system. *Ultrasound (Seoul, Korea).* 2015;34: 195-205.
230. Zhang L, Wan W, Wang Y, et al. Evaluation of Elastic Stiffness in Healing Achilles Tendon After Surgical Repair of a Tendon Rupture Using In vivo Ultrasound Shear Wave Elastography. *Med Sci Monit.* 2016;22:1186-1191. .
231. Tan S, Kudaş S, Özcan AŞ, et al. Real-time sonoelastography of the Achilles tendon: pattern description in healthy subjects and patients with surgically repaired complete ruptures. *Skeletal Radiol.* 2012;41:1067-1072..
232. Shalabi A, Kristoffersen-Wiberg M, Aspelin P, Movin T. MR evaluation of chronic Achilles tendinosis. A longitudinal study of 15 patients preoperatively and two years postoperatively. *Acta Radiol.* 2001;42:269-276.
233. Fujikawa A, Kyoto Y, Kawaguchi M, et al. Achilles tendon after percutaneous surgical repair: serial MRI observation of uncomplicated healing. *AJR Am J Roentgenol.* 2007;189:1169-1174.
234. Karjalainen PT, Aronen HJ, Pihlajamäki HK, et al. Magnetic Resonance Imaging During Healing of Surgically Repaired Achilles Tendon Ruptures. *Am J Sports Med.* 1997;25:164-171.
235. Hahn F, Meyer P, Maiwald C, et al. Treatment of Chronic Achilles Tendinopathy and Ruptures with Flexor Hallucis Tendon Transfer: Clinical Outcome and MRI Findings. *Foot Ankle Int.* 2008;29:794-802.
236. Sölveborn S-A, Moberg A. Immediate Free Ankle Motion After Surgical Repair of Acute Achilles Tendon Ruptures. *Am J Sports Med.* 1994;22:607-610.
237. Karjalainen PT, Ahovuo J, Pihlajamäki HK, et al. Postoperative MR Imaging and Ultrasonography of Surgically Repaired Achilles Tendon Ruptures. *Acta radiol.* 1996;37:639-646.
238. Sarman H, Atmaca H, Cakir O, et al. Assessment of Postoperative Tendon Quality in Patients With Achilles Tendon Rupture Using Diffusion Tensor Imaging and Tendon Fiber Tracking. *J Foot Ankle Surg.* 2015;54:782-786.
239. Valkering KP, Aufwerber S, Ranuccio F, Lunini E, Edman G, Ackermann PW. Functional weight-bearing mobilization after Achilles tendon rupture enhances early healing response: a single-blinded randomized controlled trial. *Knee Surg Sports Traumatol Arthrosc.* 2017 Jun;25(6):1807-1816
240. Lantto I, Heikkinen J, Flinkkila T, Ohtonen P, Kangas J, Siira P, Leppilahti J. Early functional treatment versus cast immobilization in tension after Achilles rupture repair: results of a prospective randomized trial with 10 or more years of follow-up. *Am J Sports Med.* 2015 Sep;43(9):2302-2309.
241. Suchak AA, Bostick GP, Beaupré LA, Durand DC, Jomha NM. The influence of early weight-bearing compared with non-weight-bearing after surgical repair of the Achilles tendon. *J Bone Joint Surg Am.* 2008 Sep;90(9):1876-1883.
242. Costa ML, MacMillan K, Halliday D, Chester R, Shepstone L, Robinson AH, Donell ST. Randomised controlled trials of immediate weight-bearing mobilization for rupture of the tendo Achillis. *J Bone Joint Surg Br.* 2006 Jan;88(1):69-77.
243. Kangas J, Pajala A, Siira P, Hämäläinen M, Leppilahti J. Early functional treatment versus early immobilization in tension of the musculotendinous unit after Achilles rupture repair: a prospective, randomized, clinical study. *J Trauma.* 2003 Jun;54(6):1171-1180.
244. Kerkhoffs GM, Struijs PA, Raaymakers EL, Marti RK. Functional treatment after surgical repair of acute Achilles tendon rupture: wrap vs walking cast. *Arch Orthop Trauma Surg.* 2002 Mar;122(2):102-105.
245. Mortensen HM, Skov O, Jensen PE. Early motion of the ankle after operative treatment of a rupture of the Achilles tendon. A prospective, randomized clinical and radiographic study. *J*

- Bone Joint Surg Am. 1999 Jul;81(7):983-990.
246. De la Fuente C, Peña y Lillo R, Carreño G, Marambio H. Prospective randomized clinical trial of aggressive rehabilitation after acute Achilles tendon ruptures repaired with Dresden technique. *Foot (Edinb)*. 2016 Mar;26:15-22.
 247. Groetelaers RP, Janssen L, van der Velden J, Wieland AW, Amendt AG, Geelen PH, Janzing HM. Functional Treatment or Cast Immobilization After Minimally Invasive Repair of an Acute Achilles Tendon Rupture: Prospective, Randomized Trial. *Foot Ankle Int*. 2014 Aug;35(8):771-778.
 248. Majewski M, Schaeren S, Kohlhaas U, Ochsner PE. Postoperative rehabilitation after percutaneous Achilles tendon repair: early functional therapy versus cast immobilization. *Disabil Rehabil*. 2008;30(20-22):1726-1732.
 249. Kader D, Mosconi M, Benazzo F, Maffulli N. Achilles Tendon Rupture. *Tendon Injuries*. 2005:187-200.
 250. Sharma P, Maffulli N. Biology of tendon injury: healing, modeling and remodeling. *J Musculoskelet Neuronal Interact*. 2006; 6:181-190.
 251. Kannus P, Jozsa L, Jarvinen M. Basic science of tendons. In: Garrett WJ, Speer K, Kirkendall DT (eds). *Principles and Practice of Orthopaedic Sports Medicine*. Lippincott Williams & Wilkins, Philadelphia. 2000:21-37.
 252. Williams JG. Achilles tendon lesions in sport. *Sports Med*. 1986;3:114-135.
 253. Maffulli N, Moller HD, Evans CH. Tendon healing: can it be optimized? *Br J Sports Med*. 2002;36:315-16.
 254. Oakes BW. Tissue healing and repair: tendons and ligaments. In: Frontera WR (ed). *Rehabilitation of Sports Injuries: Scientific Basis*. Blackwell Science, Oxford. 2003:56-98.
 255. Tillman LJ, Chasan NP. Properties of dense connective tissue and wound healing. In: Hertling D, Kessler RM (eds). *Management of Common Musculoskeletal Disorders*. Lippincott, Philadelphia. 1996:8-21.
 256. Hooley CJ, Cohen RE. A model for the creep behavior of tendon. *Int J Biol Macromol*. 1979;1:123-132.
 257. Abrahamsson SO. Matrix metabolism and healing in the flexor tendon. Experimental studies on rabbit tendon. *Scand J Plast Reconstr Surg Hand Surg Suppl*. 1991;23:1-51.
 258. Amiel D, Akeson W, Harwood FL, Frank CB. Stress deprivation effect on metabolic turnover of medial collateral ligament collagen. *Clin Orthop*. 1987;172:25-27.
 259. Enwemeka CS. Functional loading augments the initial tensile strength and energy absorption capacity of regenerating rabbit Achilles tendons. *Am J Phys Med Rehabil*. 1992;71:31-38.
 260. Kannus P, Jozsa L, Natri A, Jarvinen M. Effects of training, immobilization and remobilization on tendons. *Scand J Med Sci Sports*. 1997;7:67-71.
 261. Bring DK, Reno C, Renstrom P, Salo P, Hart DA, Ackermann PW. Joint immobilization reduces the expression of sensory neuropeptide receptors and impairs healing after tendon rupture in a rat model. *J Orthop Res*. 2009;27:274-280.
 262. Maffulli N, King JB. Effects of physical activity on some components of the skeletal system. *Sports Med* 1992;13:393-407.
 263. Akeson WH, Amiel D, Mechanic GL, Woo SL, Harwood FL, Hamer ML. Collagen cross-linking alterations in joint contractures: changes in the reducible cross-links in periarticular connective tissue collagen after nine weeks of immobilization. *Connect Tissue Res*. 1977;5:15-19.
 264. Maganaris CN, Reeves ND, Rittweger J, Sargeant AJ, Jones DA, Gerrits K, DeHaan A. Adaptive response of human tendon to paralysis. *Muscle Nerve*. 2006;33:85-92.
 265. Chiodo CP, Glazebrook M, Bluman EM, et al. Diagnosis and treatment of acute Achilles tendon rupture. *J Am Acad Orthop Surg*. 2010;18:503-510.
 266. Jones MP, Khan R, Smith R. Surgical Interventions for treating acute Achilles tendon rupture: key findings from a recent Cochrane review. *J Bone Joint Surg*. 2012;94:881-886.
 267. Suchak AA, Spooner C, Reid DC, Jomha NM. Postoperative rehabilitation protocols for Achilles tendon ruptures: a meta-analysis. *Clin Orthop Relat Res*. 2006;445:216-221.
 268. Saleh M, Marshall PD, Senior R, MacFarlane A. The Sheffield splint for controlled early mobilisation after rupture of the calcaneal tendon. A prospective, randomised comparison with plaster treatment. *J Bone Joint Surg Br*. 1992;74:206-209.
 269. Petersen OF, Nielsen MB, Jensen KH, Solgaard S. Randomized comparison of CAM walker and light-weight plaster cast in the treatment of first-time Achilles tendon rupture. *Ugeskr Laeger*. 2002;164:3852-3855.
 270. Hutchison AM, Topliss C, Beard D, Evans RM, Williams P. The Treatment of a rupture of the Achilles tendon using a dedicated management programme. *Bone Joint J*. 2015 Apr;97-B(4):510-515.
 271. Barfod KW, Brencke J, Lauridsen HB, Ban I, Ebskov L, Troelsen A. Nonoperative dynamic treatment of acute Achilles tendon rupture: the influence of early weight-bearing on clinical outcome: a blinded, randomized controlled trial. *J Bone Joint Surg Am*. 2014;96:1497-1503.
 272. Calder JD, Saxby TS. Early, active rehabilitation following mini-open repair of Achilles tendon rupture: a prospective study. *Br J Sports Med*. 2005 Nov;39(11):857-859.
 273. Bossi P. *Conoscere le apparecchiature elettromedicali*. McGraw-Hill Education (1/11/2004) ISBN-10: 8838616531 ISBN-13: 978-8838616532 , Italian. 2004.
 274. Notarnicola A, Pesce V, Vicenti G, Tafuri S, Forcignano M, Moretti B. SWAAT study: extracorporeal shock wave therapy and arginine supplementation and other nutraceuticals for insertional Achilles tendinopathy. *Advances in Therapy*. 2012; 29(9):799-814.
 275. Balius R, Álvarez G, Baró F, Jiménez F, Pedret C, Costa E, Martínez-Puig D. A 3-Arm Randomized Trial for Achilles Tendinopathy: Eccentric Training, Eccentric Training Plus a Dietary Supplement Containing Mucopolysaccharides, or Passive Stretching Plus a Dietary Supplement Containing Mucopolysaccharides. *Curr Ther Res- Clin Exp* 2016;78:1-7.
 276. Hai Binh B, Ramirez P, Martinez-Puig D. A randomized, placebo-controlled study to evaluate efficacy and safety of a dietary supplement containing mucopolysaccharides, collagen type I and vitamin C for management of different tendinopathies. *Ann Rheum Dis*. 2014;73.
 277. Nadal F, Bové T, Sanchis D, Martinez-Puig D. 473 Effectiveness of treatment of tendinitis and plantar fasciitis by tendoactive™. *Osteoarthritis and Cartilage*. 2009;17:S253.
 278. Arquer A, García M, Laucirica JA, et al. The efficacy and safety of oral mucopolysaccharide, type I collagen and vitamin C treatment in tendinopathy patients. *Apuntes Medicina de l'Esport*. 2014;49(182):31-36.
 279. Mavrogenis S, Johannessen E, Jensen P, Sindberg C. The effect of essential fatty acids and antioxidants combined with physiotherapy treatment in recreational athletes with chronic tendon disorders. A randomised, double-blind, placebo-controlled study. *Phys Ther Sport*. 2004;5(4):194-199.
 280. Ahmad J, Repka M, Raikin SM. Treatment of myotendinous Achilles ruptures. *Foot Ankle Int*. 2013;34:1074-1078.
 281. Al-Mouazzen L, Rajakulendran K, Najefi A, et al. Percutaneous repair followed by accelerated rehabilitation for acute Achilles tendon ruptures. *J Orthop Surg (Hong Kong)*. 2015; 23:352-356.
 282. Amin NH, Old AB, Tabb LP, et al. Performance outcomes after repair of complete Achilles tendon ruptures in national basket-

- ball association players. *Am J Sports Med.* 2013;41:1864-1868.
283. Amlang MH, Christiani P, Heinz P, et al. Die perkutane Naht der Achillessehne mit dem Dresdner Instrument. *Oper Orthop Traumatol.* 2006;18:287-299.
284. Ateschrang A, Gratzner C, Weise K. Incidence and effect of calcifications after open-augmented Achilles tendon repair. *Arch Orthop Trauma Surg.* 2008;128:1087-1092.
285. Bassi JL, Mahindra P. A modified flap technique as an alternate procedure for open Achilles tendon repair (the Bassi method). *Oper Orthop Traumatol.* 2006;18:171-181.
286. Bevoni R, Angelini A, D'Apote G, et al. Long term results of acute Achilles repair with triple-bundle technique and early rehabilitation protocol. *INJURY.* 2014;45:1268-1274.
287. Bostick GP, Jomha NM, Suchak AA, et al. Factors associated with calf muscle endurance recovery 1 year after Achilles tendon rupture repair. *J Orthop Sports Phys Ther.* 2010;40:345-351.
288. Boyden EM, Kitaoka HB, Cahalan TD, et al. Late versus early repair of Achilles tendon rupture. Clinical and biomechanical evaluation. *Clin Orthop Relat Res.* 1995;317:150-158.
289. Carmont MR, Grävare Silbernagel K, Brorsson A, et al. The Achilles tendon resting angle as an indirect measure of Achilles tendon length following rupture, repair, and rehabilitation. *Asia-Pacific J Sport Med Arthrosc Rehabil Technol.* 2015;2:49-55.
290. Ceccarelli F, Berti L, Giuriati L, et al. Percutaneous and minimally invasive techniques of Achilles tendon repair. *Clin Orthop Relat Res.* 2007;458:188-193.
291. Chandrakant V, Lozano-Calderon S, McWilliam J. Immediate weight bearing after modified percutaneous Achilles tendon repair. *Foot Ankle Int.* 2012;33:1093-1097.
292. Chen Z, Wei J, Hou Z, et al. Application of internal fixation of steel-wire limited loop in early Achilles tendon rupture. *Asian Pac J Trop Med.* 2013;6:902-907.
293. Coutts A, MacGregor A, Gibson J, et al. Clinical and functional results of open operative repair for Achilles tendon rupture in a non-specialist surgical unit. *J R Coll Surg Edinb.* 2002;47:753-762.
294. Cretnik A, Frank A. Incidence and outcome of rupture of the Achilles tendon. *Wien Klin Wochenschr.* 2004;116:33-38.
295. Cretnik A, Kosir R, Kosanović M. Incidence and outcome of operatively treated Achilles tendon rupture in the elderly. *Foot Ankle Int.* 2010;31:14-18.
296. Demirel M, Turhan E, Dereboy F, et al. Augmented repair of acute tendo Achilles ruptures with gastrosoleus turn down flap. *Indian J Orthop.* 2011;45:45-52.
297. Doral MN. What is the effect of the early weight-bearing mobilisation without using any support after endoscopy-assisted Achilles tendon repair? *Knee Surg Sport Traumatol Arthrosc.* 2013;21:1378-1384.
298. Eames MHA, Eames NWA, McCarthy KR, et al. An audit of the combined non-operative and orthotic management of ruptured tendo Achillis. *Injury.* 1997;28:289-292.
299. Feldbrin Z, Hendel D, Lipkin A, et al. Achilles tendon rupture and our experience with the Achillon device. *Isr Med Assoc J.* 2010;12:609-612.
300. Fernández-Fairén M, Gimeno C. Augmented repair of Achilles tendon ruptures. *Am J Sports Med.* 1997;25:177-181.
301. Fortis AP, Dimas A, Lamprakis AA. Repair of Achilles tendon rupture under endoscopic control. *Arthrosc J Arthrosc Relat Surg.* 2008;24:683-688.
302. Garabito A, Martinez-Miranda J, Sanchez-Sotelo J. Augmented repair of acute Achilles tendon ruptures using gastrocnemius-soleus fascia. *Int Orthop.* 2005;29:42-46.
303. Garrido IM, Deval JC, Bosch MN, et al. Treatment of acute Achilles tendon ruptures with Achillon device: clinical outcomes and kinetic gait analysis. *Foot Ankle Surg.* 2010;16:189-194.
304. Gorschewsky O, Vogel U, Schweizer A, et al. Percutaneous tenodesis of the Achilles tendon. A new surgical method for the treatment of acute Achilles tendon rupture through percutaneous tenodesis. *Injury.* 1999;30:315-321.
305. Gorschewsky O, Pitzl M, Pütz A, et al. Percutaneous repair of acute Achilles tendon rupture. *Foot Ankle Int.* 2004;25:219-224.
306. Guillo S, Del Buono A, Dias M, et al. Percutaneous repair of acute ruptures of the tendo Achillis. *Surg.* 2013;11:14-19.
307. Halasi T, Tállay A, Berkes I. Percutaneous Achilles tendon repair with and without endoscopic control. *Knee Surg Sports Traumatol Arthrosc.* 2003;11:409-414.
308. Hohendorf B, Siepen W, Spiering L, et al. Long-term results after operatively treated Achilles tendon rupture: fibrin glue versus suture. *J Foot Ankle Surg.* 2008;47:392-399.
309. Hufner TM, Brandes DB, Thermann H, et al. Long-term results after functional nonoperative treatment of Achilles tendon rupture. *Foot Ankle Int.* 2006;27:167-171.
310. Jaakkola JI, Beskin JL, Griffith LH, et al. Early ankle motion after triple bundle technique repair vs. casting for acute Achilles tendon rupture. *Foot Ankle Int.* 2001;22:979-984.
311. Jacob KM, Paterson R. Surgical repair followed by functional rehabilitation for acute and chronic Achilles tendon injuries: excellent functional results, patient satisfaction and no reruptures. *ANZ J Surg.* 2007;77:287-291.
312. Jennings AG, Sefton GK, Newman RJ. Repair of acute rupture of the Achilles tendon: a new technique using polyester tape without external splintage. *Ann R Coll Surg Engl.* 2004;86:445-448.
313. Josey RA, Marymont J V, Varner KE, et al. Immediate, full weight bearing cast treatment of acute Achilles tendon ruptures: a long-term follow-up study. *Foot Ankle Int.* 2003;24:775-779.
314. Jung HG, Lee KB, Cho SG, et al. Outcome of Achilles tendon ruptures treated by a limited open technique. *Foot Ankle Int.* 2008;29:803-807.
315. Kakiuchi M. A combined open and percutaneous technique for repair of tendo Achillis: comparison with open repair. *J Bone Joint Surg Br.* 1995;77:60-63.
316. Karkhanis S, Mumtaz H, Kurdy N. Functional management of Achilles tendon rupture: a viable option for non-operative management. *Foot Ankle Surg.* 2010;16:81-86.
317. Knobe M, Gradl G, Klos K, et al. Is percutaneous suturing superior to open fibrin gluing in acute Achilles tendon rupture? *Int Orthop.* 2015;39:535-542.
318. Korkmaz M, Erkoc MF, Yolcu S, et al. Weight bearing the same day versus non-weight bearing for 4 weeks in Achilles tendon rupture. *J Orthop Sci.* 2015;20:513-516.
319. Kraus R, Stahl J-P, Meyer C, et al. Frequency and effects of intratendinous and peritendinous calcifications after open Achilles tendon repair. *Foot Ankle Int.* 2004;25:827-832.
320. Labib SA, Hoffer CE, Shah JN, et al. The gift box open Achilles tendon repair method: a retrospective clinical series. *J Foot Ankle Surg.* 2016;55:39-44.
321. Lansdaal JR, Goslings JC, Reichart M, et al. The results of 163 Achilles tendon ruptures treated by a minimally invasive surgical technique and functional after treatment. *Injury.* 2007;38:839-844.
322. Lee DK. A preliminary study on the effects of acellular tissue graft augmentation in acute Achilles tendon ruptures. *J Foot Ankle Surg.* 2008;47:8-12.

323. Leppilahti J, Forsman K, Puranen J, et al. Outcome and prognostic factors of Achilles rupture repair using a new scoring method. *Clin Orthop Relat Res.* 1998;346:152-161.
324. Macquet AJ, Christensen RJ, Debenham M, et al. Open repair of the acutely torn Achilles tendon under local anaesthetic. *ANZ J Surg.* 2011;81:619-623.
325. Maffulli N, Longo UG, Maffulli GD, et al. Achilles tendon ruptures in elite athletes. *Foot Ankle Int.* 2011;32:9-15.
326. Maffulli N, Longo UG, Ronga M, et al. Favorable outcome of percutaneous repair of Achilles tendon ruptures in the elderly. *Clin Orthop Relat Res.* 2010;468:1039-1046.
327. Majewski M, Rohrbach M, Czaja S, et al. Avoiding sural nerve injuries during percutaneous Achilles tendon repair. *Am J Sports Med.* 2006;34:793-798.
328. Mandelbaum BR, Myerson MS, Forster R. Achilles tendon ruptures: a new method of repair, early range of motion, and functional rehabilitation. *Am J Sports Med.* 1995;23:392-395.
329. Maniscalco P, Bertone C, Bonci E, et al. Titanium anchors for the repair of distal Achilles tendon ruptures: preliminary report of a new surgical technique. *J Foot Ankle Surg.* 1998;36:96-100.
330. Martinelli B. Percutaneous repair of the Achilles tendon in athletes. *Bull Hosp Jt Dis.* 2000;59:149-152.
331. McCormis GP, Nawoczenski DA, DeHaven K. Functional bracing for rupture of the Achilles tendon: clinical results and analysis of ground-reaction forces and temporal data. *J Bone Jt Surg.* 1997;29A:1799-1808.
332. Metz R, van der Heijden GJ, Verleisdonk EJ, et al. Effect of complications after minimally invasive surgical repair of acute Achilles tendon ruptures: report on 211 cases. *Am J Sports Med.* 2011;39:820-824.
333. Mortensen N, Saether J, Steinke M, et al. Separation of tendon ends after Achilles tendon repair: a prospective, randomized, multicenter study. *Orthopedics.* 1992;15:899-903.
334. Motta P, Errichiello C, Pontini I. Achilles tendon rupture: a new technique for easy surgical repair and immediate movement of the ankle and foot. *Am J Sports Med.* 1997;25:172-176.
335. Nestorson J, Movin T, Moller M, et al. Function after Achilles tendon rupture in the elderly: 25 patients older than 65 years followed for 3 years. *Scand J Med Sci Sports.* 2000;71:64-68.
336. Orr J, McCriskin B, Dutton J. Achillon mini-open Achilles tendon repair: early outcomes and return to duty results in U.S. military service members. *J Surg Orthop Adv.* 2013;22:23-29.
337. Ozsoy M, Cengiz B, Ozsoy A, et al. Minimally invasive Achilles tendon repair: a modification of the Achillon technique. *Foot Ankle Int.* 2013;34:1683-1688.
338. Pajala A, Kangas J, Siira P, et al. Augmented compared with nonaugmented surgical repair of a fresh total Achilles tendon rupture. A prospective randomized study. *J Bone Joint Surg Am.* 2009;91:1092-100.
339. Parekh SG, Wray WH, Brimmo O, et al. Epidemiology and outcomes of Achilles tendon ruptures in the National Football League. *Foot Ankle Spec.* 2009;2:283-286.
340. Park HG, Moon DH, Yoon JM. Limited open repair of ruptured Achilles tendons with Bunnel-type sutures. *Foot Ankle Int.* 2001;22:985-987.
341. Rajasekar K, Gholve P, Faraj A, et al. A subjective outcome analysis of tendo-Achilles rupture. *J Foot Ankle Surg.* 2005;44:32-36.
342. Rettig AC, Liotta FJ, Klootwyk TE, et al. Potential risk of rerupture in primary Achilles tendon repair in athletes younger than 30 years of age. *Am J Sports Med.* 2005;33:119-123.
343. Richardson LC, Reitman R, Wilson M. Achilles tendon ruptures: functional outcome of surgical repair with a "pull-out" wire. *Foot Ankle Int.* 2003;24:439-443.
344. Schepull T, Kvist J, Andersson C, et al. Mechanical properties during healing of Achilles tendon ruptures to predict final outcome: a pilot Roentgen stereophotogrammetric analysis in 10 patients. *BMC Musculoskelet Disord.* 2007;8:116.
345. Silbernagel KG, Steele R, Manal K. Deficits in heel-rise height and Achilles tendon elongation occur in patients recovering from an Achilles tendon rupture. *Am J Sports Med.* 2012;40:1564-1571.
346. Soldatis J, Goodfellow D, Wilber J. End-to-end operative repair of Achilles tendon rupture. *Am J Sports Med.* 1997;25:90-95.
347. Solveborn S, Moberg A. Immediate free ankle motion after surgical repair of acute Achilles tendon ruptures. *Am J Sports Med.* 1994;22:607-610.
348. Sorrenti S. Achilles tendon rupture: effect of early mobilization in rehabilitation after surgical repair. *Foot Ankle Int.* 2006;27:407-410.
349. Speck M, Klaue K. Early full weightbearing and functional treatment after surgical repair of acute Achilles tendon rupture. *Am J Sports Med.* 1998;26:789-793.
350. Stein BE, Stroh DA, Schon LC. Outcomes of acute Achilles tendon rupture repair with bone marrow aspirate concentrate augmentation. *Int Orthop.* 2015;39:901-905.
351. Strauss E, Ishak C, Jazrawi L, et al. Operative treatment of acute Achilles tendon ruptures: an institutional review of clinical outcomes. *Injury.* 2007;38:832-838.
352. Talbot J, Williams G, Bismil Q, et al. Results of accelerated postoperative rehabilitation using novel 'suture frame' repair of Achilles tendon rupture. *J Foot Ankle Surg.* 2012;51:147-151.
353. Troop RL, Losse GM, Lane JG, et al. Early motion after repair of Achilles tendon ruptures. *Foot Ankle Int.* 1995;16:705-709.
354. Uchiyama E, Nomura A, Takeda Y, et al. A modified operation for Achilles tendon ruptures. *Am J Sports Med.* 2007;35:1739-1743.
355. Valente M, Crucul M, Alecci V, et al. Minimally invasive repair of acute Achilles tendon rupture with Achillon device. *Musculoskelet Surg.* 2012;96:35-39.
356. Wagnon R, Akayi M. The Webb-Bannister percutaneous technique for acute Achilles tendon ruptures: a functional and MRI assessment. *J Foot Ankle Surg.* 2005;44:437-444.
357. Wallace RGH, Heyes GJ, Michael ALR. The non-operative functional management of patients with a rupture of the tendo Achillis leads to low rates of re-rupture. *J Bone Joint Surg Br.* 2011;93:1362-1366.
358. Wallace RGH, Traynor IER, Kernohan WG, et al. Combined conservative and orthotic management of acute ruptures of the Achilles tendon. *J Bone Joint Surg Am.* 2004;86-A:1198-1202.
359. Cappozzo A. The forces and couples in the human trunk during level walking. *J Biomech.* 1983;16(4):265-277.
360. Hof AL. An explicit expression for the moment in multibody systems. *J Biomech.* 1992;25(10):1209-1211.
361. Cappozzo A, Felici F, Figura F, Gazzani F. Lumbar spine loading during half-squat exercises. *Med Sci Sports Exerc.* 1985;17(November):613-620.
362. Kernozek T, Gheidi N, Ragan R. Comparison of estimates of Achilles tendon loading from inverse dynamics and inverse dynamics-based static optimisation during running. *J Sports Sci.* 2017;35(21):2073-2079.
363. Rajagopal A, Dembia CL, DeMers MS, Delp DD, Hicks JL, Delp SL. Full-Body Musculoskeletal Model for Muscle-Driven Simulation of Human Gait. *IEEE Trans Biomed Eng.* 2016;63(10):2068-2079.
364. Delp SL, Loan JP, Hoy MG, Zajac FE, Topp EL, Rosen JM. An

- Interactive Graphics-Based Model of the Lower Extremity to Study Orthopaedic Surgical Procedures. *IEEE Trans Biomed Eng.* 1990;37(8):757-767.
365. Fukashiro S, Komi P V., Järvinen M, Miyashita M. Comparison between the directly measured Achilles tendon force and the tendon force calculated from the ankle joint moment during vertical jumps. *Clin Biomech.* 1993;8(1):25-30.
366. Robertson G, Caldwell G, Hamill J, Kamen G, Whittlesey S. *Research Methods in Biomechanics.* Human Kinetics 2004.
367. Camomilla V, Cereatti A, Cutti AG, Fantozzi S, Stagni R, Vannozzi G. Methodological factors affecting joint moments estimation in clinical gait analysis: a systematic review. *Biomed Eng Online.* 2017;16(1):106.
368. Lamberto G, Martelli S, Cappozzo A, Mazzà C. To what extent is joint and muscle mechanics predicted by musculoskeletal models sensitive to soft tissue artefacts? *J Biomech.* 2016.
369. Martelli S, Valente G, Viceconti M, Taddei F. Sensitivity of a subject-specific musculoskeletal model to the uncertainties on the joint axes location. *Comput Methods Biomech Biomed Engin.* 2015;18(14):1555-1563.
370. Bosmans L, Valente G, Wesseling M, et al. Sensitivity of predicted muscle forces during gait to anatomical variability in musculotendon geometry. *J Biomech.* 2015;48(10):2116-2123.
371. Lichtwark GA, Wilson AM. In vivo mechanical properties of the human Achilles tendon during one-legged hopping. *J Exp Biol.* 2005;208(24):4715-4725.
372. Gerus P, Rao G, Berton E. Subject-Specific Tendon-Aponeurosis Definition in Hill-Type Model Predicts Higher Muscle Forces in Dynamic Tasks. *PLoS One.* 2012;7(8).
373. Franz JR, Thelen DG. Imaging and simulation of Achilles tendon dynamics: Implications for walking performance in the elderly. *J Biomech.* 2016;49(9):1403-1410.
374. Ralston EL, Schmidt ER Jr. Repair of the ruptured Achilles tendon. *J Trauma.* 1971;81:1019-1036.
375. Eidelman M, Nachtigal A, Katzman A, Bialik V. Acute rupture of Achilles tendon in a 7-year-old girl. *J Pediatr Orthop B.* 2004; 13:32-33.
376. Tudisco C, Bisicchia S. Reconstruction of neglected traumatic Achilles tendon rupture in a young girl. *J Orthopaed Traumatol.* 2012;13:163-166.
377. Vasileff WK, Moutzourous V. Unrecognized pediatric partial Achilles tendon injury followed by traumatic completion: a case report and literature review. *J Foot Ankle Surg.* 2014;53:485-488.
378. Padulo J, Oliva F, Frizziero A, Maffulli N. *Muscles, Ligaments and Tendons Journal - Basic principles and recommendations in clinical and field science research: 2016 update.* MLTJ. 2016;6(1):1-5.