

## TOWARDS A MAP OF THE UPPER PLEISTOCENE LOESS OF THE PO PLAIN LOESS BASIN (NORTHERN ITALY)

**Andrea Zerboni<sup>1</sup>, Rivka Amit<sup>2</sup>, Carlo Baroni<sup>3</sup>, Mauro Coltorti<sup>4</sup>, M. Francesca Ferrario<sup>5</sup>,  
Gianfranco Fioraso<sup>6</sup>, M. Gabriella Forno<sup>7</sup>, Chiara Frigerio<sup>5</sup>, Franco Gianotti<sup>7</sup>,  
Andrea Irace<sup>6</sup>, Franz Livio<sup>5</sup>, Guido S. Mariani<sup>1</sup>, Alessandro M. Michetti<sup>5</sup>, Giovanni Monegato<sup>6</sup>,  
Paolo Mozzi<sup>8</sup>, Giuseppe Orombelli<sup>9</sup>, Alessandro Perego<sup>1</sup>, Naomi Porat<sup>2</sup>, Ivano Rellini<sup>10</sup>,  
Luca Trombino<sup>1</sup>, Mauro Cremaschi<sup>1</sup>**

<sup>1</sup> Dipartimento di Scienze della Terra "A. Desio", Università degli Studi di Milano, Milano, Italy

<sup>2</sup> Geological Survey of Israel, Jerusalem, Israel

<sup>3</sup> Dipartimento di Scienze della Terra, Università degli Studi di Pisa, Pisa, Italy

<sup>4</sup> Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente, Università di Siena, Siena, Italy

<sup>5</sup> Dipartimento di Scienza e Alta Tecnologia, Università degli Studi dell'Insubria, Como, Italy

<sup>6</sup> IGG - CNR, Torino, Italy

<sup>7</sup> Dipartimento di Scienze della Terra, Università degli Studi di Torino, Torino, Italy

<sup>8</sup> Dipartimento di Geoscienze, Università degli Studi di Padova, Padova, Italy

<sup>9</sup> Dipartimento di Scienze dell'Ambiente e della Terra, Università degli Studi di Milano-Bicocca, Milano, Italy

<sup>10</sup> Dipartimento del Territorio e delle sue Risorse, Università di Genova, Genova, Italy

*Corresponding author:* A. Zerboni <[andrea.zerboni@unimi.it](mailto:andrea.zerboni@unimi.it)>

**ABSTRACT:** Upper Pleistocene (MIS 4–2) loess sequences occur in most of continental Europe and in Northern Italy along the Po Plain Loess Basin. Loess is distributed along the flanks of the Po Plain and was deposited on glacial deposits, fluvial terraces, uplifted isolated hills, karst plateaus, slopes and basins of secondary valleys. Loess bodies are generally tiny and affected by pedogenesis, being locally slightly reworked by slope processes and bioturbation. Notwithstanding, loess in the Po Plain is an important archive of paleoenvironmental record and its mapping provides new insights in paleoenvironmental and palaeoseismic reconstructions of Northern Italy.

**KEYWORDS:** Loess, Po Plain loess basin, upper Pleistocene, mapping

### 1. INTRODUCTION

In the framework of European terrestrial palaeo-records, loess is one of the best archives for Quaternary palaeoenvironmental reconstruction (e.g. Smalley & Leach, 1978; Coudé-Gaussen, 1990; Pye, 1995; Cremaschi, 2004). Maps published in the last decade that present the distribution of loess in Europe eliminate loess outcrops in Northern Italy or, alternatively, define them as loess derivatives (Haase et al., 2007; Bertran et al., 2016; Lindner et al., 2017). As a consequence, despite systematic studies carried out on the Upper Pleistocene loess sequences at the margin of the Po Plain (see Cremaschi, 1987, 2004 and references therein), loess in Northern Italy is overlooked in comparison to the classical sequences described in continental Europe (Amit & Zerboni, 2013), and crucial data are excluded from the interpretation of loess formation at a continental scale.

For this reason, in the last few years we revised the sedimentological, paleoenvironmental, and geochronological data regarding the loess from the Po Plain Loess Basin (PPLB), including new data derived from studying some classical loess outcrops (e.g., Ferraro, 2009; Wacha et al., 2011; Cremaschi et al., 2015; Cremaschi & Zerboni, 2016), as well as freshly identified sequences (e.g., Rellini et al., 2009; Zerboni et al., 2015; Frigerio et al., 2017). Moreover, in a region - the

Po Plain foredeep - of moderate seismicity loess is a stratigraphic marker of increasing interest for earthquake hazard characterization (e.g., Livio et al., 2014). To provide the necessary spatial context for these tasks, a systematic mapping of loess outcrops along the Po Plain is on-going and preliminary results are here discussed.

### 2. THE PO PLAIN LOESS BASIN

Loess is a wind-blown, mainly silty-sized sediment. It is widespread in Northern Italy, where it is typical of glacial periods, when sediment grinding (weathering), transportation, and sedimentation are enhanced by dry environmental conditions with poor vegetation cover and characterized by strong winds (Cremaschi, 1990; Orombelli, 1990; Pye, 1995).

Except for a few thick sequences of unweathered loess deposits (as the Val Sorda sequence) and/or complex pedosequences (as at Monte Netto), most of the outcrops in Northern Italy consist of thin loess sheets located on dissected fluvial terraces, glacial deposits, uplifted isolated hills, and karst plateaus (Fig. 1). These are for the most part set along the southern margin of the Alps and the northern margin of the Apennines. Loess sequences can also be found on top of polygenetic paleosols, inside sinkholes, and intercalated in rock-shelter deposits. Furthermore, loess deposits in Northern Italy are often associated with Middle Palaeolithic archaeo-




Fig. 1 - Some loess outcrops investigated in the last few years. A) The Val Sorda sequence (Cavaion Veronese, VR): loess lies on a glacial deposit. B) Loess at the margin of the Apennine (Cavriago, RE) (Cremaschi et al., 2015). C) The Monte Netto (Capriano del Colle, BS) sequence (Zerboni et al., 2015), displaced by an array of normal faults (i.e., gravity graben). D) Loess sequence at Pecetto di Valenza (AL) (Frigerio et al., 2017).

logical sites, showing that the Po Plain Loess Basin (PPLB) was a suitable environment for Mousterian hunter-gatherers.

Available chronology, based on a number of radiocarbon, TL, and OSL dating suggests that the formation of loess occurred during the Upper Pleistocene, between at least 75 and 16 ka BP (MIS 4-2) (Cremaschi, 2004; Zerboni et al., 2015). Thicker sequences (as at Ghiardo and Monte Netto sites) also record weak to moderate soil-forming processes dating to the Upper Pleistocene interstadials and the Holocene. At many sites, the uppermost part of loess sequences likely underwent moderate to strong reworking due to post-Last Glacial Maximum (post-LGM) reactivation of slope processes, enhanced Holocene bioturbation, and, more recently, human-induced processes (tillage). As a result, colluvial and bioturbated units/soil horizons were formed, whose  $^{14}\text{C}$  and OSL dating provided mid-Holocene ages that are interpreted as due respectively

to the mixing of soil humic acids over a long time and rejuvenation of dose signal. This has been demonstrated at Monte Netto and Ghiardo sequences (Cremaschi et al., 2015; Zerboni et al., 2015).

### 3. DRAWING THE LOESS MAP

The available geological maps present loess deposits only fragmentarily, as layers inside or at the top of Quaternary formations, without providing any information on the sedimentary and post-sedimentary (pedogenetic) processes affecting loess covers. On the other hand, the available pedological maps suggest the occurrence of silt-bearing soil horizons, but interpretations of sedimentary processes ruling the accretion of silt are lacking. A few attempts to reconstruct the spatial distribution of loess in the PPLB have been preliminary published by Cremaschi (2004), Cremaschi et al. (2015), Zerboni et al. (2015); moreover, the Climex Map

of the LGM (Vai & Cantelli, 2004) offers a good representation of many loess outcrops. For these reasons, the publication of a new map presenting the spatial distribution of loess is necessary and relies on field surveys in key regions, integrated by observations on (i) geological contexts and (ii) active and inactive surface processes.

At the first stage, we will identify and study landforms acting as sedimentary traps for loess aggradation (Cremaschi, 2004), such as: (i) glacial sediments (moraines) and fluvio-glacial deposits at the southern fringe of the Pre-Alps; (ii) karst plateaus (and sinkholes) between Lombardia and the Dalmatian coast, and in the Conero promontory (Marche region); (iii) isolated hills in the middle of the Po Plain, actively growing through repeated coseismic uplift during the Late Quaternary; (iv) fluvial terraces of different ages on the Apennine foothills between southern Piemonte, Emilia Romagna and Marche regions; (v) glacial and erosional surfaces in the Apennine range of Marche, Emilia Romagna, and Liguria regions, where sedimentation can be influenced by the pre-existing topography. In the map, loess outcrops will be represented distinguishing between: (i) unweathered loess; (ii) weathered loess and loessial soils displaying different degrees of weathering and characterized by distinctive soil-forming processes; and (iii) colluvial/reworked loessic sediments (loess derivatives). At a second stage, a more detailed classification of loess units will be under evaluation. This will offer a better comprehensive palaeoenvironmental context in the light of the maximum expansion of Alpine and Apennine glaciers and the shoreline of the Adriatic Sea during the LGM, which will be included in our map as reconstructed in the Climex Map (Vai & Cantelli, 2004).

#### 4. PRELIMINARY CONCLUSIONS

Because loess deposits in Northern Italy are generally thin and can be easily affected by soil-forming processes, sometimes their characteristics do not fully meet the standard requirements generally accepted for classical loess (Haase et al., 2007). Although loess deposits and loessial soils in the Po Plain are slightly different, the main sedimentological and pedological characteristics are similar in their field appearance, particle size distribution, and mineralogy, thus suggesting a similar source. The source area was identified as the Po flood plain, which was exposed during glacial periods and was subject to intense winds at that time. The investigation of several key sequences and the systematic mapping of outcrops are demonstrating the importance of North Italian loess for the reconstruction of Upper Pleistocene palaeoclimates of the northern Mediterranean region and the paleoseismic record of the Po Plain fore-deep.

#### ACKNOWLEDGEMENTS

This research has been carried out within the activities of the INQUA GEODUST International Focus Group. Financial support from Università degli Studi di Milano, Progetto Linea 2 (2016 and 2017) entrusted to AZ.

#### REFERENCES

- Amit R., Zerboni A. (2013) - Report on the INQUA-AEOMED field-trip workshop 'Reconsidering Loess in Northern Italy' (Po Plain, 1–3 July 2013). *Alpine and Mediterranean Quaternary*, 26, XI-XV.
- Bertran P., Liard M., Sitzia L., Tissoux H. (2016) - A map of Pleistocene aeolian deposits in Western Europe, with special emphasis on France. *Journal of Quaternary Science*, 31, 844-856.
- Coudé-Gaussen G. (1990) - The loess and loess-like deposits along the sides of the western Mediterranean Sea: genetic and palaeoclimatic significance. *Quaternary International*, 5, 1-8.
- Cremaschi M. (1987) - Loess deposits of the Plain of the Po and of the adjoining Adriatic basin (Northern Italy). In: Pecci M., French H.M. (Eds.), *Loess and Periglacial phenomena*. Akademiai Kiado, Budapest, Hungary, 125-140.
- Cremaschi M. (1990) - The Loess in Northern and Central Italy: a Loess Basin between the Alps and the Mediterranean Sea. C.N.R., Centro di Studio per la Stratigrafia e Petrografia delle Alpi Centrale, Milano, Italy, pp. 187.
- Cremaschi M. (2004) - Late Pleistocene loess. In: Antonioli F., Vai G.B. (Eds.), *Litho-palaeoenvironmental Maps of Italy During the Last Two Climatic Extremes*. Climex Maps Italy, Museo Geologico Giovanni Cappellini, Bologna, Italy, 34-37.
- Cremaschi M., Zerboni A. (2016) - La questione dell'apparato morenico del Garda alla luce delle recenti ricerche sul Torrion della Val Sorda. *Geologia Insubrica*, 12/1, 103.
- Cremaschi M., Zerboni A., Nicosia C., Negrino F., Rodnight H., Spötl C. (2015) - Age, soil-forming processes, and archaeology of the loess deposits at the Apennine margin of the Po plain (northern Italy): new insights from the Ghiardo area. *Quaternary International*, 376, 173-188.
- Ferraro F. (2009) - Age, sedimentation, and soil formation in the Val Sorda loess sequence, Northern Italy. *Quaternary International*, 204, 54-64.
- Frigerio C., Bonadeo L., Zerboni A., Livio F., Ferrario M.F., Fioraso G., Irace A., Brunamonte F., Michetti A.M. (2017) - First evidence for Late Pleistocene to Holocene earthquake surface faulting in the Eastern Monferrato Arc (Northern Italy): geology, pedomorphology and structural study of the Pecetto di Valenza site. *Quaternary International*, 451, 143-164.
- Haase D., Fink J., Haase G., Ruske R., Pécsi M., Richter H., Altermann M., Jäger K.D. (2007) - Loess in Europe - its spatial distribution based on a European loess map, scale 1:2,500,000. *Quaternary Science Reviews*, 26, 1301-1312.
- Lindner H., Lehmkuhl F., Zeeden C. (2017) - Spatial loess distribution in the eastern Carpathian Basin: a novel approach based on geoscientific maps and data. *Journal of Maps*, 13, 173-181.

- Livio F., Berlusconi A., Zerboni A., Trombino L., Sileo G., Michetti A.M., Rodnight H., Spötl C. (2014) - Progressive offset and surface deformation along a seismogenic blind thrust in the Po Plain foredeep (Southern Alps, Northern Italy). *Journal of Geophysical Research: Solid Earth*, 119, 7701-7721.
- Orombelli G. (1990) - Loess in Italy: an introduction. In: Cremaschi, M. (Ed.), *The Loess in Northern and Central Italy: a Loess Basin between the Alps and the Mediterranean Sea*. C.N.R., Centro di Studio per la Stratigrafia e Petrografia delle Alpi Centrale, Milano, Italy, 11-12.
- Pye K. (1995) - The nature, origin and accumulation of loess. *Quaternary Science Reviews*, 14, 653-667.
- Rellini I., Trombino L., Firpo M., Rossi P.M. (2009) - Extending westward the loess basin between the Alps and the Mediterranean region: micromorphological and mineralogical evidences from the slope of the Ligurian Alps (northern Italy). *Geografia Fisica e Dinamica Quaternaria*, 32, 103-116.
- Smalley I.J., Leach A.J. (1978) - The origin and distribution of the loess in the Danube basin and associated regions of East-Central Europe - a review. *Sedimentary Geology*, 21, 1-26.
- Vai G.B., Cantelli L. (2004) - Litho-Palaeoenvironmental Maps of Italy During the Last Two Climatic Extremes, Map 1 - Last Glacial Maximum, Map 2 - Holocene Climatic Optimum, 1:1,000,000 scale, 32<sup>nd</sup> IGC Florence 2004, Museo Geologico Giovanni Cappellini, Bologna.
- Wacha L., Pavlakovic S.M., Novothny A., Crnjakovic M., Frechen M. (2011) - Luminescence dating of Upper Pleistocene Loess from the Island of Susak in Croatia. *Quaternary International*, 234, 50-61.
- Zerboni A., Trombino L., Frigerio C., Livio F., Berlusconi A., Michetti A.M., Rodnight H., Spötl C. (2015) - The loess-paleosol sequence at Monte Netto: a record of climate change in the Upper Pleistocene of the central Po Plain, northern Italy. *Journal of Soils and Sediments*, 15, 1329-1350.

*Ms. received: May 4, 2018*  
*Final text received: May 14, 2018*