

Review

TERT promoter mutations in head and neck squamous cell carcinoma: A systematic review and meta-analysis on prevalence and prognostic significance

Paolo Boscolo-Rizzo^{a,*}, Giancarlo Tirelli^a, Jerry Polesel^b, Egidio Sia^a, Veronica Phillips^c, Daniele Borsetto^d, Anita De Rossi^{e,f}, Silvia Giunco^{e,f}

^a Department of Medical, Surgical and Health Sciences, Section of Otolaryngology, University of Trieste, Trieste, Italy

^b Unit of Cancer Epidemiology, Centro di Riferimento Oncologico di Aviano (CRO) IRCCS, Aviano, Italy

^c University of Cambridge Medical Library, Cambridge, United Kingdom

^d Department of ENT, Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, United Kingdom

^e Department of Surgery, Oncology and Gastroenterology, Section of Oncology and Immunology, University of Padova, Padova, Italy

^f Immunology and Diagnostic Molecular Oncology Unit, Veneto Institute of Oncology IOV- IRCCS, Padova, Italy



ARTICLE INFO

Keywords:

TERT promoter mutations
Head and neck squamous cell carcinoma (HNSCC)
Oral cavity squamous cell carcinoma
Prognostic biomarkers
Survival
Systematic review
Meta-analysis
Telomerase
Telomere

ABSTRACT

Objectives: To estimate the prevalence of two most common and mutually exclusive $-124 C > T$ and $-146 C > T$ TERT promoter mutations in HNSCC and analyse their prognostic role.

Materials and methods: The databases Medline (via Ovid), Embase (via Ovid), Cochrane Library, Scopus, and Web of Science (Core Collection) were searched from inception to December 2022 to identify studies analysing TERT promoter mutations in HNSCC. Pooled prevalence of TERT promoter mutations and hazard ratio (sHR) of death/progression, with corresponding confidence intervals (CI), were estimated.

Results: The initial search returned 6416 articles, of which 17 studies, including 1830 patients, met the criteria for prevalence meta-analysis. Among them, 8 studies fitted the inclusion criterion to analyse the prognostic impact of TERT promoter mutations. Overall, 21% (95% CI: 12%-31%) of HNSCCs harboured TERT promoter mutation. TERT promoter mutations were more commonly found in oral cavity cancer (prevalence = 47%, 95% CI: 33%-61%), followed by laryngeal/hypopharyngeal cancer (prevalence = 12%, 95% CI: 4%-25%), while they were quite rare in oropharyngeal cancer (prevalence = 1%, 95% CI: 0%-4%). TERT promoter mutation $-124 C > T$ was associated with a higher risk of death (sHR = 2.01, 95% CI: 1.25-3.23) and progression (sHR = 2.79, 95% CI: 1.77-4.40), while $-146 C > T$ TERT promoter mutation did not show any significant correlation neither to overall nor progression-free survival.

Conclusion: TERT promoter mutations were mainly topographically restricted to oral cavity cancer. $-124 C > T$ was the most common TERT promoter mutation and was significantly associated to worse outcome in HNSCC.

Introduction

Worldwide, approximately 880,000 people are diagnosed with a head and neck cancer each year [1]. The majority of these (i.e., about 375,000) are localized in the oral cavity, with about 185,000 in the larynx, 185,000 in the oropharynx and hypopharynx, and 135,000 in the nasopharynx. Most of head and neck cancers are head and neck squamous cell carcinomas (HNSCC) [2].

Tobacco smoking and alcohol abuse are the major established risk

factors for HNSCC [1]. The role of high-risk alpha human papillomaviruses (HR α -HPVs) infection in the aetiology of oropharyngeal SCC has increased in the last decades, being the strongest independent prognostic factor for this subset of cancers [3,4]. Conversely, no robust biological biomarkers are available for HNSCC arising from other head and neck sites including oral cavity and larynx [1]. There is, therefore, a strong need to have biomarkers able to stratify the risk in these patients.

Among molecular markers proposed for risk stratification of HNSCC patients, the detection of telomerase reverse transcriptase (TERT)

* Corresponding author at: Department of Medical, Surgical and Health Sciences, Section of Otolaryngology, University of Trieste, Strada di Fiume 447, 34149 Trieste, Italy.

E-mail address: paolo.boscolorizzo@units.it (P. Boscolo-Rizzo).

<https://doi.org/10.1016/j.oraloncology.2023.106398>

Received 3 February 2023; Received in revised form 27 March 2023; Accepted 11 April 2023

Available online 17 April 2023

1368-8375/© 2023 Published by Elsevier Ltd.

promoter mutations has attracted considerable interest given that most HNSCCs express high levels of *TERT* transcripts that have proved to be associated with worse responses to treatment and high risk of progression [5,6]. Telomerase is an enzyme that avoids the loss of telomeres at each cell replication. It presents a catalytic subunit with reverse transcriptase activity, *TERT*, and an RNA component which primes DNA synthesis from telomere repeats (TERC, telomerase RNA component). Somatic cells shorten the telomeres each cell cycle; this prevents the unlimited cell division, whereas cells that have active telomerase possess unlimited proliferative potential [7]. Since the acquisition of unlimited proliferation capacity represents a critical hallmark required for cell malignant transformation, telomere/telomerase complex is a pivotal component in the neoplastic process [8]. *TERT* is normally expressed in adult humans only in germ cells, transit-amplifying stem-like cells, and proliferating/stimulated B and T cells, but it is estimated that in 85% of cancers telomerase are reactivated during the process of carcinogenesis [9]. While the mechanisms that lead *TERT* to be reactivated in cancer cells have still to be completely understood, *TERT* reactivation can be explained by the genetic and epigenetic factors, such as *TERT* amplifications, *TERT* structural variants, rearrangements, promoter methylation, and mutations within *TERT* promoter region [9,10]. *TERT* promoter mutations mainly appear at nucleotides 1,295,228 (−124 C > T) and 1,295,250 (−146 C > T) and represent the most common non-coding mutations in solid tumours being recorded in several cancer types with a broad spectrum of prevalence [11–15]. These two mutations occur in a mutually exclusive manner, with the −124 C > T showing greater prevalence than −146 C > T. Both mutations increase *TERT* promoter activity and augment *TERT* gene transcription by creating *de novo* binding sites for E-twenty-six (ETS) transcription factors family [11,12,16]. In addition, recent evidence suggests that *TERT* polymorphisms can also play an important role in oncogenesis [17–20]. By interacting with the Wnt/ β -catenin and the NF- κ B signalling pathways, telomerase may play non-canonical functions directly linked with tumour progression, making it a possibly appealing prognostic marker [21,22]. Recently, *TERT* promoter mutations were observed to be associated with a highly aggressive behaviour in several cancers including thyroid, bladder, and melanoma skin cancers [23–25]. However, results from studies evaluating the association between *TERT* promoter mutations, cancer biology, and outcome in HNSCC were often inconsistent [17,26–29].

Thus, the aim of this systematic review and meta-analysis was to estimate the pooled prevalence of *TERT* promoter mutations in HNSCC and to investigate their prognostic significance.

Materials and methods

Search strategy

The protocol for this systematic review was registered with PROSPERO with the code CRD42022338251. The databases Medline (via Ovid), Embase (via Ovid), Cochrane Library, Scopus, and Web of Science (Core Collection) were searched from inception to December 2022. The search strategy was developed in consultation with an experienced medical librarian (VP) using the PRESS checklist and reported according to the PRISMA-S guidelines [30,31] (eMethod in Supplementary Online Content). Databases were searched separately, rather than multiple databases being searched simultaneously on the same platform. The search syntax was adapted for each database to account for variation between thesaurus terms/controlled vocabulary across each database. Exact search terms used in each database are present in supplementary materials. Results were deduplicated using Endnote 20 software. Endnote was set to identify articles as duplicates if they matched in the Author, Year, Title, Short Title, and Reference Type fields, and was also set to ignore differences in item record spacing and punctuation in these fields when identifying duplicates. The reference lists of articles included in this review, as well as narrative reviews published in the last

10 years, were also manually searched to minimize the risk of missing data. Two authors (PBR, ES) independently screened all titles and abstracts generated by the search and then evaluated the full texts of all the relevant articles identified against the inclusion criteria (Fig. 1); a third author (DB) settled discordances when present.

Outcome measures

The primary outcome of this meta-analysis was the prevalence of *TERT* promoter mutations in HNSCCs, measured as the proportion of patients carrying the mutation, distinguishing between −124 C > T, −146 C > T, or other non-specified *TERT* promoter mutations. The secondary outcome was the influence of *TERT* promoter mutations on overall survival (OS) or progression-free survival (PFS); the measure effect of the secondary outcome was the hazard ratio (HR) of death or recurrence, according to the mutational status.

Selection criteria

All observational studies that analysed *TERT* promoter mutations in HNSCCs were included. Research letters were also considered. Inclusion criteria were: 1) studies investigating *TERT* promoter mutations in patients with HNSCC; 2) patients who underwent treatment with curative intent (i.e., surgery and/or chemo/radiotherapy); 3) studies reporting *TERT* promoter mutation prevalence or HR for death or progression, with corresponding confidence interval; 4) studies evaluating *TERT* promoter mutation in surgical specimen.

The exclusion criteria were: 1) reviews and editorials; 2) studies with fewer than five patients; 3) non-English language studies; 4) studies containing aggregated and non-extractable data, or duplicated data from previously published work; 5) studies including distant metastatic cancers.

Data extraction

An electronic data-collection form was used to extract the following data: 1) study information: first author, year of publication, cohort characteristics, total number of patients; 2) clinical data: cancer site and subsite, number of cancers for each location, pathology stage, treatment; 3) analysis of the mutations: total number of *TERT* mutations, number of wild typed cancer, number of the single mutations studied; 4) data on prognosis, when available: HR of PFS and/or OS with corresponding 95% CI. Two authors (JP, ES) independently assessed the quality of the included studies with the Newcastle-Ottawa Scale (NOS) [32].

Statistical analysis

The number of total cases and of those carrying *TERT* promoter mutation, HR of death/progression and corresponding 95% CI were extracted from each study; the standard error of the log HR was derived from the log CIs. Summary estimate of *TERT* mutation prevalence and hazard ratio (sHR), with corresponding 95% CI, were calculated according to random-effects models of DerSimonian and Laird [33] as a weighted average, giving each study a weight proportional to its precision and incorporating both within- and between-study variability. Analyses were conducted separately for *TERT* promoter mutations −124 C > T and −146 C > T and for non-specified *TERT* promoter mutations. Statistical heterogeneity among studies was evaluated using the I^2 and τ^2 statistics.[33] Prevalence analysis were stratified by cancer site and, for oral cancer, by cancer subsite. Influence analysis was performed when summary estimate was estimated from five or more studies: summary estimate was calculated by omitting one study at a time. Publication bias was assessed through a funnel plot [34]. The results of the meta-analysis were presented graphically using forest plots, reporting the estimates from individual studies, the summary estimates and corresponding 95% CI. Statistical significance was claimed for $p < 0.05$.




Fig. 1. PRISMA flow chart of study selection process.

Results

Search results and study selection

Once the duplicates were eliminated, 4467 items were screened, excluding 4219 articles based on the title. The full text of the remaining 248 articles was further reviewed, and 17 articles met the inclusion criteria for the meta-analysis, as described in Fig. 1. The included articles [13–15,17,26–29,35–43] involved 1830 patients and were published between 2013 and 2021 (eTable in Supplementary Online Content). As data for oral cavity cancers were included in Giunco *et al.* [17], only findings for non-oral cavity HNSCCs by Boscolo-Rizzo *et al.* [35] were included in the meta-analysis. Among the selected studies, 11 articles evaluated the prognostic role of *TERT* promoter mutations in HNSCC, but only 8 of those, including 1102 patients, reported HR with 95% CI that fitted the requirements for this meta-analysis.

Characteristics and quality of the included studies

Nine articles focused on the prevalence of *TERT* promoter mutations in a specific head and neck site [15,17,26,27,36–39,43]. In the seven

studies that included subjects with oropharyngeal SCC [13,14,28,29,35,37,43], HPV status was available in 287 out of 311 cases (92.3%). Among them, 155 cases (54.0%) were HPV-positive based on p16 overexpression and/or presence of HPV-DNA. Among the 11 studies that analysed prognostic impact of *TERT* promoter mutations, seven studies analysed the OS [17,26,28,29,35,37,39] and four studies focused on the PFS [17,29,37,39]. Overall, quality was satisfactory, with seven out of eight studies investigating OS or PFS with NOS ≥ 7 (eTable in Supplementary Online Content).

Prevalence of *TERT* promoter mutations in HNSCC

Overall, 21% (95% CI: 12%–31%) of HNSCCs harboured *TERT* promoter mutation (Fig. 2). Differences emerged according to cancer site ($p < 0.01$): *TERT* promoter mutations were more frequent for cancer arising in the oral cavity (47%, 95% CI: 33%–61%) and larynx/hypopharynx (12%, 95% CI: 4%–25%), while *TERT* promoter mutations were quite uncommon in oropharyngeal SCC (1%, 95% CI: 0%–4%). HNSCCs harbouring –124 C > T *TERT* promoter mutation (12%; 95% CI: 4%–23%; Fig. 3A) were twice as frequent than those carrying –146 C > T *TERT* promoter mutation (6%; 95% CI: 2%–13%; Fig. 3B). Prevalence of




Fig. 2. Forest plot of prevalence of any *TERT* promoter mutations according to the site of HNSCC.

TERT promoter mutations was not statistically significant across different oral cavity subsites ($p = 0.64$), though *TERT* promoter mutations were more frequent in cancers of the buccal mucosa (62%; 95% CI: 21%-95%) and tongue (49%; 95% CI: 24%-74%; Fig. 4).

An analysis of *TERT* promoter mutations in oral cavity SCC was conducted differentiating the studies based on the geographic area of the included patients (eFigure in Supplementary Online Content). The meta-analysis indicated that *TERT* promoter mutations were more frequently reported in studies from Asia (59%, 95% CI 38%-77%), followed by North America (49%, 95% CI 1%-99%), Europe (40%, 95% CI 28%-52%), and South America (32%, 95% CI 21%-44%). A significant

heterogeneity was found among the articles ($I^2 = 95%$; $p < 0.01$).

Prognostic significance of TERT promoter mutations

TERT promoter mutation -124C > T was associated with a significant higher risk of death (sHR = 2.01, 95% CI 1.25-3.23 - Fig. 5A) and progression (sHR = 2.79, 95% CI 1.77-4.40 - Fig. 5B). Conversely, -146 C > T *TERT* promoter mutation did not show a significant association with outcomes.




Fig. 3. Forest plot of prevalence of mutation -124 C > T (A) and 146 C > T TERT promoter mutations (B) according to cancer site.

Discussion

From this systematic review and meta-analysis, it emerged that TERT promoter mutations were identified in 21% of HNSCCs. However, these mutually exclusive mutations, i.e. -124 C > T and -146 C > T, were by far more prevalent in SCCs of the oral cavity compared to the other head and neck sites. Nearly half of oral cavity carcinomas harboured, indeed, mutations in the TERT promoter. Conversely, only 1% and 12% of the SCCs arising from the oropharynx and larynx/hypopharynx harboured TERT mutations respectively. Confirming data collected by analysing different types of malignancies and human cell lines [44], -124 C > T mutation was the most commonly found. Despite the high heterogeneity across studies, all but one of the those analysing the prevalence of TERT mutations in the oral cavity consistently report rates ≥ 30%, while all but one of the studies investigating the prevalence of these mutations in laryngeal/hypopharyngeal cancers reported prevalence rates ≤ 15%. No heterogeneity was observed in prevalence data of TERT promoter mutations in oropharyngeal cancers with most of studies consistently reporting no mutations in these malignancies. Furthermore, although not statistically significant, we observed differences in prevalence according to the subsite of origin of the tumour within the oral cavity and according to the geographical area.

We have no mechanistic explanations for these observations. Reactivation of TERT through promoter mutations has been observed to occur more frequently in tumours originating from tissues with relative low rates of self-renewal [15,44]. Highly proliferative tissues retain, indeed, a pool of telomerase positive cells necessary for their long-term self-renewal capacity and ability to maintain tissue homeostasis [45]. It has been postulated that mutations of the TERT promoter in tumours originating from these tissues do not confer a direct proliferative benefit as telomeres are still long enough or telomerase is active [45]. However, the epithelium lining the oral cavity has a relatively high rate of self-renewal with this rate being not significantly different from that of other head and neck sites [46] and is provided by a compartment of stem cells expressing detectable levels of TERT mRNA [47].

Another hypothesis is that exposure to particular risk factors may account for sites and subsites topographic differences, as well as for geographical differences. A substantial and increasing proportion of oropharyngeal SCCs, around 60%-70% in Western populations [48,49], are known to be caused by transforming HR α-HPV infection, mainly HPV type 16 [50]. HR α-HPV-driven oropharyngeal SCCs express very

high TERT levels [47]. It has been demonstrated that HPV16 E6 oncoprotein physically and functionally interacts with telomerase complex and increases TERT catalytic activity, thus contributing to cell immortalization and transformation [51]. Consequently, in HPV-driven oropharyngeal SCCs, TERT reactivation mechanisms would be independent of the promoter mutations and there would be no selective pressure towards neoplastic clones possibly harbouring TERT promoter mutations. Most of cases of oropharyngeal SCC included in the present series were tested for HPV with 54% of them being HPV-positive. Thus, HPV-induced carcinogenesis in the oropharynx may only partially explain the very low rate of TERT promoter mutations found in these tumours.

While laryngeal, hypopharyngeal, and non-HPV-driven oropharyngeal SCCs are mostly attributable to exposure to tobacco smoke and alcohol, several other risk factors, including smokeless tobacco, betel quid chewing, areca nut, poor dentition, poor oral health, and trauma due to sharp or broken tooth were described for cancer of the oral cavity [1]. Interestingly, a study conducted in a case series from Taiwan identified betel nut chewing as the main risk factor for TERT promoter mutations [38]. Thus, specific risk factors could target the TERT promoter region and explain both the peculiar topographical restriction and the different geographic distribution of tumours harbouring TERT promoter mutations. An indirect support to this hypothesis also comes from the observation that, while skin melanomas frequently harbour mutations of the TERT promoter, uveal melanomas do not show such mutations [44]. As an alternative explanation, retaining a sufficient proliferative capability in response to chronic damage related to persistent exposure to particular risk factors could be challenged by the telomere-dependent proliferative barrier, despite the high rate of self-renewal of the oral cavity epithelium. In this context, TERT promoter mutations could provide an immediate and strong proliferative advantage on these cells [45].

With respect to oncological outcomes, an important finding emerging from this meta-analysis is that HNSCC patients with tumours harbouring -124 C > T TERT promoter mutation had a poor prognosis showing a more than doubled risk of death and progression than patients with tumours not harbouring this mutation. To our knowledge, this is the first meta-analysis that has attempted to summarize the scientific evidence on the association between the presence of a specific somatic mutation of the TERT gene promoter, i.e., -124 C > T, and the oncological clinical outcomes in HNSCC patients. The importance of our




Fig. 4. Forest plot of prevalence of any *TERT* promoter mutations in oral SCC according to oral cavity subsite.

finding emerged when the association analyses were carried out considering non-specified *TERT* promoter mutations. In this case, the presence of -124 C > T, or -146 C > T or other less frequent and functionally not characterized *TERT* promoter mutations, such as -124 C > A [28], had no predictive effect on OS and PFS, highlighting the relevance of evaluating the impact of -124 C > T *TERT* hotspot on clinical behaviour independently.


Through the creation *de novo* binding site for transcription factors, the presence of -124 C > T or -146 C > T *TERT* promoter mutations is considered a reliable indicator of sustained telomerase expression that drives cancer cell immortalization and progression and has been proposed as a potential biomarker for cancer prognosis associated with clinically aggressive behaviour [16]. However, the clinicopathological association of *TERT* promoter mutations is cancer-dependent, and

studies on different tumour types, including head and neck cancers, have reported contradicting clinical effects of *TERT* promoter mutations, ranging from poorer survival associated with the -124 C > T or -146 C > T *TERT* promoter mutation to unchanged clinical outcome [17,25-29,38]. Although both the -124 C > T and -146 C > T mutations create an identical 11-base sequence for binding the ETS transcription factor, promoting a similar increase of *TERT* transcription *in vitro* [52,53], previous reports demonstrated that these mutations are functionally distinct. The -146 C > T mutation, unlike -124 C > T, activated *TERT* transcription by binding the p52/ETS complex, thereby stimulating *TERT* expression via non-canonical NF- κ B signalling [54,55]. In addition, *in vivo*, the -124 C > T mutation was associated with higher *TERT* expression/telomerase activity compared to -146 C > T [53,56]. These functional differences might partially account for the

A. Overall survival


-124 C>T *TERT* promoter mutation

Study	HR	(95% CI)
Giunco et al, 2021 [17]	2.71	(1.30-5.65)
Kim et al, 2021 [37]	0.98	(0.23-4.14)
Arantes et al, 2020 [29]	2.35	(1.00-5.56)
Dogan et al, 2019 [43]	1.16	(0.37-3.66)
Total (Random effects)	2.01	(1.25-3.23)
Heterogeneity: $I^2=0\%$, $\tau^2=0.00$, $p=0.46$		




-146 C>T *TERT* promoter mutation

Study	HR	(95% CI)
Giunco et al, 2021 [17]	1.64	(0.53-5.08)
Arantes et al, 2020 [29]	0.89	(0.46-1.71)
Total (Random effects)	1.04	(0.59-1.83)
Heterogeneity: $I^2=0\%$, $\tau^2=0.00$, $p=0.36$		



Non specified *TERT* promoter mutation


Study	HR	(95% CI)
Arantes et al, 2020 [29]	1.24	(0.71-2.18)
Boscolo-Rizzo et al, 2020 [35]	0.53	(0.18-1.60)
Yilmaz et al, 2020 [28]	1.19	(0.81-1.76)
Dogan et al, 2019 [43]	1.48	(0.54-4.02)
Mundi et al, 2019 [39]	0.59	(0.30-1.17)
Qu et al, 2014 [26]	1.66	(1.13-2.44)
Total (Random effects)	1.13	(0.82-1.57)
Heterogeneity: $I^2=45\%$, $\tau^2=0.07$, $p=0.11$		



B. Progression-free survival


-124 C>T *TERT* promoter mutation

Study	HR	(95% CI)
Giunco et al, 2021 [17]	2.71	(1.42-5.17)
Kim et al, 2021 [37]	3.22	(1.20-8.64)
Arantes et al, 2020 [29]	3.37	(1.17-9.73)
Dogan et al, 2019 [43]	1.68	(0.41-6.97)
Total (Random effects)	2.79	(1.77-4.40)
Heterogeneity: $I^2=0\%$, $\tau^2=0.00$, $p=0.87$		



-146 C>T *TERT* promoter mutation

Study	HR	(95% CI)
Giunco et al, 2021 [17]	1.35	(0.50-3.65)
Arantes et al, 2020 [29]	1.00	(0.41-2.46)
Total (Random effects)	1.15	(0.59-2.23)
Heterogeneity: $I^2=0\%$, $\tau^2=0.00$, $p=0.67$		



Non specified *TERT* promoter mutation

Study	HR	(95% CI)
Arantes et al, 2020 [29]	1.56	(0.73-3.33)
Mundi et al, 2019 [39]	0.80	(0.40-1.58)
Dogan et al, 2019 [43]	1.68	(0.41-6.97)
Total (Random effects)	1.13	(0.70-1.83)
Heterogeneity: $I^2=0\%$, $\tau^2=0.00$, $p=0.37$		




Fig. 5. Forest plot visualizing the association of *TERT* promoter mutations and overall survival (A) and progression-free survival (B).

conflicting results obtained when specified or non-specified *TERT* promoter mutations were evaluated in clinicopathological associations. In particular, the higher *TERT* expression conferred by the -124 C > T *TERT* promoter mutation could explain its association with the more aggressive clinical phenotype that emerged from this meta-analysis, considering that consistent evidence stemming from a substantial number of studies reports worse clinical course and/or shorter survival of HNSCC patients with high tumour *TERT* mRNA expression and/or telomerase activity [5,6,35,47]. In accordance with this line of reasoning, it is not surprising that a recent meta-analysis evaluating the prognostic role of *TERT* upregulation alterations in patients with SCC of the oral cavity failed to find any association between non-specified *TERT* promoter mutations with OS or PFS while *TERT* protein overexpression resulted as a prognostic indicator of poor survival in these patients [57].

The biological bridge between high *TERT*/telomerase expression and the more aggressive tumour phenotype is still partially unidentified and seems not to be attributable only to *TERT*'s ability to maintain telomere length. Indeed, growing evidence indicates that *TERT* may promote tumorigenesis through telomere-length independent functions, including enhancement of proliferation, resistance to apoptosis, inflammation, invasion and metastasis [22,58] ultimately contributing to all the major characteristics of the cancer phenotype [8].

Finally, it must also be kept in mind that the effect of the *TERT* promoter mutations on *TERT* expression, and in turn on clinical prognosis, may be further complicated by the presence of the rs2853669 single nucleotide polymorphism (SNP). The minor C-variant allele disrupts a pre-existing ETS binding site at -245 bp in the *TERT* promoter region, thus counteracting the transactivation activity of the *TERT* hotspots [59,60]. Only one study among those included in our meta-analysis examined the clinical impact and the prognostic importance of the rs2853669 polymorphism [17]. The results of this study showed that patients' non-carrier of the SNP had an increased risk of disease progression and the coexistence of the T/T genotype of rs2853669 and the -124 C > T *TERT* promoter mutation increased the risk of adverse clinical outcome conferred by this mutation [17]. Greater attention to

this SNP might substantially improve the HNSCC patients' risk stratification allowing a greater personalization of care for these patients in terms of planning follow-up protocols and selecting patients more at risk of disease progression.

Conclusion

In conclusion, *TERT* promoter mutations are mainly topographically restricted to oral cavity SCC. The presence of -124 C > T mutation in the *TERT* gene promoter is associated with a worse prognosis of HNSCC patients. Therefore, this mutation appears as a promising biomarker to stratify the prognosis in patients with oral SCC for whom no other biomarkers capable of predicting outcome are currently available. However, these findings should be taken with caution as they are based on a still limited number of studies. We recommend that future research focuses on the different prognostic impact of each *TERT* promoter mutations to obtain a more precise risk stratification in HNSCC, thus helping clinicians to better predict patient outcome and consequently tailor treatment decisions.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.oraloncology.2023.106398>.

References

[1] Johnson DE, Burtneis B, Leemans CR, Lui VWY, Bauman JE, Grandis JR. Head and neck squamous cell carcinoma. Nat Rev Dis Primers 2020;6:92. <https://doi.org/10.1038/s41572-020-00224-3>.

- [2] Cancer today n.d. <http://gco.iarc.fr/today/home> (accessed May 29, 2022).
- [3] Mody MD, Rocco JW, Yom SS, Haddad RI, Saba NF. Head and neck cancer. *Lancet* 2021;398:2289–99. [https://doi.org/10.1016/S0140-6736\(21\)01550-6](https://doi.org/10.1016/S0140-6736(21)01550-6).
- [4] Tosi A, Parisatto B, Menegaldo A, Spinato G, Guido M, Del Mistro A, et al. The immune microenvironment of HPV-positive and HPV-negative oropharyngeal squamous cell carcinoma: a multiparametric quantitative and spatial analysis unveils a rationale to target treatment-naïve tumors with immune checkpoint inhibitors. *J Exp Clin Cancer Res* 2022;41(1). <https://doi.org/10.1186/s13046-022-02481-4>.
- [5] Boscolo-Rizzo P, Rampazzo E, Polesel J, Giunco S, Menegaldo A, Mantovani M, et al. Predictive and prognostic significance of telomerase levels/telomere length in tissues and peripheral blood in head and neck squamous cell carcinoma. *Sci Rep* 2019;9(1). <https://doi.org/10.1038/s41598-019-54028-x>.
- [6] Boscolo-Rizzo P, Da Mosto MC, Rampazzo E, Giunco S, Del Mistro A, Menegaldo A, et al. Telomeres and telomerase in head and neck squamous cell carcinoma: from pathogenesis to clinical implications. *Cancer Metastasis Rev* 2016;35(3):457–74.
- [7] Huffman KE, Levene SD, Tesmer VM, Shay JW, Wright WE. Telomere Shortening Is Proportional to the Size of the G-rich Telomeric 3'-Overhang *. *J Biol Chem* 2000;275:19719–22. <https://doi.org/10.1074/jbc.M002843200>.
- [8] Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. *Cell* 2011;144:646–74. <https://doi.org/10.1016/j.cell.2011.02.013>.
- [9] Akincilar SC, Unal B, Tergaonkar V. Reactivation of telomerase in cancer. *Cell Mol Life Sci* 2016;73:1659–70. <https://doi.org/10.1007/s00018-016-2146-9>.
- [10] Leão R, Apolónio JD, Lee D, Figueiredo A, Tabori U, Castelo-Branco P. Mechanisms of human telomerase reverse transcriptase (hTERT) regulation: clinical impacts in cancer. *J Biomed Sci* 2018;25:22. <https://doi.org/10.1186/s12929-018-0422-8>.
- [11] Horn S, Figl A, Rachakonda PS, Fischer C, Sucker A, Gast A, et al. TERT promoter mutations in familial and sporadic melanoma. *Science* 2013;339(6122):959–61.
- [12] Huang FW, Hodis E, Xu MJ, Kryukov GV, Chin L, Garraway LA. Highly Recurrent TERT Promoter Mutations in Human Melanoma. *Science* 2013;339:957–9. <https://doi.org/10.1126/science.1229259>.
- [13] Annunziata C, Pezzuto F, Greggi S, Ionna F, Losito S, Botti G, et al. Distinct profiles of TERT promoter mutations and telomerase expression in head and neck cancer and cervical carcinoma. *Int J Cancer* 2018;143(5):1153–61.
- [14] Yu Y, Fan D, Song X, Zakeri K, Chen L, Kang J, et al. TERT Promoter Mutations Are Enriched in Oral Cavity Cancers and Associated With Locoregional Recurrence. *JCO Precis Oncol* 2021;5(5):1259–69.
- [15] Killela PJ, Reitman ZJ, Jiao Y, Bettegowda C, Agrawal N, Diaz LA, et al. TERT promoter mutations occur frequently in gliomas and a subset of tumors derived from cells with low rates of self-renewal. *Proc Natl Acad Sci U S A* 2013;110(15):6021–6.
- [16] Hafezi F, Jaxel L, Lemaire M, Turner JD, Perez-Bercoff D. TERT Promoter Mutations Increase Sense and Antisense Transcription from the TERT Promoter. *Biomedicines* 2021;9:1773. <https://doi.org/10.3390/biomedicines9121773>.
- [17] Giunco S, Boscolo-Rizzo P, Rampazzo E, Tirelli G, Alessandrini L, Di Carlo R, et al. TERT Promoter Mutations and rs2853669 Polymorphism: Useful Markers for Clinical Outcome Stratification of Patients With Oral Cavity Squamous Cell Carcinoma. *Front Oncol* 2021;11:782658. <https://doi.org/10.3389/fonc.2021.782658>.
- [18] Zhang X, Chen Y, Yan D, Han J, Zhu L. TERT Gene rs2736100 and rs2736098 Polymorphisms are Associated with Increased Cancer Risk: A Meta-Analysis. *Biochem Genet* 2022;60:241–66. <https://doi.org/10.1007/s10528-021-10097-0>.
- [19] Liu Z, Ma H, Wei S, Li G, Sturgis EM, Wei Q. Telomere length and TERT functional polymorphisms are not associated with risk of squamous cell carcinoma of the head and neck. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol* 2011;20:2642–5. <https://doi.org/10.1158/1055-9965.EPI-11-0890>.
- [20] Yu J, Li X, Zhou B, Yan A. Polymorphisms of the TERT-CLPTM1L Gene Are Associated with Pharynx-Larynx Cancer. *DNA Cell Biol* 2019;38:915–21. <https://doi.org/10.1089/dna.2019.4744>.
- [21] Li Y, Tergaonkar V. Noncanonical functions of telomerase: implications in telomerase-targeted cancer therapies. *Cancer Res* 2014;74:1639–44. <https://doi.org/10.1158/0008-5472.CAN-13-3568>.
- [22] Ségal-Bendirdjian E, Geli V. Non-canonical Roles of Telomerase: Unraveling the Imbroglío. *Front Cell Dev Biol* 2019;7:332. <https://doi.org/10.3389/fcell.2019.00332>.
- [23] Wan S, Liu X, Hua W, Xi M, Zhou Y, Wan Y. The role of telomerase reverse transcriptase (TERT) promoter mutations in prognosis in bladder cancer. *Bioengineered* 2021;12:1495–504. <https://doi.org/10.1080/21655979.2021.1915725>.
- [24] Mao J, Huang X, Okla MK, Abdel-Maksoud MA, Mubarak A, Hameed Z, et al. Risk Factors for TERT Promoter Mutations with Papillary Thyroid Carcinoma Patients: A Meta-Analysis and Systematic Review. *Comput Math Methods Med* 2022;2022:1–11.
- [25] Del Bianco P, Stagni C, Giunco S, Fabozzi A, Elefanti L, Pellegrini S, et al. TERT Promoter Mutations Differently Correlate with the Clinical Outcome of MAPK Inhibitor-Treated Melanoma Patients. *Cancers* 2020;12(4):946.
- [26] Qu Y, Dang S, Wu K, Shao Y, Yang Qi, Ji M, et al. TERT promoter mutations predict worse survival in laryngeal cancer patients. *Int J Cancer* 2014;135(4):1008–10.
- [27] Vinothkumar V, Arunkumar G, Revathidevi S, Arun K, Manikandan M, Rao AKDM, et al. TERT promoter hot spot mutations are frequent in Indian cervical and oral squamous cell carcinomas. *Tumor Biol* 2016;37(6):7907–13.
- [28] Yilmaz I, Erkul BE, Ozturk Sari S, Issin G, Tural E, Terzi Kaya Terzi N, et al. Promoter region mutations of the telomerase reverse transcriptase (TERT) gene in head and neck squamous cell carcinoma. *Oral Surg Oral Med Oral Pathol. Oral Radiol* 2020;130(1):63–70.
- [29] Arantes LMRB, Cruvinel-Carlioni A, de Carvalho AC, Sorroche BP, Carvalho AL, Scapulatempo-Neto C, et al. TERT Promoter Mutation C228T Increases Risk for Tumor Recurrence and Death in Head and Neck Cancer Patients. *Front Oncol* 2020;10:1275. <https://doi.org/10.3389/fonc.2020.101275>.
- [30] McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol* 2016;75:40–6. <https://doi.org/10.1016/j.jclinepi.2016.01.021>.
- [31] Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4(1). <https://doi.org/10.1186/2046-4053-4-1>.
- [32] Wells G, Shea B, O'Connell D, Peterson je, Welch V, Losos M, et al. The Newcastle–Ottawa Scale (NOS) for Assessing the Quality of Non-Randomized Studies in Meta-Analysis. 2000.
- [33] Normand S-LT. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat Med* 1999;18:321–59. [https://doi.org/10.1002/\(SICI\)1097-0258\(19990215\)18:3<321::AID-SIM28>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1097-0258(19990215)18:3<321::AID-SIM28>3.0.CO;2-P).
- [34] Funnel plots for detecting bias in meta-analysis - Journal of Clinical Epidemiology n.d. [https://www.jclinepi.com/article/S0895-4356\(01\)00377-8/fulltext](https://www.jclinepi.com/article/S0895-4356(01)00377-8/fulltext) (accessed July 21, 2022).
- [35] Boscolo-Rizzo P, Giunco S, Rampazzo E, Brutti M, Spinato G, Menegaldo A, et al. TERT promoter hotspot mutations and their relationship with TERT levels and telomere erosion in patients with head and neck squamous cell carcinoma. *J Cancer Res Clin Oncol* 2020;146(2):381–9.
- [36] Moreira A, Poulet A, Masliyah-Planchon J, Lecerf C, Vacher S, Larbi Chérif L, et al. Prognostic value of tumor mutational burden in patients with oral cavity squamous cell carcinoma treated with upfront surgery. *ESMO Open* 2021;6(4):100178.
- [37] Kim H, Kwon MJ, Park B, Choi HG, Nam ES, Cho SJ, et al. Negative Prognostic Implication of TERT Promoter Mutations in Human Papillomavirus-Negative Tonsillar Squamous Cell Carcinoma Under the New 8th AJCC Staging System. *Indian J Surg Oncol* 2021;12(S1):134–43.
- [38] Chang K-P, Wang C-I, Pickering CR, Huang Y, Tsai C-N, Tsang N-M, et al. Prevalence of promoter mutations in the TERT gene in oral cavity squamous cell carcinoma: Promoter mutations in TERT gene in oral cavity SCC. *Head Neck* 2017;39(6):1131–7.
- [39] Mundi N, Prokopec SD, Ghasemi F, Warner A, Patel K, MacNeil D, et al. Genomic and human papillomavirus profiling of an oral cancer cohort identifies TP53 as a predictor of overall survival. *Cancers Head Neck* 2019;4(1). <https://doi.org/10.1186/s41199-019-0045-0>.
- [40] Morris LGT, Chandramohan R, West L, Zehir A, Chakravarty D, Pfister DG, et al. The Molecular Landscape of Recurrent and Metastatic Head and Neck Cancers: Insights From a Precision Oncology Sequencing Platform. *JAMA Oncol* 2017;3(2):244.
- [41] Cheng KA, Kurtis B, Babayeva S, Zhuge J, Tantchou I, Cai D, et al. Heterogeneity of TERT promoter mutations status in squamous cell carcinomas of different anatomical sites. *Ann Diagn Pathol* 2015;19(3):146–8.
- [42] Schwaederle M, Krishnamurthy N, Daniels GA, Piccioni DE, Kesari S, Fanta PT, et al. Telomerase reverse transcriptase promoter alterations across cancer types as detected by next-generation sequencing: A clinical and molecular analysis of 423 patients: TERT Promoter Alterations in Cancers. *Cancer* 2018;124(6):1288–96.
- [43] Dogan S, Xu B, Middha S, Vanderbilt CM, Bowman AS, Migliacci J, et al. Identification of prognostic molecular biomarkers in 157 HPV-positive and HPV-negative squamous cell carcinomas of the oropharynx. *Int J Cancer* 2019;145(11):3152–62.
- [44] Vinagre J, Almeida A, Pópulo H, Batista R, Lyra J, Pinto V, et al. Frequency of TERT promoter mutations in human cancers. *Nat Commun* 2013;4(1). <https://doi.org/10.1038/ncomms3185>.
- [45] Chiba K, Johnson JZ, Vogan JM, Wagner T, Boyle JM, Hockemeyer D. Cancer-associated TERT promoter mutations abrogate telomerase silencing. *Elife* 2015;4:e07918.
- [46] Tomasetti C, Li L, Vogelstein B. Stem cell divisions, somatic mutations, cancer etiology, and cancer prevention. *Science* 2017;355:1330–4. <https://doi.org/10.1126/science.aaf9011>.
- [47] Boscolo-Rizzo P, Rampazzo E, Perissinotto E, Piano MA, Giunco S, Baboci L, et al. Telomere shortening in mucosa surrounding the tumor: biosensor of field cancerization and prognostic marker of mucosal failure in head and neck squamous cell carcinoma. *Oral Oncol* 2015;51(5):500–7.
- [48] Faraji F, Rettig EM, Tsai H-L, El Asmar M, Fung N, Eisele DW, et al. The prevalence of human papillomavirus in oropharyngeal cancer is increasing regardless of sex or race, and the influence of sex and race on survival is modified by human papillomavirus tumor status. *Cancer* 2019;125(5):761–9.
- [49] Zamani M, Grønhoj C, Jensen DH, Carlander AF, Agander T, Kiss K, et al. The current epidemic of HPV-associated oropharyngeal cancer: An 18-year Danish population-based study with 2,169 patients. *Eur J Cancer Oxf Engl* 2020;134:52–9.
- [50] Lechner M, Liu J, Masterson L, Fenton TR. HPV-associated oropharyngeal cancer: epidemiology, molecular biology and clinical management. *Nat Rev Clin Oncol* 2022;19:306–27. <https://doi.org/10.1038/s41571-022-00603-7>.
- [51] Liu X, Dakic A, Zhang Y, Dai Y, Chen R, Schlegel R. HPV E6 protein interacts physically and functionally with the cellular telomerase complex. *Proc Natl Acad Sci* 2009;106:18780–5. <https://doi.org/10.1073/pnas.0906357106>.
- [52] Bell RJA, Rube HT, Kreig A, Mancini A, Fouse SD, Nagarajan RP, et al. The transcription factor GABP selectively binds and activates the mutant TERT promoter in cancer. *Science* 2015;348(6238):1036–9.
- [53] Huang D-S, Wang Z, He X-J, Diplas BH, Yang R, Killela PJ, et al. Recurrent TERT promoter mutations identified in a large-scale study of multiple tumour types are associated with increased TERT expression and telomerase activation. *Eur J Cancer Oxf Engl* 2015;51(8):969–76.

- [54] Li Y, Zhou Q-L, Sun W, Chandrasekharan P, Cheng HS, Ying Z, et al. Non-canonical NF- κ B signalling and ETS1/2 cooperatively drive C250T mutant TERT promoter activation. *Nat Cell Biol* 2015;17:1327–38. <https://doi.org/10.1038/ncb3240>.
- [55] Xu X, Li Y, Bharath SR, Ozturk MB, Bowler MW, Loo BZL, et al. Structural basis for reactivating the mutant TERT promoter by cooperative binding of p52 and ETS1. *Nat Commun* 2018;9(1). <https://doi.org/10.1038/s41467-018-05644-0>.
- [56] Heidenreich B, Rachakonda PS, Hosen I, Volz F, Hemminki K, Weyerbrock A, et al. TERT promoter mutations and telomere length in adult malignant gliomas and recurrences. *Oncotarget* 2015;6:10617–33. <https://doi.org/10.18632/oncotarget.3329>.
- [57] González-Moles MÁ, Moya-González E, García-Ferrera A, Nieto-Casado P, Ramos-García P. Prognostic and Clinicopathological Significance of Telomerase Reverse Transcriptase Upregulation in Oral Cancer: A Systematic Review and Meta-Analysis. *Cancers* 2022;14:3673. <https://doi.org/10.3390/cancers14153673>.
- [58] Martínez P, Blasco MA. Telomeric and extra-telomeric roles for telomerase and the telomere-binding proteins. *Nat Rev Cancer* 2011;11:161–76. <https://doi.org/10.1038/nrc3025>.
- [59] Rachakonda PS, Hosen I, de Verdier PJ, Fallah M, Heidenreich B, Ryk C, et al. TERT promoter mutations in bladder cancer affect patient survival and disease recurrence through modification by a common polymorphism. *Proc Natl Acad Sci U S A* 2013;110(43):17426–31.
- [60] Hsu C-P, Hsu N-Y, Lee L-W, Ko J-L. Ets2 binding site single nucleotide polymorphism at the hTERT gene promoter—effect on telomerase expression and telomere length maintenance in non-small cell lung cancer. *Eur J Cancer Oxf Engl* 1990;2006(42):1466–74. <https://doi.org/10.1016/j.ejca.2006.02.014>.