

16. Maggi FA, Braga-Neto P, Chien HF, Gama MTD, Rezende Filho FM, Saraiva-Pereira ML, et al. Cross-cultural adaptation and validation of the international cooperative ataxia rating scale (ICARS) to Brazilian Portuguese. *Arq Neuropsiquiatr* 2018;76(10):228–230.
17. Early treatment diabetic retinopathy study design and baseline patient characteristics: ETDRS report number 7. *Ophthalmology* 1991;98(5):741–756.
18. Aytulun A, Cruz-Herranz A, Aktas O, Balcer LJ, Balk L, Barboni P, et al. APOSTEL 2.0 recommendations for reporting quantitative optical coherence tomography studies. *Neurology* 2021;97(2):68–79.
19. Tewarie P, Balk L, Costello F, Green A, Martin R, Schippling S, et al. The OSCAR-IB consensus criteria for retinal OCT quality assessment. *PLoS One* 2012;7(4):e34823.
20. la Morgia C, Barboni P, Rizzo G, Carbonelli M, Savini G, Scaglione C, et al. Loss of temporal retinal nerve fibers in Parkinson disease: a mitochondrial pattern? *Eur J Neurol* 2013;20(1):198–201.
21. Cornelius N, Wardman JH, Hargreaves IP, Neerghen V, Bie AS, Tümer Z, et al. Evidence of oxidative stress and mitochondrial dysfunction in spinocerebellar ataxia type 2 (SCA2) patient fibroblasts: effect of coenzyme Q10 supplementation on these parameters. *Mitochondrion* 2017;34:103–114.
22. Huang L, Zhang D, Ji J, Wang Y, Zhang R. Central retina changes in Parkinson's disease: a systematic review and meta-analysis. *J Neurol* 2021;268:4646–4654.
23. Chan VTT, Sun Z, Tang S, Chen LJ, Wong A, Tham CC, et al. Spectral-domain OCT measurements in Alzheimer's disease: a systematic review and meta-analysis. *Ophthalmology* 2019;126:497–510.
24. Marianelli BF, Filho FMR, Salles MV, de Andrade JBC, Pedrosa JL, Sallum JMF, et al. A proposal for classification of retinal degeneration in spinocerebellar ataxia type 7. *Cerebellum* 2020;20(3):384–391.
25. Oertel FC, Zeitz O, Rönnefarth M, Bereuter C, Motamedi S, Zimmermann HG, et al. Functionally relevant maculopathy and optic atrophy in spinocerebellar ataxia type 1. *Mov Disord Clin Pract* 2020;7(5):502–508.
26. Isashiki Y, Kii Y, Ohba N, Nakagawa M. Retinopathy associated with Machado-Joseph disease (spinocerebellar ataxia 3) with CAG trinucleotide repeat expansion. *Am J Ophthalmol* 2001;131(6):808–810.
27. Garcia-Martin E, Larrosa JM, Polo V, Satue M, Marques ML, Alarcia R, et al. Distribution of retinal layer atrophy in patients with parkinson disease and association with disease severity and duration. *Am J Ophthalmol* 2014;157(2):470–478.e2. Available from: <https://doi.org/10.1016/j.ajo.2013.09.028>
28. Ortuño-Lizarán I, Beach TG, Serrano GE, Walker DG, Adler CH, Cuenca N. Phosphorylated α -synuclein in the retina is a biomarker of Parkinson's disease pathology severity. *Mov Disord* 2018;33(8):1315–1324.
29. Koronyo Y, Biggs D, Barron E, Boyer DS, Pearlman JA, Au WJ, et al. Retinal amyloid pathology and proof-of-concept imaging trial in Alzheimer's disease. *JCI. Insight* 2017;2(16):e93621.
30. Huynh DP, Del Bigio MR, Ho DH, Pulst SM. Expression of Ataxin-2 in Brains from Normal Individuals and Patients with Alzheimer's Disease and Spinocerebellar Ataxia 2; 1999.
31. Sundberg CA, Lakk M, Paul S, P. Figueroa K, Scoles DR, Pulst SM, et al. The RNA-binding protein and stress granule component ATAXIN-2 is expressed in mouse and human tissues associated with glaucoma pathogenesis. *J Comp Neurol* 2022;530(2):537–552.
32. Qiu K, Wang G, Zhang R, Lu X, Zhang M, Jansonius NM. Influence of optic disc-fovea distance on macular thickness measurements with OCT in healthy myopic eyes. *Sci Rep* 2018;8(1):5233.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.

Serum Neurofilament Light Chain in Replication Factor Complex Subunit 1 CANVAS and Disease Spectrum

Ilaria Quartesan, MD,^{1,2} Elisa Vegezzi, MD,^{1,2} Riccardo Currò, MD,^{1,3} Amanda Heslegrave, BSc, PhD,^{4,5} Chiara Pisciotta, MD, PhD,⁶ Pablo Iruzubieta, MD,⁷ Alessandro Salvalaggio, MD, PhD,⁸ Gorka Fernández-Eulate, MD,⁹ Natalia Dominik, MSc,³ Bianca Rugginini, MD,¹ Arianna Manini, MD,^{3,10} Elena Abati, MD,^{3,11} Stefano Facchini, PhD,² Katarina Manso, MD,¹² Ines Albajar, MD,⁷ Rhiannon Laban, MSc,⁵ Alexander M. Rossor, MRCP, PhD,³ Anna Pichiechio, MD, PhD,^{1,2} Giuseppe Cosentino, MD, PhD,^{1,2} Paola Saveri, BSc,⁶ Ettore Salsano, MD,⁶ Francesca Andreetta, PhD,⁶ Enza M. Valente, MD, PhD,^{1,2} Henrik Zetterberg, MD, PhD,^{4,5,12,13,14,15} Paola Giunti, MD, PhD,¹⁶ Tanya Stojkovic, MD,⁹ Chiara Briani, MD,⁸ Adolfo López de Munain, MD, PhD,⁷ Henry Pareyson, MD,⁶ Mary M. Reilly, MD, FRCP, FRCPI,³ Henry Houlden, MD, PhD, MRCP,³ Cristina Tassorelli, MD, PhD,^{1,2} and Andrea Cortese, MD, PhD^{1,3*}

¹Department of Brain and Behavioral Sciences, University of Pavia, Pavia, Italy ²IRCCS Mondino Foundation, Pavia, Italy ³Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, London, United Kingdom ⁴Department of Neurodegenerative Disease, UCL Queen Square Institute of Neurology, London, United Kingdom ⁵UK Dementia Research Institute at UCL, London, United Kingdom ⁶Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy ⁷Neurology Department, Donostia University Hospital, Osakidetza, and Biodonostia Health Research Institute-UPV-EHU, San Sebastián, Spain ⁸Department of Neuroscience, University of Padova, Padova, Italy ⁹Neuro-myology Department, Institut de Myologie, Pitié-Salpêtrière Hospital, APHP, Sorbonne University, Paris, France ¹⁰Department of Neurology and Laboratory of Neuroscience,

This is an open access article under the terms of the [Creative Commons Attribution](#) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

*Correspondence to: Dr. Andrea Cortese, Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, WC1N 3BG London, UK; E-mail: andrea.cortese@ucl.ac.uk; andrea.cortese@unipv.it

Relevant conflicts of interest/financial disclosures: All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or nonfinancial interest in the subject matter or materials discussed in this manuscript.

Funding agencies: This study was supported by the Medical Research Council (MR/T001712/1), Fondazione Cariplo (grant no. 2019-1836), the Inherited Neuropathy Consortium, Italian Ministry of Health (Ricerca Corrente 2021–2022), Italian Ministry for Universities and Research (MUR, 20229MMHXP).

Received: 11 August 2023; **Revised:** 7 November 2023; **Accepted:** 15 November 2023

Published online 6 December 2023 in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/mds.29680

IRCCS Istituto Auxologico Italiano, Milan, Italy ¹¹Department of Pathophysiology and Transplantation (DEPT), University of Milan, Milan, Italy ¹²Clinical Neurochemistry Laboratory, Sahlgrenska University Hospital, Mölndal, Sweden ¹³Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The Sahlgrenska Academy at the University of Gothenburg, Mölndal, Sweden ¹⁴Hong Kong Center for Neurodegenerative Diseases, Clear Water Bay, Hong Kong, China ¹⁵Wisconsin Alzheimer's Disease Research Center, University of Wisconsin School of Medicine and Public Health, University of Wisconsin-Madison, Madison, Wisconsin, USA ¹⁶Department of Clinical and Movement Neuroscience, UCL Queen Square Institute of Neurology, London, United Kingdom

ABSTRACT: Background: Biallelic intronic AAGGG repeat expansions in the replication factor complex subunit 1 (*RFC1*) gene were identified as the leading cause of cerebellar ataxia, neuropathy, vestibular areflexia syndrome. Patients exhibit significant clinical heterogeneity and variable disease course, but no potential biomarker has been identified to date.

Objectives: In this multicenter cross-sectional study, we aimed to evaluate neurofilament light (NfL) chain serum levels in a cohort of *RFC1* disease patients and to correlate NfL serum concentrations with clinical phenotype and disease severity.

Methods: Sixty-one patients with genetically confirmed *RFC1* disease and 48 healthy controls (HCs) were enrolled from six neurological centers. Serum NfL concentration was measured using the single molecule array assay technique.

Results: Serum NfL concentration was significantly higher in patients with *RFC1* disease compared to age- and sex-matched HCs ($P < 0.0001$). NfL level showed a moderate correlation with age in both HCs ($r = 0.4353$, $P = 0.0020$) and patients ($r = 0.4092$, $P = 0.0011$). Mean NfL concentration appeared to be significantly higher in patients with cerebellar involvement compared to patients without cerebellar dysfunction (27.88 vs. 21.84 pg/mL, $P = 0.0081$). The association between cerebellar involvement and NfL remained significant after controlling for age and sex ($\beta = 0.260$, $P = 0.034$).

Conclusions: Serum NfL levels are significantly higher in patients with *RFC1* disease compared to HCs and correlate with cerebellar involvement. Longitudinal studies are warranted to assess its change over time. © 2023 The Authors. *Movement Disorders* published by Wiley Periodicals LLC on behalf of International Parkinson and Movement Disorder Society.

Key Words: ataxia; biomarkers; cerebellar ataxia neuropathy vestibular areflexia syndrome; neurofilament light chain; replication factor complex subunit 1

Biallelic AAGGG repeat expansions in intron 2 of the gene encoding replication factor complex subunit 1 (*RFC1*) were identified as the cause of cerebellar ataxia with neuropathy and vestibular areflexia

syndrome (CANVAS) and disease spectrum (here shortened as *RFC1* disease).¹

Affected individuals exhibit significant clinical heterogeneity, starting with an isolated sensory neuropathy, with or without chronic cough, and progressing to a more complex ataxia with cerebellar dysfunction in later disease stages.^{2,3} Bilateral vestibular areflexia is also often present but can be easily overlooked if not tested for. The clinical manifestations of *RFC1* disease have expanded beyond the classical CANVAS to encompass dysautonomia, parkinsonism, and cognitive impairment.⁴⁻⁶

Multiple studies have established that neurofilament light (NfL), a component of the axonal cytoskeleton released into the cerebrospinal fluid (CSF) and blood stream after neuronal damage,⁷ is a promising fluid biomarker for neurodegenerative conditions, including Charcot–Marie–Tooth disease, hereditary transthyretin (ATTR) amyloidosis, and hereditary ataxias.⁸⁻¹³ Increased CSF or serum NfL levels in patients carrying biallelic *RFC1* expansions have been observed in two single cases.^{14,15} However, there is still scant evidence of its diagnostic role or its ability to reflect progression of the condition over time.

In this multicenter cross-sectional study, we aimed to assess whether serum NfL may represent a potential biomarker for *RFC1* disease using ultrasensitive single molecule array (Simoa) immunoassay technology. Furthermore, we sought to evaluate the correlation between NfL concentration and clinical phenotype and severity.

Patients and Methods

Serum samples ($n = 50$) were prospectively collected, after informed consent, from patients with a genetically confirmed diagnosis of *RFC1* disease¹ and consecutively attending neurology clinics in six neurological centers (London: 10; Pavia: 16; Milan: 1; Paris: 7; Padova: 7; San Sebastian: 9). In 11 patients, serum was retrieved from the biorepository (Milan).

Patients' neurological history and clinical signs, including the presence of sensory neuropathy, cerebellar dysfunction (defined by the presence of one or more of the following signs: broken pursuits, dysmetric saccades, gaze-evoked or downbeat nystagmus, dysarthria, and dysphagia), vestibular areflexia, dysautonomia, cognitive impairment, and parkinsonism, were collected based on a standard template. Patients with other neurologic diseases were excluded. In prospectively collected data, neurological examinations were performed at serum collection. In the 11 patients whose serum was retrieved from biorepositories, examination was performed at a mean of 15 months from serum sampling

(ranging from 24 months before sampling to 12 months after sampling).

Based on the known progression of the disease from an isolated sensory neuropathy to a complex neuropathy with cerebellar dysfunction or full CANVAS,³ patients were divided into two clinical subtypes as a proxy of disease severity: (1) *RFC1* disease without cerebellar involvement, which includes patients with isolated sensory neuropathy/neuropathy with bilateral vestibular areflexia, and (2) *RFC1* disease with cerebellar involvement, which includes patients with complex sensory and cerebellar ataxia or full CANVAS. Therefore, neuronal damage occurring within the cerebellum has been previously shown to be associated with higher elevation of NfL serum levels compared to neuropathies.^{8–16} Loss of independent ambulation was also considered as a marker of advanced disease.

Serum samples from age-matched healthy controls (HC) were also collected from each participating center (London: 1 sample; Pavia: 13; Milan: 14; Paris: 4; Padova: 8; San Sebastian: 8). The assessment of HCs was based on medical history (through participants' interviews).

Blood sampling and storage were conducted following a standard operating procedure at each of the six different centers. In particular, blood was collected into serum separating tubes and centrifuged at 20°C at 3500 rpm for 10 min. Serum was then aliquoted and stored at –20°C. Samples were anonymized and sent blinded for clinical details to the University College London (A.H., H.Z.) for analysis of NfL levels. Serum NfL concentration was measured using the Simoa NfL assay on an HD-X analyzer (Quanterix, Billerica, MA, USA) in one round of experiments with one batch of reagents. Four quality control samples were run in duplicate; the mean intra-assay coefficient of variation of duplicate determinations for concentration was 6.9%.

Statistical analysis was performed using SPSS, version 26.00 (IBM, Armonk, NY, USA) and GraphPad Prism 9.0.0 for Windows (GraphPad Software, San Diego, CA, USA).

Demographic and clinical data were described as mean (SD) or median (interquartile range) if normally or nonnormally distributed, respectively. Data normality was assessed using Q–Q plots and analytical tests (Kolmogorov–Smirnov test, Shapiro–Wilk test, and Anderson–Darling test), requiring consistent results from all tests to confirm normality.

Means between groups were compared using the *t* test for normally distributed data and the Mann–Whitney *U* test for nonnormally distributed data. Correlations were assessed using Spearman's or Pearson's coefficients, as appropriate for data distribution. We conducted a multiple linear regression analysis to

examine the relationship between serum NfL (dependent variable), cerebellar involvement, age at time of blood collection, and sex (independent variables).

The study protocol was approved by the local institutional review boards and ethics committees. Written informed consent was obtained from all patients and HCs. The study adhered to all applicable ethical regulations.

Results

A total of 61 patients and 48 HCs were enrolled in the study. The demographics and clinical characteristics of participants are summarized in Table 1. There was no significant difference in the mean age at sample collection of the two groups ($P = 0.13$) or sex distribution ($P = 0.823$). Mean age at blood collection in patients with *RFC1* disease was 67.08 years (± 10.70), with a mean age of onset of 55.16 years (± 10.85) and a median disease duration of 11 years (6–15). Overall, 42 patients (69%) had signs and/or symptoms of cerebellar involvement. At the time of evaluation, 27 patients (44%) required walking aids. In 7 patients (11%) vestibular function was not assessed.

No significant difference in serum NfL levels was observed between stored samples and samples collected prospectively (23.18 vs. 26.62 pg/mL, $P = 0.22$).

Serum NfL concentration was significantly higher in patients with *RFC1* disease (24.34 pg/mL [IQR: 19.44–31.60]) compared to age-matched HCs (12.40 pg/mL ± 4.64) ($P < 0.0001$; Fig. 1A).

Receiver operating characteristic analysis showed that serum NfL levels could discriminate patients from controls with great accuracy (AUC of 0.9262, 95% CI [confidence interval]: 0.88–0.97) (Fig. 1B). A concentration of 15.86 pg/mL can effectively identify individuals with *RFC1* disease with a sensitivity of 92% and a specificity of 81%.

Serum NfL concentration showed a moderate correlation with age in both HCs ($r = 0.4353$, $P = 0.0020$) and patients ($r = 0.4092$, $P = 0.0011$) (Fig. 1C).

The significant difference in serum NfL concentration compared to controls was maintained for individual comparisons of controls versus patients without cerebellar involvement ($P < 0.0001$), and of controls versus patients with cerebellar involvement ($P < 0.0001$).

Also, mean NfL concentrations were significantly higher in patients with cerebellar involvement compared to patients without cerebellar dysfunction (27.88 vs. 21.84 pg/mL, $P = 0.0081$) (Fig. 1D), and the association remained significant after controlling for age and sex in a multiple linear regression model ($\beta = 0.260$, 95% CI: 0.411002, $P = 0.034$).

Conversely, there was no significant difference in NfL levels between patients without signs of vestibular

TABLE 1 Demographic details of patients and healthy control cohorts

Patient group	N	Age, y (min–max)	Sex, F (%)	NfL (pg/mL)
Controls	48	63.68 (37–79)	21 (44)	12.40 ± 4.64
<i>RFC1</i> disease	61	67.08 (38–86)	28 (46)	24.34 ± 9.24
<i>P</i> -value		0.13	0.82	<0.0001
Without cerebellar involvement	19	64.11 (46–86)	6 (31)	21.84 ± 6.99
Isolated sensory neuropathy	11	66.91 (47–86)	4 (36)	24.48 ± 6.07
Sensory neuropathy with BVA	8	60.25 (46–77)	2 (25)	18.22 ± 6.84
With cerebellar involvement	42	68.42 (38–83)	22 (52)	27.88 ± 9.58
Complex sensory and cerebellar ataxia	9	65 (48–81)	6 (67)	26.69 ± 8.60
CANVAS	33	69.36 (38–83)	16 (48)	28.20 ± 9.94
<i>P</i> -value		0.14	0.14	0.0081

Abbreviations: NfL, neurofilament light; BVA, bilateral vestibular areflexia; CANVAS, cerebellar ataxia, neuropathy, vestibular areflexia syndrome.




FIG. 1. (A) Increased serum NfL concentration in patients with *RFC1* disease compared to healthy controls (HC). (B) Receiver operator curve of serum NfL concentration for detecting patients with *RFC1* CANVAS and disease spectrum. (C) Correlation between serum NfL concentration and age in both patients carrying *RFC1* expansions and HCs. (D) Increased serum NfL concentration in patients with cerebellar involvement compared to patients without cerebellar dysfunction. Both subgroups show higher concentrations compared to HCs. Line is at mean. Error bars: standard deviation. *P*-value less than 0.05 is flagged with one star (*), *P*-value less than 0.01 is flagged with two stars (**), and *P*-value less than 0.001 is flagged with three stars (***). NfL: neurofilament light. CANVAS: cerebellar ataxia, neuropathy, vestibular areflexia syndrome. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

dysfunction ($n = 13$) and patients with vestibular impairment ($n = 41$) (25.28 vs. 26.25 pg/mL, $P = 0.7061$). Serum NfL levels did not appear to correlate with disease duration ($r = 0.014$, $P = 0.917$) and did not differ between patients with independent walking ($n = 42$) and patients using walking aid ($n = 27$) (24.12 vs. 28.36 pg/mL, $P = 0.0820$).

One patient had clinically manifest parkinsonism, with an NfL value (31.78 pg/mL) above the 75th percentile, and 3 patients presented with clinical signs of dysautonomia (mean NfL: 28.82 ± 6.26 pg/mL). Cognitive impairment was not reported in any patients.

Discussion

This study is the first to investigate serum NfL levels as a biomarker in *RFC1* disease using ultrasensitive Simoa technology. We found significantly higher serum NfL levels in patients with biallelic *RFC1* expansions compared to HCs of the same age and sex. Elevated NfL levels were observed in various clinical phenotypes, including isolated sensory neuropathy, which is more common in early disease stages. NfL levels demonstrated excellent discriminatory power, supporting its potential as a reliable biomarker in various neurodegenerative ataxias,^{12,13,17-19} as well as in hereditary neuropathies.^{8,10}

NfL is an axonal cytoskeletal protein released after axonal damage. Abnormal NfL serum levels in our patients reflect the pathology and progression of *RFC1* disease. Postmortem examination revealed loss of sensory neurons in the dorsal root ganglia and Scarpa's ganglion, as well as marked loss of cerebellar Purkinje and granular cells.^{14,20-22} A recent brain magnetic resonance imaging study also showed basal ganglia and brainstem volumetric reduction and involvement of the cerebral white matter in cases with advanced disease,²³ suggesting a widespread cerebral neurodegeneration.

We have also demonstrated that patients with cerebellar damage have significantly higher levels of serum NfL than those without cerebellar involvement. This correlation persisted after correcting for age and gender. This may be explained by the high density of neurons in cerebellum, which would lead to a significantly increased release of NfL in the bloodstream when this structure is affected.

Conversely, serum NfL did not appear to correlate with disease duration and the need for walking aids. An explanation of this phenomenon is that NfL may increase rapidly in the initial stages of the disease and then reach a plateau above a certain degree of severity, as observed in other genetic ataxias.^{12,13,18} Other possible explanations entail the difficulty in accurately defining the onset of the disease, because neuropathy symptoms may remain unnoticed for a long time, or the chronologically variable involvement of the

cerebellum in early or late disease stages relative to onset, due to factors yet to be explored, including the repeat size and additional genetic modifiers. Also, no significant difference in serum NfL levels was found based on vestibular involvement. This could be due to the limited quantitative relevance of Scarpa's ganglion, which contains around 20,000 neurons, compared to the cerebellum's 50 billion neurons.

This study has some limitations. First, the small sample size limits the statistical power of the study. Second, the relatively wide range in the time interval between serum collection and neurological examination (within 24 months at the most) may have introduced variability. However, previous studies in patients with spinocerebellar ataxias and Charcot-Marie-Tooth disease showed no significant difference in serum NfL concentration after 1 or 2 years,^{8,24} suggesting stability of NfL levels in the short time in patients with slowly progressive neurodegenerative diseases. Finally, the cross-sectional design of the study restricts causality assumptions and does not allow us to assess changes in serum NfL levels over time.

In conclusion, we have demonstrated that serum NfL holds promise as a reliable biomarker in *RFC1* disease, as NfL levels are elevated even in the early stages of the disease and exhibit a correlation with cerebellar involvement. Given the significant interindividual variability, NfL may prove to be a valuable tool for stratifying patients, and longitudinal studies are warranted to assess its ability to monitor disease progression. ■

Acknowledgments: We thank MYOBANK-AFM (S. Vasseur, M. Chapart) from the Institute of Myologie for serum preparation. We also thank the study participants and their families for their support.

Data Availability Statement

Anonymized data from this study will be shared by request from any qualified investigator.

References

1. Cortese A, Simone R, Sullivan R, et al. Biallelic expansion of an intronic repeat in *RFC1* is a common cause of late-onset ataxia. *Nat Genet* 2019;51(4):649–658.
2. Rafahi H, Szmulewicz DJ, Bennett MF, Sobreira NLM, Pope K, Smith KR, et al. Bioinformatics-based identification of expanded repeats: a non-reference intronic pentamer expansion in *RFC1* causes CANVAS. *Am J Hum Genet* 2019;105(1):151–165.
3. Cortese A, Tozza S, Yau WY, et al. Cerebellar ataxia, neuropathy, vestibular areflexia syndrome due to *RFC1* repeat expansion. *Brain* 2020;143(2):480–490.
4. Kulshreshtha D, Ganguly J, Jog M. Expanding the clinical Spectrum of *RFC1* gene mutations. *J Mov Disord* 2022;15(2):167–170.
5. Record CJ, Alsukhni RA, Currò R, et al. Severe distinct dysautonomia in *RFC1*-related disease associated with parkinsonism. *J Peripher Nerv Syst* 2022;27(4):311–315.
6. Ylikotila P, Sipilä J, Alapirtti T, et al. Association of biallelic *RFC1* expansion with early-onset Parkinson's disease. *Eur J Neurol* 2023; 30(5):1256–1261.

7. Ashton NJ, Janelidze S, Al Khleifat A, et al. A multicentre validation study of the diagnostic value of serum neurofilament light. *Nat Commun* 2021;12:3400.
8. Sandelius Å, Zetterberg H, Blennow K, et al. Serum neurofilament light chain concentration in the inherited peripheral neuropathies. *Neurology* 2018;90(6):e518–e524.
9. Millere E, Rots D, Simrén J, et al. Plasma neurofilament light chain as a potential biomarker in Charcot-Marie-tooth disease. *Eur J Neurol* 2021;28(3):974–981.
10. Ticau S, Sridharan GV, Tsour S, et al. Neurofilament light chain as a biomarker of hereditary transthyretin-mediated amyloidosis. *Neurology* 2021;96(3):e412–e422.
11. Maalmi H, Strom A, Petrera A, et al. Serum neurofilament light chain: a novel biomarker for early diabetic sensorimotor polyneuropathy. *Diabetologia* 2023;66(3):579–589.
12. Peng L, Wang S, Chen Z, et al. Blood neurofilament light chain in genetic ataxia: a meta-analysis. *Mov Disord* 2022;37(1):171–181.
13. Wilke C, Mengel D, Schöls L, et al. Levels of neurofilament light at the Preataxic and ataxic stages of spinocerebellar ataxia type 1. *Neurology* 2022;98(20):e1985–e1996.
14. Träschütz A, Wilke C, Haack TB, et al. Sensory axonal neuropathy in RFC1-disease: tip of the iceberg of broad subclinical multisystemic neurodegeneration. *Brain* 2022;145(3):e6–e9.
15. Schoeberl F, Abicht A, Kuepper C, et al. Sensory neuropathy due to RFC1 in a patient with ALS: more than a coincidence? *J Neurol* 2022;269(5):2774–2777.
16. Chelban V, Nikram E, Perez-Soriano A, et al. Neurofilament light levels predict clinical progression and death in multiple system atrophy. *Brain* 2022;145(12):4398–4408.
17. Yang L, Shao YR, Li XY, et al. Association of the Level of neurofilament light with disease severity in patients with spinocerebellar ataxia type 2. *Neurology* 2021;97(24):e2404–e2413.
18. Tezenas du Montcel S, Petit E, Olubajo T, et al. Baseline clinical and blood biomarker in patients with Preataxic and early-stage disease spinocerebellar ataxia 1 and 3. *Neurology* 2023;100(17):e1836–e1848.
19. Donath H, Woelke S, Schubert R, et al. Neurofilament light chain is a biomarker of neurodegeneration in ataxia telangiectasia. *Cerebellum* 2022;21(1):39–47.
20. Montaut S, Diedhiou N, Fahrer P, et al. Biallelic RFC1-expansion in a French multicentric sporadic ataxia cohort. *J Neurol* 2021;268(9):3337–3343.
21. Huin V, Coarelli G, Guemy C, et al. Motor neuron pathology in CANVAS due to RFC1 expansions. *Brain* 2022;145(6):2121–2132.
22. Reyes-Leiva D, Aldecoa I, Gelpi E, Rojas-García R. Motor neuron involvement expands the neuropathological phenotype of late-onset ataxia in RFC1 mutation (CANVAS). *Brain Pathol* 2022;32(4):e13051.
23. Matos PCAAP, Rezende TJR, Schmitt GS, et al. Brain structural signature of RFC1-related disorder. *Mov Disord* 2021;36(11):2634–2641.
24. Coarelli G, Darios F, Petit E, et al. Serum neurofilament light chain predicts cerebellar atrophy and clinical progression in spinocerebellar ataxia. *Neurobiol Dis* 2021;153:105311.

Supporting Data

Additional Supporting Information may be found in the online version of this article at the publisher's web-site.