

1     **Exercise addiction in athletes: comparing two assessment instruments and willingness**  
2                                 **to stop exercise after medical advice**

3                                 **Abstract**

4     Exercise is overwhelmingly beneficial for physical and mental health, but for some people exercise  
5     addiction (EA) can develop and negatively impact an individual. This study sought to 1) compare the  
6     latent structure of two instruments assessing EA and 2) examine differences in attitudes towards  
7     stopping exercise, if required to on medical grounds, among exercise-addicted and non-addicted  
8     athletes.

9     In a cross-sectional study, 1,011 athletes competing at different levels completed an anonymous on-  
10    line survey. The survey contained Exercise Dependence Scale-Revised (EDS-R), Exercise  
11    Addiction Inventory (EAI), and questions on adherence to medical prescriptions to stop exercise.  
12    We tested the latent structure of EDS-R and EAI with multigroup confirmatory factor analyses  
13    (CFA), across gender and competition level. Finally, we measured the difference of athletes'  
14    attitudes towards stopping exercise, if prescribed by a physician. Both instruments showed good fit  
15    indexes, even across gender. CFAs on EAI scores showed some violations of measurement  
16    invariance across competition level ( $\Delta\text{CFI}=.03$ ;  $\Delta\text{RMSEA}=.02$ ). On the contrary, CFAs on EDS-R  
17    scores did not show invariance violations across competition level ( $\Delta\text{CFI}<.01$ ;  $\Delta\text{RMSEA}<$   
18     $.01$ ). Finally, athletes who reached thresholds for exercise addiction, by means of EDS-R, were  
19    more prone to not follow medical prescriptions to cease exercise, independently of the competition  
20    level. These results suggest that athletes' answers on the EDS-R seems to be less affected by  
21    competition level, compared to EAI. Moreover, EDS-R outcomes could be used to identify  
22    individuals who may be unlikely to cease exercise for medical reasons, independently of their  
23    competition level.

24

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 **Public Significance Statement.** The present study suggests that the answers to Exercise Dependence  
2 Scale-Revised are not influenced by either the gender or the competition level of an athlete. On the  
3 contrary, the answers to Exercise Addiction Inventory can be influenced by athletes' competition  
4 level. Furthermore, Exercise Dependence Scale-Revised appears to be more effective than the  
5 Exercise Addiction Inventory in detecting athletes' attitude toward medical prescription to stop  
6 exercise.

7

8 **Keywords:** Exercise Dependence Scale; Exercise Addiction Inventory; Measurement invariance;  
9 Multigroup CFA; Psychometric Comparison.

## 1 **Exercise addiction in athletes: comparing two assessment instruments and** 2 **willingness to stop exercise after medical advice**

3  
4 There is overwhelming evidence that exercise is beneficial for physical and mental health  
5 (Ashdown-Franks et al., 2020; Vina et al., 2012). Moderate levels of exercise have been shown to  
6 **have a protective effect on** myocardial infarction and several types of cancer (Kim et al., 2019).  
7 Exercise is also effective in improving a wide range of symptomatic outcomes ranging from primary  
8 symptoms in mental disorders to cognition (Ashdown-Franks et al., 2020; Stubbs et al., 2018).  
9 However, exercise can also become obsessive and compulsive to an extent in which it can negatively  
10 impact on a person's physical and mental health. Such an obsessive-compulsive relationship with  
11 exercise has been termed exercise addiction (Szabo et al., 2015).

12 Exercise addiction was first described by Frederick Baekeland who found that people  
13 exercising for more than three days per week suffered from sleep and psychological symptoms akin  
14 to substance withdrawal symptoms upon cessation of exercise (Baekeland, 1970). Exercise addiction  
15 can cause an impairment in physical (e.g., spine injuries or anemia), and social functioning even **in**  
16 **the absence of** injuries (Hausenblas et al., 2017; Wouthuyzen-Bakker & van Assen, 2015). In the  
17 general population, estimates of prevalence of exercise addiction are extremely heterogeneous, with  
18 reports ranging from 4.0% in adolescent school athletes (Lichtenstein et al., 2018) to 42% in selected  
19 at-risk populations, such as those attending gyms (Lejoyeux et al., 2008).

20 A common component of every type of addiction, either to substances or to specific behaviors,  
21 is the difficulty to cease the activity despite negative consequences. For athletes, such difficulty **is a**  
22 **delicate topic** (Vina et al., 2012), for which the tradeoff **between the amount of** exercise to maintain  
23 appropriate preparation and excessive exercise is not clearly defined. Among athletes, several  
24 arrhythmogenic conditions including cardiomyopathies and coronary artery disease can  
25 contraindicate exercise due to the risk of sudden cardiac death (Maron, 2007; Zorzi et al., 2020). In  
26 several sports for example, all competitive athletes **must undergo an annual** medical examination

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 to screen for such conditions by law (Vessella et al., 2020), and if a potentially fatal cardiac anomaly  
2 is detected, the athlete is disqualified from competitive events and she/he is recommended to stop or  
3 reduce the sporting activity. Athletes with exercise addiction, however, would struggle to follow this  
4 prescription. In a previous study, people assessed as exercise addicted reported to be more likely to  
5 continue practicing despite a potentially worsening heart disease, compared to non-exercise addicted  
6 athletes (Zorzi et al., 2020). Hence, it is clinically relevant to use valid tools detecting, or at least  
7 screening, exercise addiction among competitive athletes, to have a broader picture of their sport  
8 attitude and their resistances to stop or reduce it. To date, the most widely used questionnaires to  
9 measure exercise addiction are the Exercise Addiction Inventory (EAI; Terry et al., 2004) and the  
10 Exercise Dependence Scale- Revised (EDS-R; Costa et al., 2012; Hausenblas & Symons Downs,  
11 2002a, 2002c). These tools have been extensively validated and the psychometric properties tested,  
12 especially during the last decade. For instance, Mónok et al., (2012) examined the two tools'  
13 properties on a nationally representative population and concluded that both instruments are valid to  
14 screen potential exercise addiction (Mónok et al., 2012). On the other hand, in a recent systematic  
15 review (Di Lodovico et al., 2019), the EAI was found to be preferable to the EDS-R in screening the  
16 risk of exercise addiction in sport exercisers, because it identifies higher proportions of at-risk  
17 individuals. Despite the steps forward, both examples suggest how the literature base is still  
18 developing, since several issues still need to be addressed to provide a definitive consensus on which  
19 instrument should be used to screen for exercise addiction (Di Lodovico et al., 2019; Mónok et al.,  
20 2012). For instance, the above examples refer to the general population or specific types of exercisers  
21 (e.g., endurance exercisers, bodybuilders or fitness attendees), however the responses to both  
22 instruments should be analyzed comparing amatorial and professional athletes or, more in general,  
23 on differing types of competition levels. In the same way, it should be tested if the response trends of  
24 male and female athletes could be different (Di Lodovico et al., 2019). The structural and  
25 psychometric validity of EA questionnaires is a relevant issue in specific athletic populations and has  
26 not been extensively explored to date. Furthermore, exercise addiction can be a serious problem for

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 athletes, since it could push athletes beyond their physiological limits, being unable to modulate  
2 training or to stop it when indicated on medical grounds. To date, no study has assessed how available  
3 questionnaires are valid to predict adherence to medical prescriptions to stop exercise, and  
4 importantly, how available questionnaires perform across competition level, and gender.

5 Therefore, the present study has two main aims:

- 6 1. To test the latent structure of EAI and EDS-R across competition level and gender, and to  
7 understand if one of both tools are influenced by such features.
- 8 2. To test the differences between addicted versus non-addicted exercisers (measured by  
9 means EAI and EDS-R) on adherence to medical prescription to stop exercise, even across  
10 competition level.

## 11 **Methods**

### 12 *The sample*

13 A sample of athletes were invited (via social media groups dedicated to athletes) to fill an  
14 anonymous questionnaire asking their opinion both on the importance of sports activity for  
15 themselves and about preparticipation evaluation. Athletes were defined as those individuals engaged  
16 in an organized sports program requiring regular training and competition. According to the Italian  
17 law (Decree of Italian Ministry of Health: Rules for the Health Care of Competitive Sport Activities,  
18 1982), these individuals are required to undergo mandatory preparticipation evaluation by a  
19 specialized sports medicine physician. At the beginning of the questionnaire, participants were asked  
20 if they were currently practicing competitive sports requiring mandatory medical evaluation: those  
21 who replied that they did not need the certificate of fitness were excluded from the analysis.  
22 Moreover, another exclusion criterion was applied to define participants as athletes: those exercising  
23 less than 4 hours per week were excluded. This classification was made accordingly the recent  
24 guidelines applied in sport cardiology (Pelliccia et al., 2020). The final sample consisted of 1011  
25 athletes (mean age  $33.9 \pm 13.75$  years; 23.83% females). The link to the survey was posted in social

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 networks for one month. Participants gave their informed consent to participate to the study, that was  
2 carried out according to the Declaration of Helsinki. The Institutional Review Board of the Padua  
3 Hospital, after a preliminary consultation, specified that ethical approval was not necessary for this  
4 study<sup>1</sup>, since it consisted in an observational study being carried out using anonymous questionnaires  
5 on the general population. Participants did not receive any form of payment to participate.

**Measures**

7 Gender, age, the level of competition (amateur, local, regional, national, international), years  
8 of experience, hours of weekly practice and kind of sport practiced were collected. Table 1 displays  
9 the descriptive statistics of the present study. Categories of practised sport are in Supplmenetary  
10 material (Table S1).

11 [PLEASE, INSERT TABLE 1 HERE]

12 Exercise Dependence Scale- Revised (EDS-R) and Exercise Addiction Inventory (EAI) were  
13 used to measure exercise addiction. The EDS-R is a questionnaire composed of 21 questions rated on  
14 a six-point Likert response scale (1= Never; 6 =Always). The scale is divided into seven sub-  
15 categories (three questions per scale), based on the DSM IV (American Psychiatric Association  
16 [APA], 2000) criteria for substance addiction. EDS-R categories are: withdrawal, continuance,  
17 tolerance, control loss, decrease of other activities, time, and effect of intention. For each category, a  
18 total score is obtained by summing the ratings to each item. In general, higher score suggest higher  
19 chance to be at risk of exercise addiction. A score greater than 14 suggests an at-risk individual,  
20 between 7 and 14 a nondependent-symptomatic individual, and less than 7 denotes a nondependent-  
21 asymptomatic individual. According to the EDS-R manual (Hausenblas & Symons Downs, 2002b),  
22 an exercise dependent profile is suggested whenever at least three subscales present an ‘at-risk’ score;  
23 a nondependent-asymptomatic profile whenever a nondependent-asymptomatic score occurs in at  
24 least four subscales; a nondependent-symptomatic profile otherwise.

---

<sup>1</sup> Consequently, a protocol number was not provided, since the *Institutional Review Board (IRB)* judged it as unnecessary.

## EXERCISE ADDICTION AND COMPETITION LEVEL

1           The EAI is a short questionnaire composed of six items rated on a five-point Likert scale (1 =  
2 strongly disagree, 5= strongly agree), and investigates six main addiction domains: salience, mood  
3 modification, tolerance, withdrawal symptoms, conflict and relapse (Griffiths et al., 2015; Mónok et  
4 al., 2012; Sicilia et al., 2013; Terry et al., 2004). Table 2 contains the items of both instruments. As  
5 with the EDS-R, higher scores indicate a higher risk of exercise addiction. A score greater than 23  
6 suggests at risk individuals; a score between 13 and suggests a potentially symptomatic person while  
7 a score less than 13 suggests an asymptomatic person (Griffiths et al., 2015). Table S2 of  
8 supplementary material contains the frequencies of each level of response, for each item of both  
9 instruments.

10           The following questions were used to assess participants adherence to hypothetical medical  
11 prescriptions made by a *physician*<sup>2</sup>: “*How would you react to a medical prescription to temporarily*  
12 *interrupt exercise (Q1 in the sequel); “After a medical investigation, it emerged that you have a*  
13 *critical cardiac anomaly that could cause a deadly heart attack. Unfortunately, it is necessary to*  
14 *permanently stop exercise. How would you react*<sup>3</sup>?” (Q2 in the sequel). Both questions were rated on  
15 a dichotomous scale indicating the intention to stop (0) or to continue sport activities (1) despite  
16 medical prescription.

**17 Analytic plan**

18           We performed confirmatory factor analysis (CFA) on each instrument. Considering that data  
19 were skewed (see Table S2), we considered them as ordinal and adopted a Diagonally Weighted  
20 Least Squares (Li, 2016) robust estimator in all the CFAs. As suggested by several authors (Forero  
21 et al., 2009; Kyriazos, 2018), the sample size for studies including confirmatory factor analysis

---

<sup>2</sup> The fact that the medical prescriptions were made by a physician was an information that participants derived from (a) the instruction of the section containing the questions within the online form and (b) from the word used. In particular, the adjective “medical” in Italian means that the recommendation or the exam is made by a physician, that is the only one that can make a diagnosis or prescribe to stop exercise.

<sup>3</sup> This question was the last of a series of question investigating the reaction to medical prescription. The question “how would you react?” was a short for of the question “*How would you react to this medical prescription?*”

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 should range from 200 to 500 participants. Considering that we performed two multigroup CFAs,  
2 we found it reasonable to collect at least 1000 participants (Comrey & Lee, 2013).

3 We applied a listwise deletion of missing data (i.e., only participants who both reported the  
4 gender/competition level and responded to all the items of the scale were included). In this way, we  
5 used the same set of complete cases to run all the CFAs. **Furthermore, we checked if missing data  
6 were equally and randomly distributed across items or if systematic patterns of missing data  
7 occurred for specific items. After the data-elimination, we also checked if there were differences  
8 in age between included and excluded participants. This gave us the chance to consider also  
9 potential influences of age on the CFAs results.** As suggested by several authors (Mónok et al.,  
10 2012; Schermelleh-Engel et al., 2003; Weston & Gore, 2006), we used multiple goodness-of-fit  
11 statistics in order to interpret the models' fit. For the Comparative Fit Index (CFI: Bentler, 1990)  
12 values  $\geq .95$  represent a good fit, values  $\geq .90$  for an adequate fit, while values  $< .90$  a not acceptable  
13 fit. **For Root Mean Square Error of Approximation's (RMSEA: Steiger, 1990) values  $\leq .05$   
14 indicate a good fit, values between .05 and .10 an adequate fit while values  $> .10$  a not acceptable  
15 fit .** For the Standardized Root Mean Square Residual (SRMR: Hu & Bentler, 1999), values  $\leq .05$   
16 represent a good fit, values between 0.5 and 0.8 an adequate fit while values  $> .8$  a not acceptable fit.  
17 We used the R statistical environment (R Core Team, 2020) to run all analyses and the *lavaan* package  
18 (Rosseel, 2012) for the CFAs. Furthermore, we used Cronbach's alpha to test the internal consistency  
19 of both questionnaires, by means of the *ltm* package (Rizopoulos, 2006).

20 We tested measurement invariance of both questionnaires' items score to understand if some  
21 external characteristics could affect the participants answers. Considering that our main goal was to  
22 test the EDS-R and the EAI latent structures across competition level and gender, we tested a series  
23 of multigroup CFAs on these variables, separately. As recommended by several authors (Putnick &  
24 Bornstein, 2016; Vandenberg & Lance, 2000), we focused on the configural, metric, scalar and  
25 residual invariances. **As a first step, we performed an omnibus test to both fit a baseline model  
26 and to test the general structure of both instruments.** After testing separate model for each group,



## EXERCISE ADDICTION AND COMPETITION LEVEL

1 we tested the configural invariance, to understand if the factor structure was equal across groups (van  
2 de Schoot et al., 2012). Configural invariance was supported if the model presented (at least) adequate  
3 goodness-of-fit indices and the pattern of loadings were significant in all groups. Then, we  
4 constrained the factor loadings to be equal across groups to test the metric invariance. Such test allows  
5 to understand if the contribution (i.e., the loading) of each item to its factor is equivalent among  
6 groups. Metric invariance was supported if the model fit was not worse than the fit of configural  
7 model. As suggested by several authors (Cheung & Rensvold, 2002; Gilson et al., 2013) we focused  
8 on two criteria to compare both models fit and, therefore, to discuss about the presence of  
9 measurement invariance: (1), the difference ( $\Delta$ ) between fit indices of the models, where we  
10 considered a  $\Delta CFI > .01$  and a  $\Delta RMSEA > .015$  as indicative of invariance violations; (2), the overall  
11 goodness of fit of the singular models: not acceptable fit suggested us potential non-invariance  
12 (Beaujean et al., 2011). As a fourth step, we constrained both the factor loadings and the **item**  
13 **intercepts/thresholds**<sup>4</sup> to be equal across groups, to test scalar invariance. In this way, it was possible  
14 to understand if both the starting level toward the construct and the threshold of response were  
15 equivalent among groups. We compared the overall fit of this model model with the metric invariance  
16 one. Finally, we tested the residual invariance (i.e., by constraining factor loadings, item  
17 intercepts/thresholds and residual variances to be equal across groups) to understand if the specific  
18 and error variances of the items were equivalent among groups (Putnick & Bornstein, 2016). In this  
19 case, model fit was compared with the scalar invariance one. Among the comparisons, whenever  
20 measurement non-invariance was found, we decided to test also partial measurement invariance. In  
21 particular, we constrained the parameter of the invariant items, relaxing the other parameters and  
22 comparing again the models. In this way, it was possible to identify which groups could interpret the  
23 construct assessed by a specific item in different ways. We used the *lavaan* package (Rosseel, 2012)

---

<sup>4</sup> In case of ordinal scale as the ones in the present study, an item threshold can be defined as the level of latent trait required to endorse a point of the response scale more likely than the previous response point. For ordinal and Likert-like scale, each item will have n-1 thresholds. Since item belonging to EAI are rated on a 5-point Likert scale, they present four thresholds. Items belonging to EDS-R present five thresholds.

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 to test measurement invariance. The procedure we used to define and compare models followed the  
2 recommendations of Hirschfeld and von Brachel (2014).

3 As a further comparison between instruments, we also tested the specificity and sensitivity of  
4 both tool by means of receiver operating characteristic (ROC) curve. This analysis also yielded  
5 potential cutoffs for both tools, comparing them with the ones emerged from the original scoring  
6 algorithms through the kappa statistics (Cohen, 1960). We used two external criteria to run ROC  
7 analysis, namely the answers to the question Q1 and the percentiles on the hours of weekly practice.  
8 Participants who both (a) expressed the will to continue the sport activities despite medical  
9 prescription and (b) were over the 95% percentile along the distribution of the training hours were  
10 considered as at risk of addiction. We determined the cut off on the total scores of both instruments,  
11 in line with previous works testing the structure of the instruments (Müller et al., 2013). We used the  
12 pROC package (Robin et al., 2011).

13 To understand whether the EDS-R and EAI scores were influenced by gender, years of  
14 practice and average time devoted to exercise every week, we adopted multiple linear regressions.  
15 We also considered both gender  $\times$  years of practice and gender  $\times$  average time devoted to exercise  
16 every week interactions. We used partial eta squared values ( $\eta^2$ ) to quantify the effect size, by means  
17 of the *heplots* package (Friendly, 2010).

18 Finally, based on EDS-R and EAI outcomes, we tested the difference between addicted versus  
19 non addicted exercisers on the adherence to medical prescription. We performed a multiple logistic  
20 regression, setting Q2 responses as dependent variable and EDS-R/EAI outcomes and the competition  
21 level as categorical predictors. Regarding the EDS-R/EAI outcomes, we were interested in comparing  
22 participants considered at risk of exercise addiction to participants who were either symptomatic-not  
23 addicted or asymptomatic. We also considered this comparison across competition levels. Moreover,  
24 odds ratios were calculated for each coefficient. **About the sample size used for this last group of**  
25 **analyses (i.e., logistic, multiple regressions and ROC), we used only those participants who**  
26 **provided data on the target categorical variables. For instance, in the multiple regressions, we**

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 used all those individuals that provided complete data on gender, years of practice and average  
2 practice time. To calculate scores in case of missing data on specific items of both EAI and EDS-  
3 R, we applied a mean imputation method, but only for those subjects who had at least the ~90%  
4 of items responses. As can be seen in Table S2, we imputed a very small set of cases, considering  
5 that the amount of missing data (per item) was negligible. This strategy allowed us to reduce  
6 the loss of useful data.

## 8 Results

### 9 *Prevalence and descriptive statistics*

10 The calculated prevalence of exercise addiction was 6% and 11% when measured with the EDS-R  
11 and EAI respectively. Considering mean differences on scales and subscales' scores, we observed  
12 that, in general, male participants obtained lower scores than female ones in all the overall  
13 scales and subscales, except for tolerance ( $p= .54$ ), lack of control ( $p= .07$ ), reduction in other  
14 activities ( $p=.21$ ) and intention effects EDS-R subscales ( $p=.59$ ). Furthermore, we observed that  
15 participants who reported an intention to continue exercising (i.e., data taken from answer to  
16 both questions Q1 and Q2), despite medical prescriptions or suggestions, obtained higher scores  
17 in all the overall scales and subscales. We did not observe mean differences on the withdrawal  
18 subscale between respondents to the Q1 question ( $p= .31$ ). Finally, we observed that  
19 participants competing at international level obtained the highest scores (especially compared  
20 to amatorial participants) in all the scales and subscale and, such result, was statistically  
21 significant for both EAI and EDS-R total score and almost all the EDS-R subscales (see Table  
22 1).

### 23 *Latent structure and measurement invariance*

24 The sample used to run all the CFAs consisted of 943 participants (i.e., composed by participants  
25 who reported their gender, level of competition and all the responses to EAI and EDS-R). **Descriptive**

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 **statistics of this subsample can be found within supplementary materials (see Table S3). We did**  
2 **not find systematic patterns of missingness among items (all the items presented very low and**  
3 **similar rates of missingness). We also did not find age differences among participants who were**  
4 **included or excluded.** Regarding the EAI, the CFA suggested a good fit of the entire model to the  
5 data ( $\chi^2(9) = 29.33$ ; CFI = .98; SRMR = .03; RMSEA = .05[.01-.05]). Standardized factor loadings  
6 ranged from .33 to .69 and are reported in Table 2, and internal consistency was  $\alpha = .68$ . Independent  
7 CFAs on each competition type obtained good to acceptable fit indexes, (range CFI: .95-.99; range  
8 SRMR: .04-.06; range RMSEA: .05-.09) except for the RMSEA of amatorial athletes (.09; see Table  
9 3). As for stratified models on competition levels, CFAs on gender revealed good fit indices (female:  
10  $\chi^2(9) = 6.79$ ; CFI = 1; SRMR = .03; RMSEA = .00[.00-.06]; male:  $\chi^2(9) = 28$ ; CFI = .98; SRMR = .03;  
11 RMSEA = .05[.03-.07]). Multigroup CFA to test measurement invariance showed that the model on  
12 competition level was good in terms of configural and metric invariance, but some invariance  
13 violations emerged comparing the metric with the scalar model ( $\Delta\text{CFI} = .03$ ;  $\Delta\text{RMSEA} = .02$ ). **Once**  
14 **we found the violation, we applied the Lagrange Multiplier test (by jointly using the**  
15 ***lavTestScore()* and the *parTable()* functions of the lavaan package), for releasing single**  
16 **constraints. This test allowed us to test if item thresholds statistically differed among groups.**  
17 Examining such items thresholds of the scalar model, we found that the threshold four of the item  
18 three of EAI (i.e., “I use exercise as a way of changing my mood”) of athletes competing at  
19 international level was the lowest (0.14) compared to the thresholds of athletes competing at different  
20 levels (ranging from 0.22 to 0.25). Likewise, the threshold three for of item four of EAI (i.e., “Over  
21 time I have increased the amount of exercise I do in a day”) of athletes competing at regional level  
22 was the lowest (|-0.57|) compared to the thresholds of athletes competing at different levels (ranging  
23 from |-0.61| to |-0.69|). Therefore, we relaxed the parameters of those thresholds and tested the partial  
24 scalar invariance. As shown in Table 3, the goodness-of-fit indices and the deltas improved ( $\Delta\text{CFI} <$   
25  $.01$ ;  $\Delta\text{RMSEA} < .01$ ).

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 On the contrary, the model across gender suggested configural invariance, with good indices of fit  
 2 ( $\chi^2(18) = 32.44$ ; CFI = .99; SRMR = .03; RMSEA = .04[.02-.06]), suggesting the same factor structure  
 3 across female and male participants. Similar results emerged in terms of metric, scalar and residual  
 4 invariance. Overall, the results indicate that the EAI showed the same factor loadings, the same item  
 5 intercepts and the same residual variances across female and male athletes.

6 [PLEASE, INSERT TABLE 2 HERE]

7 Regarding the EDS-R, as reported in Table 3, the fit of the entire model to the data resulted as good  
 8 ( $\chi^2(168) = 805.06$ ; CFI = .97; SRMR = .05; RMSEA = .06[.06-.07]). Standardized factor loadings  
 9 ranged from .54 to .97, as shown in Table 2. EDS-R showed very good internal consistency both as  
 10 an entire scale ( $\alpha = .83$ ), and within each subscale ( $\alpha$  withdrawal = .92;  $\alpha$  continuity = .80;  $\alpha$  tolerance =  
 11 .82;  $\alpha$  control loss = .85;  $\alpha$  reduction = .57;  $\alpha$  time = .79;  $\alpha$  intention = .82) except for the reduction in  
 12 other activities subscale. In the multigroup CFAs, all the models on competition levels ranged from  
 13 good to acceptable indices (range CFI: .97-.98; range SRMR: .06-.08; range RMSEA: .06-.07).  
 14 Similarly, we found acceptable goodness-of-fit indices on gender models (female:  $\chi^2(168) = 348.47$ ;  
 15 CFI = .97; SRMR = .06; RMSEA = .07[.06-.08] male:  $\chi^2(168) = 634.08$ ; CFI = .98; SRMR = .05;  
 16 RMSEA = .06[.06-.07]). We observed configural invariance both on competition level and gender,  
 17 with also metric, scalar and residual invariances supported by small changes in models fit (see Table  
 18 3).

19 [PLEASE, INSERT TABLE 3 HERE]

20 On the ROC analysis ( $n = 1006$ ), we observed that a cutoff of 19.5 for the EAI was sufficient  
 21 to determine people at risk to develop exercise addiction. The specificity was .56, the sensitivity was  
 22 .67 and the area under the curve (AUC) was equal to .67 (**Figure 1, upper panel**). Based on this new  
 23 cut off, 439 participants were screened as at risk (43.22%). The kappa coefficient between the new  
 24 cut off and the one suggested by Hausenblas and Symons Downs (2002b) was .28, suggesting  
 25 minimal agreement (Cohen, 1960). In regard to the EDS-R, we observed that participants with a score  
 26 higher than 76.5 were screened as at risk, with a specificity of .82, a sensitivity of .67 and an AUC of

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 .73 (**Figure 1, lower panel**). Based on this new cutoff, 181 participants were screened as at risk (i.e.,  
2 17.90%). In this case, the agreement between the new and the original cutoffs was weak ( $\kappa = .41$ ).

3 [PLEASE, INSERT FIGURE 1 HERE]

4 *Effects of gender, years of practice and average practice time on exercise addiction*

5 The multiple regression (**n= 999**) on EAI scores yielded no significant effects on gender, years  
6 of practice and the interaction effects. The effect of the average time per week spent in training  
7 emerged as small ( $\eta^2 = .03$ ) but statistically significant: a one-unit increase in hours per week spent  
8 in training led to an increase of .11 in EAI score.

9 [PLEASE, INSERT TABLE 4 HERE]

10 The multiple regression on the EDS-R total scores (**n= 999**) also yielded no significant effects  
11 across years of practice or interaction effects. There was, however, a small but significant effect of  
12 gender ( $\eta^2 < .01$ ). Moreover, we found that a one-unit increase in hours per week spent in training  
13 led to an increase of .39 in EDS-R score ( $\eta^2 = .047$ ).

14 Finally, logistic regression (**n= 976**) suggested that only the EDS-R showed a statistically  
15 significant difference between participants at risk of exercise addiction and non-addicted. We  
16 observed that the odds of continuing exercising (even in case of a critical cardiac anomaly) in  
17 participants who were screened at risk of exercise addiction were 2.91 times greater than the odds of  
18 continuing exercising in participants who were screened as exercise non-addicted (OR=2.91 CI=1.01-  
19 8.94  $p = .04$ ; see Table 5), a results that was not found in the EAI. In this case, the odds of continuing  
20 exercising (even in case of a critical cardiac anomaly) in participants who competed at local level  
21 were 0.62 times lower than the odds of continuing exercising in participants who competed at  
22 amatorial level (OR=0.62 CI=0.4-0.96  $p = .03$ ).

23 [PLEASE, INSERT TABLE 5 HERE]

24

## EXERCISE ADDICTION AND COMPETITION LEVEL

**Discussion**

During the past ten years, exercise addiction has received increasing attention (Di Lodovico et al., 2019). There is the possibility that sport could be compulsive to the point that it may modify the adherence to medical prescription to cease exercise, if required. Athletes may find this particularly difficult, considering that most of the time athletes are guided by a strong passion that could moderate the relevance of any exercise addiction (Vega et al., 2016). Both main and differential diagnoses can be difficult (Freimuth et al., 2011), as exercise addiction has been reported to exist in a continuum, from beneficial exercise to exercise addiction. Furthermore, a number of comorbid disorders, such as eating disorders in particular, and other addictive or impulsive behaviors can co-exist (Berczik et al., 2012; Freimuth et al., 2011; Lichtenstein et al., 2018; Pinna et al., 2015; Trott, Jackson, Firth, Jacob, et al., 2020). Therefore, it is fundamental to build and test assessment instruments, whose structure must be stable not only in general populations, but also in different athletic populations across several different competition levels. Moreover, a tradeoff between brevity and exhaustivity must be reached for such assessment tools. Currently there are two well-validated questionnaires, the EDS-R and the EAI, with the EDS-R having the advantage of investigating more aspects related to exercise addiction, and the EAI being faster to administer. Although some studies have compared these tools in terms of validity, usability and factorial structure, these validation studies were not stratified by gender or athletic level of competition. The present study tried to add new insight onto such issues debated in literature (Di Lodovico et al., 2019; Mónok et al., 2012), retesting the latent structures of EAI and EDS-R across competition level and gender. Furthermore, the present study tried to understand if, based on those tools' outcome, those identified as at risk of exercise addiction would be less adherent the medical prescriptions, compared to non-addicted individuals.

The difference in prevalence rates was large, and in general agreement with previous studies that have found the EAI to yield higher prevalence rates than the EDS-R (Trott, Jackson, Firth, Fisher, et al., 2020; Zorzi et al., 2020), possibly because of the differing number of items and

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 underlying theoretical structures the tools are based on. The ratio between prevalence rates was  
2 confirmed also by the ROC analyses, which suggested that EAI screened more participants as “at  
3 risk” than the EDS-R, even using new external criteria. These results suggest that the EDS-R should  
4 be preferred to the EAI in cases where more conservative screening for exercise addiction is  
5 necessary.

6 We observed that both instruments showed a good overall factorial structure, in line with  
7 previous works (Mónok et al., 2012; Terry et al., 2004). Moreover, we found that both the EAI and  
8 EDS-R can be independently used by female and male athletes **(even if male participant reported,  
9 in descriptive terms, lower scores in almost all the tested scales and subscales)**, as suggested by  
10 their measurement invariance. These results suggest, therefore, that both instruments can be used in  
11 scenarios where screening of exercise addiction is required, controlling for gender. Some potential  
12 issues in terms of measurement invariance emerged for the EAI where violations of scalar  
13 invariance emerged. **Recalling that an item threshold can be defined as the level of latent trait  
14 necessary to endorse a point of the response scale more likely than the previous response  
15 point, we found that the fourth threshold of item three (i.e., “I use exercise as a way of  
16 changing my mood”) and the third threshold of item four (i.e., “Over time I have increased  
17 the amount of exercise I do in a day”) were the two sources of such scalar non-invariance. In  
18 the former, we observed that athletes competing at international level had the lowest fourth  
19 threshold, suggesting for these athletes it is necessary for a lower level of the latent trait to  
20 answer a “5” (the top score) to this item over a “4”. In other words, such athletes seem to be  
21 more prone to score the maximum on this item than other athletes.** In line with previous works  
22 with similar results (e.g. Rhudy et al., 2020), this result could reflect either a real difference toward  
23 the tendency of using exercise as a way to change mood or a response bias caused by different  
24 endorsement of the response among groups. In the latter, we observed that the lowest third  
25 threshold belonged to the athletes competing at regional level, suggesting the same dichotomy  
26 found for item three. On the contrary, the structure of the EDS-R emerged as invariant across



## EXERCISE ADDICTION AND COMPETITION LEVEL

1 competition levels. Although two thresholds of 24 (four thresholds  $\times$  6 items of the EAI) could be  
2 interpreted as trivial from a measurement point of view (Rhudy et al., 2020), it could be preferable  
3 to use an instrument that has fewer critical points. Therefore, it appears that the EDS-R is preferable  
4 to the EAI in this population.

5 A slight difference between the instruments emerged in the ROC analysis. Based on the two  
6 external criteria, we observed that the EAI yielded more participants at risk of exercise addiction  
7 than the EDS-R did, but with less sensitivity. The ratio between our new prevalence rates were  
8 broadly in line with the ratio found by calculating the prevalence rates using the methods proposed  
9 by the instrument's authors. It appears that the EAI consistently yields almost double the prevalence  
10 of participants at risk of EA than the EDS-R. It is important to stress, however, that the results of  
11 the ROC analyses should be taken with caution. Both minimal (for EAI) and weak (for EDS-R)  
12 agreements with the original scoring algorithm suggested that the measure used as external criteria  
13 are not free from bias, due to the self-reporting nature of the tools. This constitutes a limitation of  
14 the study but also paves also the way for future studies aimed at finding more objective criteria.

15 Another aspect highlighting a difference between the EAI and the EDS-R came after testing  
16 the difference on adherence to medical prescriptions. **In general, we observed that participants**  
17 **who reported an intention to continue exercising obtained higher scores in both EAI and**  
18 **EDS-R scales and almost all EDS-R subscales. Beyond such descriptive results,** the analysis the  
19 logistic regression suggested that, when the medical prescription was to permanently stop  
20 exercising due to a potential (and hypothetical) cardiac anomaly, athletes at risk of addiction were  
21 less prone to adhere to the prescription, but only when exercise addiction was measured by the  
22 EDS-R. This result was independent of competition level. Further study should aim to replicate  
23 these results that, currently, are merely exploratory.

24 Such results have several potential clinical implications, beyond the psychometric aspects.  
25 First, the risk of exercise addiction appears to be not so infrequent, and should be routinely assessed  
26 in competitive athletes, given that it can lead to excessive exercising with higher risk of injuries,

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 impairment in occupational, and social functioning (Hausenblas et al., 2017). Considering the  
2 method of exercise addiction measurement, the present results suggest that, in case of competitive  
3 athletes, the EDS-R is preferable to the EAI, mainly because the responses to the questions in this  
4 tool appear to not be influenced by the competition level of an athlete. Levels of exercise among  
5 athletes are of course higher than those in the general population, and it is reasonable that anaerobic  
6 and cardio-respiratory training of the athlete be judged as excessive from the general population  
7 point of view. Nonetheless, when occupational and social functioning becomes impaired due to  
8 excessive time and mental focus dedicated to exercise, which is often the criteria for other forms of  
9 addiction, exercising can assume a pathological entity in this population (Hausenblas et al., 2017).  
10 Reducing exercise and replacing this with other stimulating activities should be attempted once a  
11 person is judged as addicted to exercise. Interventions to prevent occupational, social functioning  
12 impairment and recurrent injuries occurring from exercise addiction should be ideally implemented  
13 systemically, from the lowest level of competitions, since more engrained addictive behaviours  
14 could emerge when the athlete career improves.

15 Secondly, we observed differences between addicted and non-addicted athletes in terms of  
16 adherence to medical prescriptions. When the medical prescription was to permanently stop  
17 exercise due to a cardiac anomaly, this difference was detected only by assessing the risk of  
18 exercise addiction with the EDS-R, in line with a previous research (Zorzi et al., 2020). Such result  
19 suggests that the EDS-R could detect athletes' attitude more effectively than the EAI. This result is  
20 particularly relevant among subjects who may receive a contraindication to continue exercise. If  
21 they continue exercising despite the presence of a cardiac disease at risk of sudden cardiac death,  
22 the risk goes beyond social and occupational impairment or injuries. Hence, one further clinical  
23 implication of the present findings is that exercise addiction should be assessed in people with  
24 cardiac anomalies.

25 Thirdly, our results regarding the psychometric properties of the EDS-R across gender and  
26 competition level, provide new data in the literature comparing it with EAI (Di Lodovico et al.,

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 2019; Mónok et al., 2012). It appears that the EDS-R is a valuable candidate to assess exercise  
2 addiction, even if administered in non-psychological contexts. This aspect is not trivial: in general,  
3 the medical assessment to allow or not allow a person practicing competitive exercise is performed  
4 by a medical doctor specialized in cardiology or sport medicine. Both may not be very familiar with  
5 questionnaires or scales and need to assess several biological variables during an often-small  
6 amount of time. Hence, they need an exhaustive tool to routinely administer it as a part of the whole  
7 evaluation. Our results showed that EDS-R can provide useful results, and information less  
8 influenced by external features, although it contains a greater number of items and dimensions than  
9 EAI.

10         The present work has some limitations. The operational definition of athletes we used in this  
11 work follows Italian law (Decree of Italian Ministry of Health: Rules for the Health Care of  
12 Competitive Sport Activities, 1982), and guidelines derived from sport cardiology (Pelliccia et al.,  
13 2020). Therefore, such a definition could be context-specific, namely it does not completely overlap  
14 with the ones that can be found in literature (Araújo & Scharhag, 2016). It would be interesting to  
15 understand if our results could be replicated adopting more international definitions. Furthermore,  
16 this study is cross-sectional, hence no causal inference can be made. The question on adherence to  
17 medical prescription assessed an intention not to comply, asked in a self-report modality. Given all  
18 the limitations of such a self-report tool, other and more objective measures could be used in future  
19 studies. Moreover, multigroup CFAs of competition type either across or separately for gender  
20 would have been beneficial, but it would require a sample fitting 10 groups (5 competition levels ×  
21 2). Our sample size, although consisting of 943 participants with complete data, would have needed  
22 to be higher to provide reliable results. The present sample was also unbalanced across both gender  
23 and competition types, increasing the risk of mis specified and underfitted models. To address this,  
24 future studies should collect more responses from female athletes at different competition levels.  
25 Finally, it would be interesting to screen athletes for sub-threshold OCD or eating disorders  
26 considering that such disorders are often comorbid or primary to exercise addiction (Berczik et al.,

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 2012; Freimuth et al., 2011; Lichtenstein et al., 2018; Pinna et al., 2015; Trott, Jackson, Firth,  
2 Jacob, et al., 2020).

3 In summary, this study presents new useful findings. Our results indicate that the usefulness  
4 of the EAI may be limited to screening of exercise addiction requiring responses not influenced by  
5 gender. Furthermore, the structure of the EDS-R appears to be less influenced by gender or  
6 competition level of the athlete. The EAI identifies more individuals reaching thresholds of exercise  
7 addiction (Di Lodovico et al., 2019), possibly overestimating exercise addiction. The EDS-R can  
8 provide a comprehensive and multidimensional picture of addiction related symptoms, and appears  
9 to be less affected by athletes' characteristics. Finally, the outcome provided by EDS-R can be used  
10 to detect differences on athletes' attitude to comply to medical prescription to stop exercise,  
11 independently of their competition level.

12 We therefore conclude that the EDS-R is a good candidate to be administered in several  
13 contexts, including offices of general practitioners (Di Lodovico et al., 2019; Griffiths et al., 2015;  
14 Mónok et al., 2012).

**15 Acknowledgments:**

16 Brendon Stubbs is supported by a Clinical Lectureship (ICA-CL-2017-03-001) jointly  
17 funded by Health Education England (HEE) and the National Institute for Health Research (NIHR).  
18 Brendon Stubbs is part funded by the NIHR Biomedical Research Centre at South London and  
19 Maudsley NHS Foundation Trust. Brendon Stubbs also holds active grants with the Medical  
20 Research Council (GCRF and multimorbidity calls) and Guys and St Thomas Charity (GSTT).  
21 Brendon Stubbs has received consultancy fees from ASICS Europe BV. The views expressed are  
22 those of the author(s) and not necessarily those of the (partner organisation), the NHS, the NIHR,  
23 the Department of Health and Social Care, the MRC or GSTT.

24 Joseph Firth is supported by a University of Manchester Presidential Fellowship (P123958)  
25 and a UK Research and Innovation Future Leaders Fellowship (MR/T021780/1).

26 All other authors have nothing to declare.

EXERCISE ADDICTION AND COMPETITION LEVEL

1

## EXERCISE ADDICTION AND COMPETITION LEVEL

## References

- 1  
2 American Psychiatric Association. (2000). Diagnostic and Statistical Manual of Mental Disorders-  
3 IV Text Revision, APA. In *Washington, DC*.
- 4 Araújo, C. G. S., & Scharhag, J. (2016). Athlete: a working definition for medical and health  
5 sciences research. *Scandinavian Journal of Medicine & Science in Sports*, *26*(1), 4–7.  
6 <https://doi.org/10.1111/sms.12632>
- 7 Ashdown-Franks, G., Firth, J., Carney, R., Carvalho, A. F., Hallgren, M., Koyanagi, A.,  
8 Rosenbaum, S., Schuch, F. B., Smith, L., Solmi, M., Vancampfort, D., & Stubbs, B. (2020).  
9 Exercise as Medicine for Mental and Substance Use Disorders: A Meta-review of the Benefits  
10 for Neuropsychiatric and Cognitive Outcomes. *Sports Medicine*, *50*(1), 151–170.  
11 <https://doi.org/10.1007/s40279-019-01187-6>
- 12 Baekeland, F. (1970). Exercise Deprivation: Sleep and Psychological Reactions. *Archives of*  
13 *General Psychiatry*, *22*(4), 365–369. <https://doi.org/10.1001/archpsyc.1970.01740280077014>
- 14 Beaujean, A. A., Freeman, M. J., Youngstrom, E., & Carlson, G. (2011). The Structure of Cognitive  
15 Abilities in Youths With Manic Symptoms: A Factorial Invariance Study. *Assessment*, *19*(4),  
16 462–471. <https://doi.org/10.1177/1073191111399037>
- 17 Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, *107*(2),  
18 238–246. <https://doi.org/10.1037/0033-2909.107.2.238>
- 19 Berczik, K., Szabó, A., Griffiths, M. D., Kurimay, T., Kun, B., Urbán, R., & Demetrovics, Z.  
20 (2012). Exercise Addiction: Symptoms, Diagnosis, Epidemiology, and Etiology. *Substance*  
21 *Use & Misuse*, *47*(4), 403–417. <https://doi.org/10.3109/10826084.2011.639120>
- 22 Cheung, G. W., & Rensvold, R. B. (2002). Evaluating Goodness-of-Fit Indexes for Testing  
23 Measurement Invariance. *Structural Equation Modeling: A Multidisciplinary Journal*, *9*(2),  
24 233–255. [https://doi.org/10.1207/S15328007SEM0902\\_5](https://doi.org/10.1207/S15328007SEM0902_5)

## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1 Cohen, J. (1960). A Coefficient of Agreement for Nominal Scales. *Educational and Psychological*  
2 *Measurement, 20*(1), 37–46. <https://doi.org/10.1177/001316446002000104>
- 3 Comrey, A. L., & Lee, H. B. (2013). *A first course in factor analysis*. Psychology press.
- 4 Costa, S., Cuzzocrea, F., Hausenblas, H. A., Larcán, R., & Oliva, P. (2012). Psychometric  
5 examination and factorial validity of the Exercise Dependence Scale-Revised in Italian  
6 exercisers. *Journal of Behavioral Addictions JBA, 1*(4), 186–190.  
7 <https://doi.org/10.1556/jba.1.2012.009>
- 8 Decree of Italian Ministry of Health: Rules for the health care of competitive sport activities,  
9 (1982).
- 10 Di Lodovico, L., Poultais, S., & Gorwood, P. (2019). Which sports are more at risk of physical  
11 exercise addiction: A systematic review. *Addictive Behaviors, 93*, 257–262.  
12 <https://doi.org/https://doi.org/10.1016/j.addbeh.2018.12.030>
- 13 Forero, C. G., Maydeu-Olivares, A., & Gallardo-Pujol, D. (2009). Factor Analysis with Ordinal  
14 Indicators: A Monte Carlo Study Comparing DWLS and ULS Estimation. *Structural Equation*  
15 *Modeling: A Multidisciplinary Journal, 16*(4), 625–641.  
16 <https://doi.org/10.1080/10705510903203573>
- 17 Freimuth, M., Moniz, S., & Kim, S. R. (2011). Clarifying Exercise Addiction: Differential  
18 Diagnosis, Co-occurring Disorders, and Phases of Addiction. *International Journal of*  
19 *Environmental Research and Public Health, 8*(10), 4069–4081.  
20 <https://doi.org/10.3390/ijerph8104069>
- 21 Friendly, M. (2010). HE Plots for Repeated Measures Designs. *Journal of Statistical Software; Vol*  
22 *1, Issue 4 (2010)* . <https://doi.org/10.18637/jss.v037.i04>
- 23 Gilson, K.-M., Bryant, C., Bei, B., Komiti, A., Jackson, H., & Judd, F. (2013). Validation of the  
24 Drinking Motives Questionnaire (DMQ) in older adults. *Addictive Behaviors, 38*(5), 2196–

## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1 2202. <https://doi.org/https://doi.org/10.1016/j.addbeh.2013.01.021>
- 2 Griffiths, M. D., Urbán, R., Demetrovics, Z., Lichtenstein, M. B., de la Vega, R., Kun, B., Ruiz-  
3 Barquín, R., Youngman, J., & Szabo, A. (2015). A cross-cultural re-evaluation of the Exercise  
4 Addiction Inventory (EAI) in five countries. *Sports Medicine - Open*, 1(1), 5.  
5 <https://doi.org/10.1186/s40798-014-0005-5>
- 6 Hausenblas, H. A., Schreiber, K., & Smoliga, J. M. (2017). Addiction to exercise. *BMJ*, 357, j1745.  
7 <https://doi.org/10.1136/bmj.j1745>
- 8 Hausenblas, H. A., & Symons Downs, D. (2002a). Exercise dependence: a systematic review.  
9 *Psychology of Sport and Exercise*, 3(2), 89–123. <https://doi.org/https://doi.org/10.1016/S1469->  
10 [0292\(00\)00015-7](https://doi.org/10.1016/S1469-0292(00)00015-7)
- 11 Hausenblas, H. A., & Symons Downs, D. (2002b). Exercise Dependence Scale-21 Manual.  
12 *Recuperado de Http://Www. Personal. Psu. Edu/Dsd11/EDS/EDS21Manual. Pdf.*
- 13 Hausenblas, H. A., & Symons Downs, D. (2002c). How Much is Too Much? The Development and  
14 Validation of the Exercise Dependence Scale. *Psychology & Health*, 17(4), 387–404.  
15 <https://doi.org/10.1080/0887044022000004894>
- 16 Hirschfeld, G., & Von Brachel, R. (2014). Improving Multiple-Group confirmatory factor analysis  
17 in R—A tutorial in measurement invariance with continuous and ordinal indicators. *Practical*  
18 *Assessment, Research, and Evaluation*, 19(1), 7.
- 19 Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis:  
20 Conventional criteria versus new alternatives. *Structural Equation Modeling: A*  
21 *Multidisciplinary Journal*, 6(1), 1–55. <https://doi.org/10.1080/10705519909540118>
- 22 Kim, Y., Sharp, S., Hwang, S., & Jee, S. H. (2019). Exercise and incidence of myocardial  
23 infarction, stroke, hypertension, type 2 diabetes and site-specific cancers: prospective cohort  
24 study of 257 854 adults in South Korea. *BMJ Open*, 9(3), e025590.



## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1        <https://doi.org/10.1136/bmjopen-2018-025590>
- 2        Kyriazos, T. A. (2018). Applied Psychometrics: Sample Size and Sample Power Considerations in  
3        Factor Analysis (EFA, CFA) and SEM in General. *Psychology*, *09*(08), 2207–2230.  
4        <https://doi.org/10.4236/psych.2018.98126>
- 5        Lejoyeux, M., Avril, M., Richoux, C., Embouazza, H., & Nivoli, F. (2008). Prevalence of exercise  
6        dependence and other behavioral addictions among clients of a Parisian fitness room.  
7        *Comprehensive Psychiatry*, *49*(4), 353–358.  
8        <https://doi.org/https://doi.org/10.1016/j.comppsy.2007.12.005>
- 9        Li, C.-H. (2016). Confirmatory factor analysis with ordinal data: Comparing robust maximum  
10        likelihood and diagonally weighted least squares. *Behavior Research Methods*, *48*(3), 936–  
11        949. <https://doi.org/10.3758/s13428-015-0619-7>
- 12        Lichtenstein, M. B., Griffiths, M. D., Hemmingsen, S. D., & Støving, R. K. (2018). Exercise  
13        addiction in adolescents and emerging adults – Validation of a youth version of the Exercise  
14        Addiction Inventory. *Journal of Behavioral Addictions J Behav Addict*, *7*(1), 117–125.  
15        <https://doi.org/10.1556/2006.7.2018.01>
- 16        Maron, B. J. (2007). Hypertrophic Cardiomyopathy and Other Causes of Sudden Cardiac Death in  
17        Young Competitive Athletes, with Considerations for Preparticipation Screening and Criteria  
18        for Disqualification. *Cardiology Clinics*, *25*(3), 399–414.  
19        <https://doi.org/https://doi.org/10.1016/j.ccl.2007.07.006>
- 20        Mónok, K., Berczik, K., Urbán, R., Szabo, A., Griffiths, M. D., Farkas, J., Magi, A., Eisinger, A.,  
21        Kurimay, T., Kökönyei, G., Kun, B., Paksi, B., & Demetrovics, Z. (2012). Psychometric  
22        properties and concurrent validity of two exercise addiction measures: A population wide  
23        study. *Psychology of Sport and Exercise*, *13*(6), 739–746.  
24        <https://doi.org/https://doi.org/10.1016/j.psychsport.2012.06.003>

## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1 Müller, A., Claes, L., Smits, D., Gefeller, O., Hilbert, A., Herberg, A., Müller, V., Hofmeister, D.,  
2 & de Zwaan, M. (2013). Validation of the German Version of the Exercise Dependence Scale.  
3 *European Journal of Psychological Assessment*, 29(3), 213–219. <https://doi.org/10.1027/1015->  
4 [5759/a000144](https://doi.org/10.1027/1015-5759/a000144)
- 5 Pelliccia, A., Sharma, S., Gati, S., Bäck, M., Börjesson, M., Caselli, S., Collet, J.-P., Corrado, D.,  
6 Drezner, J. A., Halle, M., Hansen, D., Heidbuchel, H., Myers, J., Niebauer, J., Papadakis, M.,  
7 Piepoli, M. F., Prescott, E., Roos-Hesselink, J. W., Graham Stuart, A., ... Wilhelm, M. (2020).  
8 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular  
9 disease. *European Heart Journal*. <https://doi.org/10.1093/eurheartj/ehaa605>
- 10 Pinna, F., Dell’Osso, B., Di Nicola, M., Janiri, L., Altamura, A. C., Carpiniello, B., & Hollander, E.  
11 (2015). Behavioural addictions and the transition from DSM-IV-TR to DSM-5. *Journal of*  
12 *Psychopathology*, 21(4), 380–389.
- 13 Putnick, D. L., & Bornstein, M. H. (2016). Measurement invariance conventions and reporting: The  
14 state of the art and future directions for psychological research. *Developmental Review*, 41,  
15 71–90. <https://doi.org/https://doi.org/10.1016/j.dr.2016.06.004>
- 16 R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for  
17 Statistical Computing. <https://www.r-project.org/>
- 18 Rhudy, J. L., Arnau, R. C., Huber, F. A., Lannon, E. W., Kuhn, B. L., Palit, S., Payne, M. F.,  
19 Sturycz, C. A., Hellman, N., Guereca, Y. M., Toledo, T. A., & Shadlow, J. O. (2020).  
20 Examining Configural, Metric, and Scalar Invariance of the Pain Catastrophizing Scale in  
21 Native American and Non-Hispanic White Adults in the Oklahoma Study of Native American  
22 Pain Risk (OK-SNAP). *Journal of Pain Research*, 13, 961–969.  
23 <https://doi.org/10.2147/JPR.S242126>
- 24 Rizopoulos, D. (2006). *ltm: An R Package for Latent Variable Modeling and Item Response*

## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1        Analysis. *Journal of Statistical Software; Vol 1, Issue 5 (2007)* .
- 2        <https://doi.org/10.18637/jss.v017.i05>
- 3        Robin, X., Turck, N., Hainard, A., Tiberti, N., Lisacek, F., Sanchez, J.-C., & Müller, M. (2011).
- 4        pROC: an open-source package for R and S+ to analyze and compare ROC curves. *BMC*
- 5        *Bioinformatics*, 12(1), 77. <https://doi.org/10.1186/1471-2105-12-77>
- 6        Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Journal of Statistical*
- 7        *Software; Vol 1, Issue 2 (2012)* . <https://doi.org/10.18637/jss.v048.i02>
- 8        Schermelleh-Engel, K., Moosbrugger, H., & Müller, H. (2003). Evaluating the fit of structural
- 9        equation models: Tests of significance and descriptive goodness-of-fit measures. *Methods of*
- 10        *Psychological Research Online*, 8(2), 23–74.
- 11        Sicilia, A., Alías-García, A., Ferriz, R., & Moreno-Murcia, J. A. (2013). Adaptación y validación al
- 12        español del Inventario de Adicción al Ejercicio (EAI). *Psicothema*, 25(3), 377–384.
- 13        Steiger, J. H. (1990). Structural Model Evaluation and Modification: An Interval Estimation
- 14        Approach. *Multivariate Behavioral Research*, 25(2), 173–180.
- 15        [https://doi.org/10.1207/s15327906mbr2502\\_4](https://doi.org/10.1207/s15327906mbr2502_4)
- 16        Stubbs, B., Vancampfort, D., Hallgren, M., Firth, J., Veronese, N., Solmi, M., Brand, S., Cordes, J.,
- 17        Malchow, B., Gerber, M., Schmitt, A., Correll, C. U., De Hert, M., Gaughran, F., Schneider,
- 18        F., Kinnafick, F., Falkai, P., Möller, H.-J., & Kahl, K. G. (2018). EPA guidance on physical
- 19        activity as a treatment for severe mental illness: a meta-review of the evidence and Position
- 20        Statement from the European Psychiatric Association (EPA), supported by the International
- 21        Organization of Physical Therapists in Mental . *European Psychiatry*, 54, 124–144.
- 22        [https://doi.org/DOI: 10.1016/j.eurpsy.2018.07.004](https://doi.org/DOI:10.1016/j.eurpsy.2018.07.004)
- 23        Szabo, A., Griffiths, M. D., de La Vega Marcos, R., Mervó, B., & Demetrovics, Z. (2015).
- 24        Methodological and Conceptual Limitations in Exercise Addiction Research. *The Yale Journal*

## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1        *of Biology and Medicine*, 88(3), 303–308. <https://pubmed.ncbi.nlm.nih.gov/26339214>
- 2 Terry, A., Szabo, A., & Griffiths, M. (2004). The exercise addiction inventory: a new brief  
3 screening tool. *Addiction Research & Theory*, 12(5), 489–499.  
4 <http://10.0.4.56/16066350310001637363>
- 5 Trott, M., Jackson, S. E., Firth, J., Fisher, A., Johnstone, J., Mistry, A., Stubbs, B., & Smith, L.  
6 (2020). Exercise Addiction Prevalence and Correlates in the Absence of Eating Disorder  
7 Symptomology: A Systematic Review and Meta-analysis. *Journal of Addiction Medicine*,  
8 *Publish Ah*.  
9 [https://journals.lww.com/journaladdictionmedicine/Fulltext/9000/Exercise\\_Addiction\\_Prevale](https://journals.lww.com/journaladdictionmedicine/Fulltext/9000/Exercise_Addiction_Prevale)  
10 [nce\\_and\\_Correlates\\_in.99216.aspx](https://journals.lww.com/journaladdictionmedicine/Fulltext/9000/Exercise_Addiction_Prevale_nce_and_Correlates_in.99216.aspx)
- 11 Trott, M., Jackson, S. E., Firth, J., Jacob, L., Grabovac, I., Mistry, A., Stubbs, B., & Smith, L.  
12 (2020). A comparative meta-analysis of the prevalence of exercise addiction in adults with and  
13 without indicated eating disorders. *Eating and Weight Disorders - Studies on Anorexia*,  
14 *Bulimia and Obesity*. <https://doi.org/10.1007/s40519-019-00842-1>
- 15 van de Schoot, R., Lugtig, P., & Hox, J. (2012). A checklist for testing measurement invariance.  
16 *European Journal of Developmental Psychology*, 9(4), 486–492.  
17 <https://doi.org/10.1080/17405629.2012.686740>
- 18 Vandenberg, R. J., & Lance, C. E. (2000). A Review and Synthesis of the Measurement Invariance  
19 Literature: Suggestions, Practices, and Recommendations for Organizational Research.  
20 *Organizational Research Methods*, 3(1), 4–70. <https://doi.org/10.1177/109442810031002>
- 21 Vega, R. de la, Parastatidou, I. S., Ruíz-Barquín, R., & Szabo, A. (2016). Exercise Addiction in  
22 Athletes and Leisure Exercisers: The Moderating Role of Passion. *Journal of Behavioral*  
23 *Addictions J Behav Addict*, 5(2), 325–331. <https://doi.org/10.1556/2006.5.2016.043>
- 24 Vessella, T., Zorzi, A., Merlo, L., Pegoraro, C., Giorgiano, F., Trevisanato, M., Viel, M.,

## EXERCISE ADDICTION AND COMPETITION LEVEL

- 1 Formentini, P., Corrado, D., & Sarto, P. (2020). The Italian preparticipation evaluation  
2 programme: diagnostic yield, rate of disqualification and cost analysis. *British Journal of*  
3 *Sports Medicine*, 54(4), 231 LP – 237. <https://doi.org/10.1136/bjsports-2018-100293>
- 4 Vina, J., Sanchis-Gomar, F., Martinez-Bello, V., & Gomez-Cabrera, M. C. (2012). Exercise acts as  
5 a drug; the pharmacological benefits of exercise. *British Journal of Pharmacology*, 167(1), 1–  
6 12. <https://doi.org/10.1111/j.1476-5381.2012.01970.x>
- 7 Weston, R., & Gore, P. A. (2006). A Brief Guide to Structural Equation Modeling. *The Counseling*  
8 *Psychologist*, 34(5), 719–751. <https://doi.org/10.1177/0011000006286345>
- 9 Wouthuyzen-Bakker, M., & van Assen, S. (2015). Exercise-induced anaemia: a forgotten cause of  
10 iron deficiency anaemia in young adults. *British Journal of General Practice*, 65(634), 268 LP  
11 – 269. <https://doi.org/10.3399/bjgp15X685069>
- 12 Zorzi, A., Cardaioli, F., Cipriani, A., D'Ascenzi, F., Corrado, D., & Solmi, M. (2020). Correlation  
13 between exercise addiction and attitude of competitive athletes towards cardiac  
14 preparticipation evaluation. *Journal of Cardiovascular Medicine (Hagerstown, Md.)*.  
15 <https://doi.org/10.2459/jcm.0000000000000997>

## EXERCISE ADDICTION AND COMPETITION LEVEL

1

**Tables**2 **Table 1**3 *Descriptives of the sample*

4

| Variable                                   | EDS-R (M; SD) |                      |                  |                    |                     |                  |                    |                     |                    | EAI (M; SD)         |
|--|---------------|----------------------|------------------|--------------------|---------------------|------------------|--------------------|---------------------|--------------------|---------------------|
|  | %             | Total                | Withdrawal       | Continuity         | Tolerance           | Control          | Reduction          | Time                | Intention          | Total               |
| <b>Gender</b>                              |               |                      |                  |                    |                     |                  |                    |                     |                    |                     |
| Missing                                    | 0             |                      |                  |                    |                     |                  |                    |                     |                    |                     |
| Females <sup>a</sup>                       | 23.83         | 65.5<br>(15.28)      | 11.93<br>(4.3)   | 8.77<br>(4.1)      | 9.76<br>(3.72)      | 7.53<br>(3.72)   | 7.57<br>(2.65)     | 12.83<br>(3.01)     | 7.1<br>(3.28)      | 19.58<br>(4.41)     |
| Males                                      | 76.17         | 62.15<br>(14.16)**   | 11.16<br>(4.39)* | 7.85<br>(3.99)**   | 9.58<br>(3.48)      | 7.03<br>(3.39)   | 7.3<br>(2.66)      | 12<br>(3.01)**      | 7.22<br>(3.11)     | 18.39<br>(4.63)**   |
| <b>Level of practiced sport</b>            |               |                      |                  |                    |                     |                  |                    |                     |                    |                     |
| Missing                                    | 3.59          | 63.74<br>(17.62)     | 11.83<br>(5.02)  | 8.56<br>(3.95)     | 9.77<br>(3.41)      | 7.31<br>(3.13)   | 7.44<br>(2.45)     | 11.8<br>(3.8)       | 7.03<br>(3.49)     | 17.23<br>(5.06)     |
| Amateur <sup>a</sup>                       | 26.23         | 60.55<br>(14.8)      | 11.6<br>(4.34)   | 7.5<br>(3.86)      | 9.11<br>(3.6)       | 6.71<br>(3.28)   | 7.14<br>(2.7)      | 11.26<br>(3.04)     | 7.22<br>(3.22)     | 17.78<br>(4.52)     |
| Local                                      | 18.65         | 61.68<br>(13.66)     | 11.85<br>(4.17)  | 8.53<br>(4.02)*    | 8.87<br>(3.22)      | 6.76<br>(3.22)   | 7.22<br>(2.52)     | 11.57<br>(3.1)      | 6.89<br>(3.18)     | 18.47<br>(4.7)      |
| Regional                                   | 19.98         | 62.45<br>(12.16)     | 11.31<br>(4.49)  | 7.96<br>(3.93)     | 9.32<br>(3.4)       | 6.94<br>(3.18)   | 7.66<br>(2.66)     | 12.39<br>(2.65)**** | 6.86<br>(2.83)     | 19.17<br>(4.46)**   |
| National                                   | 24.49         | 64<br>(14.96)*       | 10.96<br>(4.34)  | 8.13<br>(4.1)      | 9.92<br>(3.39)*     | 7.56<br>(3.78)*  | 7.39<br>(2.69)     | 12.73<br>(2.81)**** | 7.3<br>(3.2)       | 18.65<br>(4.46)     |
| International                              | 10.66         | 69.51<br>(15.37)**** | 10.69<br>(4.44)  | 8.63<br>(4.45)*    | 12.05<br>(3.49)**** | 8.3<br>(4)****   | 7.54<br>(2.77)     | 14.19<br>(2.62)**** | 8.11<br>(3.19)*    | 20.87<br>(4.36)**** |
| <b>Answer to Q1</b>                        |               |                      |                  |                    |                     |                  |                    |                     |                    |                     |
| Missing                                    | 0             |                      |                  |                    |                     |                  |                    |                     |                    |                     |
| “I would stop exercising” (0) <sup>a</sup> | 64.39         | 62.31<br>(14.1)      | 11.32<br>(4.36)  | 7.95<br>(4)        | 9.53<br>(3.49)      | 7.02<br>(3.42)   | 7.27<br>(2.56)     | 12.14<br>(3.02)     | 7.07<br>(3.08)     | 18.55<br>(4.57)     |
| “I would not stop exercising” (1)          | 35.61         | 75.26<br>(16.6)****  | 11.96<br>(4.74)  | 10.47<br>(3.8)**** | 11.27<br>(4)***     | 9.4<br>(3.9)**** | 9.27<br>(3.62)**** | 13.43<br>(3.15)***  | 9.46<br>(3.63)**** | 21.19<br>(4.7)****  |

## EXERCISE ADDICTION AND COMPETITION LEVEL

**Answer to Q2**

|   |       |                                  |                              |                                |                           |                                |                               |                                 |                                |                                 |
|---|-------|----------------------------------|------------------------------|--------------------------------|---------------------------|--------------------------------|-------------------------------|---------------------------------|--------------------------------|---------------------------------|
| Missing                                       | 0     |                                  |                              |                                |                           |                                |                               |                                 |                                |                                 |
| “I would stop<br>exercising” (0) <sup>a</sup> | 65.88 | 60.83<br>(13.59)                 | 11.1<br>(4.37)               | 7.64<br>(3.87)                 | 9.42<br>(3.5)             | 6.84<br>(3.36)                 | 6.99<br>(2.52)                | 11.9<br>(2.99)                  | 6.94<br>(3.04)                 | 18.21<br>(4.62)                 |
| “I would not stop<br>exercising” (1)          | 34.12 | 67.14<br>(15.34) <sup>****</sup> | 11.83<br>(4.35) <sup>*</sup> | 8.93<br>(4.22) <sup>****</sup> | 10<br>(3.57) <sup>*</sup> | 7.75<br>(3.63) <sup>****</sup> | 8.1<br>(2.78) <sup>****</sup> | 12.81<br>(3.03) <sup>****</sup> | 7.69<br>(3.32) <sup>****</sup> | 19.61<br>(4.45) <sup>****</sup> |

1

2

Note. EAI, exercise addiction inventory; EDS-R, exercise dependence scale, 21 items. <sup>a</sup>= **this level is assumed as the reference one for analyses testing mean differences among groups.** \*= p< .05; \*\*= p< .01; \*\*\*= p< .001; \*\*\*\*= p< .0001. P values are adjusted by applying a False Discovery Rate correction.

3

4





## EXERCISE ADDICTION AND COMPETITION LEVEL

|   |  |     |
|---|--|-----|
| 2 | Conflicts have arisen between me and my family<br>and/or my partner<br>about the amount of exercise                        | .48 |
| 3 | I use exercise as a way of changing my mood  | .63 |
| 4 | Over time I have increased the amount of exercise<br>I do in a day   | .65 |
| 5 | If I have to miss an exercise session, I feel moody<br>and irritable   | .69 |
| 6 | If I cut down the amount of exercise I do, and then<br>start again, I always<br>end up exercising as often as I did before | .33 |

---

1

2 Note. EAI: Exercise Addiction Inventory, 6 items; EDS-R: Exercise Dependence Scale- Revised, 21 items.

## EXERCISE ADDICTION AND COMPETITION LEVEL

**Table 3***Multigroup confirmatory factor analyses for EDS21 and EAI across gender and competition type*

| #                         | Model                | $\chi^2$ | df   | SRMR | CFI | $\Delta$ CFI | RMSEA | $\Delta$ RMSEA |
|---------------------------|----------------------|----------|------|------|-----|--------------|-------|----------------|
| <i>EAI: Competition</i>   |                      |          |      |      |     |              |       |                |
| 1                         | Omnibus              | 29.33    | 9    | .03  | .98 |              | .05   |                |
| 2                         | Amateur              | 29.58    | 9    | .05  | .95 |              | .09   |                |
| 3                         | Local                | 11.57    | 9    | .04  | .99 |              | .04   |                |
| 4                         | Regional             | 18.80    | 9    | .05  | .96 |              | .08   |                |
| 5                         | National             | 17.67    | 9    | .04  | .97 |              | .06   |                |
| 6                         | International        | 12.09    | 9    | .06  | .97 |              | .06   |                |
| 7                         | Conf. inv.           | 55.71    | 45   | .05  | .99 |              | .04   |                |
| 8                         | Metric i. (7 vs 8)   | 79.55    | 65   | .06  | .99 | < .01        | .03   | < .01          |
| 9                         | Scalar i. (8 vs 9)   | 202.75   | 133  | .06  | .96 | .03          | .05   | .02            |
| 10                        | Par.s.i. (8 vs 10)   | 155.66   | 125  | .06  | .98 | < .01        | .04   | < .01          |
| 11                        | Resid.i. (9 vs 11)   | 202.75   | 133  | .06  | .96 | 0            | .05   | 0              |
| 12                        | Par.r.i. (10 vs 12)  | 155.66   | 125  | .06  | .98 | 0            | .04   | 0              |
| <i>EAI: Gender</i>        |                      |          |      |      |     |              |       |                |
| 13                        | Female               | 6.79     | 9    | .03  | 1   |              | 0     |                |
| 14                        | Male                 | 28       | 9    | .03  | .98 |              | .05   |                |
| 15                        | Conf. inv.           | 33.99    | 18   | .03  | .99 |              | .04   |                |
| 16                        | Metric i. (15 vs 16) | 31.90    | 23   | .03  | .99 | < .01        | .04   | < .01          |
| 17                        | Scalar i. (16 vs 17) | 55.53    | 40   | .03  | .99 | < .01        | .03   | < .01          |
| 18                        | Resid.i. (17 vs 18)  | 55.53    | 40   | .03  | .99 | 0            | .03   | 0              |
| <i>EDS-R: Competition</i> |                      |          |      |      |     |              |       |                |
| 19                        | Omnibus              | 805.06   | 168  | .05  | .97 |              | .06   |                |
| 20                        | Amateur              | 365.88   | 168  | .06  | .97 |              | .07   |                |
| 21                        | Local                | 284.42   | 168  | .07  | .97 |              | .06   |                |
| 22                        | Regional             | 287.32   | 168  | .07  | .97 |              | .06   |                |
| 23                        | National             | 311.01   | 168  | .06  | .98 |              | .06   |                |
| 24                        | International        | 264.64   | 168  | .08  | .97 |              | .07   |                |
| 25                        | Conf. inv.           | 1000.12  | 840  | .06  | .99 |              | .03   |                |
| 26                        | Metric i. (25 vs 26) | 1203.28  | 896  | .07  | .99 | < .01        | .04   | .01            |
| 27                        | Scalar i. (26 vs 27) | 1346.17  | 1156 | .07  | .99 | < .01        | .03   | .01            |
| 28                        | Resid.i. (27 vs 28)  | 1354     | 1172 | .07  | .99 | 0            | .03   | < .01          |
| <i>EDS-R: Gender</i>      |                      |          |      |      |     |              |       |                |
| 29                        | Female               | 348.47   | 168  | .06  | .97 |              | .07   |                |
| 30                        | Male                 | 634.08   | 168  | .05  | .98 |              | .06   |                |
| 31                        | Conf. inv.           | 957.72   | 336  | .05  | .98 |              | .06   |                |
| 32                        | Metric i. (31 vs 32) | 908.68   | 350  | .05  | .98 | < .01        | .06   | < .01          |
| 33                        | Scalar i. (32 vs 33) | 1010.32  | 424  | .05  | .98 | < .01        | .05   | < .01          |
| 34                        | Resid.i. (33 vs 34)  | 1010.70  | 424  | .05  | .98 | 0            | .05   | 0              |

Note. EAI: Exercise Addiction Inventory, 6 items; EDS-R: Exercise Dependence Scale- Revised, 21 items. Conf. Inv.: Configural invariance. Metric i. : Metric invariance. Scalar i. : Scalar invariance. Resid i. : Residual invariance. Par.s.i: partial scalar invariance Par.r.i.: partial residual invariance.

## EXERCISE ADDICTION AND COMPETITION LEVEL

1 **Table 4**

2

3 *Multiple regression results.*

4

| <b>Predictors</b>              | <b>Response variable: EAI</b> |          |                            | <b>Response variable: EDS-R</b> |          |                            |
|--------------------------------|-------------------------------|----------|----------------------------|---------------------------------|----------|----------------------------|
|                                | <b><math>\beta</math></b>     | <b>p</b> | <b><math>\eta^2</math></b> | <b><math>\beta</math></b>       | <b>p</b> | <b><math>\eta^2</math></b> |
| Gender (Male)                  | -1.06                         | .12      | .01                        | -4.98                           | .02      | .01                        |
| Years of practice              | -0.02                         | .3       | .01                        | -0.03                           | .73      | < .01                      |
| Average practice time          | 0.11                          | < .01    | .03                        | 0.34                            | < .01    | .05                        |
| Gender X Years of practice     | -0.02                         | .58      | < .01                      | 0.01                            | .98      | < .01                      |
| Gender X Average practice time | 0.02                          | .58      | < .01                      | 0.14                            | .2       | < .01                      |

5 Note: p: p value;  $\eta^2$ : partial eta squared

## EXERCISE ADDICTION AND COMPETITION LEVEL

**Table 5**

*Results of multiple logistic regressions of EAI and EDS-R on question evaluating the adherence to medical prescriptions*

| Predictors                                       | Response variable: Q2 |            |                      |
|--|-----------------------|------------|----------------------|
|  | $\beta$               | p value    | OR(CI)               |
| <b>EAI</b><br>(Reference: Addicted)              | 0.64                  | .19        | 1.9(.72-5.05)        |
| <b>Competition level</b><br>(Reference: Amateur) |                       |            |                      |
| Local  | -0.48                 | <b>.03</b> | <b>.62(.4-.96)</b>   |
| Regional   | -0.2                  | .34        | .82(.53-1.24)        |
| National   | -0.06                 | .77        | .94(.64-1.39)        |
| International                                    | 0.28                  | .29        | 1.32(.79-2.2)        |
| <b>EDS-R</b><br>(Reference: Addicted)            | 1.07                  | <b>.04</b> | <b>2.9(1.01-8.9)</b> |
| <b>Competition level</b><br>(Reference: Amateur) |                       |            |                      |
| Local  | -0.35                 | .1         | .7(.45-.107)         |
| Regional   | -0.12                 | .53        | .88(.58-1.32)        |
| National   | -0.01                 | .98        | .99(.67-1.46)        |
| International                                    | 0.13                  | .6         | 1.14(.68-1.9)        |

Notes. Q2: "A medical investigation revealed that you have a critical cardiac anomaly that could cause a deadly heart attack. Unfortunately, it is necessary to permanently stop exercise. How would you react?". Statistically significant effects are reported in bold.


EXERCISE ADDICTION AND COMPETITION LEVEL

**Figures**

1  
2  
3  
4  
5  
6

**Figure 1**

*ROC curves for EAI (upper panel) and EDS-R (lower panel).*



7