


pca

europaean journal of
postclassicalarchaeologies

Volume 14
2024



pca

european journal of
postclassicalarchaeologies

volume 14/2024

SAP Società Archeologica s.r.l.

Mantova 2024

EDITORS

Alexandra Chavarría (chief editor)

Gian Pietro Brogiolo (executive editor)

EDITORIAL BOARD

Paul Arthur (Università del Salento)

Alicia Castillo Mena (Universidad Complutense de Madrid)

Margarita Díaz-Andreu (ICREA - Universitat de Barcelona)

Enrico Cirelli (Alma Mater Studiorum - Università di Bologna)

José M. Martín Civantos (Universidad de Granada)

Caterina Giostra (Università Cattolica del Sacro Cuore, Milano)

Matthew H. Johnson (Northwestern University of Chicago)

Vasco La Salvia (Università degli Studi G. D'Annunzio di Chieti e Pescara)

Bastien Lefebvre (Université Toulouse - Jean Jaurès)

Alberto León (Universidad de Córdoba)

Tamara Lewit (University of Melbourne)

Yuri Marano (Scuola Archeologica Italiana di Atene)

Federico Marazzi (Università degli Studi Suor Orsola Benincasa di Napoli)

Maurizio Marinato (Università degli Studi di Padova)

Johannes Preiser-Kapeller (Österreichische Akademie der Wissenschaften)

Andrew Reynolds (University College London)

Mauro Rottoli (Laboratorio di archeobiologia dei Musei Civici di Como)

Colin Rynne (University College Cork)

Marco Valenti (Università degli Studi di Siena)

Giuliano Volpe (Università degli Studi di Foggia)

Post-Classical Archaeologies (PCA) is an independent, international, peer-reviewed journal devoted to the communication of post-classical research. PCA publishes a variety of manuscript types, including original research, discussions and review articles. Topics of interest include all subjects that relate to the science and practice of archaeology, particularly multidisciplinary research which use specialist methodologies, such as zooarchaeology, paleobotany, archaeometallurgy, archaeometry, spatial analysis, as well as other experimental methodologies applied to the archaeology of post-classical Europe.

Submission of a manuscript implies that the work has not been published before, that it is not under consideration for publication elsewhere and that it has been approved by all co-authors. Authors must clear reproduction rights for any photos or illustration, credited to a third party that they wish to use (including content found on the Internet). For more information about **ethics** (including plagiarism), copyright practices and guidelines please visit the website www.postclassical.it.

PCA is published once a year in May. Manuscripts should be submitted to editor@postclassical.it in accordance to the guidelines for contributors in the webpage <http://www.postclassical.it>.

Post-Classical Archaeologies' manuscript **review process** is rigorous and is intended to identify the strengths and weaknesses in each submitted manuscript, to determine which manuscripts are suitable for publication, and to work with the authors to improve their manuscript prior to publication.

This journal has the option to publish in **open access**. For more information on our open access policy please visit the website www.postclassical.it.

How to **quote**: please use "PCA" as abbreviation and "European Journal of Post-Classical Archaeologies" as full title.

Cover image: the production of past lives layers at present society (by Gonzalo Ruiz Zapatero).

"Post-Classical Archaeologies" is indexed in Scopus and classified as Q3 by the Scimago Journal Rank (2022). It was approved on 2015-05-13 according to ERIH PLUS criteria for inclusion and indexed in Carhus+2018. Classified A by ANVUR (Agenzia Nazionale di Valutazione del sistema Universitario e della Ricerca).

DESIGN:

Paolo Vedovetto

PUBLISHER:

SAP Società Archeologica s.r.l.

Strada Fienili 39/a, 46020 Quingentole, Mantua, Italy

www.saplibri.it

Authorised by Mantua court no. 4/2011 of April 8, 2011

For subscription and all other information visit the website www.postclassical.it.

Volume funded by the
University of Padova
Department of Cultural Heritage



CONTENTS PAGES

EDITORIAL

5

RESEARCH - WHY AND FOR WHOM ARE WE DOING ARCHAEOLOGY TODAY?

G. Ruiz Zapatero Archaeology today: people, knowledge and dissemination

7

C. Lewis Who cares? Why and for whom are we now doing archaeology in the UK?

31

M. Nevell The public benefit of archaeology during an era of financial austerity: three British case studies from the commercial, university, and museum sectors

55

J.-P. Demoule French archaeology: for whom, but also against whom?

67

G.P. Brogiolo Archeologia e 'coscienze generazionali'. Quali prospettive in Italia?

79

BEYOND THE THEME

C. Croci Da Giove Capitolino a San Pietro: Costantino e il tramonto della Roma antica

89

H. Tůmová, E. Cirelli Ravenna surrounded by waters. Landscape changes and urban transformations in Late Antiquity

121

L. Pugliese Indagini GPR e LiDAR sulla rocca di Garda. Una finestra sull'abitato del *castrum* tardoantico e altomedievale

145

G.P. Brogiolo, F. Malaspina Storie di un territorio di frontiera tra i fiumi Serio e Oglio (VI-XIII secolo)

163

M. Fecchio, M. Marinato Change in diet or visibility problem? Observations on the marine isotopic values of early medieval populations in coastal Croatia 195

G. Vannini The urban fabric as a source for an archaeological history of the topography of medieval Florence 217

DOSSIER - CITIZEN SCIENCE IN ARCHAEOLOGY

F. Pinna, M. Sanna Montanelli Citizen Archaeology. Un contributo alla costruzione di una archeologia pubblica come "scienza normale" 237

F. Frandi, G. Pastura Cittadinanza dell'archeologia e infrastrutture ferroviarie. Sostenibilità sociale di un rapporto necessario 257

M. Paladini L'utilizzo delle interviste qualitative non strutturate nell'archeologia partecipata: il caso studio di Lio Piccolo (laguna nord di Venezia) 267

RETROSPECT

C. Rynne The development of post-medieval and industrial archaeology in Ireland 287

PROJECT

A. Arrighetti, M. Repole Expeditious archaeoseismological analysis of a medieval town centre. The case of Siena and the PROTECT project 302

REVIEWS

Slavko Ciglènečki, *Between Ravenna and Constantinople: Rethinking Late Antique Settlement Patterns* - by **A. Chavarría Arnau**

Rafael Hidalgo Prieto, Inmaculada Carrasco Gómez, Adalberto Ottati (eds), *Palatia. Ville e palazzi imperiali di época romana (secoli I-V d.C.)/Imperial villas and palaces of the Roman period (1st-5th centuries AD)* - by **J.A. Pintado**

Isabel María Sánchez, Yolanda Peña Cervantes (eds), *Fundus Ecclesiae. Evidencias materiales de las actividades económicas de la iglesia en Hispania durante la Antigüedad Tardía y Alta Edad Media* - by **V. Amorós-Ruiz**

Jana Kopáčková, *Olevm et Vinvm Adriaticvm. Production of Olive Oil and Wine in Roman Histria and Dalmatia* - by **T. Lewit**

Beth Munro, *Recycling the Roman Villa. Material Salvage and the Medieval Circular Economy* - by **A. Chavarría Arnau**

Cristina Godoy Fernández, Andreu Muñoz Melgar (eds), *El monument tardoromà de Centelles. Dades, context, propostes* - by **G.P. Brogiolo**

Roberta Conversi (ed), *Il sito della Piana di San Martino a Pianello Val Tidone. Dal castrum tardoantico all'abbandono nel XVIII secolo. Scavi e ricerche 2018-2021* - by **G.P. Brogiolo**

Riccardo Santangeli Valenzani (ed), *Roma Altomedievale. Paesaggio urbano, società e cultura (secoli V-X)* - by **A. Chavarría Arnau**

Mirko Fecchio, Maurizio Marinato*

Change in diet or visibility problem? Observations on the marine isotopic values of early medieval populations in coastal Croatia

1. Introduction

The island of Rab is in the Kvarner archipelago of the Croatian littoral-mountain region. It has been the subject of archaeological investigations for over a decade by the Department of Cultural Heritage of the University of Padua and the Faculty of Humanities and Social Sciences of the University of Zagreb (fig. 1). The archaeological investigations have covered numerous sites both in the countryside (churches, monasteries, castles, and villas, as well as landscape analysis) (Brogiolo *et al.* 2013; Jurković *et al.* 2012; Brogiolo *et al.* 2017) and within the urban landscape of ancient Rab, a city with protohistoric origins that saw Roman and Late Antique-Byzantine phases, followed by significant Late Medieval reorganization shaping the current architectural structure (Brogiolo *et al.* 2023).

Between 2015 and 2018, archaeological campaigns were carried out in Dinka Dokule Street (fig. 1), documenting the evolution of this urban area, characterized initially by a Late Antique funerary zone, followed by a fortification from the late 6th to early 7th centuries that overlays part of the tombs (some in sarcophagi), and a series of subsequent metallurgical activities (Brogiolo *et al.* 2023).

This paper focuses on the faunal remains uncovered at the construction site for a city wall (late 6th-early 7th century AD) and in a building adjacent to the wall, dated between the 7th and 8th centuries AD; and three burials, one of which is multiple, from the initial phase. The data have been supplemented with analyses from a burial, likely dating to the early medieval period, contemporary with the

* University of Padova, Department of Cultural Heritage, mirko.fecchio@phd.unipd.it, maurizio.marinato@unipd.it.




Fig. 1. Island of Rab and the excavations conducted at Dinka Dokule and St. Lawrence in Banjol.

second phase of the church of Saint Lawrence, which is three kilometres far from the Dinka Dokule site, discovered during the 2015 and 2016 excavations (Brogiolo *et al.* 2017).

This research aims to reveal, through zooarchaeological study and stable isotope analyses ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$), the aspects that characterized animal management and the diet of the early medieval population of an *emporion* of the Byzantine Empire.

2. Historical and environmental background

The history of the island of Rab dates back to the Middle Paleolithic. During the Bronze Age, several hillforts were strategically built for territorial control. Evidence of burials at Gromačica and Pečina confirms the presence of these fortified villages until the Iron Age. By the end of the Bronze Age, a tribe unified some of the hillforts, forming larger centers such as Rab and Kastelina, and building new ones. Archaeological findings indicate commercial exchanges between Liburnia, Greece, and Italy (Batovič 1987, pp. 147-169). According to Strabo, the Liburni, part of the Illyrian population, arrived on the northern Adri-

atic coasts after being expelled from Corcyra by Corinthian colonists. Pliny the Elder, in his *Naturalis Historia*, described the Liburni as savage tribes dedicated to piracy, who were defeated by Consul *Gaius Sempronius Tuditanus* in 129 BC (Wilkes 1977, p. 160). Pliny also mentioned several cities in the Gulf of Kvarner, including *Arba*, which received *jus italicum* (G. Plinio S., O.C., L. III, 140), becoming fortified military ports that controlled trade across the North of the Adriatic Sea.

With the advent of the Imperial age, the island of Rab and the entire regions of Liburnia and Dalmatia became the Province *Superior Illyricum*, renamed Dalmatia under Emperor *Titus Flavius Domitian* in 96 AD, and linked to Salona (Tomaz 2003, pp. 103-104). During the Gothic War (535-553 AD), Justinian reconquered the islands of the Kvarner, placing them under the Eastern Roman Empire. In the 7th century, while much of the Italian peninsula fell under Lombard control, Dalmatia faced Slavic invasions but retained control of the coastal areas, including Rab, which remained strongly Latin. The Dalmatian islands remained under the exarchate of Ravenna until 726 AD, then became connected to Constantinople after Ravenna fell to the Lombards in 751 AD (Tomaz 2003, pp. 240-241). In 812, the *Pax Nicephori* established imperial boundaries, placing the Dalmatian coasts and islands under Eastern dominion. Constantine VII's *De administrando imperio* (950) referred to the citizens of various Adriatic cities, including *Arba*, as Romans.

Information on the diet of Croatian populations in the Early Middle Ages is scarce, mainly due to the limited presence of written sources on the subject. Most of the available data therefore comes from biochemical analyses conducted on skeletal remains from the period, which will be examined for comparison, or from recent paleoenvironmental studies carried out on specific case studies within the archipelago.

Through the analysis of lake sediments from Vrana and Osor, the IATECA research program (Interdisciplinary Approach to the Territorial Evolution of the Kvarner Archipelago) has highlighted the evolution of ancient landscapes and socio-cultural changes on the island of Cres, located a few kilometers from Rab. Despite the chronological imprecision of sedimentary deposits over short time scales, the dating model for Vrana is more reliable than that Lake Osor's. This has allowed the reconstruction of landscape exploitation in the region from the beginning of Roman rule to the early Middle Ages.

The study on the occupation of Kvarner, focused on the southern part of Cres, highlights two significant periods. The first is characterized by the establishment of Roman power starting from the 1st century BC, followed by the territorial reorganization of northern Liburnia between the late 2nd and 4th centuries AD. During this period, the construction of urban walls and structures in cities such as Krk, Rab, and Osor, as well as the settlement of large maritime and rural estates, can

be observed. Lake sediments supplement our understanding of the region, indicating the introduction of crops such as olives in the 1st century BC, followed by walnuts and deforestation in the 2nd century AD. The 3rd century saw the introduction of cereals, vineyards, and chestnuts, reflecting an agricultural change related to regional reorganization.

In the 4th century, there was an increase in livestock around Lake Vrana, possibly linked to wool production for the Roman army. During the Byzantine period, cereal cultivation intensified, probably in response to the crisis of Mediterranean trade links from the 7th century. These transformations reflect an adaptation to socio-geopolitical and climatic dynamics, showing how the Kvarner islands were integral parts of Mediterranean exchange flows despite their apparent isolation. This study has underscored the importance of agricultural and pastoral practices in the adaptation of local communities to historical and environmental changes (Čaušević-Bully *et al.* 2021).

According to archaeobotanical studies, during the Roman and late antique periods, the diet was enriched with varieties such as spelt, barley, emmer, millet, grapes, olives, legumes, cabbage, and carrots. The cities of Naron and Vis and the areas around the city of Zara were renowned, even by the author Strabo, to produce wine and olive oil (Škegro 2006, pp.157-159; Wilkes 1992, p. 220). Although agriculture dominated, historical sources and individual archaeozoological studies in southern Dalmatia testify to the consumption of fish and animals hunted, such as deer, hares, and wild birds (Wilkes 1992, p. 222; Schwartz 1996). In Early Medieval Croatia, evidenced by the material culture of funerary contexts, a mixed agricultural model with cultivation and livestock farming of cattle, pigs, sheep, goats, and poultry emerged (Evans 1989, p. 206). The economy was also influenced by trade along the Adriatic coast to Venice, as evidenced by commercial artefacts and imported objects (Piteša 2006).

These findings outline a continuity in the management of agricultural and food resources across historical periods, showing adaptations to available resources and the commercial dynamics of the time. The few data available in the literature on the exploitation of plant and animal resources find only some common points with what emerged from the present study.

3. Zooarchaeological evidence

The analysed archaeozoological record comprises 1403 bone fragments, of which over 72% were taxonomically identified thanks to the excellent preservation of the remains. Approximately 380 fragments, mainly ribs, and skulls of small to medium-sized herbivores, could not be identified to the species level.



Fig. 2. Percentage distribution of the Number of Remains (NR) for each animal taxa documented in the investigated faunal sample.

The analysis¹ has provided a comprehensive overview of animal exploitation between the 6th and 8th centuries AD, including both major domestic species (sheep and goats, pigs, and cattle) and fish resources, which were an essential food source for the island's inhabitants. The high number of butchery marks on the bones, along with the study of the age at slaughter and the sex of the individuals, has led to a deeper understanding of meat consumption by the inhabitants of ancient *Arba*.

The count of remains (NR) and the determination of the Minimum Number of Individuals (MNI) revealed that caprines (*Ovis aries* and *Capra hircus*) were the most prevalent species at the site, followed by pigs (*Sus domesticus*) and domestic chickens (*Gallus gallus*) (tab. 1). Large herbivores such as horses and cattle were much less common, with cattle represented by only 50 bone elements. Although present in minimal percentages, the faunal assemblage also includes wild mammals (*Cervus elaphus* and *Lepus europaeus*), wild birds, and tortoises, with uncertainty remaining as to whether they are *Testudo hermanni* or *Testudo graeca*, both terrestrial species. In contrast, fish resources were more frequently identified, determined to the species level and, in some sporadic cases, to the genus level (fig. 2).

4. Rearing animals on an island: the recovery of primary and secondary products

The study of slaughter age (fig. 3) and the sex determination of the main domestic species, coupled with a meticulous taphonomic analysis aimed at record-

¹ For the methodological protocol applied to the study of animal bones, reference is made to FECCHIO, TECCHIATI 2021, pp. 203-225, and to the zooarchaeological manuals by DE GROSSI MAZZORIN 2008 and GIFFORD-GONZALEZ 2018.

	<i>Bos taurus</i>	<i>Ovis aries</i>	<i>Capra hircus</i>	<i>Ovis vel Capra</i>	<i>Sus domesticus</i>	<i>Equus caballus</i>	<i>Gallus gallus</i>	<i>Cervus elaphus</i>	<i>Lepus europaeus</i>	Galliformes	Testudo	Small to medium sized herbivores	Large herbivores	ND
<i>Calvaria</i>	-	6	6	-	-	-	-	5	-	-	-	-	-	
<i>Cranium</i>	1	-	-	12	15	-	-	-	-	-	-	12	2	17
<i>Maxilla</i>	-	-	-	3	5	-	-	-	-	-	-	-	-	-
<i>Dentes</i>	7	-	-	41	24	1	-	1	-	-	-	-	-	1
<i>Mandibula</i>	2	6	-	23	4	-	-	-	-	-	-	-	-	-
<i>Costae</i>	13	-	-	19	10	-	1	-	-	-	-	66	16	61
<i>Vertebrae</i>	5	-	3	67	32	-	-	-	-	-	-	10	2	8
<i>Sternum</i>	-	-	-	-	3	-	4	-	-	-	-	-	-	-
<i>Coracoide</i>	-	-	-	-	-	-	3	-	-	-	-	-	-	-
<i>Furcula</i>	-	-	-	-	-	-	1	-	-	-	-	-	-	-
<i>Scapula</i>	-	5	6	12	9	-	4	-	-	-	-	-	-	-
<i>Humerus</i>	1	8	6	17	16	-	8	1	-	-	1	-	-	-
<i>Radius</i>	4	7	4	23	5	-	2	-	-	-	-	1	-	-
<i>Ulna</i>	1	1	1	9	6	-	7	-	1	-	-	-	-	-
<i>Carpalia</i>	1	-	-	2	1	-	-	-	-	-	-	-	2	-
<i>Metacarpus</i>	1	7	6	3	5	-	-	-	-	-	-	-	-	-
<i>Carpometacarpus</i>	-	-	-	-	-	-	1	-	-	-	-	-	-	-
<i>Pelvis</i>	1	3	4	10	6	1	1	-	-	-	-	1	1	2
<i>Femur</i>	1	3	2	20	6	-	8	-	2	1	-	1	-	3
<i>Tibia</i>	2	1	-	41	11	-	-	-	1	-	-	-	-	-
<i>Tibiotarsus</i>	-	-	-	-	-	-	9	-	-	1	-	-	-	-
<i>Fibula</i>	-	-	-	-	1	-	-	-	-	-	-	-	-	-
<i>Talus</i>	2	3	4	1	5	-	-	-	-	-	-	-	-	-
<i>Os sacrum</i>	1	-	-	-	-	-	-	-	-	-	-	2	-	-
<i>Calcaneus</i>	1	7	1	2	11	-	-	-	-	-	-	-	-	-
<i>Tarsalia</i>	1	-	-	1	3	-	-	-	-	-	-	-	-	-
<i>Tarsometatarsus</i>	-	-	-	-	-	-	26	-	-	-	-	-	-	-
<i>Metatarsus</i>	1	11	3	6	10	1	-	-	-	-	-	-	-	-
<i>Metapodia</i>	-	-	-	17	1	-	-	-	-	-	-	-	-	-
<i>Phalanges</i>	5	12	11	6	20	-	4	1	-	-	-	-	-	-
<i>Diaphyses</i>	-	-	-	-	-	-	-	-	1	1	-	27	8	5
ND	-	-	-	-	-	-	-	-	-	-	-	-	-	134
Total	51	80	57	335	209	3	79	8	5	3	1	120	31	

Tab. 1. Table of terrestrial animal remains from the entire faunal assemblage.




Fig. 3. Age at death for the main domestic species based on the state of eruption, replacement, and dental wear.

ing all butchery marks on the specimens, have allowed for an overall understanding of animal management. Mortality curves for pigs, obtained from the fusion state of epiphyses to the diaphysis of long bones and the degree of dental wear², indicate that they were slaughtered at a young or subadult age when the species reached its maximum meat yield. This, combined with numerous butchery marks along the pelvic and scapular joints, demonstrates that *Sus domesticus* constituted a fundamental protein resource in the daily diet of islanders. Pigs were typically slaughtered around 18-24 months of age; in antiquity, the development and fattening process of pigs were slower, and the peak meat yield relative to the animal maintenance costs occurred later in age compared to modern times (De Grossi Mazzorin, Minniti 2001, pp. 71-72). It can be inferred that the older specimens (5%) were predominantly female and were essential for species preservation on the site. The presence of all anatomical districts suggests that pig farming likely occurred near the site.

For domestic poultry³, unlike the appendicular and axial regions of the skeleton, the cranial region is completely absent, as it was removed during bird slaughter and generally has a fragile structure prone to deterioration under various taphonomic processes. Small butchery traces have been identified near the wings and hind limbs of the bird, areas with less muscle mass, indicating a hypothetical division of the animal before consumption. Considering that chickens

² The methods of PAYNE 1973 and GRANT 1982 were considered solely for comparison with the process currently employed by us to derive age classes based on the wear state of the occlusal surface of teeth.

³ COHEN, SERJEANTSON 1986 was used to distinguish domestic avifauna from wild ones.

are easily raised in urban settings and can provide food resources both through direct slaughter and secondary product recovery, a significant number of bones in the faunal assemblage is expected.

The incidence of cattle in the faunal sample is quite low, with a minimum of two fully adult individuals. Sex determination relied on the dimensions and relative osteometric measurements of a complete metacarpal, suggesting a female specimen⁴. The few butchery remains were found along major joints and on bones with the least amount of meat, such as the astragalus, calcaneus, and metapodials, likely to select large meat portions. It can only be inferred that the few *Bos taurus* specimens were primarily exploited for traction and subsequently slaughtered for primary product recovery⁵.

The use of small domestic ruminants was more intricate: the central role these species played in the island's economy is confirmed by numerous data obtained from dental erosion status and the fusion of epiphyses of long bones. The combination of both methodologies confirmed much more generalized breeding strategies. A high percentage of fully adult or senile specimens (48.7%), associated with milk and wool recovery in the case of sheep, is supplemented by a considerable number of young individuals (32%), many of whom show clear signs of butchery. The mortality curve in this case suggests a clear intention to systematically recover meat from these species.

The good frequency of anatomical parts with various diagnostic elements allowed for the distinction between sheep and goats for some fragments (Boessneck *et al.* 1964, pp. 331-358): 80 cases were attributed to *Ovis aries*, 57 to *Capra hircus*, while the remaining 335 were generic caprine remains. Taphonomic analysis of the bones of these species revealed a clear predominance of butchery cuts on goat remains relative to the number of skeletal elements. This data supports the hypothesis that, unlike *Ovis aries*, there was greater interest in meat recovery for *Capra hircus*, a fact also evidenced at other early medieval coastal sites in the Adriatic Sea (De Grossi Mazzorin, De Venuto 2009, p. 24; Powell 2004, pp. 307-313).

5. Hunting and fishing

Less than 2% of the identified remains are attributable to wild mammals and avifauna, a figure that inevitably suggests that hunting did not play an essential

⁴ The data is further confirmed by the application of the Nobis index (NOBIS 1954, pp. 155-194) ($Bp100/GL, 51.3100/192.4 = 26.6$ cm).

⁵ The paleopathological study revealed evidence of animal exploitation as a workforce, indicated by a case of extensive exostosis on a third anterior phalanx and by two instances of eburnation, respectively observed in a bovine astragalus and calcaneus (BAKER, BROTHWELL 1980).

role in the economy and diet of the island's inhabitants. This evidence could be due to intensive agriculture developed on the island, which left little room for hunting activities. Unfortunately, in the absence of studies regarding the ancient flora of the island, it is impossible to hypothesize how much territory was dedicated to agriculture and how much was left fallow.

The intensive sieving of sediment during excavation allowed the inclusion of 82 fish bone fragments in the study (tab. 2). Apparently, in the protein diet of the inhabitants of *Arba*, sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) were present, although there are also attestations of vertebrae and cranial elements of European conger, red porgy, scorpionfish, or common smooth hound, significantly increasing the variability of resources recovered from the sea⁶ (fig. 4). This data contradicts the results obtained from the isotopic signal of the examined burials. Such fish species, with demersal habits, prefer coastal habitats and only occasionally migrate to deeper waters. Therefore, fishing likely concentrated, mainly during the summer months, in the shallow waters of the bay upon which the city of *Arba* overlooked. A similar hypothesis can be advanced for the 107 remains of bivalves and gastropods related to marine malacofauna,

	<i>Conger conger</i>	<i>Dicentrarchus labrax</i>	<i>Engraulis encrasicolus</i>	<i>Mustelus mustelus</i>	<i>Pagrus pagrus</i>	<i>Scorpaenidae</i>	<i>Sparidae</i>	<i>Sparus aurata</i>	<i>Pisces</i>	<i>Caridiidae</i>	<i>Cerithidae</i>	<i>Conus mediterraneus</i>	<i>Gibbula adriatica</i>	<i>Murex trunculus</i>	<i>Patella</i>	<i>Spondylus gaederopus</i>	<i>Ostrea edulis</i>
<i>Dentale</i>	-	1	-	-	-	-	-	12	-	-	-	-	-	-	-	-	-
<i>Dentes</i>	-	-	-	-	-	-	-	4	-	-	-	-	-	-	-	-	-
<i>Premaxilla</i>	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-
<i>Maxilla</i>	-	-	-	-	1	-	-	1	-	-	-	-	-	-	-	-	-
<i>Mandibula</i>	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-
<i>lomandibula</i>	-	-	-	-	1	-	-	1	-	-	-	-	-	-	-	-	-
<i>Preoperculous</i>	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-	-	-
<i>Supracleitrum</i>	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-	-
<i>Postcleitrum</i>	-	-	-	-	-	-	-	-	1	-	-	-	-	-	-	-	-
<i>Vertebrae</i>	1	5	1	3	-	-	2	8	34	-	-	-	-	-	-	-	-
<i>Valva</i>	-	-	-	-	-	-	-	-	-	10	-	-	-	-	17	10	2
<i>Gasteropode</i>	-	-	-	-	-	-	-	-	-	-	31	2	27	10	-	-	-
Total	1	6	1	3	2	1	2	29	35	10	31	2	27	10	17	10	2

Tab. 2. Table of marine animal remains from the entire faunal assemblage.

⁶ For the determination of marine fish species, reference was made to the anatomical atlas by CANNON (1987) and the website of the University of Nottingham, available at the following link: <http://fishbone.nottingham.ac.uk/>. For acquiring osteometric measurements, MORALES, ROSEN LUND 1979 was consulted.




Fig. 4. Percentages, based on the Number of Remains, of the fish species identified.

most of which were collected in the intertidal zone, on sand, or the rocky shores. Murices (*Murex trunculus*), limpets, or the more common *Cerithidae* show no signs of breakage, indicating selection for food purposes. In the case of bivalves, only a few nicks on the edge attributable to the pressure exerted at the time of opening the molluscs were observed. Much more prized was the consumption of the spiny oyster (*Spondylus gaederopus*), present in almost 10% of the entire investigated malacofauna.

6. Anthropological and stable carbon and nitrogen isotope analyses

In the case of the Rab Island context, four burials were considered for stable carbon and nitrogen isotope analysis: three from the urban context of Dinka Dokule and one from the rural site of the Church of Saint Lawrence. A total of six individuals were studied anthropologically (Casolin 2020/2021). From the urban context, a sarcophagus containing three subjects was identified: two mature adult males (individuals A and B) and one adolescent (individual C) approximately 18-20 years old. The second tomb from this context is a stone coffin burial of a subadult subject approximately six months to one year old. The last tomb from the urban context was a pit burial of an elderly adult woman. Finally, in the Saint Lawrence context, a single pit burial of a mature adult female was identified. The small number of burials does not allow for conclusions or hypotheses about the island's population, but it provides an informative basis for future work starting from other sites. Regarding the health status of the studied individuals, none exhibited pathological signs indicative of malignant neoplasms or infectious diseases. Only common skeletal alterations were found, such as periosteal reactions, osteoarthritis, nonspecific stress indicators, nonspecific infections, and dental pathologies. Specifically, only the infant dis-

played *cribra orbitalia* (Walker *et al.* 2009) as a nonspecific stress indicator, with a mild occurrence in the roof of the right orbit. Hypoplasia (Ortner 2003; Hillson 2005) was identified in three cases from the city: the senile adult female, the mature adult male (individual A), and the adolescent, with an onset age ranging between 3.5 and 5 years.

Osteoarthritis (Ortner 2003; Waldron 2009), attributable to age and physical activity, was documented in the analyzed sample, with the vertebral column being the most affected skeletal region. Specifically, several cases involving the vertebral bodies were observed among the studied adults. Schmorl's nodes and a herniated disc were present in the two mature adult males (A and B) from the city. Additionally, there was fusion of two thoracic vertebral arches in the mature adult female from San Lorenzo, and slight eburnation on the superior and inferior facets of the lumbar vertebrae in the senile adult female buried in the city. Other skeletal regions, though less affected, also exhibited arthritic alterations, such as the shoulder girdle, particularly the clavicles, ribs, and hands. Dental pathologies (Hillson 2005) were extensively documented in all adult subjects and the adolescent. Several cases of widespread periodontopathy and dental wear on both dental arches were found. Adults and adolescents from the city exhibited small amounts of tartar. Only two individuals (the senile female and adolescent from the town) presented early stages of caries (respectively: on the right third maxillary molar and the right first maxillary molar). The mature adult female from San Lorenzo was the only one showing antemortem loss of all mandibular molars.

Except for the senile female, all adults exhibited periostitis of the lower limbs (primarily *tibiae* and *fibulae*) as a nonspecific infection. This alteration involves ossification of the periosteum following trauma or nonspecific infectious processes (Ortner 2003). A unique case identified in the analyzed sample was a metabolic endocranial pathology of uncertain etiology found in a senile female from the city. This case involved internal frontal hyperostosis, characterized by a subcircular apposition of new bone, approximately 4.4x3.5 mm, on the endocranial surface near the frontal crest.

For the isotopic study⁷, two bone fragments (usually from femurs and ribs) and the dentin of the first permanent molar were sampled from the six individuals, for eleven samples. Sampling from these two tissues allows for the reconstruction of dietary changes during life stages, as collagen from femurs provides information

⁷ The standard collagen extraction protocol (RICHARDS, HEDGES 1999; PRIVAT *et al.* 2002) was performed on bone and teeth samples. Analyses were conducted using an Elementar Analyzer EA – Flash 2000 connected via ConFlo to a Thermo Fisher Delta V Advantage mass spectrometer, located in the Department of Geosciences (Unipd). CH6 and CH7, N-1 and N-2, UREA, and ALANINE were used as international standards during the analysis phase. All twenty-three analyzed samples met the C:N ratio within the range of 2.9 and 3.6.

about the last 10-20 years of life and ribs about the last 5-10 years, while teeth provide information from childhood (Hakenbeck 2013). To avoid variations in isotopic composition, bone collagen was extracted from non-affected bone areas (Olsen *et al.* 2014). To calibrate and analyse the human diet, contemporaneous fauna samples were collected from the burials. The sampling sought to include the highest representativeness of different taxa, including herbivores, omnivores, birds, and fish. Carnivores were the only absent group. In total, eighteen samples were taken, with only one failing during collagen extraction. The data obtained shows that individuals from both urban and rural contexts had a similar diet. Carbon values are plotted on the x-axis, while nitrogen is on the y-axis (fig. 5). The value of carbon in collagen, between -19.43‰ and -17.40‰, is relatively narrow, indicating a diet based on the consumption of C₃ plants. Carbon values in dentin, although there are only three samples from the urban collective tomb, have an even narrower range but are entirely similar to those of bone collagen (-19.01‰ to -19.65‰). Therefore, individuals' diets did not change significantly over their lifetimes. Nitrogen values in collagen have ranged from 9.07‰ to 12.03‰, indicating that individuals had a diet primarily based on terrestrial C3 resources with a good protein intake. Dentin data is in line with collagen, ranging from 9.48‰ to 10.10‰. Comparing isotopic results with anthropological and paleopathological data, no difference in diet based on sex of adults was observed. The highest nitrogen value corresponds to the infant, mainly due to information recorded in their bones during breastfeeding (Richards *et al.* 2002). The second male from the urban collective tomb has the highest nitrogen values among adults, while the adolescent buried in the same sarcophagus has the lowest values among all analysed subjects, suggesting limited access to protein resources.

Regarding fauna, all samples have a C₃-based diet except for the two sea bream samples, which show much higher carbon values, indicating a C₄-based diet (average carbon: -10.20‰, average nitrogen: 6.72‰). Herbivores have the lowest nitrogen values⁸, while birds have the highest carbon and nitrogen values. Sheep and goats⁹ have higher nitrogen values compared to other herbivores, and pigs¹⁰ have similar values to them. Birds have a much more varied diet¹¹. Variations within individual species may be due to different dating or dietary stresses experienced by different specimens during life.

⁸ On average, cattle show carbon values of -19.91‰ and nitrogen values of 4.70‰. Horses exhibit carbon values of -20.46‰ and nitrogen values of 3.95‰. Deer, on the other hand, displays carbon values of -21.24‰ and nitrogen values of 4.63‰. Finally, hares demonstrate values of -20.27‰ for carbon and 4.36‰ for nitrogen.

⁹ Sheep and goats exhibit an average carbon value of -20.81‰ and a nitrogen value of 5.50‰.

¹⁰ Pigs exhibit an average carbon value of -20.55‰ and a nitrogen value of 6.22‰.

¹¹ Roosters and *Galliformes* exhibit an average carbon value of -19.32‰ and a nitrogen value of 7.05‰.




Fig. 5. The graph compares the values of stable carbon and nitrogen isotopes in bone collagen, tooth dentin of the analyzed individuals, and collagen from the selected fauna.

7. Tracing dietary changes in coastal Croatian sites between the Roman and Early Medieval periods

The results obtained from archaeozoological and isotopic analyses of carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) on human and animal bone samples from two sites located in the island of Rab have been compared with data from fourteen different Croatian sites, ascertained through a meticulous survey of existing literature (tab. 3). All these sites are either adjacent to or near the Adriatic Sea, thereby ensuring geographical consistency in the comparisons (fig. 6). These sites cover a considerable chronological range, from the Roman period to the end of the Early Middle Ages. This temporal breadth has not only allowed us to contextualize the results obtained for the interred individuals and fauna of Rab within a specific ge-

	Site	Chronology	Bibliography
1	Rab	6 th -8 th century AD	-
2	Hvar - Radoševic	5 th -early 7 th century AD	Zagorc <i>et al.</i> 2024
3	Velim - Velištak	7 th -mid 9 th century AD	Lightfoot <i>et al.</i> 2012
4	Podvršje	5 th -early 7 th century AD	
5	Glavice Gluvine	8 th -early 10 th century AD	
6	Radašinovci - Vinogradine	9 th -early 10 th century AD	
7	Šibenik – Sveti Lovre	9 th -early 11 th century AD	
8	Zadar-Relje	3 rd -4 th century AD	
9	Vis-Bandirica	2 nd century BC-2 nd century AD	
10	Dubravice	8 th -9 th century AD	Gunjaca 1987; Novak <i>et al.</i> 2016
11	Stranče - Gorica	8 th -mid 9 th century AD	Novak <i>et al.</i> 2016
12	Konjsko polje – Livade	9 th century AD	Petrinec 2005; Novak <i>et al.</i> 2016
13	Omišalj - Mirine	5 th -7 th century AD	Čaušević-Bully <i>et al.</i> 2023
14	Martinšćica	5 th -7 th century AD	Čaušević-Bully <i>et al.</i> 2023
15	Vačani	8 th -10 th century AD	Novak <i>et al.</i> 2016

Tab. 3. Sites considered for comparing stable isotope analyses with chronology and reference bibliography.



Fig. 6. Distribution of the sites considered for comparing stable isotope analyses.

ographic framework but also to observe potential dietary evolutions from the 2nd century BC to the 11th century AD.

In all the contexts examined, stable isotopic analyses were conducted on human and animal remains to calibrate the local trophic level. Among the fourteen sites considered, two present burial phases are attributable to the Roman occupation (Zadar-Relja - 3rd-4th century AD; Vis-Bandirica - 2nd century BC-2nd century AD), while another four are assignable to the Late Antique phase between the 5th and early 7th centuries AD (Hvar-Radošević; Podvršje; Omišalj-Mirine; Martinščica). More numerous are the Early Medieval sites (Velim-Velištak; Glavice Gluvine; Radašinovci-Vinogradine; Šibenik-Sveti Lovre; Dubravice; Stranče-Gorica; Konjsko polje-Livade), which have allowed us to obtain a local isotopic signal from the 7th to the 11th centuries AD.

The isotopic averages of 65 human bone samples from the Roman period, 68 from the Late Antique period, and 293 from the Early Medieval period, as well as 90 contemporaneous faunal samples, have provided the opportunity to verify possible variations in human and animal diets over the centuries¹². These variations possibly reflect socioeconomic, environmental, and cultural changes that have characterized the area over time.

The comparison of carbon and nitrogen values across the various analyzed populations reveals a notable similarity, with some local peculiarities and variations related to the different historical periods considered (fig. 7).

During the Roman period, analyses conducted by Lightfoot (2009) and Lightfoot *et al.* (2012) on the two examined contexts show differences. However, a diet predominantly based on C₃ terrestrial resources with a variable contribution of marine proteins emerges. For instance, at Vis-Bandirica, the diet was primarily composed of C₃ plants with a variable intake of marine proteins, while at Zadar-Relja, a C₃-based diet was observed, but with the consumption of C₄ plants and/or low trophic level marine foods.

In the Late Antiquity period, a similar situation to the Roman period is observed, with a diet predominantly based on the C₃ ecosystem and limited use of marine resources. Specifically, Podvršje presents a range of values similar to Vis-Bandirica (Lightfoot 2009; Lightfoot *et al.* 2012). Samples from Mirine and Martinščica indicate a diet based on C₃ plants with a pronounced intake of C₄-type vegetables. Individuals from Martinščica seem to have consumed a higher quantity of C₄ food, which could indicate the consumption of algae and other marine foods (Čaušević-Bully *et al.* 2023). On the island of Hvar, the analyzed individuals reveal a diet similar to that of the two insular sites of Cres and Krk, with a C₃-based diet and minimal consumption of marine foods (Čaušević-Bully *et al.* 2023; Novak *et al.* 2016).

¹² In the count, human and animal bone samples from the site of Rab are also included.




Fig. 7. The graph shows the comparison of carbon and nitrogen isotopic data of the analyzed subjects from the Island of Rab with Roman, Late Antique, and Early Medieval contexts from the islands, coast, and inland of Croatia.

During the Early Medieval period, the eight contexts considered show values indicating a relatively uniform diet among the inhabitants of the eastern Adriatic coast, based on C₃ terrestrial resources, with a variable intake of C₄, mainly due to millet consumption, and a very low contribution of marine resources (Novak *et al.* 2016). At Velim-Velištak, subjects presented a diet that included a variable intake of C₄ or low trophic level marine proteins and a good consumption of animal proteins. At Glavice Gluvine, most of the population consumed a higher quantity of C₄ plants, such as millet, while $\delta^{15}\text{N}$ values were less varied, suggesting less diversified use of animal resources. At Radašinovci-Vinogradine, millet consumption varied within the population, with a high intake of animal proteins and a certain variety in the quantity consumed. At Šibenik-Sveti Lovre, compared to the previous three contexts, a greater consumption of C₃ resources and small amounts of freshwater resources, visible from high trophic levels, is observed, given the proximity to the mouth of the Krka River (Lightfoot 2009; Lightfoot *et al.* 2012).

Regarding the remaining four sites – Dubravice, Stranče-Gorica, Konjsko polje-Livade, Omišalj-Mirine, and Vaćani-Laluše – it was possible to detect a diet mainly based on C₃ terrestrial resources, with some intake of C₄ foods or possibly marine origin (Novak *et al.* 2016). These results are entirely similar to those of the other Early Medieval contexts analyzed by Lightfoot *et al.* 2012.

Another notable finding from this comparison is the absence of distinctions based on sex and age at death among adults. However, for subadult subjects, a certain variety is observed: although they fall within adult ranges, in some cases, they show greater variability, especially in $\delta^{13}\text{C}$ values, covering a broader range, indicating higher consumption of C₄ or marine resources (Novak *et al.* 2016; Čaušević-Bully *et al.* 2023; Zagorc *et al.* 2024). The variability in $\delta^{15}\text{N}$ values in subadults is primarily linked to the weaning phase.

In comparison with archaeological data and possible indicators of social status, most contexts do not show diet variations attributable to the burial location, tomb type, or presence of grave goods. The only exception involves two adult males from Vaćani, who present high $\delta^{15}\text{N}$ values indicative of greater animal resource consumption and were buried in sarcophagi with rich grave goods (Novak *et al.* 2016).

The Early Medieval individuals from Rab exhibit data similar to other contemporary sites but with a diet entirely and predominantly based on C₃ terrestrial resources and good consumption of animal proteins.

Regarding the fauna, a comparison was made with that analyzed in the study by Lightfoot *et al.* (2012), both for the Roman and Early Medieval periods and with the only sample from the Early Medieval context of Vaćani (Novak *et al.* 2016). Different taxa generally fall within the range of a diet primarily based on C₃ plants, except for the Early Medieval pigs and dogs studied by Lightfoot *et al.*

(2012), which show greater variability and partial consumption of C_4 resources. The observed differences, mostly at the trophic level, can be correlated with chronological differences between the analyzed samples. As for fish, at the current state of research, comparisons are not possible, as the only samples analyzed are those present in this study.

8. Conclusion

The analysis of faunal remains has provided a detailed overview of animal exploitation practices from the 6th to the 8th century AD on the island of Rab. This study examined the use of major domestic species (goats, pigs, and cattle) and aquatic resources in the local diet. Pigs, constituting approximately 15% of the faunal sample, were slaughtered and butchered on-site, with a particular focus on major joints for meat distribution. Swine husbandry primarily targeted meat production, with slaughtering typically occurring between 18 and 24 months of age. Poultry, crucial to Rab's urban economy, provided quick access to protein resources through meat and eggs. Sheep and goats, predominant in the sample, played a central role in the local economy, both for secondary product recovery and meat consumption. Analysis of fish and bivalve remains highlighted their contribution to the island's protein intake, featuring prized species like *Ostrea edulis* and *Spondylus gaederopus* among marine finds. Unlike fishing, hunting played a minor role in the island's economy and diet, possibly due to intensive agriculture and deforestation for goat husbandry. With limited studies on plant species and cultivated fruits, the relationship between cultivated and uncultivated areas on the island remains uncertain.

The analysis of carbon and nitrogen values across various populations reveals significant similarities, with local peculiarities and historical period variations. During the Roman period, diets were mainly based on C_3 terrestrial resources with variable marine protein intake, as seen in Vis-Bandirica and Zadar-Relja. In Late Antiquity, diets remained predominantly C_3 -based with limited marine resources, as observed in Podvršje, Mirine, and Martinščica. In the Early Medieval period, diets were relatively uniform along the eastern Adriatic coast, primarily based on C_3 terrestrial resources with variable C_4 intake, mainly from millet, and minimal marine resources. Notable sites include Velim-Velištak, Glavice Gluvine, Radašinovci-Vinogradine, and Šibenik-Sveti Lovre. The remaining sites (Dubravice, Stranče-Gorica, Konjsko polje-Livadine, Omišalj-Mirine, and Vaćani-Laluše) also showed similar dietary patterns. No significant diet differences based on sex or adult age were found, but subadults showed greater variability in $\delta^{13}C$ values, indicating higher C_4 or, possibly, marine resource intake. Archaeological data mostly did not indicate social status-based dietary differences, ex-

cept for two adult males from Vaćani. Faunal comparisons showed diets primarily based on C_3 plants, with some variability in pigs and dogs. The most prominent finding from the comparison of both diachronic and contemporaneous contexts is that, despite originating from islands, coastal areas, and inland zones not far from the coast, the consumption of marine proteins is very low. In many contexts, the positive carbon values have been more closely associated with millet consumption. In the absence of strontium isotope analysis, we cannot be certain whether the individuals buried at Rab and Banjol are local residents, a consideration that could isotopically explain the absence of marine resources in their diet. This hypothesis, in our view, is contradicted by the results of isotopic analyses obtained for the comparison sites. To achieve greater discrimination between consumers of terrestrial and marine resources in the study area, future analyses utilizing stable oxygen ($\delta^{18}O$) and sulphur ($\delta^{34}S$) isotopes would be desirable. These, along with the already obtained carbon and nitrogen isotopic signals, would highlight the actual exploitation and consumption of marine resources (Craig *et al.* 2006).

Acknowledgments

Although the paper is the result of joint work between the two authors, Mirko Fecchio wrote paragraphs 2-5, Maurizio Marinato wrote paragraphs 1 and 6. Paragraphs 7 and 8 were written jointly.

The excavations at the sites of Dinka Dokule and Banjol were directed by Professors Alexandra Chavarría Arnau (Department of Cultural Heritage at the University of Padua) and Miljenko Jurkovic (University of Zagreb); the zooarchaeological and anthropological study was supported by the Conservation Department in Rijeka, while the isotopic analyses were conducted under the supervision of Professor Manuel Rigo (Department of Geosciences at the University of Padua). We wish to thank them for their invaluable support to this research.

Abstract

The zooarchaeological analysis conducted at the archaeological site of Dinka Dokule in the city of Rab (Croatia) outlined an overall picture of animal exploitation during the early medieval phase of occupation of this building (6th-8th century AD). This includes both the main domestic species and the fish resources. The reconstruction of the complex animal butchery process, combined with stable isotope analyses ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) on both fauna and human remains from contemporary burials, has enhanced the understanding of the consumption of plant and, above all, animal products. Comparison with other Croatian sites reveals similar dietary patterns with some changes from the Roman to the Early Medieval period. Despite the presence of various species of marine fish among the meal remains stable isotope analyses have not indicated significant consumption of marine resources.

Keywords: Late Antiquity - Early Middle Ages, zooarchaeology, stable isotopes, Island of Rab.

L'analisi zooarcheologica condotta nel sito archeologico di Dinka Dokule, nella città di Rab (Croazia), ha delineato un quadro complessivo dello sfruttamento animale durante la fase di occupazione altomedievale di questo edificio (VI-VIII secolo d.C.). Questo comprende sia le principali specie domestiche sia le risorse ittiche. La ricostruzione del complesso processo di macellazione degli animali, combinata con le analisi degli isotopi stabili ($\delta^{13}\text{C}$ e $\delta^{15}\text{N}$) sia sulla fauna che sui resti umani provenienti da sepolture contemporanee, ha migliorato la comprensione del consumo di prodotti vegetali e, soprattutto, animali. Il confronto con altri siti croati rivela modelli alimentari simili, con alcuni cambiamenti dal periodo romano a quello altomedievale. Nonostante la presenza di varie specie di pesci marini tra i resti dei pasti, le analisi degli isotopi stabili non hanno indicato un consumo significativo di risorse marine.

Parole chiave: Tarda Antichità - Alto Medioevo, zooarcheologia, isotopi stabili, isola di Rab.

References

- ARCHAEOLOGICAL FISH RESOURCE <http://fishbone.nottingham.ac.uk/> (ultimo accesso 26 marzo 2024).
- J. BAKER, D. BROTHWELL 1980, *Animal Diseases in Archaeology*, London.
- S. BATOVIĆ 1987, *Prapovijesni ostaci na otoku Rabu*, in M. GROSS, I. KAMPUŠ, T. RAUKAR, P. STRČIĆ (eds), *Rapski Zbornik*, Zagreb, pp. 147-169.
- J. BOESSNECK, H.H. MÜLLER, M. TEICHERT 1964, *Osteologische Unterscheidungsmerkmale zwischen Schaf und Ziege*, "Julius-Kühn-Archiv", 78, pp. 1-129.
- G.P. BROGIOLO, A. CHAVARRÍA ARNAU, F. CERES, M. FECCHIO, M. MARINATO, L. MINZON 2023, *Lo scavo in via Dinka Dokule n. 5 di Rab nel quadro della storia della città*, "Hortus Artium Medievalium", 29, pp. 59-75.

- G.P. BROGIOLO, A. CHAVARRÍA ARNAU, F. GIACOMELLO, M. JURKOVIĆ, G. BILOGRIVIĆ 2017, *The Late Antique church of Saint Lawrence, Banjol (island of Rab, Croatia). Results of the first two archaeological campaigns (2015-2016)*, "Hortus Artium Medievalium", 23, pp. 666-673.
- G.P. BROGIOLO, M. NEBBIA, F. BENETTI 2013, *Ricerche sul paesaggio dell'isola di Arbe*, in G. CUSCITO (ed), *Le modificazioni del paesaggio nell'altoadriatico tra pre-protostoria ed altomedioevo*, Atti della XLIII Settimana di Studi Aquileiesi (10-12 maggio 2012), Trieste, pp. 217-229.
- A. CASOLIN 2020/2021, *Studio antropologico e paleopatologico del materiale antropologico proveniente dagli scavi di via Dinka Dokule n. 5 (2015-2017-2018) e San Lorenzo in Banjol (2015-2016), (Isola di Rab, Croazia)*, Tesi triennale, Università degli Studi di Padova.
- M. ČAUŠEVIĆ-BULLY, M. NOVAK, A. DELLISTE, S. BULLY, M. CARIĆ 2023, *Insular population of the Kvarner Archipelago in the 5th century: Mirine-Fulium and Martinšćica church complex case studies*, "Antiquité Tardive", 31, pp. 63-82.
- M. ČAUŠEVIĆ-BULLY, C. MASSA, M. ROUSSEAU, V. BICHET, H. RICHARD, M. NOVAK 2021, *L'homme et son environnement. Approche interdisciplinaire sur l'évolution des territoires insulaires de l'archipel du Kvarner (Croatie)*, "Antiquité Tardive", 29, pp. 57-68.
- A. COHEN, D. SERJEANTSON 1986, *A manual for the identification of bird bones from archaeological sites*, London.
- O. CRAIG, R. ROSS, S.H. ANDERSEN, N. MILNER, G. N. BAILEY 2006, *Focus: sulphur isotope variation in archaeological marine fauna from northern Europe*, "Journal of Archaeological Science", 33, pp. 1642-1646.
- J. DE GROSSI MAZZORIN 2008, *Archeozoologia: lo studio dei resti animali in archeologia*, Bari.
- J. DE GROSSI MAZZORIN, C. MINNITI 2001, *L'allevamento e l'approvvigionamento alimentare di una comunità urbana. L'utilizzazione degli animali a Roma tra il VII e il X secolo*, in M.S. ARENA, P. DELOGU, L. PAROLI, M. RICCI, L. SAGUI, L. VENDITELLI (eds), *Roma dall'antichità al medioevo. Archeologia e storia nel museo nazionale romano Crypto Balbi*, Milano, pp. 69-78.
- J. DE GROSSI MAZZORIN, G. DE VENUTO 2009, *L'economia produttiva animale: i risultati delle analisi archeozoologiche*, in P. ARTHUR, M. LEO IMPERIALE, M. TINELLI (eds) *Apigliano. Un villaggio bizantino e medioevale in Terra d'Otranto. I reperti*, Lecce, pp. 23-27.
- H.M.A. EVANS 1989, *The Early Mediaeval Archaeology of Croatia: A.D. 600-900*, Oxford.
- M. FECCHIO, U. TECCHIATI 2021, *I resti faunistici altomedievali della torre*, in A. CHAVARRÍA ARNAU, G.P. BROGIOLO (eds), *Torba (VA). Scavi 2013-2019*, Mantova, pp. 203-225.
- D. GIFFORD-GONZALEZ 2018, *An introduction to zooarchaeology*, Switzerland.
- A. GRANT 1982, *The use of tooth wear as a guide to the age of domestic ungulates*, London.
- Z. GUNJAČA 1987, *Dubravice - ranosrednjovjekovno groblje*, "Arheološki pregled", 28, pp. 148-150.
- S. HAKENBECK 2013, *Potentials and limitations of isotope analysis in Early Medieval archaeology*, "European Journal of Post-Classical Archaeologies", 3, pp. 95-111.
- S. HILLSON 2005, *Teeth*, Cambridge.
- M. JURKOVIĆ, G.P. BROGIOLO, T. TURKOVIĆ, A. CHAVARRÍA ARNAU, I. MARIĆ 2012, *Kaštelina na otoku Rabu. Od rimske vile do ranobizantske utvrde*, in J. ANDRIĆ, R. LONČARIĆ (eds), *Rapski Zbornik 2*, Rab, pp. 1-13.
- E. LIGHTFOOT 2009, *Bioarchaeological analyses of archaeological populations from Croatia: a comparison of isotopic and archaeological results*, PhD thesis, University of Cambridge.
- E. LIGHTFOOT, M. ŠLAUS, T.C. O'CONNELL 2012, *Changing cultures, changing cuisines: Cultural transitions and dietary change in iron age, roman, and early medieval Croatia*, "American Journal of Physical Anthropology", 148, 4, pp. 543-556.

- A. MORALES, K. ROSENBLUND 1979, *Fish Bone Measurements. An attempt to standardize the Measuring of fish bones from archaeological sites*, Copenhagen.
- G. NOBIS 1954, *Zur Kenntnis der ur- und frühgeschichtlichen Rinder Nord- und Mitteldeutschlands*, "Zeitschrift für Tierzucht und Züchtungsbiologie", 63, pp. 155-194.
- M. NOVAK, R. HOWCROFT, R. PINHASI, M. ŠLAUS 2016, *Dietary trends in early medieval Croatia as evidenced by stable isotope analysis*, "American Journal of Physical Anthropology", 62, p. 242.
- K.C. OLSEN, C. D. WHITE, F. J. LONGSTAFFE, K. VON HEYKING, G. MCGLYNN, G. GRUPE, F. J. RUHLI 2014, *Intraskeletal Isotopic Compositions ($\delta^{13}C$, $\delta^{15}N$) of Bone Collagen: Nonpathological and Pathological Variation*, "American Journal of Physical Anthropology", 153, pp. 598-604.
- D.J. ORTNER 2003, *Identification of Pathological Conditions in Human Skeletal Remains*, Washington.
- S. PAYNE 1973, *Kill-off Patterns in Sheep and Goat: The Mandibles from Aşvan Kale*, "Anatolian Studies", 23, pp. 281-303.
- M. PETRINEC 2005, *Ranosrednjovjekovno groblje na položaju Livade u Konjskom polju*, "Starohrvatska prosvjeta", 32, pp. 21-52.
- A. PITEŠA 2006, *The Slavs and the Early Croatian State*, in D. DAVISON, V. GAFFNEY, E. MARIN (eds), *Dalmatia: Research in the Roman Province, 1970-2001*, Oxford, pp. 193-212.
- A. POWELL 2004, *The faunal remains*, in R. HODGES, W. BOWDEN, K. LAKO (eds) *Byzantine Butrint, Excavations and Surveys 1994-1999*, Oxford, pp. 305-320.
- K.L. PRIVAT, T.C. O'CONNELL, M.P. RICHARDS 2002, *Stable Isotope Analysis of Human and Faunal Remains from the Anglo-Saxon Cemetery at Berinsfield, Oxfordshire: Dietary and Social Implications*, "Journal of Archaeological Science", 29, 7, pp. 779-90.
- M.P. RICHARDS 2020, *Isotope analysis for diet studies*, in M.P. RICHARDS, K. BRITTON (eds), *Archaeological Science. An introduction*, Cambridge, pp. 125-143.
- M.P. RICHARDS, R.E.M. HEDGES 1999, *Stable Isotope Evidence for Similarities in the Types of Marine Foods Used by Late Mesolithic Humans at Sites Along the Atlantic Coast of Europe*, "Journal of Archaeological Science", 26, pp. 717-722.
- M.P. RICHARDS, S. MAYS, B.T. FULLER 2002, *Stable Carbon and Nitrogen isotope values of bone and teeth reflect weaning age at the Medieval Wharram Percy Site, Yorkshire, UK*, "American Journal of Physical Anthropology", 119, pp. 205-210.
- C. SCHWARTZ 1996, *The Faunal Remains*, in J. CHAPMAN, R. SHIEL, S. BATOVIĆ (eds), *The Changing Face of Dalmatia: Archaeological and Ecological Studies in a Mediterranean Landscape*, London, pp. 176-178.
- A. ŠKEGRO 2006, *The Economy of Roman Dalmatia*, in D. DAVISON, V. GAFFNEY, E. MARIN (eds), *Dalmatia: Research in the Roman Province, 1970-2001*, Oxford, pp. 149-173.
- L. TOMAZ 2003, *In Adriatico nell'Antichità e nell'Alto Medioevo. Da Dionigi di Siracusa ai Dogi Orseolo*, Venezia.
- P.L. WALKER, R.R. BATHURST, R. RICHMAN, T. GJERDRUM, A. ANDRUSHKO 2009, *The causes of porotic hyperostosis and cribra orbitalia: a reappraisal of the iron-deficiency-anaemia hypothesis*, "American Journal of Physical Anthropology", 139, pp. 109-125.
- T. WALLDRON 2009, *Paleopathology*, Cambridge.
- J.J. WILKES 1992, *The Illyrians*, Oxford.
- J.J. WILKES 1977, *The Population of Roman Dalmatia*, "Aufstieg und Niedergang der römischen Welt", 2(6), pp. 732-766.
- B. ŽAGORC, M. BLANZ, P. GELABERT, S. SAWYER, V. OBERREITER, O. CHERONET, H. SHAN CHEN, M. CARIĆ, E. VISKOVIĆ, I. OLALDE, M. IVANOVA-BIEG, M. NOVAK, D. REICH, R. PINHASI 2024, *Bioarchaeological Perspectives on Late Antiquity in Dalmatia: Paleogenetic, Dietary, and Population Studies of the Hvar – Radošević burial site*, Preprints and early-stage research, <https://www.biorxiv.org/content/10.1101/2024.05.14.594056v1>.