

ON THE LOCAL MAXIMIZERS OF HIGHER CAPACITY RATIOS

LUCA BARACCO, OLGA BERNARDI,
CHRISTIAN LANGE, AND MARCO MAZZUCHELLI

ABSTRACT. We prove an analogue of the 4-dimensional local Viterbo conjecture for the higher Ekeland-Hofer capacities: on the space of 4-dimensional smooth star-shaped domains of unitary volume, endowed with the C^3 topology, the local maximizers of the k -th Ekeland-Hofer capacities are those domains symplectomorphic to suitable rational ellipsoids.

CONTENTS

1. Introduction	1
1.1. The Viterbo conjecture	1
1.2. Main result	2
1.3. Open questions on higher capacity ratios	3
1.4. Outline of the proof of Theorem A	6
1.5. Organization of the paper	8
2. The Clarke action functional	8
2.1. Smooth convex bodies	8
2.2. Spectral invariants	10
2.3. Finite dimensional reduction	11
2.4. Besse convex bodies	12
2.5. Local maximizers of the spectral ratio	16
3. The Ekeland-Hofer capacities	17
3.1. Definition of the capacities	17
3.2. Capacities vs spectral invariants	19
3.3. Local maximizers of the capacity ratio	24
4. Maximizers of the capacity ratios among ellipsoids	27
5. Maximizers of the capacity ratios among toric domains	28
5.1. Concave toric domains	29
5.2. Convex toric domains	29
Appendix A. The Fadell-Rabinowitz index	36
Acknowledgments	40
References	40

1. INTRODUCTION

1.1. The Viterbo conjecture. Symplectic capacities are fundamental invariants that govern many rigidity phenomena in symplectic and contact topology. In this

Date: March 23, 2023.

2020 Mathematics Subject Classification. 37J39, 53D05, 53D35.

Key words and phrases. Viterbo conjecture, symplectic capacities, Ekeland-Hofer capacities.

Marco Mazzucchelli is partially supported by the ANR grants CoSyDy (ANRCE40-0014) and COSY (ANR-21-CE40-0002). Olga Bernardi and Marco Mazzucchelli are partially supported by the PRIN project 2017S35EHN 003 2019-2021 “Regular and stochastic behaviour in dynamical systems”.

paper, we consider symplectic capacities c defined on subsets B of the symplectic vector space (\mathbb{C}^n, ω) , for $n \geq 2$. If B is a domain (i.e. the closure of a non-empty open subset) of finite volume, its associated *capacity ratios* are defined as

$$\widehat{c}(B) = \frac{c(B)}{\text{vol}(B, \omega)^{1/n}}.$$

Here, the volume is the integral of ω^n on B . In his seminal paper [Vit00], Viterbo conjectured that $\widehat{c}(B) \leq \widehat{c}(B^{2n})$ for any convex body $B \subset \mathbb{C}^n$ and any symplectic capacity c , where B^{2n} denotes any round ball in \mathbb{C}^n . This conjecture, which is currently open, is particularly hard as it combines symplectic invariants with the non-symplectic notion of convexity. Its validity would have far reaching consequences well beyond symplectic geometry: it would imply the validity of the long-standing Mahler conjecture from convex geometry, as it was pointed out by Artstein-Avidan, Karasev, and Ostrover [AAKO14].

By a $2n$ -dimensional *smooth convex body*, we mean a compact subset $B \subset \mathbb{C}^n$ with non-empty interior and smooth positively-curved boundary ∂B ; for the sake of convenience, we shall always require that B contains the origin in its interior. We consider the C^k topology on the space of smooth convex bodies: two smooth convex bodies $B_0, B_1 \subset \mathbb{C}^n$ are C^k -close when their boundaries $\partial B_0, \partial B_1$ are C^k -close submanifolds. A theorem of Abbondandolo and Benedetti [AB19], extending an earlier result of Abbondandolo, Bramham, Hryniewicz, and Salomao [ABHS18] for the Ekeland-Hofer capacity in dimension 4, asserts the validity of a local version of the Viterbo conjecture: on a sufficiently C^3 -small neighborhood of the round ball B^{2n} , the local maximizers of the function $B \mapsto \widehat{c}(B)$ are precisely the smooth convex bodies symplectomorphic to B^{2n} . In dimension 4, this theorem has been recently strengthened by Edtmair [Edt22]: on a sufficiently C^3 -small neighborhood of the round ball B^{2n} , all symplectic capacities coincide.

1.2. Main result. The goal of this paper is to investigate questions analogous to the (local and global) Viterbo conjecture for higher symplectic capacities, whose definition is the following. Each tuple $\mathbf{a} = (a_1, \dots, a_n) \in (0, \infty]^n$ defines a $2n$ -dimensional domain

$$E(\mathbf{a}) = \left\{ z = (z_1, \dots, z_n) \in \mathbb{C}^n \mid \sum_{j=1}^n \frac{|z_j|^2}{a_j} = \frac{1}{\pi} \right\}.$$

If all the entries a_i are finite, $E(\mathbf{a})$ is an ellipsoid. A *symplectic k -capacity* on \mathbb{C}^n is a function $c_k : 2^{\mathbb{C}^n} \rightarrow [0, \infty]$, where $2^{\mathbb{C}^n}$ is the collection of subsets of \mathbb{C}^n , satisfying the following three properties:

- **(Monotonicity)** $c_k(U) \leq c_k(V)$ if there exists $\psi \in \text{Symp}(\mathbb{C}^n, \omega)$ such that $\psi(U) \subset V$.
- **(Conformality)** $c_k(bU) = b^2 c_k(U)$ for each $b \in \mathbb{R}$ and $U \subset \mathbb{C}^n$.
- **(Normalization)** For each $\mathbf{a} \in (0, \infty]^n$ with $a_1 < \infty$, if we enumerate the set of positive multiples of the entries a_j in increasing order as $\tau_1 < \tau_2 < \tau_3 < \dots$, the value $c_k(E(\mathbf{a}))$ is the k -th element in the sequence

$$\underbrace{\tau_1, \dots, \tau_1}_{\times m_1}, \underbrace{\tau_2, \dots, \tau_2}_{\times m_2}, \underbrace{\tau_3, \dots, \tau_3}_{\times m_3}, \dots$$

where m_h is the number of parameters a_j having τ_h as a positive multiple.

Ordinary symplectic capacities on \mathbb{C}^n , as appearing in the Viterbo conjecture, are 1-capacities¹. While there are many 1-capacities in the literature, to the best of the authors' knowledge there are only few constructions of symplectic k -capacities: the Ekeland-Hofer capacities c_k^{EH} [EH89, EH90] defined via an infinite dimensional linking argument (see Section 3.1), the Viterbo capacities c_k^{V} [Vit89] defined via finite dimensional reductions, the Gutt-Hutchings capacities c_k^{GH} [GH18] defined via S^1 -equivariant symplectic homology, and the very recent Zhang capacities [Zha21] defined via the microlocal theory of sheaves. At least the three k -capacities c_k^{EH} , c_k^{V} , and c_k^{GH} are expected to coincide².

We say that $B \subset \mathbb{C}^n$ is a $2n$ -dimensional *smooth star-shaped domain* when it is diffeomorphic to a compact $2n$ -dimensional ball and has a smooth boundary ∂B that is transverse to all radial lines on \mathbb{C}^n . Notice that the smooth convex bodies, and in particular the ellipsoids, are smooth star-shaped domains. Once again, we consider the C^k topology on the space of smooth star-shaped domains: B_0 and B_1 are C^k -close when their boundaries $\partial B_0, \partial B_1$ are C^k -close submanifolds.

An ellipsoid $E = E(a_1, \dots, a_n)$ is called *rational* when all the ratios a_j/a_h are rational numbers. In this case, we denote by $\tau(E)$ the least common multiple of the parameters a_1, \dots, a_n . For each integer $m \geq 1$, we define $k_m(E)$ to be the minimal integer $k \geq 1$ such that $c_k(E) = m \tau(E)$. Here, c_k is any symplectic k -capacity, and the values $k_m(E)$ are clearly independent of the choice of such c_k 's. Finally, we set $K(E) := \{k_m(E) \mid m \geq 1\}$. Our main result is a confirmation of a version of the local Viterbo conjecture for the higher Ekeland-Hofer capacities in dimension 4.

Theorem A. *On the space of 4-dimensional smooth star-shaped domains endowed with the C^3 topology, the local maximizers of $B \mapsto \widehat{c}_k^{\text{EH}}(B)$ are precisely those domains symplectomorphic to a 4-dimensional rational ellipsoid E with $k \in K(E)$.*

1.3. Open questions on higher capacity ratios. In the Viterbo conjecture, the convexity assumptions on the domains cannot be removed: in an unpublished note [Her98], Hermann showed that, for any symplectic 1-capacity c_1 on \mathbb{C}^n , the associated capacity ratio \widehat{c}_1 is unbounded over the space of star-shaped domains. In contrast, a theorem of Artstein-Avidan, Milman, and Ostrover [AAMO08] asserts that any 1-capacity ratio is uniformly bounded over the space of convex bodies by a constant that is independent of the dimension (previously, Viterbo [Vit00] showed that there is a uniform bound growing linearly in the dimension). A minor modification in their proof provides the following uniform bound for all k -capacity ratios.

Theorem 1.1 (Artstein-Avidan, Milman, Ostrover). *There exists a constant $a > 0$ such that, for each integer $n \geq 1$, for each symplectic k -capacity c_k on \mathbb{C}^n , and for each $2n$ -dimensional convex body B , we have $\widehat{c}_k(B) \leq ak$.*

Proof. By [AAMO08, Theorem 1.6], there exists a constant $a > 0$ such that, for any $n \geq 1$ and for any convex body $B \subset \mathbb{C}^n$, we have

$$\frac{c_{\text{lin}}(B)}{\text{vol}(B)^{1/n}} \leq a,$$

¹In the literature, and in particular in the statement of the Viterbo conjecture, a stronger form of monotonicity is sometimes assumed for the symplectic capacities: $c(U) \leq c(V)$ if there exists a symplectic embedding of U into V . Capacities and, more generally, k -capacities satisfying the weaker monotonicity stated above are sometimes called relative or non-intrinsic [MS17, Sect. 12.1].

²The equality $c_k^{\text{EH}} = c_k^{\text{GH}}$ was recently announced by Gutt and Ramos [GR23].

where

$$c_{\text{lin}}(B) := \inf \left\{ b > 0 \mid \psi(B) \subset \underbrace{E(b) \times \mathbb{C}^{n-1}}_{=E(b, \infty, \dots, \infty)} \text{ for some } \psi \in \text{Sp}(2n) \right\}.$$

The normalization of the symplectic k -capacity implies $c_k(E(b) \times \mathbb{C}^{n-1}) = kb$, and together with the monotonicity property we obtain $c_k(B) \leq k c_{\text{lin}}(B)$. Therefore

$$\widehat{c}_k(B) \leq k \frac{c_{\text{lin}}(B)}{\text{vol}(B)^{1/n}} \leq ak. \quad \square$$

In view of Theorem 1.1, it is natural to raise the following question, which for 1-capacities would be answered by the Viterbo conjecture if confirmed.

Question B. *For any symplectic k -capacity c_k , what are the global maximizers of the function $B \mapsto \widehat{c}_k(B)$ over the space of convex bodies in \mathbb{C}^n ?*

Simple algebraic computations allow to detect the local and global maximizers of \widehat{c}_k over the space of ellipsoids.

Proposition C. *Let c_k be any symplectic k -capacity on \mathbb{C}^n .*

- (i) *The local maximizers of the capacity ratio \widehat{c}_k on the space of $2n$ -dimensional ellipsoids are precisely those rational ellipsoids E such that $k \in K(E)$.*
- (ii) *The global maximum of the capacity ratio \widehat{c}_k on the space of $2n$ -dimensional ellipsoids is*

$$\max_{\mathbf{a}} \widehat{c}_k(E(\mathbf{a})) = (q+1)^{\frac{n-r+1}{n}} (q+2)^{\frac{r-1}{n}},$$

where $q \geq 0$ and $r \in \{1, \dots, n\}$ are integers such that $k = qn + r$. The global maximizers are precisely those rational ellipsoids $E(\mathbf{a})$ of the form

$$\frac{a_i}{a_n} = \begin{cases} \frac{q+1}{q+2}, & \text{if } r > 1 \text{ and } 1 \leq i \leq r-1, \\ 1 & \text{if } r \leq i \leq n. \end{cases}$$

One might be tempted to conjecture that each capacity ratio \widehat{c}_k achieves its maximum over the space of ellipsoids. While for $k = 1$ this is implied by the Viterbo conjecture, for large k this turns out to be false!

Example 1.2. Proposition C implies that the 4-dimensional ellipsoid maximizing any k -capacity ratio \widehat{c}_k is $E(\lceil \frac{k}{2} \rceil, \lceil \frac{k+1}{2} \rceil)$, and

$$\widehat{c}_k(E(\lceil \frac{k}{2} \rceil, \lceil \frac{k+1}{2} \rceil)) = \sqrt{\lceil \frac{k}{2} \rceil \lceil \frac{k+1}{2} \rceil},$$

where $\lceil x \rceil = \min\{y \in \mathbb{Z} \mid y \geq x\}$ denotes the ceiling function. For any k -capacity $c_k \in \{c_k^{\text{EH}}, c_k^{\text{V}}, c_k^{\text{GH}}\}$, the polydisks $P(a, b) = E(a) \times E(b)$ have capacity ratios

$$\widehat{c}_k(P(a, b)) = k \frac{\min\{a, b\}}{\sqrt{2ab}},$$

as it was computed in [EH90, Moa94, GH18], see Example 5.1; in particular $P(1, 1)$ maximizes all capacity ratios \widehat{c}_k among 4-dimensional polydisks:

$$\widehat{c}_k(P(1, 1)) = \frac{k}{\sqrt{2}}.$$

For $k = 2$, the maximizer polydisk $P(1, 1)$ and the maximizer ellipsoid $E(1, 2)$ have the same capacity ratio

$$\widehat{c}_2(P(1, 1)) = \widehat{c}_2(E(1, 2)) = \sqrt{2},$$

whereas for any $k \geq 3$ we have

$$\widehat{c}_k(P(1, 1)) > \widehat{c}_k(E(\lceil \frac{k}{2} \rceil, \lceil \frac{k+1}{2} \rceil)), \quad \forall k \geq 3. \quad \square$$

Example 1.2 suggests that $P(1, 1)$ may have relevant maximality properties for the higher capacity ratios. This is the case within a suitable class of toric domains in \mathbb{C}^n , which we now introduce. Let $S^1 \subset \mathbb{C}$ be the unit circle in the complex plane, and consider the action of the torus $\mathbb{T}^n = S^1 \times \dots \times S^1$ on \mathbb{C}^n given by

$$(e^{i\theta_1}, \dots, e^{i\theta_n}) \cdot (z_1, \dots, z_n) = (e^{i\theta_1} z_1, \dots, e^{i\theta_n} z_n).$$

A toric domain is a \mathbb{T}^n -invariant subset of \mathbb{C}^n given by the closure of some non-empty open set. Any domain $\Omega \subset [0, \infty)^n$ is the profile of an associated toric domain

$$X_\Omega := \{(z_1, \dots, z_n) \in \mathbb{C}^n \mid (\pi|z_1|^2, \dots, \pi|z_n|^2) \in \Omega\}.$$

Two classes of toric domains are particularly relevant in symplectic geometry:

- The *convex* toric domains X_Ω , which are those toric domains such that $\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid (|x_1|, \dots, |x_n|) \in \Omega\}$ is compact and convex.
- The *concave* toric domains X_Ω , which are those toric domains whose profile Ω is compact and such that $[0, \infty)^n \setminus \Omega$ is convex.

Polydisks $P(\mathbf{a}) = E(a_1) \times \dots \times E(a_n)$ are examples of convex toric domains, and ellipsoids $E(\mathbf{a})$ are the only toric domains that are simultaneously convex and concave. We stress that “convex” and “concave” here do not refer to the convexity of X_Ω as a subset of \mathbb{C}^n : all convex toric domains are convex subsets of \mathbb{C}^n , but some concave toric domains are convex subsets of \mathbb{C}^n as well. In his Ph.D. thesis [Moa94], Moatty provided combinatorial formulas to compute the Viterbo capacities c_k^\vee of convex or concave toric domains. Analogous formulas were provided by Gutt and Hutchings [GH18] for their capacities c_k^{GH} .

If $B \subset \mathbb{C}^n$ is a smooth star-shaped domain, its boundary ∂B admits a canonical contact form $\lambda|_{\partial B}$, which is the restriction of the 1-form on \mathbb{C}^n given by

$$\lambda = \frac{1}{2} \sum_{j=1}^n (x_j dy_j - y_j dx_j). \quad (1.1)$$

Here, $x_1, y_1, \dots, x_n, y_n$ are the standard Darboux coordinates on \mathbb{C}^n . As pointed out in [GH18], the formulas for convex and concave toric domains hold for all symplectic k -capacities satisfying the following extra condition:

- **(Closed Reeb orbits)** For all smooth star-shaped domains $B \subset \mathbb{C}^n$ such that $(\partial B, \lambda)$ has non-degenerate closed Reeb orbits, $c_k(B)$ is the period of a closed Reeb orbit with Conley-Zehnder index $2k + n - 1$.

Moatty-Gutt-Hutchings’ formulas may be employed to study the global maximizers of the higher capacity ratios over convex and concave toric domains. The combinatorics becomes rather involved. We worked out the details in dimension 4.

Proposition D. *For any symplectic k -capacity on \mathbb{C}^2 satisfying the extra “closed Reeb orbits” assumption, the following points hold.*

- (i) On the space of 4-dimensional concave toric domains, \widehat{c}_k achieves its global maximum on the ellipsoid $E(\lceil \frac{k}{2} \rceil, \lceil \frac{k+1}{2} \rceil)$.
- (ii) On the space of 4-dimensional convex toric domains, \widehat{c}_k achieves its maximum on:
 - the round balls $E(a, a)$, $a > 0$, if $k = 1$,
 - the ellipsoids $E(a, 2a)$ and the polydisks $P(a, a)$, $a > 0$, if $k = 2$,
 - the polydisks $P(a, a)$, $a > 0$, if $k \geq 3$.

The situation is largely unexplored in higher dimensions. Let $P_{2n} := P(1, \dots, 1)$ be the $2n$ -dimensional polydisk with unitary parameters, and E_{2n} the global maximizer of \widehat{c}_k over the space of $2n$ -dimensional ellipsoids (see Proposition C(ii)). Simple computations show that there exists a minimal integer $\kappa(n) \geq 2$ such that

$$\widehat{c}_k(P_{2n}) \geq \widehat{c}_k(E_{2n}), \quad \forall k \geq \kappa(n).$$

Moreover, $\kappa(n) \rightarrow \infty$ as $n \rightarrow \infty$. This raises the following questions.

Question E. For each dimension $2n \geq 4$ and for each $k \geq \kappa(n)$, is the polydisk P_{2n} a global maximizers of the k -capacity ratios \widehat{c}_k over the space of $2n$ -dimensional convex toric domains?

Question F. In some dimension $2n \geq 4$, do there exist arbitrarily large integers k and $2n$ -dimensional convex bodies $B \subset \mathbb{C}^n$ such that $\widehat{c}_k(B) > \widehat{c}_k(P_{2n})$?

1.4. Outline of the proof of Theorem A. The local Viterbo conjecture was established as a consequence of a systolic characterization of Zoll contact manifolds, which we now briefly recall. Let (N, α) be a closed contact manifold, X its Reeb vector field defined by $\alpha(X) \equiv 1$ and $d\alpha(X, \cdot) \equiv 0$, and $\phi^t : N \rightarrow N$ its Reeb flow. The Weinstein conjecture, which is a theorem for several classes of closed contact manifolds including all 3-dimensional ones, asserts that the Reeb flow ϕ^t has at least one closed orbit. If this is the case, we denote by $\tau_1(N, \alpha) = \inf\{t > 0 \mid \text{fix}(\phi^t) \neq \emptyset\}$ the systole, namely the minimum among the periods of the closed Reeb orbits; if the Weinstein conjecture fails for (N, α) , we instead set $\tau_1(N, \alpha) = 0$. We consider the associated systolic ratio

$$\widehat{\tau}_1(N, \alpha) = \frac{\tau_1(N, \alpha)}{\text{vol}(N, \alpha)^{1/n}}, \quad (1.2)$$

where $n = \frac{1}{2}(\dim(N) + 1)$, and $\text{vol}(N, \alpha)$ is the integral of $\alpha \wedge (d\alpha)^{n-1}$. On the space of contact forms on N endowed with the C^3 -topology, the local maximizers of the function $\alpha \mapsto \widehat{\tau}_1(N, \alpha)$ are precisely the *Zoll* contact forms: those contact forms whose associated Reeb orbits are all closed and have the same minimal period (namely $\phi^{\tau_1(N, \alpha)} = \text{id}$ and $\text{fix}(\phi^t) = \emptyset$ for all $t \in (0, \tau_1(N, \alpha))$). This remarkable theorem was established in full generality by Abbondandolo and Benedetti [AB19], and previously in dimension 3 in a series of papers [APB14, ABHS18, BK21].

A closed contact manifold (N, α) , or just the contact form α , is called *Besse* when all its Reeb orbits are closed, and in this case they automatically have a common period according to a theorem of Wadsley [Wad75]. The simplest examples of Besse closed contact manifolds are the boundaries of the rational ellipsoids. Actually, in dimension 4, for each smooth star-shaped domain $B \subset \mathbb{C}^2$ with a Besse boundary there exists a symplectomorphism $\psi \in \text{Symp}(\mathbb{C}^n, \omega)$ such that $\psi(B)$ is a rational ellipsoid; this is a consequence of the classification of 3-dimensional Besse contact

spheres [MR23, Theorem 1.1], together with a straightforward generalization of [ABHS18, Prop. 4.3].

Assume now that (N, α) is a closed contact 3-manifold. For each $k \geq 1$, we denote by $\tau_k(N, \alpha)$ the infimum of the values $\tau > 0$ such that there exist at least k closed Reeb orbits (counting iterates as well) of period less than τ ; in formulas:

$$\tau_k(N, \alpha) := \inf \left\{ \tau > 0 \mid \sum_{t \in (0, \tau)} \#(\text{fix}(\phi^t)/\sim) \geq k \right\},$$

where \sim is the equivalence relation identifying $z \sim \phi^t(z)$ for all $z \in N$ and $t \in \mathbb{R}$. Extending the notation (1.2), we write

$$\widehat{\tau}_k(N, \alpha) := \frac{\tau_k(N, \alpha)}{\text{vol}(N, \alpha)^{1/2}}.$$

If (N, α) is Besse, we denote by $k_1 := k_1(N, \alpha)$ the minimal positive integer such that τ_{k_1} is a common period of the Reeb orbits, i.e. $\phi^{\tau_{k_1}} = \text{id}$. This notation is consistent with the one employed in Section 1.2 with 4-dimensional rational ellipsoids E : we have $k_1(E) = k_1(\partial E, \alpha)$. A result of Abbondandolo, Lange, and Mazzucchelli [ALM22] generalized the above mentioned systolic characterization of Zoll contact 3-manifolds as follows: on the space of contact forms on a closed 3-manifold N , endowed with the C^3 topology, the local maximizers of the function $\alpha \mapsto \widehat{\tau}_k(N, \alpha)$ are precisely the Besse contact forms α with $k_1(N, \alpha) = k$.

Consider now a $2n$ -dimensional smooth star-shaped domain B . Stokes theorem implies that $\text{vol}(B, \omega) = \text{vol}(\partial B, \lambda)$. Moreover, if B is a smooth convex body and c_1 denotes any of the above mentioned 1-capacities c_1^{EH} , c_1^{V} , or c_1^{GH} , we have $c_1(B) = \tau_1(\partial B, \lambda)$, as it was proven by Sikorav [Sik90], Viterbo [Vit89], and Abbondandolo and Kang [AK22] for the three respective capacities. Therefore $\widehat{c}_1(B) = \widehat{\tau}_1(\partial B, \lambda)$, and the systolic Zoll characterization implies the local Viterbo conjecture.

The proof of Theorem A employs the Clarke action functional associated with a smooth convex body B , whose set of critical values is precisely the action spectrum $\sigma(\partial B)$, that is, the set of periods of the closed Reeb orbits of ∂B . Ekeland and Hofer introduced spectral invariants $s_k(B)$, which are critical values of the Clarke action functional selected by suitable min-max procedures. A result of Ginzburg, Gürel, and Mazzucchelli [GGM21] asserts that a $2n$ -dimensional smooth convex body B_0 has Besse boundary if and only if $s_k(B_0) = s_{k+n-1}(B_0)$ for some $k \geq 1$. As a first crucial ingredient for the proof of Theorem A, we show that, for any smooth convex body B_1 sufficiently C^2 -close to such a B_0 , the spectral invariant $s_{k_1}(B_1)$ is the minimum among the element in the action spectrum $\sigma(\partial B_1)$ that are close to $s_{k_1}(B_0)$ (Lemma 2.2).

Assume now that B_0 as above has dimension $2n = 4$. We denote by $K(B_0)$ the collection of values k such that $s_k(B_0) = s_{k+1}(B_0)$, and denote by $k_m = k_m(B_0)$ the m -th smallest element of $K(B_0)$. Once again, this notation agrees with the one employed in Section 1.2 for the rational ellipsoids. For each $m \geq 1$, Lemma 2.2 together with some Morse theory for the Clarke action functional implies that $s_{k_m}(B_1) \leq m\tau_{k_1}(B_1)$ for all smooth convex bodies B_1 sufficiently C^2 -close to B_0 , and equality holds if $B_1 = B_0$ (Lemma 2.3). Therefore, by the above mentioned result of Abbondandolo, Lange, and Mazzucchelli [ALM22], we conclude in Theorem 2.4 that any such B_0 is a local maximizer, with respect to the C^3

topology, of the spectral ratios

$$\widehat{s}_{k_m}(B) := \frac{s_{k_m}(B)}{\text{vol}(B)^{1/2}}.$$

It is well known that $s_1(B) = \tau_1(\partial B, \lambda) = c_1^{\text{EH}}(B)$ for all $2n$ -dimensional smooth convex bodies B . Conjecturally, $s_k(B) = c_k^{\text{EH}}(B)$ for all $k \geq 2$ as well. We shall show that at least the inequality $c_k^{\text{EH}}(B) \leq s_k(B)$ holds for all $k \geq 2$ (Proposition 3.1). Moreover, the equality $c_k^{\text{EH}}(B) = s_k(B)$ holds for all $k \geq 2$ and for all 4-dimensional Besse convex bodies, using the already mentioned fact that every such B is symplectomorphic to a rational ellipsoid. By employing these facts together with Theorem 2.4, we infer that the Besse 4-dimensional smooth convex bodies B_0 with associated integers $k_m = k_m(B_0)$ are local maximizers, with respect to the C^3 topology, of the capacity ratios $\widehat{c}_k^{\text{EH}}$.

Finally, building on a well-known argument originally due to Alvarez Paiva and Balacheff [APB14], we show that any local maximizer B of the capacity ratios $\widehat{c}_k^{\text{EH}}$ over the space of $2n$ -dimensional smooth star-shaped domains must have Besse boundary, and the Reeb orbits therein must have common period $c_k(B)$. In particular every such B is symplectomorphic to a rational ellipsoid E that is a local maximizer of $\widehat{c}_k^{\text{EH}}$, and our sharp result for ellipsoids (Proposition C(i)) implies that $k \in K(E)$.

1.5. Organization of the paper. In Section 2, after recalling the general properties of the Clarke action functional, we prove that Besse convex bodies are local maximizers of suitable spectral ratios (Theorem 2.4). In Section 3 we prove that the Ekeland-Hofer capacities of smooth convex bodies are bounded from above by the corresponding spectral invariants of the Clarke action functional (Proposition 3.1), and employ this result and Theorem 2.4 to prove Theorem A. The proof of Proposition 3.1 requires some subtle properties of the Fadell-Rabinowitz index that are either stated under slightly different assumptions or not explicitly stated in the literature, and we included the details in Appendix A. In Section 4 we prove Proposition C. Finally, in Section 5, we prove Proposition D.

2. THE CLARKE ACTION FUNCTIONAL

2.1. Smooth convex bodies. Let $B \subset \mathbb{C}^n$ be a $2n$ -dimensional smooth convex body, and as usual we assume that B contains the origin in its interior. We consider the Hamiltonian $h : \mathbb{C}^2 \rightarrow [0, \infty)$ that is 2-homogeneous and is identically equal to 1 on the boundary ∂B , i.e.

$$h(cz) = c^2, \quad \forall c \geq 0, z \in \partial B.$$

Its Hamiltonian vector field is defined as usual by $\omega(X_h, \cdot) = dh$, where $\omega = d\lambda$ is the standard symplectic form of \mathbb{C}^n , and λ is its primitive (1.1). The Hamiltonian flow ϕ_h^t is 1-homogeneous, i.e. $\phi_h^t(cz) = c\phi_h^t(z)$, and its restriction $\phi_h^t|_{\partial B}$ coincides with the Reeb flow of ∂B (with respect to the canonical contact form $\lambda|_{\partial B}$). In particular, there is a one-to-one correspondence between closed Reeb orbits of ∂B and cylinders of periodic orbits of ϕ_h^t . The action spectrum $\sigma(\partial B)$ is the set of periods of the closed Reeb orbits, i.e.

$$\sigma(\partial B) = \{t > 0 \mid \text{fix}(\phi_h^t) \neq \emptyset\}.$$

Since the Hamiltonian h has positive definite Hessian everywhere outside the origin, it admits a Legendre dual $h^* : \mathbb{C}^n \rightarrow [0, \infty)$ given by

$$h^*(w) = \max_{z \in \mathbb{C}^n} (\langle w, z \rangle - h(z)), \quad (2.1)$$

which is also 2-homogeneous. We set $S^1 = \mathbb{R}/\mathbb{Z}$, and consider the Hilbert space

$$L_0^2(S^1, \mathbb{C}^n) = \left\{ u \in L^2(S^1, \mathbb{C}^n) \mid \int_{S^1} u(t) dt = 0 \right\}.$$

Its elements are precisely the maps of the form $u = \dot{\gamma}$, where $\gamma \in W^{1,2}(S^1, \mathbb{C}^n)$. We consider the functionals $\mathcal{A} : L_0^2(S^1, \mathbb{C}^n) \rightarrow \mathbb{R}$ and $\mathcal{H} : L_0^2(S^1, \mathbb{C}^n) \rightarrow [0, \infty)$, given by

$$\mathcal{A}(\dot{\gamma}) = \frac{1}{2} \int_{S^1} \langle J\gamma(t), \dot{\gamma}(t) \rangle dt, \quad \mathcal{H}(\dot{\gamma}) = \int_{S^1} h^*(-J\dot{\gamma}(t)) dt.$$

Notice that the expression of \mathcal{A} involves a primitive of $\dot{\gamma}$, but is actually independent of the choice of such a primitive. The functional \mathcal{A} is a non-degenerate quadratic form, and we consider its positive open cone $\mathcal{A}^{-1}(0, \infty)$. The circle S^1 acts on it by translation, i.e. $t \cdot u = u(t + \cdot)$ for all $t \in S^1$ and $u \in \mathcal{A}^{-1}(0, \infty)$, and both \mathcal{A} and \mathcal{H} are S^1 -invariant.

The *Clarke action functional* $\tilde{\Psi} : \mathcal{A}^{-1}(0, \infty) \rightarrow (0, \infty)$ is defined by

$$\tilde{\Psi}(u) = \frac{\mathcal{H}(u)}{\mathcal{A}(u)}.$$

Let us summarize its properties (and refer the reader to Ekeland-Hofer's [EH87] for a proof of the non-trivial ones):

- **(\mathbb{C}_* -invariance)** It is 0-homogeneous and S^1 -invariant, i.e. $\tilde{\Psi}(cu) = \tilde{\Psi}(u) = \tilde{\Psi}(t \cdot u)$ for all $c > 0$, $t \in S^1$, and $u \in \mathcal{A}^{-1}(0, \infty)$. Overall, $\tilde{\Psi}$ is \mathbb{C}_* -invariant, where $\mathbb{C}_* = \mathbb{C} \setminus \{0\} \cong (0, \infty) \times S^1$.
- **(Regularity)** It has the same regularity as \mathcal{H} : it is $C^{1,1}$ and admits a Hessian in the sense of Gateaux at every point.
- **(Clarke variational principle)** There is a one-to-one correspondence

$$\begin{array}{ccc} \text{crit}(\tilde{\Psi}) \cap \tilde{\Psi}^{-1}(\tau) & \xleftrightarrow{1:1} & \text{fix}(\phi_h^\tau) \\ u_z & \longleftrightarrow & z \end{array}$$

where $u_z(t) = \frac{d}{dt} \phi_H^{\tau t}(z)$ and, conversely, $z = \tau^{-1} \nabla h^*(-Ju_z(0))$.

- **(Morse indices)** The Morse index and nullity³ of any critical point $u_z \in \text{crit}(\tilde{\Psi}) \cap \tilde{\Psi}^{-1}(\tau)$ are finite. Indeed, the Morse index is equal (up to a conventional additive constant) to the Maslov index of the associated τ -periodic orbit $t \mapsto \phi_h^t(z)$, and the nullity is equal to $\dim \ker(d\phi_h^\tau(z) - I) - 2$.
- **(Palais-Smale condition)** Being 0-homogeneous, $\tilde{\Psi}$ cannot satisfy the Palais-Smale condition. Nevertheless, let V be the rescaled version of the anti-gradient of $\tilde{\Psi}$ given by

$$V(u) := -\|u\|_{L^2} \nabla \tilde{\Psi}(u),$$

³Here, we keep into account the \mathbb{C}_* -invariance of $\tilde{\Psi}$: the nullity of a critical point u of $\tilde{\Psi}$ is defined as $\dim \ker(d^2 \tilde{\Psi}(u)) - 2$.

whose flow $g_t : \mathcal{A}^{-1}(0, \infty) \rightarrow \mathcal{A}^{-1}(0, \infty)$ is 1-homogeneous and preserves the L^2 norm, i.e. $\frac{d}{dt} \|g_t(u)\|_{L^2} = 0$. On an L^2 -sphere of any radius $r > 0$, the Clarke action functional $\tilde{\Psi}$ satisfies the Palais-Smale condition with respect to V : any sequence $u_k \in \mathcal{A}^{-1}(0, \infty)$ such that $V(u_k) \rightarrow 0$, $\|u_k\|_{L^2} = r$, and $\sup \tilde{\Psi}(u_k) < \infty$ admits a converging subsequence.

Due to its 0-homogeneity, in the literature the Clarke action functional $\tilde{\Psi}$ usually appears restricted to S^1 -invariant hypersurfaces of $\mathcal{A}^{-1}(0, \infty)$ transverse to the radial directions: for instance $\mathcal{A}^{-1}(1)$ or $\mathcal{H}^{-1}(1)$. In this paper, we equivalently restrict $\tilde{\Psi}$ to the smooth hypersurface $\Lambda = \{u \in \mathcal{A}^{-1}(0, \infty) \mid \|u\|_{L^2} = 1\}$, and denote the restriction by

$$\Psi := \tilde{\Psi}|_{\Lambda}.$$

We still call Ψ the Clarke action functional. Clearly, Ψ is S^1 -invariant, and we have $S^1 \cdot u \subset \text{crit}(\Psi) \cap \Psi^{-1}(\tau)$ if and only if $\mathbb{C}_* \cdot u \subset \text{crit}(\tilde{\Psi}) \cap \tilde{\Psi}^{-1}(\tau)$. Overall, Ψ satisfies the classical properties required by S^1 -equivariant Morse theory, except for its lack of C^2 regularity, which nevertheless will be easily circumvented later on.

2.2. Spectral invariants. In their seminal work [EH87], Ekeland and Hofer introduced and studied spectral invariants for the Clarke action functional. We briefly recall the construction.

Consider the subspace Λ_+ given by all $u \in \Lambda$ having the form

$$u(t) = \sum_{k=1}^{\infty} e^{i2\pi kt} u_k,$$

for some $u_k \in \mathbb{C}^n$. Notice that Λ_+ is the unit sphere of an infinite dimensional Hilbert subspace of $L_0^2(S^1, \mathbb{C}^n)$, and is S^1 -invariant. Since Λ_+ is contractible, the Gysin sequence of the S^1 -bundle $\Lambda_+ \times ES^1 \rightarrow \Lambda_+ \times_{S^1} ES^1$ readily implies that its S^1 -equivariant cohomology with rational coefficients is given by $H_{S^1}^*(\Lambda_+; \mathbb{Q}) = \mathbb{Q}[e]$, where e is a generator of $H_{S^1}^2(\Lambda_+; \mathbb{Q}) \cong \mathbb{Q}$. From now on, all cohomology rings will be assumed to have rational coefficients, and we will suppress \mathbb{Q} from the notation. The inclusion $\Lambda_+ \hookrightarrow \Lambda$ is an S^1 -equivariant homotopy equivalence, and therefore induces a ring isomorphism in S^1 -equivariant cohomology. Summing up, we have

$$H_{S^1}^*(\Lambda) = \mathbb{Q}[e],$$

where e now denotes a generator of $H_{S^1}^2(\Lambda) \cong \mathbb{Q}$. In particular, $e^k \neq 0$ in $H_{S^1}^*(\Lambda)$ for all $k \geq 0$; in terms of the Fadell-Rabinowitz index (see Appendix A), this means $\text{ind}_{\text{FR}}(\Lambda) = \infty$.

For each $k \geq 1$, the k -th spectral invariant of the smooth convex body B is defined by

$$s_k(B) := \inf \{c > 0 \mid \text{ind}_{\text{FR}}(\{\Psi < c\}) \geq k\},$$

where $\{\Psi < \tau\}$ denotes the sublevel set $\Psi^{-1}(0, \tau)$. We recall some classical properties of these values:

- **(Spectrality)** Every $s_k(B)$ is a critical value of the Clarke action functional associated with B , that is, an element of the action spectrum $\sigma(\partial B)$.
- **(Systole)** $s_1(B) = \min \Psi = \sigma(\partial B)$.

- **(Lusternik-Schnirelmann)** For all $k \geq 1$ we have $s_k(B) \leq s_{k+1}(B)$. If $s_k(B) = s_{k+j}(B)$ for some $j \geq 1$, then $\text{ind}_{\text{FR}}(\mathcal{U}) > j$ for any S^1 -invariant neighborhood $\mathcal{U} \subset \Lambda$ of $\text{crit}(\Psi) \cap \Psi^{-1}(s_k(B))$.
- **(C^0 continuity)** For any sequence of smooth convex bodies B_j converging to a smooth convex body B in the C^0 topology, we have $s_k(B_j) \rightarrow s_k(B)$.

2.3. Finite dimensional reduction. The $C^{1,1}$ regularity of the Clarke action functional Ψ is not sufficient to apply those results of Morse theory involving the Hessian, such as the Morse lemma. Nevertheless, Ψ becomes smooth as soon as one restricts it to the subspace of $W^{1,2}$ curves that do not go through the origin of \mathbb{C}^n . With such restriction, however, one loses compactness properties such as the Palais-Smale condition. In order to both improve the regularity of Ψ and retain the compactness properties, we will apply the finite dimensional reduction introduced by Ekeland and Hofer, which we now briefly recall. We refer the reader to [EH87, Sec. II.2] for the proofs.

Let $b > \min \Psi$ be a fixed value, and let us focus on the sublevel set $\{\Psi < b\}$ of the Clarke action functional. For each integer $N \geq 1$, let $F = F_N \subset L_0^2(S^1, \mathbb{C}^n)$ be the S^1 -invariant finite dimensional Hilbert subspace given by those $u \in L_0^2(S^1, \mathbb{C}^n)$ of the form

$$u(t) = \sum_{0 < |k| \leq N} e^{i2\pi kt} u_k,$$

where $u_k \in \mathbb{C}^n$. We consider the open subset $V = \{u \in F \mid W_u \neq \emptyset\}$, where

$$W_u := \{v \in F^\perp \mid \mathcal{A}(u+v) > 0, \tilde{\Psi}(u+v) < b\}.$$

The function $W_u \rightarrow (0, \infty)$, $v \mapsto \tilde{\Psi}(u+v)$ has a unique non-degenerate global minimizer $\nu(u) \in F^\perp$. We denote by $S \subset F$ the unit sphere of the Hilbert space F , and set $U := V \cap S$. The *reduced Clarke action functional* is defined as

$$\psi : U \rightarrow (0, b), \quad \psi(u) = \tilde{\Psi}(u + \nu(u)) = \Psi(\iota(u)),$$

where

$$\iota : U \hookrightarrow \Lambda, \quad \iota(u) := \frac{u + \nu(u)}{\|u + \nu(u)\|_{L^2}}.$$

The properties of this finite dimensional setting are the following.

- **(S^1 -invariance)** The map ν is \mathbb{C}_* -equivariant, and therefore the reduced Clarke action functional ψ is S^1 -invariant.
- **(Critical points)** Both ν and ψ are $C^{1,1}$, and there is a one-to-one correspondence

$$\begin{aligned} \text{crit}(\psi) &\xleftrightarrow{1:1} \text{crit}(\Psi) \cap \Psi^{-1}(0, b) \\ u &\longleftrightarrow \iota(u). \end{aligned}$$

Both ν , ψ , and the S^1 action are smooth on a sufficiently small neighborhood of the critical set $\text{crit}(\psi)$. Moreover, at every $u \in \text{crit}(\psi)$, the Hessians $d^2\psi(u)$ and $d^2\Psi(\iota(u))$ have the same Morse index and the same nullity.

- **(Compactness)** For each $c \in (0, b)$, the closed sublevel set $\{\psi \leq c\}$ is compact.

- **(Approximation)** For each $a \in (0, b)$, the map $\iota : \{\psi < a\} \hookrightarrow \{\Psi < a\}$ is an S^1 -equivariant homotopy equivalence.

2.4. Besse convex bodies. We say that a $2n$ -dimensional smooth convex body B is *Besse* when its boundary ∂B , equipped with the canonical contact form $\lambda|_{\partial B}$ of Equation (1.1), is a Besse contact manifold: all its Reeb orbits are closed, and therefore have a common period according to a theorem of Wadsley [Wad75]. A theorem of Ginzburg, Gürel, and Mazzucchelli [GGM21] implies that the Besse property is detected by the spectral invariants:

- **(Spectral characterization)** We have $c := s_k(B) = s_{k+n-1}(B)$ for some $k \geq 1$ if and only if B is Besse, and c is a common period for the Reeb orbits on ∂B (i.e. $\phi_h^c = \text{id}$, where $h : \mathbb{C}^n \rightarrow [0, \infty)$ is the 2-homogeneous Hamiltonian such that $h^{-1}(1) = \partial B$). In this case, the critical manifold $\text{crit}(\Psi) \cap \Psi^{-1}(c)$ has Morse index $2k - 2$ and nullity $2n - 2$. Moreover, $c < s_{k+n}(B)$ and, when $k \geq 2$, $c > s_{k-1}(B)$.

The Clarke action functional Ψ associated with a Besse convex body B also satisfies the following properties, which were established by Mazzucchelli and Radeschi [MR23]. Analogous properties in the different setting of geodesic flows were established earlier by Radeschi and Wilkings [RW17].

- **(Critical manifolds)** For each $c \in \sigma(\partial B)$, the critical manifold

$$\mathcal{K} := \text{crit}(\Psi) \cap \Psi^{-1}(c) \cong \text{fix}(\phi_h^c|_{\partial B})$$

is a non-degenerate (i.e. $\ker d^2\Psi(u) = T_u\mathcal{K}$ for all $u \in \mathcal{K}$), connected, odd dimensional, rational homology sphere.

- **(Perfectness)** The Clarke action functional Ψ is perfect for the S^1 -equivariant Morse theory with rational coefficients: for each $a < b$ the inclusions induce short exact sequences

$$0 \rightarrow H_{S^1}^*(\{\Psi < b\}, \{\Psi < a\}) \rightarrow H_{S^1}^*(\{\Psi < b\}) \rightarrow H_{S^1}^*(\{\Psi < a\}) \rightarrow 0.$$

Remark 2.1. For a rational ellipsoid E (which is a Besse convex body) the perfectness of the associated Clarke action functional, together with the Lusternik-Schnirelmann property, readily implies that $s_k(E) = c_k^{\text{EH}}(E)$ for all $k \geq 1$. Since $\mathbf{a} \mapsto s_k(E(\mathbf{a}))$ and $\mathbf{a} \mapsto c_k^{\text{EH}}(E(\mathbf{a}))$ are continuous, the latter identity holds for irrational ellipsoids as well. \square

In order to prove our main Theorem A, we shall prove analogous statements for the spectral invariants $s_k(B)$: Theorems 2.4 and, ultimately, 3.11. A crucial ingredient for these statements is the following perturbation result for the spectral invariants. We recall that two $2n$ -dimensional smooth convex bodies B_0 and B_1 are C^k -close if their boundaries are C^k -close embedded hypersurfaces of \mathbb{C}^n .

Lemma 2.2. *Let B_0 be a $2n$ -dimensional Besse convex body, $k \geq 1$ an integer such that $s_k(B_0) = s_{k+n-1}(B_0)$, and $[a, b]$ a compact neighborhood of $s_k(B_0)$ such that $[a, b] \cap \sigma(\partial B_0) = s_k(B_0)$. Then, for any smooth convex body B_1 that is sufficiently C^2 -close to B_0 , we have*

$$s_k(B_1) = \min(\sigma(\partial B_1) \cap [a, b]).$$

Proof of Lemma 2.2. The statement is straightforward when $k = 1$. Indeed, $B \mapsto s_k(B)$ is a continuous function with respect to the C^0 topology on the space of

smooth convex bodies, and $s_1(B) = \min \sigma(\partial B)$ for each smooth convex body B . From here on, we shall assume $k \geq 2$.

Let $h_0 : \mathbb{C}^n \rightarrow [0, \infty)$ be the 2-homogeneous Hamiltonian such that $h_0^{-1}(1) = \partial B_0$, and $\Psi_0 : \Lambda \rightarrow (0, \infty)$ the corresponding Clarke action functional. Consider the critical value $c := s_k(B_0) = s_{k+n-1}(B_0)$. The spectral Besse characterization implies that the critical manifold $\mathcal{K} := \text{crit}(\Psi_0) \cap \Psi_0^{-1}(c)$ is diffeomorphic to ∂B_0 , non-degenerate, and with Morse index $d := 2k - 2$.

Let $[a, b]$ be a compact neighborhood of c such that $[a, b] \cap \sigma(\partial B_0) = c$, and consider the reduced Clarke action functional $\psi_0 = \Psi_0 \circ \iota_0 : U_0 \rightarrow (0, \infty)$, where the reduction is chosen so that ψ_0 approximates Ψ_0 up to a level larger than b (see Section 2.3). We denote by $K := \iota_0^{-1}(\mathcal{K}) = \text{crit}(\psi_0) \cap \psi_0^{-1}(c)$ the corresponding critical manifold of ψ_0 , and recall that K and \mathcal{K} have the same Morse index d . The normal bundle $N \subset TU_0|_K$ of K admits an S^1 -invariant splitting $N = N^+ \oplus N^-$ such that $\pm d^2 \psi_0(u)$ is positive definite on N_u^\pm for each $u \in K$. The rank of N^- is the Morse index d . For each $u \in K$ and $r > 0$, we denote by $D_u^\pm(r) \subset N_u^\pm$ the closed ball of radius r centered at the origin, measured with the Riemannian metric of U_0 . The union $D^\pm(r) := \cup_{u \in K} D_u^\pm(r)$ is the closed neighborhood of radius r of the zero-section. By the Morse-Bott lemma, there exists a closed S^1 -invariant neighborhood $D \subset U_0$ of K small enough so that $\psi_0|_D$ is smooth, and an S^1 -equivariant diffeomorphism identifying $D \equiv D^+(r_+) \oplus D^-(r_-)$, for some $r_- > r_+ > 0$, so that K is identified with the zero-section and

$$\psi_0|_D(u; x, y) = c + \|x\|^2 - \|y\|^2, \quad \forall u \in K, \quad x \in D_u^+(r_+), \quad y \in D_u^-(r_-).$$

In these local coordinates, we have $\partial_{yy}^2 \psi_0|_D \equiv -I$. We require the radius r_+ to be small enough so that

$$\max_D \psi_0 = c + r_+^2 < b.$$

From now on, in order to simplify the notation, we simply write $D^\pm = D^\pm(r_\pm)$.

Since B is Besse, Ψ_0 is perfect for the S^1 -equivariant Morse theory with rational coefficients, and so is ψ_0 thanks to its approximation property (see Section 2.3). Therefore, the embedding $\iota_0 : U_0 \rightarrow \Lambda$ induces an isomorphism

$$\iota_0^* : H_{S^1}^d(\Lambda, \{\Psi_0 < c\}) \xrightarrow{\cong} H_{S^1}^d(D^-, \partial D^-).$$

Since K is simply connected, the negative normal bundle $N^- \rightarrow K$ is orientable, and therefore the inclusion and Thom isomorphism give

$$H_{S^1}^d(D^-, \partial D^-) \xrightarrow{\cong} H_{S^1}^d(D^-|_{S^1 \cdot u}, \partial D^-|_{S^1 \cdot u}) \cong H_{S^1}^0(S^1 \cdot u) \cong \mathbb{Q}, \quad \forall u \in K.$$

We consider the product $S^1 \times D_u^-$ equipped with the S^1 -action inherited from the first factor. For each $u \in K$ and $x \in D_u^+$, we consider the S^1 -equivariant map

$$j_x : S^1 \times D_u^- \rightarrow D, \quad j_x(t, y) = t \cdot (u; x, y),$$

which depends continuously on x . For $x = 0$, the map j_0 is an m -fold covering map onto $D^-|_{S^1 \cdot u}$, where $m \geq 1$ is the order of iteration of u , i.e. the largest integer m such that $\frac{1}{m} \cdot u = u$. Moreover, $\psi_0 \circ j_x|_{S^1 \circ \partial D_u^-} \leq c + r_+^2 - r_-^2 < c$. Therefore, the compositions $\iota_{0,x} := \iota_0 \circ j_x$ induce the isomorphisms

$$\iota_{0,x}^* = \iota_{0,0}^* : H_{S^1}^d(\Lambda, \{\Psi_0 < c\}) \xrightarrow{\cong} H_{S^1}^d(S^1 \times D_u^-, S^1 \times \partial D_u^-). \quad (2.2)$$

We fix $\epsilon > 0$ small enough so that

- (i) $[(1 - \epsilon)^3 c, (1 - \epsilon)^{-2} c] \subset [a, b]$.
- (ii) $c + r_+^2 - r_-^2 < (1 - \epsilon)^2 c$.

We consider a 2-homogeneous Hamiltonian $h_1 : \mathbb{C}^n \rightarrow [0, \infty)$ such that

$$\delta := \|h_1 - h_0\|_{C^2(B_0 \setminus \{0\})}$$

is sufficiently small. Namely, $B_1 := h_1^{-1}[0, 1]$ is a convex body C^2 -close to B_0 . For each $s \in [0, 1]$, the convex combinations $h_s := sh_1 + (1 - s)h_0$ define the convex bodies $B_s := h_s^{-1}[0, 1]$ interpolating between B_0 and B_1 . We require $\delta \leq \epsilon$, so that

$$(1 - \epsilon)h_s \leq h_0 \leq (1 - \epsilon)^{-1}h_s, \quad \forall s \in [0, 1], \quad (2.3)$$

We denote by $\Psi_s : \Lambda \rightarrow (0, \infty)$ the Clarke action functional associated with h_s , and $\psi_s = \Psi_s \circ \iota_s : U_s \rightarrow (0, \infty)$ its finite dimensional reduction. Up to choosing δ small enough, we can insure that all the domains U_s are open subsets of the same finite dimensional vector subspace of $L_0^2(S^1; \mathbb{C}^n)$ and contain the compact set D . The dual Hamiltonians h_s^* satisfy inequalities analogous to (2.3), and so do the Clarke action functionals Ψ_s , i.e.

$$(1 - \epsilon)\Psi_s \leq \Psi_0 \leq (1 - \epsilon)^{-1}\Psi_s, \quad \forall s \in [0, 1]. \quad (2.4)$$

We require δ to be small enough so that, for each $s \in [0, 1]$, we have:

- (iii) $\sigma(\partial B_s) \cap [a, b] = \sigma(\partial B_s) \cap [(1 - \epsilon)c, (1 - \epsilon)^{-1}c]$;
- (iv) $\psi_s|_D$ is smooth and C^2 -close to ψ_0 , and in particular $\psi_s|_D \leq (1 - \epsilon)^{-1}\psi_0|_D$, and $\partial_{yy}^2 \psi_s$ is negative definite at all points of D .
- (v) $\text{crit}(\psi_s) \cap \psi_s^{-1}[a, b] \subset D$.

The inequalities (2.4) imply that we have the inclusions of sublevel sets

$$\{\Psi_s < (1 - \epsilon)^3 c\} \hookrightarrow \{\Psi_0 < (1 - \epsilon)^2 c\} \hookrightarrow \{\Psi_s < (1 - \epsilon)c\} \hookrightarrow \{\Psi_0 < c\}. \quad (2.5)$$

Notice that $[(1 - \epsilon)^3 c, (1 - \epsilon)c]$ is an interval of regular values of Ψ_s , and $[(1 - \epsilon)^2 c, c]$ is an interval of regular values of Ψ_0 , according to properties (i) and (iii). Therefore, the composition of any two subsequent inclusions in (2.5) is an S^1 -equivariant homotopy equivalence. This implies that each of the inclusions in (2.5) induces an isomorphism in S^1 -equivariant cohomology, in particular the last one:

$$H_{S^1}^*(\{\Psi_0 < c\}) \xrightarrow{\cong} H_{S^1}^*(\{\Psi_s < (1 - \epsilon)c\}). \quad (2.6)$$

The inequalities (2.4) and property (i) further imply that we have inclusions

$$\{\Psi_0 \leq c\} \hookrightarrow \{\Psi_s \leq (1 - \epsilon)^{-1}c\} \hookrightarrow \{\Psi_0 \leq b\},$$

and since their composition is an S^1 -equivariant homotopy equivalence, the second inclusion induces an injective homomorphism

$$H_{S^1}^*(\{\Psi_0 \leq b\}) \hookrightarrow H_{S^1}^*(\{\Psi_s \leq (1 - \epsilon)^{-1}c\}). \quad (2.7)$$

Equations (2.6) and (2.7) imply that

$$s_k(\partial B_1) \in [(1 - \epsilon)c, (1 - \epsilon)^{-1}c].$$

In particular, $[(1 - \epsilon)c, (1 - \epsilon)^{-1}c]$ contains some critical values of the Clarke action functional Ψ_1 , and we denote the smallest one by

$$c_1 := \min(\sigma(\partial B_1) \cap [a, b]) \in [(1 - \epsilon)c, (1 - \epsilon)^{-1}c].$$

For each $s \in [0, 1]$ and $x \in D_u^+$, properties (ii) and (iv) imply

$$\begin{aligned} \psi_s \circ j_x|_{S^1 \times \partial D_u^-} &\leq (1 - \epsilon)^{-1} \psi_0 \circ j_x|_{S^1 \times \partial D_u^-} \\ &\leq (1 - \epsilon)^{-1} (c + r_+^2 - r_-^2) \\ &< (1 - \epsilon)c. \end{aligned}$$

Therefore, if we denote $\iota_{s,x} := \iota_s \circ j_x$, we have

$$\iota_{s,x}(S^1 \times \partial D_u^-) \subset \{\Psi_s < (1 - \epsilon)c\} \subset \{\Psi_0 < c\}.$$

Since $\iota_{s,x}$ depends continuously on s and x , it induces the same isomorphism as in (2.2) in relative cohomology, i.e.

$$\iota_{s,x}^* = \iota_{0,0}^* : H_{S^1}^d(\Lambda, \{\Psi_0 < c\}) \xrightarrow{\cong} H_{S^1}^d(S^1 \times D_u^-, S^1 \times \partial D_u^-). \quad (2.8)$$

We consider a critical circle $S^1 \cdot p \in \text{crit}(\psi_1) \cap \psi_1^{-1}(c_1)$. Property (v) guarantees that $S^1 \cdot p \subset D$, and therefore we can write p as

$$p = (u; x, y) \in D^+ \oplus D^- \equiv D.$$

Property (iv) implies that the restriction $\psi_1|_{\{x\} \oplus D_u^-}$ is a strictly concave function with a unique maximizer at y . Therefore

$$\iota_{1,x}(S^1 \times D_u^-) \subset \{\Psi_1 \leq c_1\}.$$

Therefore, the isomorphism (2.8) factors as

$$\begin{array}{ccc} H_{S^1}^d(\Lambda, \{\Psi_0 < c\}) & \xrightarrow[\cong]{\iota_{s,x}^*} & H_{S^1}^d(S^1 \times D_u^-, S^1 \times \partial D_u^-) \\ \downarrow i^* & \nearrow \iota_{s,x}^* & \\ H_{S^1}^d(\{\Psi_1 \leq c_1\}, \{\Psi_1 \leq (1 - \epsilon)c\}) & & \end{array}$$

where i^* is induced by the inclusion. The diagram implies that i^* is injective. Finally, consider the following commutative diagram, whose rows are parts of long exact sequences of inclusions, and whose vertical homomorphisms are induced by inclusions:

$$\begin{array}{ccccc} H_{S^1}^{d-1}(\{\Psi_0 < c\}) & \xrightarrow{\partial^*=0} & H_{S^1}^d(\Lambda, \{\Psi_0 < c\}) & \xleftarrow{l^*} & H_{S^1}^d(\Lambda) \\ \cong \downarrow & & \downarrow i^* & & \downarrow j^* \\ H_{S^1}^{d-1}(\{\Psi_1 < (1 - \epsilon)c\}) & \xrightarrow{\partial^*} & H_{S^1}^d(\{\Psi_1 \leq c_1\}, \{\Psi_1 < (1 - \epsilon)c\}) & \xrightarrow{m^*} & H_{S^1}^d(\{\Psi_1 \leq c_1\}) \end{array}$$

The cohomology class $e^{k-1} \in H_{S^1}^d(\Lambda)$ belongs to the image of l^* , since $c = s_k(\partial B_0)$. Let $f \in H_{S^1}^d(\Lambda, \{\Psi_0 < c\})$ be the non-zero cohomology class such that $l^*f = e^{k-1}$. The diagram readily implies that the connecting homomorphism ∂^* on the bottom line is the zero homomorphism, namely that m^* is injective. Therefore,

$$j^*e^{k-1} = m^*i^*f \neq 0,$$

which implies $s_k(\partial B_1) \leq c_1$. Since $s_k(\partial B_1) \in [(1 - \epsilon)c, (1 - \epsilon)^{-1}c]$, and since c_1 is the smaller critical value of Ψ_1 in $[(1 - \epsilon)c, (1 - \epsilon)^{-1}c]$, we conclude that $s_k(\partial B_1) = c_1$. \square

2.5. Local maximizers of the spectral ratio. From now on, we will focus on dimension 4. Let B be a 4-dimensional smooth convex body, $h : \mathbb{C}^2 \rightarrow [0, \infty)$ the 2-homogeneous Hamiltonian such that $h^{-1}(1) = \partial B$, and $\Psi : \Lambda \rightarrow (0, \infty)$ the associated Clarke action functional. We denote by $\tau_k(B)$ the infimum of the values $\tau > 0$ such that $\text{crit}(\Psi) \cap \Psi^{-1}(0, \tau)$ contains at least k critical circles. In terms of the Hamiltonian flow ϕ_h^t , this can be expressed as

$$\tau_k(B) := \inf \left\{ \tau > 0 \mid \sum_{t \in (0, \tau]} \# \left(\frac{\text{fix}(\phi_h^t) \setminus \{0\}}{\sim} \right) \geq k \right\},$$

where \sim is the equivalence relation identifying $z \sim c\phi_h^t(z)$ for all $t \in \mathbb{R}$ and $c > 0$.

Assume that B is Besse. We denote by $\tau(B)$ the minimal common period of the closed Reeb orbit on ∂B , namely the minimal $t > 0$ such that $\phi_h^t = \text{id}$. The critical set $\text{crit}(\Psi) \cap \Psi^{-1}(0, \tau(B))$ consists of a finite union of non-degenerate critical circles (indeed, since B is symplectomorphic to an ellipsoid according to the forthcoming Proposition 3.7, at most two critical circles). More generally, for each integer $m \geq 1$, $\text{crit}(\Psi) \cap \Psi^{-1}(m\tau(B), (m+1)\tau(B))$ consists of a finite union of non-degenerate critical circles: the suitable iterates of the critical circles in $\text{crit}(\Psi) \cap \Psi^{-1}(0, \tau(B))$. Since the Clarke action functional Ψ is perfect, every critical value of Ψ is equal to some spectral invariant $s_k(B)$, i.e.

$$\sigma(\partial B) = \{s_k(B) \mid k \geq 1\}.$$

For all integers $m \geq 1$, we define

$$k_m = k_m(B) := \min \{k \geq 1 \mid s_k(B) = m\tau(B)\}.$$

If $k_1 \geq 2$, for each $k \in \{1, \dots, k_1 - 1\}$ the critical set $\text{crit}(\Psi) \cap \Psi^{-1}(s_k(B))$ is a single non-degenerate critical circle. Therefore, the perfectness and the Lusternik-Schinerlmann property of Ψ imply that

$$s_k(B) = \tau_k(B), \quad \forall k \in \{1, \dots, k_1\}. \quad (2.9)$$

Moreover, the spectral characterization of Besse convex bodies (see Section 2.4) implies that $s_{k_m}(B) = s_{k_m+1}(B) < s_{k_m+2}(B)$ and, if $k_m > 1$, $s_{k_m-1}(B) < s_{k_m}(B)$. We define

$$K(B) := \{k_m(B) \mid m \geq 1\}.$$

Lemma 2.3. *Let B_0 be a 4-dimensional Besse convex body, and $m \geq 1$ an integer. For each smooth convex body B_1 that is sufficiently C^2 -close to B_0 , we have*

$$s_{k_m(B_0)}(B_1) \leq m\tau_{k_1(B_0)}(B_1).$$

Proof. Fix an integer $m \geq 1$, and consider $k_1 = k_1(B_0)$ and $k_m = k_m(B_0)$. Let $[a, b] \subset (0, \infty)$ be a compact neighborhood of $s_{k_m}(B_0)$ that is small enough so that

$$\sigma(\partial B_0) \cap [a, b] = \{s_{k_m}(B_0)\}.$$

Let $\epsilon > 0$ be small enough so that $m(s_{k_1}(B_0) - \epsilon) \geq a$. Let Ψ be the Clarke action functional associated with B_0 . Since $\text{crit}(\Psi) \cap \Psi^{-1}(\tau_k(B_0))$ is a non-degenerate critical circle for each $k \in \{1, \dots, k_1 - 1\}$, any smooth convex body B_1 that is sufficiently C^2 -close to B_0 satisfies

$$|\tau_k(B_1) - \tau_k(B_0)| < \epsilon, \quad \forall k \in \{1, \dots, k_1\}.$$

This, together with (2.9), implies $\tau_{k_1}(B_1) > s_{k_1}(B_0) - \epsilon$, and therefore

$$m\tau_{k_1}(B_1) \geq a.$$

By Lemma 2.2, we have

$$s_{k_m}(B_1) = \min(\sigma(\partial B_1) \cap [a, b]).$$

Since $m \tau_{k_1}(B_1) \in \sigma(\partial B_1)$, we conclude $s_{k_m}(B_1) \leq m \tau_{k_1}(B_1)$. \square

We can finally prove the main result of this section. For each $k \geq 1$ and for each 4-dimensional smooth convex body $B \subset \mathbb{C}^n$, we set

$$\widehat{\tau}_k(B) = \frac{\tau_k(B)}{\text{vol}(B)^{1/2}}, \quad \widehat{s}_k(B) = \frac{s_k(B)}{\text{vol}(B)^{1/2}},$$

where the volume $\text{vol}(B) = \text{vol}(B, \omega)$ is obtained by integrating $\omega \wedge \omega$.

Theorem 2.4. *Let B_0 be a 4-dimensional Besse convex body. For each integer $k \in K(B_0)$, any smooth convex body B_1 that is sufficiently C^3 -close to B_0 satisfies $\widehat{s}_k(B_1) \leq \widehat{s}_k(B_0)$.*

Proof. Consider the integers $k_m = k_m(B_0)$ associated with B_0 . The spectral invariants $s_{k_m}(B_0)$ are given by

$$s_{k_m}(B_0) = m \tau_{k_1}(B_0).$$

A theorem of Abbondandolo, Lange, and Mazzucchelli [ALM22], extending an earlier theorem due to Abbondandolo, Bramham, Hryniewicz, and Salomao [ABHS18] for the special case $k_1 = 1$, implies that any smooth convex body B_1 that is sufficiently C^3 -close to B_0 satisfies

$$\widehat{\tau}_{k_1}(B_1) \leq \widehat{\tau}_{k_1}(B_0).$$

Fix an integer $m \geq 1$. By Lemma 2.3, if B_1 is sufficiently C^2 -close to B_0 , we have

$$s_{k_m}(B_1) \leq m \tau_{k_1}(B_1).$$

Overall, we proved that $\widehat{s}_{k_m}(B_1) \leq \widehat{s}_{k_m}(B_0)$ for each smooth convex body B_1 that is sufficiently C^3 -close to B_0 . \square

In Theorem 3.11 at the end of Section 3.3, we will strengthen this theorem and fully characterize the local maximizers of the spectral ratios \widehat{s}_k .

3. THE EKELAND-HOFER CAPACITIES

3.1. Definition of the capacities. We begin by briefly recalling Ekeland and Hofer's construction [EH89, EH90] of their symplectic capacities. The functional setting involves the Sobolev space $E := W^{1/2,2}(S^1, \mathbb{C}^n)$, which is the space of all periodic curves $\gamma : S^1 \rightarrow \mathbb{C}^n$ of the form

$$\gamma(t) = \sum_{k \in \mathbb{Z}} e^{i2\pi kt} \gamma_k,$$

where $\gamma_k \in \mathbb{C}^n$, such that

$$\|\gamma\|_E^2 := \sum_{k \in \mathbb{Z}} |k| \cdot \|\gamma_k\|^2 < \infty.$$

The circle S^1 acts on E by translations, i.e. $t \cdot \gamma = \gamma(t + \cdot)$ for all $t \in S^1$ and $\gamma \in E$. The Sobolev space E admits an orthogonal direct sum decomposition

$$E = E_- \oplus E_0 \oplus E_+, \tag{3.1}$$

where E_{\pm} consists of the γ such that $\gamma_k = 0$ for all $\mp k \geq 0$, and E_0 consists of the constants $\gamma \equiv \gamma_0$. We consider the quadratic form $a : E \rightarrow \mathbb{R}$ given by

$$a(\gamma) = \pi \sum_{k \in \mathbb{Z}} k \|\gamma_k\|^2.$$

On the dense subspace $W^{1,2}(S^1, \mathbb{C}^n) \subset E$, this quadratic form is related to the quadratic form $\mathcal{A} : L_0^2(S^1, \mathbb{C}^n) \rightarrow \mathbb{R}$ employed in the previous section by

$$a(\gamma) = \mathcal{A}(\dot{\gamma}).$$

The Hamiltonians involved in Ekeland-Hofer's setting are non-negative smooth functions of the form $H : \mathbb{C}^n \rightarrow [0, \infty)$ that vanish on some non-empty open subset of \mathbb{C}^n and have the form $H(z) = r\|z\|^2$ outside a compact set, for some $r > \pi$ that is not a multiple of π ; we briefly refer to such H as to *admissible* Hamiltonians. The associated Hamiltonian action functional $\Phi_H : E \rightarrow \mathbb{R}$, given by

$$\Phi_H(\gamma) = a(\gamma) - \int_0^1 H(\gamma(t)) dt,$$

is C^∞ , S^1 -invariant, satisfies the Palais-Smale condition, and its critical circles $S^1 \cdot \gamma$ are precisely the circles of curves of the form $\gamma(t) = \phi_H^t(\gamma(0))$, where ϕ_H^t is the Hamiltonian flow of H . However, unlike for the Clarke action functional, the critical circles of Φ_H have infinite Morse index.

A preliminary step to construct the Ekeland-Hofer capacities consists in defining suitable spectral invariants for Φ_H . Since all Morse indices are infinite, the recipe is not an ordinary min-max, but a more sophisticated linking argument. We write each element of E as $\gamma = \gamma_- + \gamma_0 + \gamma_+$ according to the orthogonal decomposition (3.1). Let \mathcal{D} be the group of those S^1 -equivariant homeomorphisms $\Theta : E \rightarrow E$ of the form

$$\Theta(\gamma) = e^{\theta_-(\gamma)}\gamma_- + \gamma_0 + e^{\theta_+(\gamma)}\gamma_+ + \mathcal{K}(\gamma), \quad (3.2)$$

where $\theta_{\pm} : E \rightarrow \mathbb{R}$ and $\mathcal{K} : E \rightarrow E$ are continuous S^1 -invariant maps that send bounded sets to precompact sets, vanish on the sublevel set $\{a \leq 0\}$, and vanish as well outside a ball of sufficiently large radius in E . Let S_+ be the unit-sphere of the Hilbert subspace $E_+ \subset E$. The *Ekeland-Hofer index* of an S^1 -invariant subset $\mathcal{V} \subset E$ is defined as

$$\text{ind}_{\text{EH}}(\mathcal{V}) = \inf_{\Theta \in \mathcal{D}} \inf_{\mathcal{U}} \text{ind}_{\text{FR}}(\mathcal{U}),$$

where the inner infimum ranges over all S^1 -invariant open subsets $\mathcal{U} \subset E$ containing $\mathcal{V} \cap \Theta(S_+)$. For each integer $k \geq 1$, we define the k -th spectral invariant associated with the Ekeland-Hofer index as

$$c_k^{\text{EH}}(H) := \inf \{c \in \mathbb{R} \mid \text{ind}_{\text{EH}}(\{\Phi_H < c\}) \geq k\}.$$

These values satisfy the following properties:

- **(Finiteness)** We have $c_1^{\text{EH}}(H) \leq c_2^{\text{EH}}(H) \leq \dots \leq c_{nk}^{\text{EH}}(H) < \infty$ for all Hamiltonians $H \in \mathcal{F}(\mathcal{B})$ such that $H(z) > k\pi\|z\|^2$ outside a compact set. From now on, whenever we write $c_k^{\text{EH}}(H)$ we implicitly assume that H satisfies this latter condition.
- **(Positivity)** $c_1^{\text{EH}}(H) > 0$.
- **(Spectrality)** Every $c_k^{\text{EH}}(H)$ is a critical value of the Hamiltonian action functional Φ_H .

- **(Monotonicity)** If $H_1 \leq H_2$ pointwise, then $c_k^{\text{EH}}(H_1) \geq c_k^{\text{EH}}(H_2)$.

For a bounded subset $B \subset \mathbb{C}^n$, we denote by $\mathcal{F}(B)$ the family of admissible Hamiltonians $H : \mathbb{C}^n \rightarrow [0, \infty)$ vanishing on some neighborhood of the closure \bar{B} . The k -th Ekeland-Hofer capacity of B is defined as

$$c_k^{\text{EH}}(B) = \inf_{H \in \mathcal{F}(B)} c_k^{\text{EH}}(H).$$

For an unbounded subset $U \subset \mathbb{C}^n$, the k -th Ekeland-Hofer capacity $c_k^{\text{EH}}(U)$ is defined as the supremum of $c_k^{\text{EH}}(B)$ over all bounded subsets $B \subset U$. The c_k^{EH} 's satisfy the monotonicity, conformality, and normalization properties of symplectic k -capacities (see Section 1.2).

Throughout this section, in order to simplify the notation, we shall denote the Ekeland-Hofer spectral invariants and capacities c_k^{EH} simply by c_k .

3.2. Capacities vs spectral invariants. We say that a closed hypersurface $N \subset \mathbb{C}^n$ is of *restricted contact type* when there exists a primitive ν of the standard symplectic form ω of \mathbb{C}^n such that $\nu|_N$ is a contact form. Notice that the action spectrum $\sigma(N)$, which is the collection of the periods of the closed Reeb orbits of (N, ν) , is independent of the specific choice of the primitive ν . Ekeland and Hofer [EH90, Prop. 2] established the following remarkable facts.

- **(Hausdorff continuity)** The functions $B \mapsto c_k(B)$ are continuous over the space of connected compact domains $B \subset \mathbb{C}^n$ with smooth restricted contact type boundary, endowed with the topology induced by the Hausdorff distance.
- **(Spectrality)** For each connected compact subset $B \subset \mathbb{C}^n$ with smooth restricted contact type boundary, we have $c_k(B) \in \sigma(\partial B)$.

Notice in particular that, if $B \subset \mathbb{C}^n$ is a smooth convex body, both the Ekeland-Hofer capacities $c_k(B)$ and the spectral invariants $s_k(B)$ (see Section 2.2) are elements of the action spectrum $\sigma(\partial B)$. A result of Sikorav [Sik90, Section 6.5] shows that these invariants coincide for $k = 1$, i.e.

$$c_1(B) = s_1(B) = \min \sigma(\partial B). \quad (3.3)$$

It is not known whether $c_k(B) = s_k(B)$ for $k \geq 2$ as well. In this section, we prove that there is at least an inequality.

Proposition 3.1. *For each smooth convex body $B \subset \mathbb{C}^n$, we have*

$$c_k(B) \leq s_k(B), \quad \forall k \geq 2.$$

The proof of this proposition is technical, and will take the rest of this subsection. A reader mainly interested in the application may skip it and proceed directly to the next Section 3.3.

Let $B \subset \mathbb{C}^n$ be a smooth convex body, $h : \mathbb{C}^n \rightarrow [0, \infty)$ be the 2-homogeneous Hamiltonian such that $h^{-1}(1) = \partial B$, $\tilde{\Psi} = \mathcal{H}/\mathcal{A} : \mathcal{A}^{-1}(0, \infty) \rightarrow (0, \infty)$ the associated (unrestricted) Clarke action functional, and $\Psi = \tilde{\Psi}|_{\Lambda}$ its restriction. The following lemma is extracted from the arguments in [Sik90, Sec. 6.5].

Lemma 3.2. *For each $\gamma \in W^{1,2}(S^1, \mathbb{C}^n) \cap a^{-1}(0, \infty)$ and $\zeta \in E_- \oplus E_0 \oplus \mathbb{R}\gamma$, we have*

$$a(\zeta) \leq \tilde{\Psi}(\dot{\gamma}) \int_{S^1} h(\zeta(t)) dt.$$

Proof. Since $a(\gamma) > 0$ and $a|_{E_- \oplus E_0} \leq 0$, the polynomial

$$c \mapsto a(\zeta + c\gamma) = a(\zeta) + c \langle J\zeta, \dot{\gamma} \rangle_{L^2} + c^2 a(\gamma)$$

has at least one root. Therefore its discriminant is non-negative, i.e.

$$a(\gamma)a(\zeta) \leq \frac{\langle J\zeta, \dot{\gamma} \rangle_{L^2}^2}{4} \quad (3.4)$$

The dual Hamiltonian h^* , defined by (2.1), satisfies the Fenchel inequality

$$h(z) + h^*(w) \geq \langle w, z \rangle.$$

Therefore, for each $c > 0$, we have

$$c^2 \mathcal{H}(\dot{\gamma}) = \mathcal{H}(c\dot{\gamma}) = \int_{S^1} h^*(-Jc\dot{\gamma}) dt \geq c \langle J\zeta, \dot{\gamma} \rangle_{L^2} - \int_{S^1} h(\zeta) dt,$$

that is,

$$c \langle J\zeta, \dot{\gamma} \rangle_{L^2} - c^2 \mathcal{H}(\dot{\gamma}) \leq \int_{S^1} h(\zeta) dt.$$

The left-hand side is maximal for $c = \frac{1}{2} \langle J\zeta, \dot{\gamma} \rangle_{L^2} \mathcal{H}(\dot{\gamma})^{-1}$, and we obtain the estimate

$$\frac{\langle J\zeta, \dot{\gamma} \rangle_{L^2}^2}{4} \leq \mathcal{H}(\dot{\gamma}) \int_{S^1} h(\zeta) dt.$$

This, together with (3.4), implies the lemma. \square

The Hilbert spaces $L_0^2(S^1, \mathbb{C})$ and E admit the S^1 action by time-translations, and the $\mathbb{R}_+ = (0, \infty)$ action by scalar multiplication. Overall, they thus admit an action of $\mathbb{C}_* \equiv \mathbb{R}_+ \times S^1$. We introduce the \mathbb{C}_* -equivariant injective continuous map

$$i : L_0^2(S^1, \mathbb{C}^n) \rightarrow E_- \oplus E_+, \quad i(\dot{\gamma}) = \gamma - \int_{S^1} \gamma(t) dt.$$

For each $c > \min \sigma(\partial B)$ we consider the c -sublevel sets of the unrestricted and restricted Clarke action functionals, which are related by

$$\{\tilde{\Psi} \leq c\} = \mathbb{R}_+ \{\Psi \leq c\}.$$

The inclusion $\{\Psi \leq c\} \hookrightarrow \{\tilde{\Psi} \leq c\}$ is an S^1 -equivariant homotopy equivalence, and therefore

$$\text{ind}_{\text{FR}}(\{\tilde{\Psi} \leq c\}) = \text{ind}_{\text{FR}}(\{\Psi \leq c\}).$$

We set

$$\mathcal{U}_c := i(\{\tilde{\Psi} \leq c\}) \subset (E_- \oplus E_+) \setminus E_-.$$

Lemma 3.3. *For all $\epsilon > 0$ there exists an admissible Hamiltonian $H \in \mathcal{F}(B)$ such that*

$$\sup_{E_- \oplus E_0 \oplus \mathcal{U}_c} \Phi_H \leq c + \epsilon.$$

Proof. Since $B = h^{-1}[0, 1]$, we can choose an admissible Hamiltonian $H \in \mathcal{F}(B)$ such that $H \geq ch - c - \epsilon$. For each $\zeta \in E_- \oplus E_0 \oplus \mathcal{U}_c$, Lemma 3.2 implies

$$\Phi_H(\zeta) = a(\zeta) - \int_{S^1} H(\zeta) dt < c \int_{S^1} h(\zeta) dt - \int_{S^1} H(\zeta) dt \leq c + \epsilon. \quad \square$$

For any integer $N \geq 0$, we introduce the finite dimensional Hilbert subspace $F_N \subset E$ of all periodic curves $\gamma \in E$ of the form

$$\gamma(t) = \sum_{|k| \leq N} e^{i2\pi kt} \gamma_k. \quad (3.5)$$

We denote by $\Pi_N : E \rightarrow F_N$ the orthogonal projection.

Lemma 3.4. *For each $c > \min \sigma(\partial B)$ there exists $N_c \geq 1$ such that*

$$\Pi_N(\mathcal{U}_c) \cap E_- = \emptyset, \quad \forall N \geq N_c.$$

Proof. Consider an arbitrary $\gamma \in W^{1,2}(S^1, \mathbb{C}^n)$ such that $\gamma_0 = 0$, $a(\gamma) > 0$ and $\Pi_N \gamma \in E_-$. This latter condition implies

$$a(\gamma) = \underbrace{a(\Pi_N \gamma)}_{\leq 0} + a((\text{id} - \Pi_N)\gamma) \leq a((\text{id} - \Pi_N)\gamma) \leq \pi \sum_{k > N} k \|\gamma_k\|^2. \quad (3.6)$$

Consider the constant

$$r := \inf_{w \in \mathbb{C}^n \setminus \{0\}} \frac{h^*(w)}{\|w\|^2} > 0,$$

which we employ to bound

$$\mathcal{H}(\dot{\gamma}) \geq r \|\dot{\gamma}\|_{L^2}^2 \geq r \sum_{k > N} 4\pi^2 k^2 \|\gamma_k\|^2 \geq 4\pi^2 r N \sum_{k > N} k \|\gamma_k\|^2. \quad (3.7)$$

The estimates (3.6) and (3.7) imply

$$\tilde{\Psi}(\dot{\gamma}) = \frac{\mathcal{H}(\dot{\gamma})}{a(\gamma)} \geq 4\pi r N,$$

and the desired quantity of the lemma can be set to $N_c := c/(4\pi r)$. \square

For technical reasons, it will be useful to consider a compact S^1 -invariant subset of \mathcal{U}_c .

Lemma 3.5. *There exists an S^1 -invariant compact subset $\mathcal{V}_c \subset \mathcal{U}_c$ such that*

$$\text{ind}_{\text{FR}}(\mathbb{R}_+ \mathcal{V}_c) \geq \text{ind}_{\text{FR}}(\mathcal{V}_c) \geq \text{ind}_{\text{FR}}(\{\Psi \leq c\}).$$

Proof. Consider the reduced Clarke action functional $\psi : U \rightarrow (0, c+1)$ introduced in Section 2.3. We recall that U is an open subset of the unit-sphere $S \subset F$, where $F = F_N$ is the finite dimensional Hilbert space introduced above, and N is sufficiently large. The function ψ has the form

$$\psi(u) = \tilde{\Psi}(u + \nu(u)),$$

where $\nu : U \rightarrow L_0^2(S^1, \mathbb{C}^n)$ is an S^1 -equivariant $C^{1,1}$ map. The sublevel set

$$\mathcal{K}_c := \{u + \nu(u) \mid \psi(u) \leq c\}$$

is compact in $L_0^2(S^1, \mathbb{C}^n)$, and the inclusion $\mathcal{K}_c \hookrightarrow \{\tilde{\Psi} \leq c\}$ is an S^1 -equivariant homotopy equivalence. In particular

$$\text{ind}_{\text{FR}}(\mathcal{K}_c) = \text{ind}_{\text{FR}}(\{\tilde{\Psi} \leq c\}) = \text{ind}_{\text{FR}}(\{\Psi \leq c\}).$$

The image $\mathcal{V}_c = i(\mathcal{K}_c)$ is an S^1 -invariant compact subset of \mathcal{U}_c , and by the monotonicity property of the Fadell-Rabinowitz index (see Appendix A) we conclude

$$\text{ind}_{\text{FR}}(\mathbb{R}_+ \mathcal{V}_c) \geq \text{ind}_{\text{FR}}(\mathcal{V}_c) \geq \text{ind}_{\text{FR}}(\mathcal{K}_c). \quad \square$$

In [EH90, Prop. 1], Ekeland and Hofer proved that

$$\text{ind}_{\text{EH}}(E_- \oplus E_0 \oplus V) = \dim_{\mathbb{C}}(V) = \text{ind}_{\text{FR}}(V \setminus \{0\})$$

for every finite dimensional S^1 -invariant vector subspace $V \subset E_+$. Building on Ekeland and Hofer arguments, we need to prove the following analogous statement for the \mathbb{C}_* -invariant set $\mathcal{W}_c := \mathbb{R}_+ \mathcal{V}_c$. We shall employ the same notation as in Appendix A: if W is a subset of a vector space, we set $W_* := W \setminus \{0\}$.

Lemma 3.6. *For each $c > \min \sigma(\partial B)$, we have $\text{ind}_{\text{EH}}(E_- \oplus E_0 \oplus \mathcal{W}_c) \geq \text{ind}_{\text{FR}}(\mathcal{W}_c)$.*

Proof. Fix an arbitrary $\Theta \in \mathcal{D}$, and an arbitrary S^1 -equivariant open neighborhood $\mathcal{U} \subset E$ of $\Theta(E_- \oplus E_0 \oplus \mathcal{W}_c) \cap S_+$. We need to prove

$$\text{ind}_{\text{FR}}(\mathcal{U}) \geq \text{ind}_{\text{FR}}(\mathcal{W}_c).$$

We consider the finite dimensional Hilbert subspace $F = F_N \subset E$ introduced above, for a sufficiently large N that we will fix later. This Hilbert subspace splits as a direct sum $F = F_- \oplus F_0 \oplus F_+$, where $F_0 = E_0$ and $F_{\pm} = F_{N,\pm} := E_{\pm} \cap F$. We denote by $\Pi_{\pm} = \Pi_{N,\pm} : E \rightarrow F_{\pm}$ and $\Pi = \Pi_N : E \rightarrow F$ the orthogonal projections. We require $N \geq N_c$, so that Lemma 3.4 implies that $\Pi_+(\mathcal{W}_c)$ does not contain the origin. We set

$$W := \Pi_+(\mathcal{W}_c) \cup \{0\}, \quad W_* := \Pi_+(\mathcal{W}_c) = W \setminus \{0\},$$

so that

$$\Pi(E_- \oplus E_0 \oplus \mathcal{W}_c) = F_- \oplus F_0 \oplus W_*.$$

The monotonicity property of the Fadell-Rabinowitz index (see Appendix A) implies that

$$\text{ind}_{\text{FR}}(W_*) \geq \text{ind}_{\text{FR}}(\mathcal{W}_c).$$

The vector space F_- has complex dimension nN , and the circle S^1 acts on it by rotations without fixed points. We set $Y = Y_N := F_- \oplus W$. Lemma A.2 and $\mathcal{W}_c \neq \emptyset$ implies

$$\text{ind}_{\text{FR}}(Y_*) = \text{ind}_{\text{FR}}(W_*) + nN \geq \text{ind}_{\text{FR}}(\mathcal{W}_c) + nN > nN.$$

We claim that, if we fix $N \geq N_c$ large enough, we have

$$(\Pi \circ \Theta(Y \oplus F_0)) \cap S_+ \subset \mathcal{U}. \quad (3.8)$$

Indeed, assume by contradiction that there exists a sequence $\gamma_N \in Y_N \oplus F_0$ such that $\Pi_N \circ \Theta(\gamma_N) \in S_+ \setminus \mathcal{U}$ for arbitrarily large $N \geq N_c$. We write

$$\gamma_N = \gamma_{N,-} + \gamma_{N,0} + \gamma_{N,+},$$

where $\gamma_{N,\pm} = \Pi_{N,\pm}(\gamma_N)$. Since $(\Pi_{N,-} + \Pi_{N,0}) \circ \Theta(\gamma_N) = 0$ and Θ is of the form (3.2), we have

$$\gamma_{N,-} + \gamma_{N,0} + (e^{-\theta_-(\gamma_N)} \Pi_{N,-} + \Pi_0) \mathcal{K}(\gamma_N) = 0. \quad (3.9)$$

Since θ_- and \mathcal{K} maps bounded sets to precompact sets and vanish outside a bounded subset of E , Equation (3.9) implies that the sequence $\gamma_{N,-} + \gamma_{N,0}$ is precompact. Since Θ is the identity outside a bounded set of E , the sequence $\gamma_{N,+}$ is also bounded; otherwise, after extracting a subsequence we would have $\|\gamma_{N,+}\|_E \rightarrow \infty$

as $N \rightarrow \infty$, and for N large enough we would have $\gamma_N = \Theta(\gamma_N)$ and obtain the contradiction

$$1 = \|\Pi_N \circ \Theta(\gamma_N)\|_E = \|\gamma_N\|_E \geq \|\gamma_{N,+}\|_E \xrightarrow{N \rightarrow \infty} \infty.$$

Notice that $\gamma_{N,+}$ is contained in $\Pi_{N,+}(Y_N) = \Pi_{N,+}(\mathcal{W}_c) = \mathbb{R}_+ \Pi_{N,+}(\mathcal{V}_c)$, and we recall that $\Pi_{N,+}(\mathcal{V}_c)$ is compact (Lemma 3.5). Since the sequence $\gamma_{N,+}$ is uniformly bounded, it is precompact in $\Pi_{N,+}(\mathcal{W}_c) \cup \{0\}$. Overall, we showed that the sequence γ_N is precompact, and after extracting a subsequence we have a convergence

$$\lim_{N \rightarrow \infty} \gamma_N = \gamma \in (E_- \oplus E_0 \oplus \mathcal{W}_c \cup \{0\}) \setminus \mathcal{U}.$$

But $\Theta(\gamma) \in S_+$, and since $\Theta(0) = 0$ we must have $\gamma \neq 0$, and therefore $\gamma \in E_- \oplus E_0 \oplus \mathcal{W}_c$. This gives a contradiction, since \mathcal{U} is an open neighborhood of $\Theta(E_- \oplus E_0 \oplus \mathcal{W}_c) \cap S_+$, and completes the proof of (3.8).

We introduce the S^1 -equivariant continuous map $\phi := \Pi \circ \Theta|_F : F \rightarrow F$, which satisfies

$$\phi(Y \oplus F_0) \cap S_+ \subset \mathcal{U}$$

according to (3.8). We denote by $B' \subset F$ the closed unit ball, which is S^1 -invariant. Since the homeomorphism Θ is the identity on $E_- \oplus E_0$ and outside a ball of E of sufficiently large radius, the map ϕ is the identity on $F_- \oplus F_0$ and outside a compact set. Therefore $B := \phi^{-1}(B') \subset F$ is an S^1 -invariant compact neighborhood of the origin. We define another S^1 -equivariant continuous map

$$\psi := (\text{id} - \Pi_+) \circ \phi|_{Y \oplus F_0} : Y \oplus F_0 \rightarrow F_- \oplus F_0,$$

which satisfies $\psi(0, x) = (0, x)$ for all $x \in F_0$. The S^1 -invariant subset

$$Z := \psi^{-1}(0) \cap \partial B,$$

satisfies

$$\phi(Z) \subset \phi(Y \oplus F_0) \cap S_+ \subset \mathcal{U}.$$

Let $U \subset Y \oplus F_0$ be an S^1 -invariant open subset containing Z that is sufficiently small so that $\phi(U) \subset \mathcal{U}$. Proposition A.4 applied to the map ψ implies that $\text{ind}_{\text{FR}}(U) \geq \text{ind}_{\text{FR}}(Y_*) - nN$. Summing up, we obtained the desired lower bound

$$\text{ind}_{\text{FR}}(\mathcal{U}) \geq \text{ind}_{\text{FR}}(U) \geq \text{ind}_{\text{FR}}(Y_*) - nN \geq \text{ind}_{\text{FR}}(\mathcal{W}_c). \quad \square$$

Proof of Proposition 3.1. Assume that $s_k(B) < c$, so that

$$\text{ind}_{\text{FR}}(\mathcal{W}_c) \geq \text{ind}_{\text{FR}}(\{\Psi \leq c\}) \geq k.$$

By Lemma 3.3, for each $\epsilon > 0$ there exists an admissible Hamiltonian $H \in \mathcal{F}(B)$ such that

$$\sup_{E_- \oplus E_0 \oplus \mathcal{W}_c} \Phi_H \leq c + \epsilon.$$

Lemma 3.6 implies

$$\text{ind}_{\text{EH}}(E_- \oplus E_0 \oplus \mathcal{W}_c) \geq \text{ind}_{\text{FR}}(\mathcal{W}_c) \geq k.$$

Therefore $c_k(B) \leq c_k(H) \leq c + \epsilon$. Since $\epsilon > 0$ can be chosen arbitrarily small, we infer that $c_k(B) \leq c$. Finally, since this latter inequality holds for all $c > s_k(B)$, we conclude $c_k(B) \leq s_k(B)$. \square

3.3. Local maximizers of the capacity ratio. Let us extend the notion of Besse convex body to the class of smooth star-shaped domains. A $2n$ -dimensional smooth star-shaped domain $B \subset \mathbb{C}^n$ is called *Besse* when its boundary ∂B , equipped with the canonical contact form $\lambda|_{\partial B}$ of Equation (1.1), is a Besse contact manifold: all its Reeb orbits are closed, and therefore have a common period according to Wadsley theorem [Wad75]. Besse convex bodies, and in particular rational ellipsoids, are a subclass of Besse star-shaped domains. Actually, in dimension 4, these classes are the same up to symplectomorphisms.

Proposition 3.7. *Let B be a 4-dimensional smooth star-shaped domain. The following conditions are equivalent:*

- (i) B is Besse.
- (ii) ∂B is strictly contactomorphic to the boundary of a rational ellipsoid E , i.e. there exists a diffeomorphism $\psi : \partial B \rightarrow \partial E$ such that $\psi^* \lambda|_{\partial E} = \lambda|_{\partial B}$.
- (iii) There exists a diffeomorphism $\psi : \mathbb{C}^2 \rightarrow \mathbb{C}^2$ such that $\psi(B) = E$, $\psi^* \omega = \omega$, and $\psi|_{\partial B}^* \lambda = \lambda|_{\partial B}$.
- (iv) B is symplectomorphic to a rational ellipsoid E , i.e. there exists a diffeomorphism $\psi : B \rightarrow E$ such that $\psi^* \omega = \omega$.

Proof. Assume that B is a 4-dimensional Besse star-shaped domain. As it was pointed out in [MR23, Theorem 1.1], the contact classification of Seifert fibrations [GL18, CGM20] implies that the boundary of B is strictly contactomorphic to the boundary of a rational ellipsoid E : there exists a diffeomorphism $\psi : \partial B \rightarrow \partial E$ such that $\psi^* \lambda|_{\partial E} = \lambda|_{\partial B}$. A straightforward generalization of [ABHS18, Prop. 4.3] implies that ψ can be extended to a symplectomorphism $\psi \in \text{Symp}(\mathbb{C}^2, \omega)$. Therefore conditions (i) and (ii) are equivalent, and imply condition (iii).

Condition (iii) immediately implies condition (iv). Assume now that a 4-dimensional smooth star-shaped domain B is symplectomorphic to a rational ellipsoid E via a symplectomorphism $\psi : B \rightarrow E$. The Reeb orbits on ∂B and ∂E are leaves of the characteristic foliations $\ker(\omega|_{\partial B})$ and $\ker(\omega|_{\partial E})$ respectively. Since $\psi^* \omega = \omega$, the restriction $\psi|_{\partial B}$ maps Reeb orbits of ∂B to time-reparametrized Reeb orbits of ∂E . Since all Reeb orbits of ∂E are closed, all Reeb orbits of ∂B are closed as well. Therefore condition (iv) implies condition (i). \square

Let B be a 4-dimensional Besse star-shaped domain, and $\psi \in \text{Symp}(\mathbb{C}^2, \omega)$ a symplectomorphism mapping B to a rational ellipsoid $E := \psi(B)$, and whose restriction $\psi|_{\partial B} : \partial B \rightarrow \partial E$ is a strict contactomorphism. Since $\psi|_{\partial B}$ conjugates the Reeb flows on ∂B and ∂E , in particular $\sigma(\partial B) = \sigma(\partial E)$. The monotonicity property of the Ekeland-Hofer capacities implies that $c_k(E) = c_k(B)$. In particular, the capacities $c_k(B)$ cover the whole spectrum $\sigma(\partial B)$, i.e.

$$\sigma(\partial B) = \{c_k(B) \mid k \geq 1\},$$

since the same holds for the ellipsoid E . We denote by $\tau(B)$ the minimal common period of the closed Reeb orbit on ∂B . For all integers $m \geq 1$, we define

$$k_m(B) := \min \{k \geq 1 \mid c_k(B) = m \tau(B)\},$$

$$K(B) := \{k_m(B) \mid m \geq 1\}.$$

Clearly, $\tau(E) = \tau(B)$ and $K(E) = K(B)$. These definitions of $k_m(B)$ and $K(B)$ agree with the ones in Section 1.2 in the special case of rational ellipsoids. They

also agree with the one given in Section 2.5 for Besse convex domains, thanks to the following lemma.

Lemma 3.8. *For each 4-dimensional Besse convex body B , we have $c_k(B) = s_k(B)$ for all $k \geq 1$.*

Proof. Let $\psi \in \text{Symp}(\mathbb{C}^2, \omega)$ be a symplectomorphism mapping B to a rational ellipsoid $E := \psi(B)$, so that $\tau = \tau(B) = \tau(E)$. We already know that both the spectral invariants $s_k(B)$ and the Ekeland-Hofer capacities $c_k(B) = c_k(E)$ cover the whole action spectrum $\sigma = \sigma(\partial B) = \sigma(\partial E)$, i.e.

$$\sigma = \{s_k(B) \mid k \geq 1\} = \{c_k(E) \mid k \geq 1\}.$$

Moreover $s_1(B) = c_1(E) = \min \sigma$, $s_k(B) \leq s_{k+1}(B)$, and $c_k(E) \leq c_{k+1}(E)$. The spectral Besse characterization for the Clarke action functional (Section 2.4) implies that $s_1(B) = \tau$ if and only if $s_1(B) = s_2(B)$; moreover, for any $k \geq 2$, it implies that $s_k(B)$ is a multiple of τ if and only if either $s_{k-1}(B) = s_k(B)$ or $s_k(B) = s_{k+1}(B)$. These properties fully determine $s_k(B)$ for the Besse convex body B . The normalization property of the symplectic k -th capacities implies that $c_k(E)$ satisfy analogous properties: $c_1(E) = \tau$ if and only if $c_1(E) = c_2(E)$; for any $k \geq 2$, $c_k(E)$ is a multiple of τ if and only if either $c_{k-1}(E) = c_k(E)$ or $c_k(E) = c_{k+1}(E)$. We conclude that $s_k(B) = c_k(E)$. \square

We infer the first half of Theorem A from Theorem 2.4.

Lemma 3.9. *Let B_0 be a 4-dimensional Besse star-shaped domain. For each integer $k \in K(B_0)$, any smooth star-shaped domain B_1 that is sufficiently C^3 -close to B_0 satisfies $\widehat{c}_k(B_1) \leq \widehat{c}_k(B_0)$.*

Proof. Fix $k \in K(B_0)$, and a symplectomorphism $\psi \in \text{Symp}(\mathbb{C}^2, \omega)$ mapping B_0 to a rational ellipsoid. If a smooth star-shaped domain B_1 is C^3 -close to B_0 , then $\psi(B_1)$ is C^3 -close to $\psi(B_0)$. In particular, $\psi(B_1)$ is a smooth convex body, and Theorem 2.4 implies $\widehat{s}_k(\psi(B_1)) \leq \widehat{s}_k(\psi(B_0))$. Proposition 3.1 implies $c_k(\psi(B_1)) \leq s_k(\psi(B_1))$, and Lemma 3.8 implies $c_k(\psi(B_0)) = s_k(\psi(B_0))$. Therefore we conclude

$$\widehat{c}_k(B_1) = \widehat{c}_k(\psi(B_1)) \leq \widehat{s}_k(\psi(B_1)) \leq \widehat{s}_k(\psi(B_0)) = \widehat{c}_k(\psi(B_0)) = \widehat{c}_k(B_0). \quad \square$$

Next, we address the opposite implication in Theorem A. The following lemma holds in arbitrary dimension.

Lemma 3.10. *Let B_0 be a local maximizer of the capacity ratio \widehat{c}_k over the space of $2n$ -dimensional smooth star-shaped domains endowed with the C^∞ topology. Then B_0 is a Besse star-shaped domain and $c_k(B_0)$ is a common period for the Reeb orbits on ∂B_0 .*

Proof. The proof builds on an argument originally due to Alvarez Paiva and Balacheff [APB14]. Let B_0 be a $2n$ -dimensional smooth star-shaped domain, and $h_0 : \mathbb{C}^n \rightarrow [0, \infty)$ the 2-homogeneous Hamiltonian such that $h_0^{-1}(1) = \partial B_0$. The restriction of its Hamiltonian flow $\phi_{h_0}^t|_{\partial B_0}$ is the Reeb flow of ∂B_0 . Assume by contradiction that $c_k(B_0)$ is not a common period for the Reeb orbits on the boundary ∂B_0 . Namely, there exists $\delta > 0$ and a non-empty open subset $U \subset \partial B_0$ such that

$$\phi_h^t(z) \neq z, \quad \forall z \in U, t \in [c_k(B_0) - \delta, c_k(B_0) + \delta].$$

Let $\chi : \partial B_0 \rightarrow [0, \infty)$ be a smooth function supported in U and not identically zero. For $\epsilon > 0$, let $h_\epsilon : \mathbb{C}^n \rightarrow [0, \infty)$ be the 2-homogeneous Hamiltonian such that

$h_\epsilon|_{\partial B_0} = h_0|_{\partial B_0} + \epsilon\chi$. For all $\epsilon > 0$ small enough, the sublevel set $B_\epsilon := h_\epsilon^{-1}[0, 1]$ is a smooth star-shaped domain strictly contained in B_0 , and in particular

$$\text{vol}(B_\epsilon) < \text{vol}(B_0).$$

By our choice of χ , for all $\epsilon > 0$ small enough we have

$$\text{fix}(\phi_{h_\epsilon}^t) = \text{fix}(\phi_{h_0}^t), \quad \forall t \in [c_k(B_0) - \delta, c_k(B_0) + \delta].$$

Therefore the continuous function $\epsilon \mapsto c_k(B_\epsilon)$ takes values into the intersection $\sigma(\partial B_0) \cap [c_k(B_0) - \delta, c_k(B_0) + \delta]$ for all $\epsilon \geq 0$ small enough. Since the action spectrum $\sigma(\partial B_0)$ is nowhere dense (see, e.g., [Sik90, Prop. 7.4]), we conclude that $c_k(B_\epsilon) = c_k(B_0)$, and therefore $\widehat{c}_k(B_\epsilon) > \widehat{c}_k(B_0)$, for all $\epsilon > 0$ small enough. \square

Proof of Theorem A. We consider the capacity ratio \widehat{c}_k over the space of 4-dimensional smooth star-shaped domains, endowed with the C^3 topology. Let B be a 4-dimensional smooth star-shaped domain symplectomorphic to a rational ellipsoid E . Proposition 3.7 implies that B is Besse, and Lemma 3.9 asserts that B is a local maximizer of \widehat{c}_k for all $k \in K(B) = K(E)$.

Conversely, assume that a 4-dimensional smooth star-shaped domain B_0 is a local maximizer of \widehat{c}_k . Lemma 3.10 implies that B_0 is Besse, and therefore Proposition 3.7 implies that there exists $\psi \in \text{Symp}(\mathbb{C}^2, \omega)$ mapping B_0 to a rational ellipsoid $E(\mathbf{a})$. For each \mathbf{b} sufficiently close to \mathbf{a} , the preimage $B_1 := \psi^{-1}(E(\mathbf{b}))$ is a smooth star-shaped domain C^3 -close to B_0 , and therefore

$$\widehat{c}_k(E(\mathbf{b})) = \widehat{c}_k(B_1) \leq \widehat{c}_k(B_0) = \widehat{c}_k(E(\mathbf{a})).$$

Namely, $E(\mathbf{a})$ is a local maximizer of \widehat{c}_k over the space of 4-dimensional ellipsoids, and Proposition C(i) implies that $k \in K(E(\mathbf{a})) = K(B_0)$. \square

We can also complete Theorem 2.4, fully characterizing the local maximizers of the spectral ratios \widehat{s}_k over the space of 4-dimensional smooth convex domains.

Theorem 3.11. *On the space of 4-dimensional smooth convex domains endowed with the C^3 topology, the local maximizers of $B \mapsto \widehat{s}_k(B)$ are precisely those domains symplectomorphic to a $2n$ -dimensional rational ellipsoid E with $k \in K(E)$.*

Proof. Theorem 2.4, together with Proposition 3.7, provides one implication: the smooth convex bodies B symplectomorphic to a 4-dimensional rational ellipsoid E are local maximizers of \widehat{s}_k for all $k \in K(B) = K(E)$.

Conversely, let B_0 be a smooth convex domain that is a local maximizer of \widehat{s}_k . An analogous argument to the one in the proof of Lemma 3.10 implies that B_0 is Besse, and Proposition 3.7 provides a symplectomorphism $\psi \in \text{Symp}(\mathbb{C}^2, \omega)$ mapping B_0 to a rational ellipsoid $E(\mathbf{a})$. For each \mathbf{b} sufficiently close to \mathbf{a} , the preimage $B_1 := \psi^{-1}(E(\mathbf{b}))$ is a smooth convex body C^3 -close to B_0 , and therefore $\widehat{s}_k(B_1) \leq \widehat{s}_k(B_0)$. Proposition 3.1 implies $c_k(B_1) \leq s_k(B_1)$, and Lemma 3.8 implies $s_k(B_0) = c_k(B_0)$. Overall, we obtained

$$\widehat{c}_k(E(\mathbf{b})) = \widehat{c}_k(B_1) \leq \widehat{s}_k(B_1) \leq \widehat{s}_k(B_0) = \widehat{c}_k(B_0) = \widehat{c}_k(E(\mathbf{a})).$$

Namely, $E(\mathbf{a})$ is a local maximizer of \widehat{c}_k over the space of 4-dimensional ellipsoids. Proposition C(i) implies that $k \in K(E(\mathbf{a})) = K(B_0)$. \square

4. MAXIMIZERS OF THE CAPACITY RATIOS AMONG ELLIPSOIDS

We recall that, throughout this paper, we compute the volumes of domains in \mathbb{C}^n by integrating the volume form ω^n ; namely, for a $2n$ -dimensional domain K , the volume $\text{vol}(K) = \text{vol}(K, \omega)$ is $n!$ times the usual $2n$ -dimensional Euclidean volume. For a $2n$ -dimensional ellipsoid $E(\mathbf{a})$, with $\mathbf{a} = (a_1, \dots, a_n)$, we have

$$\text{vol}(E(\mathbf{a})) = a_1 \dots a_n$$

All symplectic k -capacities coincide over the space of ellipsoids, according to their normalization property (see Section 1.2). Throughout this section, we denote by c_k any symplectic k -capacity.

Proof of Proposition C(i). Let $\mathbf{a} = (a_1, \dots, a_n)$ be a local maximizer of the function $\mathbf{a} \mapsto \widehat{c}_k(E(\mathbf{a}))$, and assume without loss of generality that $0 < a_1 \leq \dots \leq a_n < \infty$. We first prove that every a_i divides $c_k(E(\mathbf{a}))$. Indeed, if we had $ra_i < c_k(E(\mathbf{a})) < (r+1)a_i$ for some integer $r \geq 0$ and some minimal $i \in \{1, \dots, n\}$, then for any $t < 1$ sufficiently close to 1 the configuration $\mathbf{b}(t) = (b_1(t), \dots, b_n(t))$ given by

$$b_j(t) = \begin{cases} t a_i, & \text{if } j = i, \\ a_j, & \text{if } j \neq i \end{cases}$$

would satisfy $c_k(E(\mathbf{b}(t))) = c_k(E(\mathbf{a}))$ and $\text{vol}(E(\mathbf{b}(t))) < \text{vol}(E(\mathbf{a}))$, contradicting the maximality of \mathbf{a} .

Therefore, $c_k(E(\mathbf{a})) = r_1 a_1 = \dots = r_n a_n$ for some integers $r_1 \geq r_2 \geq \dots \geq r_n \geq 1$. We now prove that

$$c_k(\mathbf{a}) = c_{k+n-1}(\mathbf{a}),$$

which readily implies $r_1 + \dots + r_n = k + n - 1$ and therefore $k \in K(E(\mathbf{a}))$. Assume by contradiction that $c_k(\mathbf{a}) < c_{k+n-1}(\mathbf{a})$. This implies that $c_k(E(\mathbf{a})) = c_{k-1}(E(\mathbf{a}))$. For any $t < 1$ sufficiently close to 1, the configuration $\mathbf{b}(t) = (ta_1, a_2, \dots, a_n)$ has capacity $c_k(E(\mathbf{b}(t))) = c_k(E(\mathbf{a}))$, but lower volume $\text{vol}(E(\mathbf{b}(t))) < \text{vol}(E(\mathbf{a}))$, contradicting the maximality of \mathbf{a} .

We are left to prove the converse implication. Let $E(\mathbf{a})$ be a rational ellipsoid. As usual, assume without loss of generality that $a_1 \leq \dots \leq a_n$. Fix any integer $k \in K(E(\mathbf{a}))$. There exist integers $r_i \geq 1$ such that $r_1 a_1 = \dots = r_n a_n$ and $k = r_1 + \dots + r_n - n + 1$. For all \mathbf{b} sufficiently close to \mathbf{a} , we have $\widehat{c}_k(E(\mathbf{b})) = f(\mathbf{b})$, where f is the 0-homogeneous function

$$f(\mathbf{b}) = \frac{\min\{r_1 b_1, \dots, r_n b_n\}}{(b_1 \dots b_n)^{1/n}}.$$

As usual, assume without loss of generality that $b_1 \leq \dots \leq b_n$. We set

$$t := \frac{r_1 a_1}{\min\{r_1 b_1, \dots, r_n b_n\}}.$$

Notice that $t b_i \geq a_i$. Therefore, for all \mathbf{b} sufficiently close to \mathbf{a} , we have

$$\widehat{c}_k(E(\mathbf{b})) = f(\mathbf{b}) = f(t\mathbf{b}) = \frac{r_1 a_1}{(t b_1 \dots t b_n)^{1/n}} \leq \frac{r_1 a_1}{(a_1 \dots a_n)^{1/n}} = f(\mathbf{a}) = \widehat{c}_k(E(\mathbf{a})). \quad \square$$

Proof of Proposition C(ii). For configurations $\mathbf{a} = (a_1, \dots, a_n)$ with

$$0 < a_1 \leq \dots \leq a_n < \infty, \quad (4.1)$$

we have

$$\widehat{c}_k(E(\mathbf{a})) \leq \frac{ka_1}{(a_1 \dots a_n)^{1/n}} \xrightarrow{a_n \rightarrow \infty} 0.$$

Therefore, the function $\mathbf{a} \mapsto \widehat{c}_k(E(\mathbf{a}))$ achieves its maximum at some \mathbf{a} , which we assume to satisfy (4.1) without loss of generality. By Proposition C(i), we have $k \in K(E(\mathbf{a}))$. Namely

$$c_k(E(\mathbf{a})) = c_{k+n-1}(E(\mathbf{a})) = s_1 a_1 = \dots = s_n a_n$$

for some integers $s_1 \geq s_2 \geq \dots \geq s_n \geq 1$.

We claim that

$$s_i - s_j \leq 1, \quad \forall 1 \leq i < j \leq n.$$

Otherwise, we can find $i < j$ such that $s_i - s_j \geq 2$, $s_i - s_{i+1} \geq 1$, and $s_{j-1} - s_j \geq 1$ (the case $i = j - 1$ is allowed). The configuration $\mathbf{b} = (b_1, \dots, b_n)$ given by

$$b_l = \begin{cases} \frac{s_i}{s_i-1} a_i, & \text{if } l = i, \\ \frac{s_j}{s_j+1} a_j, & \text{if } l = j, \\ a_l, & \text{if } l \neq i, j \end{cases}$$

has the same value $c_k(E(\mathbf{b})) = c_k(E(\mathbf{a}))$, but

$$\text{vol}(E(\mathbf{b})) = \frac{s_i s_j}{(s_i - 1)(s_j + 1)} \text{vol}(E(\mathbf{a})) = \frac{s_i s_j}{s_i s_j + \underbrace{s_i - s_j - 1}_{\geq 1}} \text{vol}(E(\mathbf{a})) < \text{vol}(E(\mathbf{a})),$$

contradicting the maximality of \mathbf{a} .

Summing up, we proved that

$$\begin{aligned} c_k(E(\mathbf{a})) &= c_{k+n-1}(E(\mathbf{a})) = (s+2)a_1 = (s+2)a_2 = \dots = (s+2)a_{i-1} \\ &= (s+1)a_i = (s+1)a_{i+1} = \dots = (s+1)a_n \end{aligned}$$

for some integers $s \geq 0$ and $i \in \{1, \dots, n\}$. Since $c_k(E(\mathbf{a})) = c_{k+n-1}(E(\mathbf{a}))$, we have

$$k + n - 1 = (s+2)(i-1) + (s+1)(n-i+1),$$

that is, $k = sn + i$, which implies that $q = s$ and $r = i$. \square

5. MAXIMIZERS OF THE CAPACITY RATIOS AMONG TORIC DOMAINS

Throughout this section, we denote by c_k any symplectic k -capacity satisfying the extra ‘‘closed Reeb orbits’’ assumption (see Section 1.2). We shall consider 4-dimensional toric domains X_Ω whose profile Ω is the subgraph of monotone non-increasing functions $f : [0, x_0] \rightarrow [0, \infty)$ such that $f|_{[0, x_0]} > 0$ and $f(x_0) = 0$, i.e.

$$\Omega = \Omega_f := \{(x, y) \in \mathbb{R}^2 \mid x \in [0, x_0], y \in [0, f(x)]\}.$$

5.1. Concave toric domains. Let X_Ω be a 4-dimensional concave toric domain. Its profile $\Omega = \Omega_f \subset [0, \infty)^2$ must be the subgraph of a (monotone decreasing) convex function $f : [0, x_0] \rightarrow [0, \infty)$ such that $f|_{[0, x_0]} > 0$ and $f(x_0) = 0$. As usual, we denote by $\text{vol}(X_\Omega) = \text{vol}(X_\Omega, \omega)$ the volume of X_Ω computed by integrating the volume form ω^2 . This volume $\text{vol}(X_\Omega)$ is twice the Euclidean area of Ω .

Gutt and Hutchings [GH18, Theorem 1.14] provided a formula for the higher capacities of $2n$ -dimensional concave toric domains. For our 4-dimensional X_Ω the formula reads:

$$c_k(X_\Omega) = \max \left\{ [v]_\Omega \mid v = (v_1, v_2) \in \{1, \dots, k\}^2, v_1 + v_2 = k + 1 \right\}, \quad (5.1)$$

where

$$[v]_\Omega = \min \{ \langle v, w \rangle \mid w \in \text{graph}(f) \}.$$

Proposition D(i) is a direct consequence of this formula and of the properties of symplectic k -capacities.

Proof of Proposition D(i). Let $v \in \{1, \dots, k\}^2$ be a vector such that $c_k(X_\Omega) = [v]_\Omega$, and $w = (w_1, w_2) \in \text{graph}(f)$ be a vector such that $[v]_\Omega = \langle v, w \rangle$. We denote by $g : \mathbb{R} \rightarrow \mathbb{R}$ an affine function such that $g(w_1) = w_2$ and $g \leq f$. We set $a_2 := g(0)$ and $a_1 := a_2/g'(0)$, so that $g(a_1) = 0$. The ellipsoid $E(a_1, a_2) = X_\Upsilon$ is a concave toric domain whose associated profile $\Upsilon = \Omega_g$ satisfies $\Upsilon \subseteq \Omega$, and therefore

$$\text{vol}(E(a_1, a_2)) = 2 \text{area}(\Upsilon) \leq 2 \text{area}(\Omega) = \text{vol}(X_\Omega).$$

Moreover, by the formula (5.1) applied to $E(a_1, a_2)$, we have

$$c_k(E(a_1, a_2)) \geq [v]_\Upsilon = [v]_\Omega = c_k(X_\Omega).$$

Overall, we obtained $\widehat{c}_k(E(a_1, a_2)) \geq \widehat{c}_k(X_\Omega)$. Finally, by Proposition C, the function $(a_1, a_2) \mapsto \widehat{c}_k(E(a_1, a_2))$ achieves its maximum at

$$(a_1, a_2) = \left(\left\lceil \frac{k}{2} \right\rceil, \left\lceil \frac{k+1}{2} \right\rceil \right). \quad \square$$

5.2. Convex toric domains. Let X_Ω be a 4-dimensional convex toric domain. Its profile $\Omega = \Omega_f \subset [0, \infty)^2$ must be the subgraph of a concave and non-increasing function $f : [0, x_0] \rightarrow [0, y_0]$ such that $f(0) = y_0 > 0$ and $f(x_0) = 0$. We consider the convex body

$$\widehat{\Omega} := \{ (x_1, x_2) \in \mathbb{R}^2 \mid (|x_1|, |x_2|) \in \Omega \}.$$

We denote by $\|\cdot\|_\Omega$ the norm on \mathbb{R}^2 whose unit ball is the polar of $\widehat{\Omega}$, i.e.

$$\|v\|_\Omega := \max_{w \in \widehat{\Omega}} \langle v, w \rangle, \quad \forall v \in [0, \infty)^2.$$

We set $V_k := \{v_0, \dots, v_k\}$, where $v_j = (j, k-j) \in [0, k]^2$. Gutt and Hutchings [GH18, Theorem 1.16] provided a formula for the higher capacities of $2n$ -dimensional convex toric domains, which in the special case of our 4-dimensional X_Ω reads:

$$c_k(\Omega) = \min_{v \in V_k} \|v\|_\Omega.$$

Example 5.1. For each $a, b > 0$, the polydisk $P(a, b) := E(a) \times E(b) = X_\Omega$ is a convex toric domain with profile $\Omega = [0, a] \times [0, b]$. Therefore $c_k(P(a, b)) = k \min\{a, b\}$, $\text{vol}(P(a, b)) = 2ab$, and

$$\widehat{c}_k(P(a, b)) = k \frac{\min\{a, b\}}{\sqrt{2ab}}.$$

The polydisk that maximizes \widehat{c}_k is (up to rescaling) $P(1, 1)$, for which $\widehat{c}_k(P(1, 1)) = k/\sqrt{2}$. \square

The proof of Proposition D(ii) requires several lemmas.

Lemma 5.2. *Over the space of 4-dimensional convex toric domains, the capacity ratios \widehat{c}_k admit a global maximizer.*

Proof. Consider a profile $\Omega = \Omega_f$ that is the subgraph of a concave and non-increasing function $f : [0, x_0] \rightarrow [0, y_0]$ such that $f(0) = y_0$ and $f(x_0) = 0$. Since the capacity ratios are scale invariant, we can always assume that $y_0 = 1$. Assume in addition that $x_0 \leq 1$. The volume of the associated convex toric domain X_Ω is bounded from below as

$$\text{vol}(X_\Omega) = 2 \text{area}(\Omega) \geq x_0.$$

The k -th capacity of X_Ω is bounded from above as

$$c_k(X_\Omega) = \min_{j \in \{0, \dots, k\}} \max_{w \in \Omega} \langle v_j, w \rangle \leq \min_{j \in \{0, \dots, k\}} (jx_0 + (k-j)) \leq kx_0.$$

Therefore $\widehat{c}_k(X_\Omega) \leq k\sqrt{x_0} < \widehat{c}_k(P(1, 1))$ if $x_0 < 1/2$. Analogously, $\widehat{c}_k(X_\Omega) \leq k/\sqrt{2}$ if $x_0 > 2$.

Let \mathcal{T} be the space of 4-dimensional convex toric domains, and $\mathcal{T}' \subset \mathcal{T}$ the subspace of those X_{Ω_f} such that $f : [0, x_0] \rightarrow [0, 1]$ is concave, non-increasing, and satisfies $f(0) = 1$ and $x_0 \in [1/2, 2]$. The conclusion of the previous paragraph implies

$$\sup_{\mathcal{T}} \widehat{c}_k = \sup_{\mathcal{T}'} \widehat{c}_k.$$

The space \mathcal{T}' is compact for the Hausdorff topology, and since $\widehat{c}_k|_{\mathcal{T}'}$ is continuous, its restriction to \mathcal{T}' achieves its maximum. \square

We set

$$\Delta_{\pm} := \{(x, y) \in (0, \infty)^2 \mid \pm x < \pm y\}, \quad \Delta := \{(x, x) \in \mathbb{R}^2 \mid x > 0\},$$

and

$$\partial_{\pm}\Omega := \Delta_{\pm} \cap \partial\Omega, \quad \partial_0\Omega := \Delta \cap \partial\Omega, \quad \partial_*\Omega := \partial_+\Omega \cup \partial_0\Omega \cup \partial_-\Omega.$$

Notice that $\partial_0\Omega$ is a singleton, and Ω is the convex set enclosed by $\partial_*\Omega$ together with two segments on the x and y axes.

Lemma 5.3. *Assume that X_Ω maximizes \widehat{c}_k over the space of convex toric domains. Then $\partial_*\Omega$ is piecewise linear with at most $k-1$ corners.*

Proof. For each $v \in V_k$ there exists $w_v \in \partial_*\Omega$ such that $\|v\|_\Omega = \langle v, w_v \rangle$. The point w_v is not necessarily unique, but for $v = (0, k)$ we choose $w_v = (0, y_0)$, and for $v = (k, 0)$ we choose $w_v = (x_0, 0)$. We denote by $\Omega' \subset \Omega$ the convex hull of the finite set

$$\{(0, 0)\} \cup \{w_v \mid v \in V_k\} \subset \partial\Omega.$$

Notice that $X_{\Omega'}$ is a convex toric domain with $c_k(\Omega') = c_k(\Omega)$. Therefore, $\widehat{c}_k(\Omega') \geq \widehat{c}_k(\Omega)$, with equality if and only if $\Omega = \Omega'$. This shows that $\partial_*\Omega$ is piecewise linear with at most $k-1$ corners. \square

Lemma 5.3, together with Proposition C, implies the claim of Proposition D(ii) in the case $k = 1$. Hence, from now on we consider $k \geq 2$. Moreover, in view of Lemma 5.3, from now on we only need to consider convex domains X_Ω whose profile Ω has piecewise linear boundary.


Lemma 5.4. *Assume that X_Ω maximizes \widehat{c}_k over the space of convex toric domains. For each corner w of $\partial_*\Omega$, there exists $v \in V_k$ such that $c_k(X_\Omega) = \|v\|_\Omega = \langle v, w \rangle$.*

Proof. If such a v does not exist, by chamfering Ω near w we would produce $\Omega' \subsetneq \Omega$ still defining a convex toric domain $X_{\Omega'}$ with the same capacity $c_k(X_{\Omega'}) = c_k(X_\Omega)$ but smaller volume $\text{vol}(X_{\Omega'}) < \text{vol}(X_\Omega)$, in contradiction to the fact that X_Ω maximizes \widehat{c}_k . \square

Lemma 5.5. *Assume that X_Ω maximizes \widehat{c}_k over the space of convex toric domains. Then $\partial_*\Omega$ is piecewise linear with at most two corners. Moreover:*

- (i) *If $\partial_*\Omega$ has no corners, then X_Ω is an ellipsoid.*
- (ii) *If $w = (x, x) \in \partial_0\Omega$ is a corner of $\partial_*\Omega$, then it is its only corner and $X_\Omega = P(x, x)$.*
- (iii) *$\partial_+\Omega$ contains at most one corner $w = (x, y)$ of $\partial_*\Omega$, and if such a corner exists then $y = y_0$.*
- (iv) *$\partial_-\Omega$ contains at most one corner $w = (x, y)$ of $\partial_*\Omega$, and if such a corner exists then $x = x_0$.*

The possible situations allowed by the last two points are described by the following pictures:



Proof. Assume that $w = (x, x) \in \partial_0\Omega$ is a corner of $\partial_*\Omega$. Then, by Lemma 5.4, there exists $v \in V_k$ such that $c_k(X_\Omega) = \|v\|_\Omega = \langle v, w \rangle$. The square $\Omega' := [0, x]^2$ is contained in Ω , and satisfies $c_k(X_{\Omega'}) = \|v\|_\Omega = c_k(X_\Omega)$. Since X_Ω maximizes \widehat{c}_k , we infer $X_\Omega = X_{\Omega'} = P(x, x)$.

Assume instead that $\partial_+\Omega$ contains a corner $w = (x, y)$. We claim that $y = y_0$ (and, in particular, that there are no other corners $w' = (x', y') \in \partial_+\Omega$ with $x' < x$). Indeed, once again by Lemma 5.4, there exists $v_j = (j, k - j) \in V_k$ such that $c_k(X_\Omega) = \|v_j\|_\Omega = \langle v_j, w \rangle$. Since $x < y$, we have

$$\begin{aligned} \langle v_h, w \rangle &> c_k(X_\Omega), & \text{if } 0 \leq h < j, \\ \langle v_h, w \rangle &< c_k(X_\Omega), & \text{if } j < h \leq k. \end{aligned}$$

This implies that the convex domain $\Omega' := (\mathbb{R} \times [0, y]) \cap \Omega$, which is contained in Ω , has the same capacity $c_k(X_{\Omega'}) = c_k(X_\Omega)$. Therefore, since X_Ω is a maximizer of \widehat{c}_k , we infer $\Omega' = \Omega$, and so $y = y_0$.

An analogous argument shows that, if $\partial_-\Omega$ contains a corner $w = (x, y)$, then $x = x_0$ (and then, in particular, there are no other corners $w' = (x', y') \in \partial_+\Omega$ with $x' > x$). \square

By Lemma 5.5 it suffices to consider regions of the form

$$\Omega = \Omega(\alpha, s, t) := \{0 \leq y \leq 1 + s\} \cap \{0 \leq x \leq 1 + t\} \cap \{1 - \alpha(x - 1) \leq y\}$$

for $\alpha \in [0, \infty]$, $s \in [0, \alpha]$ and $t \in [0, 1/\alpha]$. The corners of $\partial_*\Omega$ are

$$v^+ := (1 - \frac{s}{\alpha}, 1 + s), \quad v^- := (1 + t, 1 - \alpha t).$$

It will be convenient to parametrize α as

$$\alpha = \alpha(i, r) = \frac{i + r}{k - (i + r)}$$

where $(i, r) \in \{0, 1, \dots, k - 1\} \times [0, 1) \cup \{(k, 0)\}$. As above we set $v_j = (j, k - j)$, $j = 1, \dots, k$.

Lemma 5.6. *We have*

$$\begin{aligned} \|v_j\|_\Omega &= \langle v_j, v^+ \rangle, & \forall j \leq i, \\ \|v_j\|_\Omega &= \langle v_j, v^- \rangle, & \forall j \geq i + 1. \end{aligned}$$

Proof. The maximum $\|v_j\|_\Omega = \max_{w \in \Omega}$ is attained at v^+ or v^- . Whether it is attained at v^+ or v^- if $v^+ \neq v^-$ depends on which side of the line l through the origin perpendicular to the line through v^+ and v^- the point v_j lies. Since the slope of l is $1/\alpha$, the claim follows. \square

Lemma 5.7. *We have*

$$\begin{aligned} \|v_j\|_\Omega &\geq \|v_i\|_\Omega = k \left(1 + \frac{sr}{i + r}\right), & \forall j \leq i, \\ \|v_j\|_\Omega &\geq \|v_{i+1}\|_\Omega = k \left(1 + \frac{t(1-r)}{k - (i + r)}\right), & \forall j \geq i + 1. \end{aligned}$$

Proof. The inequalities follow from the facts that the coordinates of v_j sum up to k and that the x -coordinate of v^+ (v^-) is not larger (smaller) than its y -coordinate. For the equalities we compute using Lemma 5.6

$$\begin{aligned} \|v_i\|_\Omega &= \langle v_i, v^+ \rangle = i - \frac{s}{\alpha}i + k - i + ks - is = k \left(1 + s \left(1 - i \frac{\alpha + 1}{k\alpha}\right)\right) \\ &= k \left(1 + s \left(1 - i \frac{1}{i + r}\right)\right) = k \left(1 + \frac{sr}{i + r}\right) \end{aligned}$$

and

$$\begin{aligned} \|v_{i+1}\|_\Omega &= \langle v_{i+1}, v^- \rangle = i + it + 1 + t + k - i - 1 - \alpha tk + \alpha it + \alpha t \\ &= k + t(1 + i - \alpha(k - i - 1)) \\ &= k + t \frac{(1 + i)(k - (i + r)) - (k - (i + 1))(i + r)}{k - (i + r)} \\ &= k \left(1 + \frac{t(1 - r)}{k - (i + r)}\right). \end{aligned} \quad \square$$

Using elementary geometry we compute the Euclidean area of Ω to be

$$\text{area}(\Omega) = 1 + t + s - \frac{s^2}{2\alpha} - \frac{t^2\alpha}{2}.$$

Hence, with $\text{vol}(X_\Omega) = 2 \text{area}(\Omega)$ and Lemma 5.7 we obtain

$$2\widehat{c}_k(X_\Omega)^2 = \frac{c_k(X_\Omega)^2}{\text{area}(\Omega)} = \frac{k^2 \left(1 + \min \left\{ \frac{sr}{i+r}, \frac{t(1-r)}{k-(i+r)} \right\}\right)^2}{1+t+s - \frac{s^2}{2\alpha} - \frac{t^2\alpha}{2}} \quad (5.2)$$

We claim that $2\widehat{c}_k(X_\Omega)^2 \leq k^2$ with equality if and only if $s = t = 0$ (in which case Ω is independent of α), or $k = 2$ and $X_\Omega = E(2, 1)$. Showing this claim will complete the proof of Proposition D(ii). The claim is equivalent to

$$\left(1 + \min \left\{ \frac{sr}{i+r}, \frac{t(1-r)}{k-(i+r)} \right\}\right)^2 \leq 1+t+s - \frac{s^2}{2\alpha} - \frac{t^2\alpha}{2} \quad (5.3)$$

with equality if and only if $s = t = 0$, or $k = 2$ and $X_\Omega = E(2, 1)$.

Since we have already treated the case of rectangles in Example 5.1, we assume that $0 < i+r < k$, i.e. $\alpha \in (0, \infty)$. In particular, we have $r < 1$. Moreover, we can assume that $\frac{sr}{i+r} = \frac{t(1-r)}{k-(i+r)}$, i.e.

$$t = \frac{r}{1-r} \frac{s}{\alpha}; \quad (5.4)$$

indeed, otherwise we could decrease the volume $\text{vol}(X_\Omega)$ by decreasing t or s without changing $c_k(X_\Omega)$. It suffices to show that we have a strict inequality in (5.3) in the case $s \neq 0$ unless $k = 2$ and $X_\Omega = E(2, 1)$. For $s \neq 0$ a strict inequality in (5.3) is (using (5.4)) equivalent to

$$\begin{aligned} 2s \frac{r}{i+r} + s^2 \frac{r^2}{(i+r)^2} &< s \left(1 + \frac{r}{(1-r)\alpha}\right) - \frac{s^2}{2\alpha} \left(1 + \frac{r^2}{(1-r)^2}\right) \\ \Leftrightarrow \frac{s}{\alpha} \left(\frac{r^2\alpha}{(i+r)^2} + \frac{(1-r)^2 + r^2}{2(1-r)^2}\right) &< 1 + \frac{r}{(1-r)\alpha} - \frac{2r}{i+r}. \end{aligned} \quad (5.5)$$

Note that $\frac{s}{\alpha} \leq \min \left\{1, \frac{1-r}{r\alpha}\right\}$. Therefore, the sharp inequality (5.5) is implied both by

$$\frac{r^2\alpha}{(i+r)^2} + \frac{(1-r)^2 + r^2}{2(1-r)^2} < 1 + \frac{r}{(1-r)\alpha} - \frac{2r}{i+r} \quad (5.6)$$

and by

$$\frac{(1-r)r}{(i+r)^2} + \frac{(1-r)^2 + r^2}{2(1-r)r\alpha} < 1 + \frac{r}{(1-r)\alpha} - \frac{2r}{i+r}. \quad (5.7)$$

Let us first look at inequality (5.6).

Lemma 5.8. *For $0 < i+r < k$ inequality (5.6) is equivalent to*

$$0 < ((k-i)(k-i-2) + r)r + (i+r)(k-i-r) \frac{1-2r}{2(1-r)}. \quad (5.8)$$

Proof. Indeed, we have

$$\begin{aligned} (5.6) &\Leftrightarrow \frac{r^2}{(i+r)(k-i-r)} + \frac{2r}{i+r} < \frac{r(k-i-r)}{(1-r)(i+r)} + 1 - \frac{(1-r)^2 + r^2}{2(1-r)^2} \\ &\Leftrightarrow (r^2 + 2r(k-i-r))(1-r) < r(k-i-r)^2 + (i+r)(k-i-r) \frac{1-2r}{2(r-1)^2} \\ &\Leftrightarrow 0 < r(k^2 + i^2 - 2ik - 2k + 2i + r) + (i+r)(k-i-r) \frac{1-2r}{2(r-1)} \\ &\Leftrightarrow (5.8). \end{aligned} \quad \square$$

Let us provide sufficient conditions for inequality (5.8), and hence for (5.5), to hold.

Lemma 5.9. *For $0 < i + r < k$ inequality (5.8) holds if*

- (i) $i \leq k - 2$ and $r \in [0, \frac{1}{2}]$,
- (ii) $i = k - 1$ and $r \in [0, u_{k-1} := \frac{k-1}{2k-1})$,
- (iii) $i = 0$ and $r \in [0, u_0 := \frac{2k^2-3k}{2k^2-2k-1})$.

Proof. Statement (i) is immediate from Lemma 5.8. For $i = k - 1$, inequality (5.8) is equivalent to

$$\begin{aligned} 0 &< (-1+r)r + (k-1+r)(1-r)\frac{1-2r}{2(1-r)} \\ \Leftrightarrow 0 &< 2r^2 - 2r + k - 1 + r - 2rk + 2r - 2r^2 \\ \Leftrightarrow r &< u_{k-1} = \frac{k-1}{2k-1}, \end{aligned}$$

which shows (ii). For $i = 0$ we have $r > 0$ and inequality (5.8) is equivalent to

$$\begin{aligned} 0 &< (k(k-2)+r)r + r(k-r)\frac{1-2r}{2(1-r)} \\ \Leftrightarrow (2r-1)(k-r) &< 2(1-r)(k(k-2)+r) \\ \Leftrightarrow 2rk - 2r^2 - k + r &< 2k^2 - 4k + 2r - 2rk^2 + 4rk - 2r^2 \\ \Leftrightarrow r &< u_0 = \frac{2k^2-3k}{2k^2-2k-1}, \end{aligned}$$

which shows (iii) and completes the proof of the lemma. \square

Let us now look at inequality (5.7).

Lemma 5.10. *For $0 < i + r < k$ inequality (5.7) is equivalent to*

$$r < i^2 + \frac{(i+r)^2(2r-1)}{2(1-r)r\alpha}. \quad (5.9)$$

Proof. Indeed, we have

$$\begin{aligned} (5.7) \Leftrightarrow \frac{r}{(i+r)^2} ((1-r) + 2(i+r)) &< 1 + \frac{1}{2(1-r)r\alpha} (2r^2 - (1-r)^2 - r^2) \\ \Leftrightarrow r - r^2 + 2ir + 2r^2 &< i^2 + 2ir + r^2 + \frac{(i+r)^2(2r-1)}{2(1-r)r\alpha} \\ \Leftrightarrow (5.9). \end{aligned} \quad \square$$

Now we provide sufficient conditions for inequality (5.8) and hence further conditions for (5.5) to hold.

Lemma 5.11. *For $0 < i + r < k$, inequality (5.9) holds if*

- (i) $i > 0$ and $r \in [\frac{1}{2}, 1)$,
- (ii) $i = 0$ and $r \in (l_0 := \frac{k}{2k-1}, 1)$,
- (iii) $i = k - 1$ and $r \in (l_{k-1} := \frac{k-1}{2k^2-2k-1}, 1)$.

Proof. Statement (i) is immediate from Lemma 5.10. For $i = 0$ we have $r > 0$ and inequality (5.8) is equivalent to

$$\begin{aligned} r &< \frac{r^2(2r-1)(k-r)}{2(1-r)r^2} = \frac{(2r-1)(k-r)}{2(1-r)}. \\ \Leftrightarrow 2r - 2r^2 &< 2kr - k - 2r^2 + r \\ \Leftrightarrow l_0 = \frac{k}{2k-1} &< r, \end{aligned}$$

which shows (ii). For $i = k-1$ inequality (5.8) is equivalent to

$$\begin{aligned} r &< (k-1)^2 + \frac{(k-1+r)^2(1-r)(2r-1)}{2(1-r)r(k-1+r)} \\ \Leftrightarrow r &< (k-1)^2 + \frac{(k-1+r)(2r-1)}{2r} \\ \Leftrightarrow 2r^2 &< 2rk^2 - 4rk + 2r + 2rk - 2r + 2r^2 - k + 1 - r \\ \Leftrightarrow l_{k-1} = \frac{k-1}{2k^2-2k-1} &< r, \end{aligned}$$

which shows (iii) and completes the proof of the lemma. \square

It remains to show that the conditions in Lemma 5.9 and Lemma 5.11 imply a strict inequality in (5.3) unless $k = 2$, and to study the equality case for $k = 2$. By part (i) of Lemma 5.9 and Lemma 5.11 it suffices to look at the cases $i = k-1$ and $i = 0$. This will be done in the subsequent two lemmas.

Lemma 5.12. *We have $l_{k-1} \leq u_{k-1}$, and $l_{k-1} < u_{k-1}$ unless $k = 2$. In the case $k = 2$ we get an equality in (5.3) if and only if $\alpha = \alpha(i, r) = 2$, $s = \alpha = 2$ and $t = \frac{1}{2}$, in which case $X_\Omega = E(2, 1)$.*

Proof. A quick computation using the expressions for l_{k-1} and u_{k-1} from Lemma 5.9 and Lemma 5.11 (still assuming $k \geq 2$) confirms that $l_{k-1} \leq u_{k-1}$ with equality if and only if $k = 2$. Hence, for the equality discussion it suffices to consider the case $k = 2$, $i = 1$ and $r = l_1 = u_1 = \frac{1}{3}$. Then $\alpha = \alpha(i, r) = 2$. Plugging this into (5.5) with an equality instead of an inequality implies that $s = 2$ and by (5.4) thus $t = \frac{1}{2}$. \square

Lemma 5.13. *We have $l_0 \leq u_0$, and $l_0 < u_0$ unless $k = 2$. In the case $k = 2$, $r = l_1 = u_1 = \frac{2}{3}$ and $i = 0$ we get an equality in (5.3) if and only if $\alpha = \alpha(i, r) = \frac{1}{2}$, $s = \alpha = \frac{1}{2}$ and $t = 2$, in which case $X_\Omega = E(2, 1)$.*

Proof. A quick computation using the expressions for l_0 and u_0 from Lemma 5.9 and Lemma 5.11 (still assuming $k \geq 2$) shows that $l_0 \leq u_0$ is equivalent to $3k \leq k^2 + 2$. It follows that equality holds if and only if $k = 2$. Hence, for the equality discussion it suffices to consider the case $k = 2$, $i = 0$ and $r = l_1 = u_1 = \frac{2}{3}$. Then $\alpha = \alpha(i, r) = \frac{1}{2}$. Plugging this into (5.5) with an equality instead of an inequality implies that $s = \frac{1}{2}$ and by (5.4) thus $t = 2$. \square

This completes the proof of Proposition D(ii).

APPENDIX A. THE FADELL-RABINOWITZ INDEX

Consider the universal bundle $ES^1 \rightarrow BS^1$. The classifying space BS^1 has cohomology ring $H^*(BS^1; \mathbb{Q}) = \mathbb{Q}[e]$, where e is a generator of $H^2(BS^1; \mathbb{Q})$. From now on, all cohomology rings will be assumed to have rational coefficients, and we will suppress \mathbb{Q} from the notation. For each non-empty topological space X equipped with an S^1 action, we consider the S^1 -equivariant cohomology

$$H_{S^1}^*(X) := H^*(X \times_{S^1} ES^1).$$

Here, $X \times_{S^1} ES^1 := (X \times ES^1)/S^1$ and the circle S^1 acts diagonally on the product. If $Y \subset X$ is an S^1 -invariant subspace, the S^1 -equivariant relative cohomology is defined by

$$H_{S^1}^*(X, Y) := H^*(X \times_{S^1} ES^1, Y \times_{S^1} ES^1).$$

Let $\pi : X \times_{S^1} ES^1 \rightarrow BS^1$, $\pi([x, y]) = [y]$ be the quotient-projection. With an abuse of notation, we will still denote by e the cohomology class $\pi^*e \in H_{S^1}^2(X)$. The *Fadell-Rabinowitz index* of X , first introduced in [FR78], is defined as⁴

$$\text{ind}_{\text{FR}}(X) = \inf \{k \geq 0 \mid e^k = 0 \text{ in } H_{S^1}^*(X)\},$$

with the usual convention $\inf \emptyset = \infty$. Its elementary properties are the following:

- **(Non-triviality)** $\text{ind}_{\text{FR}}(X) = 0$ if and only if $X = \emptyset$.
- **(Monotonicity)** $\text{ind}_{\text{FR}}(X) \leq \text{ind}_{\text{FR}}(Y)$ if there exists an S^1 -equivariant continuous map $f : X \rightarrow Y$.
- **(Subadditivity)** $\text{ind}_{\text{FR}}(X) \leq \text{ind}_{\text{FR}}(Y) + \text{ind}_{\text{FR}}(Z)$ if $X = \text{int}(Y) \cup \text{int}(Z)$, where the interior is taken with respect to the subspace topology.

Remark A.1. In the literature, the cohomology employed in the definition of the Fadell-Rabinowitz index is often the Alexander-Spanier one [Spa95, Section 6.4], which has a suitable continuity property [HZ94, Appendix A.8]: if a topological space is metrizable, the cohomology of any subspace is isomorphic to the direct limit of the cohomology of its open neighborhoods. We shall equivalently employ the more common singular cohomology, which does not satisfy such a strong continuity property, and circumvent the issue by taking open neighborhoods in statements such as the Lusternik-Schnirelmann property of spectral invariants (Section 2.2), Proposition A.4 below, and in the definition of the Ekeland-Hofer index (Section 3.1).

In the proof of Proposition 3.1, we need some properties of the Fadell-Rabinowitz index that are essentially established in the original source [FR78] and in Fadell-Husseini-Rabinowitz [FHR82], but stated under slightly different assumptions or contained within the proofs of some statements. We provide a self-contained account here for the reader's convenience. For any subset B of a vector space, we denote

$$B_* := B \setminus \{0\}.$$

Throughout this appendix, we denote by S^1 the unit circle in \mathbb{C} . We shall consider vector spaces $\mathbb{C}^m \oplus \mathbb{R}^n$, with $m \geq 1$ and $n \geq 2$, equipped with the S^1 action

$$e^{it} \cdot (z_1, \dots, z_m, x_1, \dots, x_n) = (e^{ik_1 t} z_1, \dots, e^{ik_m t} z_m, x_1, \dots, x_n),$$

⁴In the literature the Fadell-Rabinowitz index is occasionally defined as $\text{ind}_{\text{FR}}(X) - 1$.

for some positive integers k_1, \dots, k_m . The fixed point set of this action is the vector subspace $\mathbb{R}^n \equiv \{0\} \oplus \mathbb{R}^n$. On $\mathbb{C}^m \oplus \mathbb{R}^n$, we also have the action of the multiplicative group $(0, \infty)$ given by the scalar multiplication, i.e.

$$c \cdot (z_1, \dots, z_m, x_1, \dots, x_n) = (cz_1, \dots, cz_m, cx_1, \dots, cx_n).$$

Overall, the S^1 and $(0, \infty)$ actions give a $\mathbb{C}_* \equiv S^1 \times (0, \infty)$ action. Direct sum of vector spaces as above will be implicitly equipped with the product \mathbb{C}_* action. We say that a subset of $\mathbb{C}^m \oplus \mathbb{R}^n$ is \mathbb{C} -invariant when it is \mathbb{C}_* -invariant and contains the origin. The first lemma is a special case of [FR78, Prop. 4.3].

Lemma A.2. *For each \mathbb{C} -invariant subset $Y \subset \mathbb{C}^m$, we have*

$$\text{ind}_{\text{FR}}((Y \oplus \mathbb{C})_*) = \text{ind}_{\text{FR}}(Y_*) + 1.$$

Proof. We set $X := Y \oplus \mathbb{C}$. Since $X_* = (Y_* \oplus \mathbb{C}) \cup (Y \oplus \mathbb{C}_*)$, by the subadditivity of the Fadell-Rabinowitz index we have

$$\begin{aligned} \text{ind}_{\text{FR}}(X_*) &\leq \text{ind}_{\text{FR}}(Y_* \oplus \mathbb{C}) + \text{ind}_{\text{FR}}(Y \oplus \mathbb{C}_*) \\ &= \text{ind}_{\text{FR}}(Y_*) + \text{ind}_{\text{FR}}(\mathbb{C}_*) = \text{ind}_{\text{FR}}(Y_*) + 1. \end{aligned}$$

Since $Y_* \subset X_*$, we have $\text{ind}_{\text{FR}}(Y_*) \leq \text{ind}_{\text{FR}}(X_*)$. Assume by contradiction that $k := \text{ind}_{\text{FR}}(Y_*) = \text{ind}_{\text{FR}}(X_*)$. We consider the commutative diagram

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^{2k-1}(X_*) & \xrightarrow{\pi_*} & H_{S^1}^{2k-2}(X_*) & \xrightarrow{\sim e} & H_{S^1}^{2k}(X_*) & \longrightarrow & \dots \\ & & \downarrow i^*=0 & & \downarrow & & \downarrow & & \\ \dots & \longrightarrow & H^{2k-1}(Y_*) & \xrightarrow{\pi'_*} & H_{S^1}^{2k-2}(Y_*) & \xrightarrow{\sim e} & H_{S^1}^{2k}(Y_*) & \longrightarrow & \dots \end{array}$$

where the vertical homomorphisms are induced by the inclusions, and both rows are Gysin exact sequences. Notice that $i^* = 0$, since the inclusion $i : Y_* \hookrightarrow X_*$ is homotopic to a constant. Since $k = \text{ind}_{\text{FR}}(X_*)$, we have $e^{k-1} = \pi_*(\mu)$ for some $\mu \in H^{2k-1}(X_*)$, and therefore $e^{k-1} = \pi'_*(i^*(\mu)) = 0$ in $H_{S^1}^{2k-2}(Y_*)$, contradicting the fact that $\text{ind}_{\text{FR}}(Y_*) = k$. \square

Since S^1 acts trivially on \mathbb{R}^n , we have the Künneth isomorphism

$$H_{S^1}^*(\mathbb{R}_*^n) = H^*(BS^1 \times \mathbb{R}_*^n) \cong H^*(BS^1) \otimes H^*(\mathbb{R}_*^n).$$

In particular, $H^*(\mathbb{R}_*^n) \equiv H^0(BS^1) \otimes H^*(\mathbb{R}_*^n)$ is naturally a subgroup of $H_{S^1}^*(\mathbb{R}_*^n)$.

Lemma A.3. *Let $Y \subset \mathbb{C}^m$ be a \mathbb{C} -invariant subset, μ a generator of $H^{n-1}(\mathbb{R}_*^n)$, and $\delta : H_{S^1}^*(\mathbb{R}_*^n) \rightarrow H_{S^1}^{*+1}((Y \oplus \mathbb{R}^n)_*, \mathbb{R}_*^n)$ the connecting homomorphism. Then*

$$\text{ind}_{\text{FR}}(Y_*) = \inf \{k \geq 0 \mid e^k \smile \delta(\mu) = 0 \text{ in } H_{S^1}^*((Y \oplus \mathbb{R}^n)_*, \mathbb{R}_*^n)\}.$$

Proof. We set $X := Y \oplus \mathbb{R}^n$, decompose X_* as the union $X_* = (Y_* \oplus \mathbb{R}^n) \cup (Y \oplus \mathbb{R}_*^n)$, and consider the associated Mayer-Vietoris sequence in S^1 -equivariant cohomology

$$\dots \longrightarrow H_{S^1}^*(X_*) \xrightarrow{a^*} H_{S^1}^*(Y_* \oplus \mathbb{R}^n) \oplus H_{S^1}^*(Y \oplus \mathbb{R}_*^n) \xrightarrow{b^*} H_{S^1}^*(Y_* \oplus \mathbb{R}_*^n) \longrightarrow \dots$$

Since Y and \mathbb{R}^n are S^1 -equivariantly contractible spaces, we have $H_{S^1}^*(Y_* \oplus \mathbb{R}^n) \cong H_{S^1}^*(Y_*)$ and $H_{S^1}^*(Y \oplus \mathbb{R}_*^n) \cong H_{S^1}^*(\mathbb{R}_*^n)$. Moreover, since S^1 acts trivially on \mathbb{R}^n , we have $H_{S^1}^*(Y_* \oplus \mathbb{R}_*^n) \cong H_{S^1}^*(Y_*) \otimes H^*(\mathbb{R}_*^n)$. Therefore, the Mayer-Vietoris sequence can be rewritten as

$$\dots \longrightarrow H_{S^1}^*(X_*) \xrightarrow{a^*} H_{S^1}^*(Y_*) \oplus H_{S^1}^*(\mathbb{R}_*^n) \xrightarrow{b^*} H_{S^1}^*(Y_*) \otimes H^*(\mathbb{R}_*^n) \longrightarrow \dots$$

Let us now consider the long exact sequence of the pair $\mathbb{R}_*^n \subset X_*$, which reads

$$\dots \longrightarrow H_{S^1}^*(X_*) \xrightarrow{j^*} H_{S^1}^*(\mathbb{R}_*^n) \xrightarrow{\delta} H_{S^1}^{*+1}(X_*, \mathbb{R}_*^n) \longrightarrow \dots$$

Let μ be a generator of $H^{n-1}(\mathbb{R}_*^n)$, and $k := \text{ind}_{\text{FR}}(Y_*)$. We have a non-zero cohomology class

$$e^{k-1} \otimes \mu \in H^*(BS^1) \otimes H^*(\mathbb{R}_*^n) = H_{S^1}^*(\mathbb{R}_*^n)$$

Let us assume by contradiction that $\delta(e^{k-1} \otimes \mu) = 0$, so that there exists a cohomology class $\nu \in H_{S^1}^*(X_*)$ such that $j^*\nu = e^{k-1} \otimes \mu$. In the above Mayer-Vietoris sequence, we have $a^*(\nu) = (\nu|_{Y_*}, e^{k-1} \otimes \mu)$, and therefore

$$0 = b^*(\nu|_{Y_*}, e^{k-1} \otimes \mu) = \nu|_{Y_*} \otimes 1 - e^{k-1}|_{Y_*} \otimes \mu,$$

which gives a contradiction, since $e^{k-1}|_{Y_*} \otimes \mu \neq 0$. This shows that

$$e^{k-1} \smile \delta(1 \otimes \mu) = \delta(e^{k-1} \smile \mu) \neq 0.$$

On the other hand, since $\text{ind}_{\text{FR}}(Y_*) = k$, we have

$$b^*(0, e^k \otimes \mu) = -e^k|_{Y_*} \otimes \mu = 0.$$

Therefore, there exists $\eta \in H_{S^1}^*(X_*)$ such that $a^*(\eta) = (0, e^k \otimes \mu)$. This implies $j^*(\eta) = e^k \otimes \mu$, and we conclude

$$e^k \smile \delta(1 \otimes \mu) = \delta(e^k \smile \mu) = \delta j^*(\eta) = 0. \quad \square$$

The last statement is a slight generalization of [FHR82, Prop. 2.2].

Proposition A.4. *Let $Y \subset \mathbb{C}^p$ be a \mathbb{C} -invariant subset, $B \subset \mathbb{C}^p \oplus \mathbb{R}^n$ an S^1 -invariant compact neighborhood of the origin, and $\psi : Y \oplus \mathbb{R}^n \rightarrow \mathbb{C}^q \oplus \mathbb{R}^n$ an S^1 -equivariant continuous map such that $\psi(0, x) = (0, x)$ for all $x \in \mathbb{R}^n$. If $\text{ind}_{\text{FR}}(Y_*) > q$, then any S^1 -invariant open subset $U \subset Y \oplus \mathbb{R}^n$ containing the intersection $\psi^{-1}(0) \cap \partial B$ has Fadell-Rabinowitz index $\text{ind}_{\text{FR}}(U) \geq \text{ind}_{\text{FR}}(Y_*) - q$.*

Proof. We set $k := \text{ind}_{\text{FR}}(Y_*)$ and $X := Y \oplus \mathbb{R}^n$. Assume by contradiction that there exists an S^1 -invariant open subset $U \subset X_*$ containing $Z := \psi^{-1}(0) \cap \partial B$ such that $\text{ind}_{\text{FR}}(U) < k - q$. In particular, since $\text{ind}_{\text{FR}}(U)$ is finite, $U \cap \mathbb{R}^n = \emptyset$. Let $V \subset X_* \setminus Z$ be an S^1 -invariant open subset containing $X \cap \partial B \setminus U$ and that is sufficiently small so that $X \cap \psi^{-1}(0) = \emptyset$. Finally, we set $W := U \cup V$, which is an S^1 -invariant open subset of X_* containing $X \cap \partial B$.

We defined the S^1 -invariant subsets $X' := (X_* \cap B) \cup W$ and $X'' := (X_* \setminus X') \cup W$, which are both open subsets of X_* , have union $X' \cup X'' = X_*$ and intersection $X' \cap X'' = W$. Notice that, for each $x \in X_*$ there exists $\epsilon > 0$ small enough so that $\epsilon x \in X'$ and $\epsilon^{-1}x \in X''$. This readily implies that both inclusions

$$(X', X' \cap \mathbb{R}_*^n) \hookrightarrow (X_*, \mathbb{R}_*^n), \quad (X'', X'' \cap \mathbb{R}_*^n) \hookrightarrow (X_*, \mathbb{R}_*^n)$$

admit an S^1 -equivariant homotopic right inverse, and therefore induce monomorphisms

$$\begin{aligned} H_{S^1}^*(X_*, \mathbb{R}_*^n) &\hookrightarrow H_{S^1}^*(X', X' \cap \mathbb{R}_*^n), \\ H_{S^1}^*(X_*, \mathbb{R}_*^n) &\hookrightarrow H_{S^1}^*(X'', X'' \cap \mathbb{R}_*^n). \end{aligned}$$

This, together with the Mayer-Vietoris exact sequence

$$\dots \rightarrow H_{S^1}^*(X_*, \mathbb{R}_*^n) \rightarrow \begin{array}{c} H_{S^1}^*(X', X' \cap \mathbb{R}_*^n) \\ \oplus \\ H_{S^1}^*(X'', X'' \cap \mathbb{R}_*^n) \end{array} \rightarrow H_{S^1}^*(W, W \cap \mathbb{R}_*^n) \rightarrow \dots$$

readily implies that the inclusion induces a monomorphism

$$H_{S^1}^*(X_*, \mathbb{R}_*^n) \hookrightarrow H_{S^1}^*(W, W \cap \mathbb{R}_*^n). \quad (\text{A.1})$$

Analogously, since both inclusions $X' \cap \mathbb{R}_*^n \hookrightarrow \mathbb{R}_*^n$ and $X'' \cap \mathbb{R}_*^n \hookrightarrow \mathbb{R}_*^n$ admit a homotopic right inverse, they induce monomorphisms in cohomology, and the Mayer-Vietoris sequence

$$\dots \rightarrow H^*(\mathbb{R}_*^n) \rightarrow H^*(X' \cap \mathbb{R}_*^n) \oplus H^*(X'' \cap \mathbb{R}_*^n) \rightarrow H^*(W \cap \mathbb{R}_*^n) \rightarrow \dots$$

readily implies that the inclusion induces a monomorphism

$$H^*(\mathbb{R}_*^n) \hookrightarrow H^*(W \cap \mathbb{R}_*^n). \quad (\text{A.2})$$

Consider the commutative diagram

$$\begin{array}{ccc} H_{S^1}^*(\mathbb{R}_*^n) & \xrightarrow{\delta} & H_{S^1}^{*+1}(X_*, \mathbb{R}_*^n) \\ \downarrow & & \downarrow \\ H_{S^1}^*(W \cap \mathbb{R}_*^n) & \xrightarrow{\delta} & H_{S^1}^{*+1}(W, W \cap \mathbb{R}_*^n) \\ \cong \downarrow & & \downarrow \\ H_{S^1}^*(V \cap \mathbb{R}_*^n) & \xrightarrow{\delta} & H_{S^1}^{*+1}(V, V \cap \mathbb{R}_*^n) \\ \uparrow & & \uparrow \psi^* \\ H_{S^1}^*(\mathbb{R}_*^n) & \xrightarrow{\delta} & H_{S^1}^{*+1}((\mathbb{C}^q \oplus \mathbb{R}^n)_*, \mathbb{R}_*^n) \end{array}$$

Here, the horizontal arrows are connecting homomorphisms, and the unspecified vertical arrows are homomorphisms induced by the inclusions. Let μ be a generator of $H^{n-1}(\mathbb{R}_*^n) \subset H_{S^1}^{n-1}(\mathbb{R}_*^n)$. By the injectivity of the homomorphism (A.2), in the above diagram μ is mapped to a non-zero element of $H_{S^1}^*(V \cap \mathbb{R}_*^n) = H_{S^1}^*(W \cap \mathbb{R}_*^n)$, which we still denote by μ . Since $\text{ind}_{\text{FR}}(Y_*) = k$, Lemma A.3 implies that

$$e^{k-1} \smile \delta(\mu) \neq 0 \text{ in } H_{S^1}^*(X_*, \mathbb{R}_*^n),$$

and therefore, by the injectivity of the homomorphism (A.1),

$$e^{k-1} \smile \delta(\mu) \neq 0 \text{ in } H_{S^1}^*(W, W \cap \mathbb{R}_*^n). \quad (\text{A.3})$$

Lemma A.2 implies $\text{ind}_{\text{FR}}(\mathbb{C}_*^q) = q$. Therefore, another application of Lemma A.3 implies that

$$e^q \smile \delta(\mu) = 0 \text{ in } H_{S^1}^*((\mathbb{C}^q \oplus \mathbb{R}^n)_*, \mathbb{R}_*^n).$$

By applying ψ^* , we obtain $e^q \smile \delta(\mu) = 0$ in $H_{S^1}^*(V, V \cap \mathbb{R}_*^n)$, and therefore

$$e^q \smile \delta(\mu) \in H_{S^1}^*(W, V).$$

Since $\text{ind}_{\text{FR}}(U) \leq k - q - 1$, we have

$$e^{k-q-1} \in H_{S^1}^*(W, U).$$

Finally

$$e^{k-1} \smile \delta(\mu) = e^{k-q-1} \smile (e^q \smile \delta(\mu)) \in H_{S^1}^*(W, U \cup V) = H_{S^1}^*(W, W) = 0,$$

which contradicts (A.3). \square

Acknowledgments. We are indebted with Alberto Abbondandolo, who first noticed while working with Lange and Mazzucchelli on [ALM22] that the higher capacity ratios of the polydisk $P(1, 1)$ were higher than the ones of 4-dimensional ellipsoids, and who suggested to improve an earlier formulation of Theorem A beyond the space of smooth convex bodies. We also thank Jean Gutt, Felix Schlenk, and Jean-Claude Sikorav for some helpful conversations on symplectic capacities.

REFERENCES

- [AAKO14] Shiri Artstein-Avidan, Roman Karasev, and Yaron Ostrover, *From symplectic measurements to the Mahler conjecture*, Duke Math. J. **163** (2014), no. 11, 2003–2022. MR3263026
- [AAMO08] Shiri Artstein-Avidan, Vitali Milman, and Yaron Ostrover, *The M -ellipsoid, symplectic capacities and volume*, Comment. Math. Helv. **83** (2008), no. 2, 359–369. MR2390048
- [AB19] Alberto Abbondandolo and Gabriele Benedetti, *On the local systolic optimality of Zoll contact forms*, 2019. arXiv:1912.04187.
- [ABHS18] Alberto Abbondandolo, Barney Bramham, Umberto L. Hryniewicz, and Pedro A. S. Salomão, *Sharp systolic inequalities for Reeb flows on the three-sphere*, Invent. Math. **211** (2018), no. 2, 687–778. MR3748316
- [AK22] Alberto Abbondandolo and Jungsoo Kang, *Symplectic homology of convex domains and Clarke’s duality*, Duke Math. J. **171** (2022), no. 3, 739–830. MR4382979
- [ALM22] Alberto Abbondandolo, Christian Lange, and Marco Mazzucchelli, *Higher systolic inequalities for 3-dimensional contact manifolds*, J. Éc. polytech. Math. **9** (2022), 807–851. MR4436034
- [APB14] J. C. Alvarez Paiva and F. Balacheff, *Contact geometry and isosystolic inequalities*, Geom. Funct. Anal. **24** (2014), no. 2, 648–669. MR3192037
- [BK21] Gabriele Benedetti and Jungsoo Kang, *A local contact systolic inequality in dimension three*, J. Eur. Math. Soc. (JEMS) **23** (2021), no. 3, 721–764. MR4210723
- [CGM20] Daniel Cristofaro-Gardiner and Marco Mazzucchelli, *The action spectrum characterizes closed contact 3-manifolds all of whose Reeb orbits are closed*, Comment. Math. Helv. **95** (2020), no. 3, 461–481. MR4152621
- [Edt22] Oliver Edtmair, *Disk-like surfaces of section and symplectic capacities*, 2022. arXiv:2206.07847.
- [EH87] I. Ekeland and H. Hofer, *Convex Hamiltonian energy surfaces and their periodic trajectories*, Comm. Math. Phys. **113** (1987), no. 3, 419–469. MR925924
- [EH89] I. Ekeland and H. Hofer, *Symplectic topology and Hamiltonian dynamics*, Math. Z. **200** (1989), no. 3, 355–378. MR978597
- [EH90] Ivar Ekeland and Helmut Hofer, *Symplectic topology and Hamiltonian dynamics. II*, Math. Z. **203** (1990), no. 4, 553–567. MR1044064
- [FHR82] E. R. Fadell, S. Y. Husseini, and P. H. Rabinowitz, *Borsuk-Ulam theorems for arbitrary S^1 actions and applications*, Trans. Amer. Math. Soc. **274** (1982), no. 1, 345–360. MR670937
- [FR78] Edward R. Fadell and Paul H. Rabinowitz, *Generalized cohomological index theories for Lie group actions with an application to bifurcation questions for Hamiltonian systems*, Invent. Math. **45** (1978), no. 2, 139–174. MR478189
- [GGM21] Viktor L. Ginzburg, Başak Z. Gürel, and Marco Mazzucchelli, *On the spectral characterization of Besse and Zoll Reeb flows*, Ann. Inst. H. Poincaré C Anal. Non Linéaire **38** (2021), no. 3, 549–576. MR4227045
- [GH18] Jean Gutt and Michael Hutchings, *Symplectic capacities from positive S^1 -equivariant symplectic homology*, Algebr. Geom. Topol. **18** (2018), no. 6, 3537–3600. MR3868228

- [GL18] Hansjörg Geiges and Christian Lange, *Seifert fibrations of lens spaces*, Abh. Math. Semin. Univ. Hambg. **88** (2018), no. 1, 1–22. MR3785783
- [GR23] Jean Gutt and Vinicius G. B. Ramos, *The equivalence of Ekeland-Hofer and equivariant symplectic homology capacities*, 2023. forthcoming.
- [Her98] David Hermann, *Non-equivalence of symplectic capacities for open sets with restricted contact type boundary*, 1998. Prépublication d’Orsay numéro 32.
- [HZ94] Helmut Hofer and Eduard Zehnder, *Symplectic invariants and Hamiltonian dynamics*, Birkhäuser Advanced Texts, Birkhäuser Verlag, Basel, 1994. MR1306732
- [Moa94] Laurent Moatzy, *Capacités et équivalence symplectique des domaines de Reinhardt*, 1994. Ph.D. Thesis, Université Toulouse III - Paul Sabatier.
- [MR23] Marco Mazzucchelli and Marco Radeschi, *On the structure of Besse convex contact spheres*, Trans. Amer. Math. Soc. **376** (2023), no. 3, 2125–2153. MR4549701
- [MS17] Dusa McDuff and Dietmar Salamon, *Introduction to symplectic topology*, Third, Oxford Graduate Texts in Mathematics, Oxford University Press, Oxford, 2017. MR3674984
- [RW17] Marco Radeschi and Burkhard Wilking, *On the Berger conjecture for manifolds all of whose geodesics are closed*, Invent. Math. **210** (2017), no. 3, 911–962. MR3735632
- [Sik90] Jean-Claude Sikorav, *Systèmes hamiltoniens et topologie symplectique*, ETS Editrice Pisa, 1990.
- [Spa95] Edwin H. Spanier, *Algebraic topology*, Springer-Verlag, New York, 1995. Corrected reprint of the 1966 original. MR1325242
- [Vit00] Claude Viterbo, *Metric and isoperimetric problems in symplectic geometry*, J. Amer. Math. Soc. **13** (2000), no. 2, 411–431. MR1750956
- [Vit89] Claude Viterbo, *Capacités symplectiques et applications (d’après Ekeland-Hofer, Gromov)*, 1989, pp. Exp. No. 714, 345–362. Séminaire Bourbaki, Vol. 1988/89. MR1040580
- [Wad75] A. W. Wadsley, *Geodesic foliations by circles*, J. Differential Geometry **10** (1975), no. 4, 541–549. MR400257
- [Zha21] Bingyu Zhang, *Capacities from the Chiu-Tamarkin complex*, 2021. arXiv:2103.05143.

LUCA BARACCO AND OLGA BERNARDI
 UNIVERSITÀ DI PADOVA, DIPARTIMENTO DI MATEMATICA “TULLIO LEVI CIVITA”
 VIA TRIESTE 63, 35121 PADOVA, ITALY
Email address: baracco@math.unipd.it, obern@math.unipd.it

CHRISTIAN LANGE
 LUDWIG-MAXIMILIANS-UNIVERSITÄT MÜNCHEN, MATHEMATISCHES INSTITUT
 THERESIENSTRASSE 39, D-80333 MUNICH, GERMANY
Email address: lange@math.lmu.de

MARCO MAZZUCHELLI
 CNRS, UMPA, ÉCOLE NORMALE SUPÉRIEURE DE LYON
 46 ALLÉE D’ITALIE, 69364 LYON, FRANCE
Email address: marco.mazzucchelli@ens-lyon.fr