

On microfunctions at the boundary along CR manifolds

ANDREA D'AGNOLO¹ and GIUSEPPE ZAMPIERI²

¹*Mathématiques; Univ. Paris 6; 4, Place Jussieu; F-75252 Paris Cedex 05*

²*Dip. di Matematica, Università di Padova, via Belzoni 7, I-35131 Padova, Italy*

Received 16 August 1995; accepted in final form 4 December 1995

Abstract. Let X be a complex analytic manifold, $M \subset X$ a C^2 submanifold, $\Omega \subset M$ an open set with C^2 boundary $S = \partial\Omega$. Denote by $\mu_M(\mathcal{O}_X)$ (resp. $\mu_\Omega(\mathcal{O}_X)$) the microlocalization along M (resp. Ω) of the sheaf \mathcal{O}_X of holomorphic functions.

In the literature (cf. [A-G], [K-S 1,2]) one encounters two classical results concerning the vanishing of the cohomology groups $H^j \mu_M(\mathcal{O}_X)_p$ for $p \in T_M^* X$. The most general gives the vanishing outside a range of indices j whose length is equal to $s^0(M, p)$ (with $s^{+, -, 0}(M, p)$ being the number of respectively positive, negative and null eigenvalues for the 'microlocal' Levi form $L_M(p)$). The sharpest result gives the concentration in a single degree, provided that the difference $s^-(M, p') - \gamma(M, p')$ is locally constant for $p' \in T_M^* X$ near p (with $\gamma(M, p) = \dim^{\mathbb{C}}(T_M^* X \cap iT_M^* X)_z$ for z the base point of p).

The first result was restated for the complex $\mu_\Omega(\mathcal{O}_X)$ in [D'A-Z 2], in the case $\text{codim}_M S = 1$. We extend it here to any codimension and moreover we also restate for $\mu_\Omega(\mathcal{O}_X)$ the second vanishing theorem.

We also point out that the principle of our proof, related to a criterion for constancy of sheaves due to [K-S 1], is a quite new one.

Key words: Solvability of the $\bar{\partial}$ -complex.

Mathematics Subject Classifications (1991): 58G, 32F.

1. Notations

Let X be a complex analytic manifold and $M \subset X$ a C^2 submanifold. One denotes by $\pi: T^* X \rightarrow X$ and $\pi: T_M^* X \rightarrow M$ the cotangent bundle to X and the conormal bundle to M in X respectively. Let $\dot{T}^* X$ be the cotangent bundle with the zero section removed, and let $\rho: M \times_X T^* X \rightarrow T^* M$ be the projection associated to the embedding $M \hookrightarrow X$.

For a subset $A \subset X$ one defines the strict normal cone of A in X by $N^X(A) := TX \setminus C(X \setminus A, A)$ where $C(\cdot, \cdot)$ denotes the normal Whitney cone (cf [K-S 1]).

Let $z_0 \in M$, $p \in (T_M^* X)_{z_0}$. We put $T_{z_0}^{\mathbb{C}} M = T_{z_0} M \cap iT_{z_0} M$; $\lambda_M(p) = T_p T_M^* X$; $\lambda_0(p) = T_p(\pi^{-1}\pi(p))$, $\nu(p)$ = the complex Euler radial field at p , and we set $\gamma(M, p) = \dim^{\mathbb{C}}(T_M^* X \cap iT_M^* X)_{z_0}$. If no confusion may arise, we will sometimes drop the indices z_0 or p in the above notations.

Let ϕ be a C^2 function in X with $\phi|_M \equiv 0$ and $p = (z_0; d\phi(z_0))$. In a local system of coordinates (z) at z_0 in X we define $L_\phi(z_0)$ as the Hermitian form with matrix $(\partial z_i \bar{\partial} z_j \phi)_{i,j}$. Its restriction $L_M(p)$ to $T_{z_0}^{\mathbf{C}}M$ does not depend on the choice of ϕ and is called the Levi form of M at p . Let $s^{+, -, 0}(M, p)$ denote the number of respectively positive, negative and null eigenvalues of $L_M(p)$.

One denotes by $\mathbf{D}^b(X)$ the derived category of the category of bounded complexes of sheaves of \mathbf{C} -vector spaces and by $\mathbf{D}^b(X; p)$ the localization of $\mathbf{D}^b(X)$ at $p \in T^*X$, i.e. the localization of $\mathbf{D}^b(X)$ with respect to the null system $\{F \in \mathbf{D}^b(X); p \notin \text{SS}(F)\}$ (here $\text{SS}(F)$ denotes the micro-support in the sense of [K-S 2], a closed conic involutive subset of T^*X).

Remark 1.1. We recall that a complex F which verifies $\text{SS}(F) \subset T_M^*X$ in a neighborhood of $p \in T_M^*X$ is microlocally isomorphic (i.e. isomorphic in $\mathbf{D}^b(X; p)$) to a constant sheaf on M . This criterion, stated in [K-S 1] for a C^2 manifold M , extends easily to C^1 manifolds (cf [D'A-Z 1]).

Let \mathcal{O}_X be the sheaf of germs of holomorphic functions on X and \mathbf{C}_A , ($A \subset X$ locally closed), the sheaf which is zero in $X \setminus A$ and the constant sheaf with fiber \mathbf{C} in A . We shall consider the complex $\mu_A(\mathcal{O}_X) := \mu\text{hom}(\mathbf{C}_A, \mathcal{O}_X)$ of microfunctions along A (where $\mu\text{hom}(\cdot, \cdot)$ is the bifunctor of [K-S 1]). Special interest lies in the complexes $\mu_M(\mathcal{O}_X)$ and $\mu_\Omega(\mathcal{O}_X)$ for Ω being an open subset of the manifold M (cf [S]).

2. Statement of the results

Let X be a complex analytic manifold of dimension n , $M \subset X$ a C^2 submanifold of codimension l , $\Omega \subset M$ an open set with C^2 boundary $S = \partial\Omega$, and set $r = \text{codim}_M S$ (we assume Ω locally on one side of S for $r = 1$). Let $z_0 \in M$, $p \in (T_M^*X)_{z_0}$. Define

$$\begin{aligned} d_M(p) &= \text{codim}_X M + s^-(M, p) - \gamma(M, p), \\ c_M(p) &= n - s^+(M, p) + \gamma(M, p). \end{aligned}$$

Let us recall the following classical results concerning the cohomology of $\mu_M(\mathcal{O}_X)$.

THEOREM A. ([A-G], [K-S 1]) *Assume $\dim^{\mathbf{R}}(\nu(p) \cap \lambda_M(p)) = 1$. Then*

$$H^j \mu_M(\mathcal{O}_X)_p = 0 \quad \text{for } j \notin [d_M(p), c_M(p)].$$

THEOREM B. ([H], [K-S 1]) *Assume $\dim^{\mathbf{R}}(\nu(p) \cap \lambda_M(p)) = 1$ and $s^-(M, p') - \gamma(M, p') \equiv \text{const}$ for $p' \in T_M^*X$ close to p . Then*

$$H^j \mu_M(\mathcal{O}_X)_p = 0 \quad \text{for } j \neq d_M(p).$$

Dealing with $\mu_\Omega(\mathcal{O}_X)$ (and choosing now $p \in S \times_M \dot{T}_M^*X$), one knows that

THEOREM C. ([D'A-Z 2]) *Assume $\text{codim}_M S = 1$ and $\dim^{\mathbf{R}}(\nu(p) \cap \lambda_M(p)) = 1$. Then*

$$H^j \mu_\Omega(\mathcal{O}_X)_p = 0 \quad \text{for } j \notin [d_M(p), c_M(p)].$$

The aim of the present note is, on the one hand, to extend Theorem C to the case of any codimension for S in M , and, on the other hand, to state the analogue of Theorem B for the complex $\mu_\Omega(\mathcal{O}_X)$. We point out that the method of our proof, based on the criterion of [K-S 1, Proposition 6.2.2] (with its C^1 -variant of [D'A-Z 1]), is a quite new one.

Our results, valid for any $r = \text{codim}_M S$, go as follows.

THEOREM 2.1. *Assume*

$$\dim^{\mathbf{R}}(\nu(p) \cap \lambda_S(p)) = 1. \tag{2.1}$$

Then

$$H^j \mu_\Omega(\mathcal{O}_X)_p = 0 \quad \text{for } j \notin [d_M(p), c_M(p) + r - 1]. \tag{2.2}$$

When M is a real analytic manifold of dimension n and X a complexification of M , then Theorem 2.1 states the concentration in degree n for $\mu_\Omega(\mathcal{O}_X)_p$. This should be proved as well by the aid of Proposition 3.1 of [S]. In fact, since Ω has C^2 -boundary, then $M \setminus \Omega$ is C^ω -convex (i.e. convex in suitable real analytic coordinates at z_o).

THEOREM 2.2. *Assume (2.1) and moreover*

$$\begin{cases} s^-(M, p') - \gamma(M, p') \text{ is constant for } p' \in \overline{\Omega} \times_M T_M^*X \text{ near } p, \\ s^-(S, p') - \gamma(S, p') \text{ is constant} \\ \quad \text{for } p' \in T_S^*X \cap \rho^{-1}(N^M(\Omega)^{\circ a}) \text{ near } p, \\ s^-(M, p) - \gamma(M, p) = s^-(S, p) - \gamma(S, p). \end{cases} \tag{2.3}$$

Then

$$H^j \mu_\Omega(\mathcal{O}_X)_p = 0 \quad \text{for } j \notin [d_M(p), d_M(p) + r - 1].$$

Remark 2.3. We notice that the sets appearing in (2.3) are very natural in this context; one has in fact

$$\begin{cases} T_M^*X \cap \text{SS}(\mathbf{C}_\Omega) = \overline{\Omega} \times_M T_M^*X, \\ T_S^*X \cap \text{SS}(\mathbf{C}_\Omega) = T_S^*X \cap \rho^{-1}(N^M(\Omega)^{\circ a}). \end{cases}$$

3. Proofs of the results

Proof of Theorem 2.1. We set $\Omega^- = M \setminus \overline{\Omega}$ and use the distinguished triangle

$$\mu_S(\mathcal{O}_X) \rightarrow \mu_M(\mathcal{O}_X) \rightarrow \mu_\Omega(\mathcal{O}_X) \oplus \mu_{\Omega^-}(\mathcal{O}_X) \xrightarrow{+1}. \quad (3.4)$$

We remark that by its own definition: $L_S(p) = L_M(p)|_{T_{z_o}^{\mathbf{C}}S}$, ($p \in S \times_M \dot{T}_M^*X$).

This gives:

$$\begin{aligned} s^{+,-}(S, p) &\leq s^{+,-}(M, p) \leq s^{+,-}(S, p) + (\dim T_{z_o}^{\mathbf{C}}M - \dim T_{z_o}^{\mathbf{C}}S) \\ &= s^{+,-}(S, p) + (r + \gamma(M, p) - \gamma(S, p)). \end{aligned}$$

Thus if the integers $c_M(p)$, $d_M(p)$ and $c_S(p)$, $d_S(p)$ are defined as in Section 2, we have at once

$$\begin{aligned} c_M(p) &\leq c_S(p) \leq c_M(p) + r, \\ d_M(p) &\leq d_S(p) \leq d_M(p) + r. \end{aligned} \quad (3.5)$$

The vanishing of (2.2) for $j > c_M(p) + r - 1$ then follows by applying Theorem A to M and S .

The vanishing of (2.2) for $j < d_M(p)$ is immediate for $d_S(p) > d_M(p)$ due to Theorem A and (3.1).

When $d_S(p) = d_M(p)$ it remains to be proven that

$$H^{d_M(p)} \mu_S(\mathcal{O}_X)_p \rightarrow H^{d_M(p)} \mu_M(\mathcal{O}_X)_p \quad \text{is injective.} \quad (3.6)$$

To this end we perform a contact transformation χ near p which interchanges (setting $q = \chi(p)$)

$$\begin{cases} T_M^*X \rightarrow T_{\widetilde{M}}^*X & \text{codim } \widetilde{M} = 1, s^-(\widetilde{M}, q) = 0, \\ T_S^*X \rightarrow T_{\widetilde{S}}^*X & \text{codim } \widetilde{S} = 1, \end{cases} \quad (3.7)$$

(cf. [D'A-Z 3]). Let \widetilde{M}^+ and \widetilde{S}^+ be the closed half spaces with boundary \widetilde{M} and \widetilde{S} and inner conormal q . We have

PROPOSITION 3.1. *Let $d_S = d_M$. Then in the above situation*

$$\begin{cases} s^-(\widetilde{S}, q) = 0, \\ \widetilde{S}^+ \subset \widetilde{M}^+. \end{cases} \quad (3.8)$$

Proof. Quantizing χ by a kernel $K \in \text{Ob}(\mathbf{D}^b(X \times X))$ we get by [K-S 1, Proposition 11.2.8]

$$\begin{cases} \phi_K(\mathbf{C}_M) \cong \mathbf{C}_{\tilde{M}^+}[d_M(p) - 1] & \text{in } \mathbf{D}^b(X; q), \\ \phi_K(\mathbf{C}_S) \cong \mathbf{C}_{\tilde{S}^+}[d_S(p) - s^-(\tilde{S}, q) - 1] & \text{in } \mathbf{D}^b(X; q). \end{cases}$$

Moreover the natural morphism $\mathbf{C}_M \rightarrow \mathbf{C}_S$ is transformed via ϕ_K to a non null morphism $\mathbf{C}_{\tilde{M}^+}[d_M(p) - 1] \rightarrow \mathbf{C}_{\tilde{S}^+}[d_S(p) - s^-(\tilde{S}, q) - 1]$. Thus

$$\begin{aligned} & \text{Hom}_{\mathbf{D}^b(X; q)}(\mathbf{C}_{\tilde{M}^+}[d_M(p) - 1], \mathbf{C}_{\tilde{S}^+}[d_S(p) - s^-(\tilde{S}, q) - 1]) \\ &= H^0(\mathbf{R}\Gamma_{\tilde{M}^+}(\mathbf{C}_{\tilde{S}^+})_y[d_S(p) - d_M(p) - s^-(\tilde{S}, q)]) \\ &\neq 0, \end{aligned}$$

where $y = \pi(q)$. Since we are assuming $d_M(p) = d_S(p)$, (3.5) follows. \square

End of the proof of Theorem 2.1. From the proof of Proposition 3.1 it follows that ϕ_K transforms the morphism (3.3) in

$$\mathcal{H}_{\tilde{S}^+}^1(\mathcal{O}_X)_y \rightarrow \mathcal{H}_{\tilde{M}^+}^1(\mathcal{O}_X)_y, \quad (3.9)$$

where $y = \pi(q)$, which is clearly injective. The proof of Theorem 2.1 is now complete. \square

Proof of Theorem 2.2. From now on we will drop p in our notations, due to the constancy assumptions (2.3).

If $r > 1$ one has $\bar{\Omega} = M$ and $N^M(\Omega)^{\circ a} = T^*M$. Thus, by (2.3), we enter the hypotheses of Theorem B for both M and S . The claim follows in this case from (3.1), (3.3) and from the inequalities (3.2).

We may then assume $r = 1$. The problem in this case is that (2.3) holds only along $\text{SS}(\mathbf{C}_\Omega)$.

Let $\chi: T^*X \rightarrow T^*X$ be a contact transformation from a neighborhood of p to a neighborhood of $q = \chi(p)$, such that

$$\begin{cases} T_M^*X \rightarrow T_{\tilde{M}}^*X & \text{codim } \tilde{M} = 1, \\ T_S^*X \rightarrow T_{\tilde{S}}^*X & \text{codim } \tilde{S} = 1, s^-(\tilde{S}, q') \equiv 0. \end{cases}$$

Notice that, for $y = \pi(q)$, $T_y\tilde{M} = T_y\tilde{S}$. Quantizing χ by a kernel K , we thus have that either $\phi_K(\mathbf{C}_{\bar{\Omega}})$ or $\phi_K(\mathbf{C}_\Omega)$ is a simple sheaf along the conormal bundle to a C^1 submanifold $Y \subset X$. Since $d_M = d_S - 1$, then $s^-(\tilde{M}, q) = 0$, $\tilde{M}^+ \subset \tilde{S}^+$ and $\phi_K(\mathbf{C}_\Omega) = \mathbf{C}_Y[d_M - 1]$. Denoting by W the open domain with boundary Y and

exterior conormal q , we have by Lemma 3.3 of [Z] that W is pseudoconvex at y , and one concludes since

$$\chi_* \mu_\Omega(\mathcal{O}_X)_q[-d_M] \cong \mathcal{H}_{X \setminus W}^1(\mathcal{O}_X)_y, \quad \square$$

References

- [A-G] Andreotti, A. and Grauert, H.: Théorèmes de finitude pour la cohomologie des espaces complexes *Bull. Soc. Math. France* 90 (1962), 193–259.
- [D'A-Z 1] D'Agnolo, A. and Zampieri, G.: A propagation theorem for a class of sheaves of microfunctions, *Rend. Mat. Acc. Lincei*, s.9 1 (1990), 53–58.
- [D'A-Z 2] D'Agnolo, A. and Zampieri, G.: Vanishing theorem for sheaves of microfunctions at the boundary on CR-manifolds, *Commun. in P.D.E.* 17 (1992), no. 5 and 6, 989–999.
- [D'A-Z 3] D'Agnolo, A. and Zampieri, G.: Generalized Levi forms for microdifferential systems (M. Kashiwara, T. Monteiro Fernandes, P. Schapira, eds.), *D-modules and microlocal geometry*, Walter de Gruyter and Co., 1992.
- [D'A-Z 4] D'Agnolo, A. and Zampieri, G.: Levi's forms of higher codimensional submanifolds, *Rend. Mat. Acc. Lincei*, s. 9, 2 (1991), 29–33.
- [H] Hörmander, L.: An introduction to complex analysis in several complex variables, Van Nostrand. Princeton (1966).
- [K-S 1] Kashiwara, M. and Schapira, P.: Microlocal study of sheaves, *Astérisque* 128 (1985).
- [K-S 2] Kashiwara, M. and Schapira, P.: Sheaves on manifolds, Springer-Verlag, 292 (1990).
- [S] Schapira, P.: Front d'onde analytique au bord II, Sem. E.D.P. Ecole Polyt., Exp. XIII (1986).
- [S-K-K] Sato, M., Kawai, T. and Kashiwara, M.: Hyperfunctions and pseudo-differential equations, *Lecture Notes in Math.*, Springer-Verlag, 287 (1973), 265–529.
- [Z] Zampieri, G.: The Andreotti-Grauert vanishing theorem for polyhedrons of \mathbb{C}^n *Journal of Math. Sci. Univ. Tokyo* 2(1) (1995), 233–246.