

Review

Regulated Transport into the Nucleus of Herpesviridae DNA Replication Core Proteins

Gualtiero Alvisi^{1*}, David A. Jans², Daria Camozzi³, Simone Avanzi¹, Arianna Loregian¹, Alessandro Ripalti⁴ and Giorgio Palù^{1*}

¹ Department of Molecular Medicine, University of Padua, Padua 35121, Italy;

E-Mails: simone.avanzi@unipd.it (S.A.); arianna.loregian@unipd.it (A.L.)

² Department of Biochemistry and Molecular Biology, Monash University, Victoria 3800, Australia;

E-Mail: david.jans@monash.au

³ National Research Council of Italy, Institute of Molecular Genetics, Unit of Bologna,

Bologna 40136, Italy; E-Mail: daria.camozzi@gmail.com

⁴ Azienda Ospedaliera Universitaria di Bologna Policlinico S. Orsola-Malpighi, Bologna 40138,

Italy; E-Mail: alessandro.ripalti@unibo.it

* Authors to whom correspondence should be addressed; E-Mails: gualtiero.alvisi@unipd.it (A.G.);

giorgio.palu@unipd.it (G.P.); Tel.: +39-049-827-2353 (A.G.); +39-049-827-2350 (G.P.);

Fax: +39-049-827-2355 (A.G. & G.P.).

Received: 22 July 2013; in revised form: 3 September 2013 / Accepted: 4 September 2013 /

Published: 16 September 2013

Abstract: The *Herpesviridae* family comprises several major human pathogens belonging to three distinct subfamilies. Their double stranded DNA genome is replicated in the nuclei of infected cells by a number of host and viral products. Among the latter the viral replication complex, whose activity is strictly required for viral replication, is composed of six different polypeptides, including a two-subunit DNA polymerase holoenzyme, a trimeric primase/helicase complex and a single stranded DNA binding protein. The study of herpesviral DNA replication machinery is extremely important, both because it provides an excellent model to understand processes related to eukaryotic DNA replication and it has important implications for the development of highly needed antiviral agents. Even though all known herpesviruses utilize very similar mechanisms for amplification of their genomes, the nuclear import of the replication complex components appears to be a heterogeneous and highly regulated process to ensure the correct spatiotemporal localization of each protein. The nuclear transport process of these enzymes is controlled by three mechanisms, typifying the main processes through which protein nuclear

import is generally regulated in eukaryotic cells. These include cargo post-translational modification-based recognition by the intracellular transporters, piggy-back events allowing coordinated nuclear import of multimeric holoenzymes, and chaperone-assisted nuclear import of specific subunits. In this review we summarize these mechanisms and discuss potential implications for the development of antiviral compounds aimed at inhibiting the Herpesvirus life cycle by targeting nuclear import of the Herpesvirus DNA replicating enzymes.

Keywords: herpesviruses; replicase; CK2; inhibitors; anti-viral therapy; nuclear transport regulation; importins; phosphorylation; hsp90

1. Introduction

The *Herpesviridae* family comprises several human pathogens of pivotal clinical importance, including the α -*Herpesvirinae* herpes simplex virus type 1 and 2 (HSV-1/-2) and Varicella Zoster virus (VZV), the β -*Herpesvirinae* human cytomegalovirus (HCMV) and human herpesvirus 6 and 7 (HHV-6/-7), as well as the γ -*Herpesvirinae* Epstein-Barr virus (EBV) and Kaposi's sarcoma-associated herpesvirus (KSHV), (see [1] for a review). Although differing in important properties such as host specificity, clinical manifestations, and *in vitro* growth kinetics, all *Herpesviridae* family members replicate their linear double stranded (ds) DNA genome in the nucleus of productively infected cells. Lytic viral DNA replication is a process that starts at defined sites on the viral genomes, the origin of DNA replication (ori_{Lyt}). Most β -herpesviruses have a single ori_{Lyt} , whereas several α - and γ -herpesviruses have two (VZV, EBV and KSHV) or three (HSV1, HSV2), respectively. Although all herpesviruses control replication initiation at ori_{Lyt} via virus-encoded DNA binding polypeptides, a *bona fide* ori-binding protein (OBP) has been identified only in α -herpesviruses and some β -herpesviruses, while in HCMV and γ -herpesviruses this function is performed by viral proteins also transactivating viral gene expression. The exact molecular events that lead to herpetic DNA replication are still mysterious, with several lines of more recent evidence putting into question the dogma of a rolling circle-based replication, instead supporting a recombination-dependent branching mechanism [2]. Despite this, there is general agreement regarding the formation of concatamers during the process, from which genomic units need to be resolved before viral genome encapsidation occurs.

Regardless of its exact mechanism, viral DNA replication takes place within the host cell nucleus and is mediated by a number of cellular and viral proteins [3–7]. The latter include an OBP, a DNA polymerase holoenzyme, composed by a DNA-dependent DNA polymerase catalytic subunit (pol) and a DNA polymerase accessory protein (PAP) conferring processivity to pol, as well as a trimeric helicase/primase complex formed by a primase, an helicase and a primase associated factor (PAF). Finally, a single stranded (ss) DNA-binding protein is also required. These factors sequentially assemble on the ori_{Lyt} and mediate viral genome replication. Therefore, herpesviral replication fork proteins need to reach the nucleus after having been translated in the cytoplasm. Here we summarize current knowledge regarding signals and transporters responsible for nuclear targeting of herpesviral

DNA replication proteins and the mechanisms regulating this process, with a focus on possible strategies to impair viral replication by interfering with it.

2. Nuclear Import

A hallmark of eukaryotic cells is the high degree of compartmentalization. The genetic information resides in the cell nucleus, separated from the cytoplasmic apparatus responsible for protein synthesis by a double membrane structure, the nuclear envelope (NE). The passage of molecules such as mRNA and proteins through the NE occurs necessarily through aqueous channels delimited by huge NE-embedded multiprotein-complexes called nuclear pore complexes (NPCs) [8]. It is generally accepted that molecules smaller than 50 kDa can freely diffuse through the NPCs [9], while the transport of larger molecules is an active, signal-dependent process. Among these molecules, cellular mRNAs and pre-miRNAs need to be exported from the nucleus after transcription to the cytoplasm in order to be translated or processed by DICER, respectively, while proteins such as histones and lamins, need to be imported into the nucleus after having been synthesized in the cytoplasm to reach the DNA or the inner NE, respectively [10,11].

Active transport of molecules across the NPC-delimited channel is a signal-mediated process effected by transporters belonging to the importin (IMP) β superfamily of cellular transporters that recognize specific targeting signals on cargoes. These signals can be divided into nuclear localization sequences (NLSs) and nuclear export sequences (NESs), responsible for targeting into and out of the nucleus, respectively (Figure 1) [12]. The study of viral proteins generally has proven to be invaluable in delineating the key molecular details of many eukaryotic cell processes [13,14], with nucleocytoplasmic transport being no exception. The first evidence of nuclear targeting signals stemmed from the work of Kalderon *et al.* with respect to simian virus 40 (SV40), that showed that SV40 large tumor antigen (T-ag), a protein normally localizing in the host cell nucleus, mislocalized to the cytoplasm upon mutation of a short stretch of basic amino acids (aas) [15]. Importantly, fusion of such a sequence to otherwise cytoplasmic proteins was sufficient to confer nuclear localization [16]. The 7-aas highly basic sequence **PKKKRKV** (single letter aa code) was delineated as an NLS, sufficient and necessary to confer T-ag nuclear localization [15,16]. Since then, a number of different types of NLS have been identified which can be classified on the basis of their sequence, and their ability to be recognized by different IMPs, and NLSs similar to that of T-ag are often referred to as “classical” NLSs (cNLSs) [17], which are highly basic, generally lysine-rich, and confer direct interaction with the adapter protein $\text{IMP}\alpha$ complexed to $\text{IMP}\beta$ 1 (see below). cNLSs can be classified as monopartite or bipartite, according to the number of basic stretches of aas composing them. Monopartite NLSs resemble the T-ag-NLS and are a single cluster of basic residues, matching the consensus (K-K/R-X-K/R), while the bipartite NLSs, with two clusters of basic residues separated by a spacer region of 10–13 aa, resemble the NLS from the *Xenopus laevis* histone assembly factor nucleoplasmin (**KR**-11 aas-**KKKK**) [17]. $\text{IMP}\alpha$ is formed by two distinct domains: a short, basic N-terminal $\text{IMP}\beta$ -binding (IBB) domain responsible for binding to $\text{IMP}\beta$ 1, and the remainder of the molecule comprising 10 Armadillo (ARM) tandem repeats, including two regions of ARM repeats representing the major and minor NLS-binding sites [18]. In the absence of $\text{IMP}\beta$ 1, the IBB domain competes with the NLSs for the major NLS-binding site, resulting in an auto-inhibitory mechanism [19]; $\text{IMP}\alpha$ - $\text{IMP}\beta$ interaction enables NLS-binding by displacing the


IBB domain. Monopartite NLSs bind preferentially to the major NLS binding site, while bipartite NLSs interact with both binding pockets. The complex is subsequently docked to the NPC and translocates through it as mediated by IMP β , before reaching the nucleus. Once inside the nucleus, binding to IMP β of the small Ras-related guanine nucleotide binding protein Ran in activated GTP-bound form, results in dissociation of IMP β from the IBB domain of IMP α , which in turn competes for the major NLS binding site on IMP α , thus effecting cargo release into the nucleus [20] (Figure 1). Importantly, the conformation of two key structural loops in Ran depends on the nucleotide bound, which thereby determines the affinity for IMP β homologues, regulating the interaction between the latter and the cargoes [21,22]. The RanGTP/RanGDP ratio is low in the cytoplasm and high in the nucleus, ultimately ensuring directionality of the nuclear transport process [23]. This RanGTP gradient is maintained by the asymmetrical distribution of two key regulators of Ran nucleotide state: Ran activating GTPase (RanGAP), which is predominantly found in the cytoplasm [24], and the nuclear Ran guanine nucleotide-exchange factor (RanGEF, also known as RCC1), which mediates substitution of GDP with GTP [25]. Cytoplasmic RanGDP is imported into the nucleus by its specific transporter nuclear transport factor 2 (NTF2) [26], and subsequently converted to RanGTP by the action of RCC1. As mentioned above, IMP α /IMP β 1-cargo complexes are dissociated in the nucleus after interaction of IMP β with RanGTP [20]. The IMP β -RanGTP complex is subsequently recycled to the cytoplasm, while IMP α is exported from the nucleus by the IMP β homologue CAS, as complexed with RanGTP [27]. Once in the cytoplasm, the action of RanGAP catalyzes the hydrolysis of GTP to GDP, resulting in the dissociation of RanGDP from the IMPs which are then available for a new round of nuclear import, while the newly generated RanGDP can be transported into the nucleus by NTF2 [26].

As mentioned above, cargoes can be imported into the nucleus through several possible pathways, in addition to the classical nuclear import pathway, whereby the IMP α / β heterodimer mediates import of a cNLS bearing cargo. Seven different IMP α homologues have been identified in humans, and can be assigned to three subgroups, on the basis of homology and cargo binding specificity [28]. Alternatively, IMP β 1 can recognize the m3G-cap of snRNAs via the adaptor Snurportin 1, to mediate nuclear import of snRNPs [29]. Additionally certain basic NLSs are recognized specifically by IMP β 1 directly [30–32]; an example is the **RQARRNRRRRWR** sequence from the human immunodeficiency virus (HIV-1) Rev protein [33]. Finally, certain atypical NLSs, such as the one recently described on KHSV LANA, can also be directly recognized by IMP β 1 [34].

On the other hand, the almost 20 IMP β 1 homologues described so far bind cargoes directly without the need for an adaptor such as IMP α [35,36]. Of these, IMP β 2 is the best characterized, with > 20 cargoes, mostly RNA binding proteins, already identified, including the RNA binding protein heterogeneous ribonucleoprotein particle A1 (hnRNP-A1), which contains the 30 aa M9-NLS (FGNYNNQSSNFGPMKGGNFGGRSSGPYGGG) [37]. IMP β 2 recognized NLSs have been identified in various other RNA binding proteins [38–41], but crystallographic and biochemical approaches were necessary to enable definition of the predictive consensus PY-NLS, consisting of highly disordered sequences of about 20–30 residues, which can be divided in *N*- and *C*-terminal motifs. The consensus sequence determined for the latter (R/K/H-X(2,5)-PY) is highly conserved in the majority of identified PY-NLSs, wherein substitution of key aas impairs recognition by IMP β 2 [42,43]. In contrast, the *N*-terminal motif has a looser consensus, formed by either a hydrophobic or basic stretch of aas, the bPY-NLS and hPY-NLS subclasses respectively. The basis of this appears to be the existence of two

NLS binding sites embedded inside the C-terminal domain of IMPβ2: the “A” site, necessary for strong and specific interaction with the R/K/H-X(2,5)-PY motif and the “B” site, which is rich in acidic residues but with hydrophobic residues scattered throughout, enabling either basic or hydrophobic linear peptides to be accommodated. Mutational analysis on either basic or hydrophobic aas within PY-NLSs results in reduced interaction with IMPβ2, confirming the requirement of mutated residues for proper interaction. Thus, it is likely that the role of the A site is essential, with the B site being required to increase IMPβ2 binding affinity [43,44]. Besides differences in target protein recognition, IMPβ2 and IMPβ1 share the same mechanisms for cargo transport and release into the nuclear compartment. In the case of IMPβ2, cargo release is through Ran-GTP binding to a disordered loop of IMPβ2 (H8 Loop), which contributes to the B site, facilitating cargo release from IMPβ2 into the nuclear compartment [43,44].

Figure 1. Cellular transporters involved in signal dependent nuclear targeting of herpesviral proteins. Transport of cargoes across the nuclear envelope (NE) occurs exclusively through channels delimited by the nuclear pore complexes (NPCs), upon recognition of specific nuclear localization complexes (NLSs) by specialized cellular transporters, with different signals being recognized by a variety of transporters: the IMPα/β1 heterodimer recognizes cNLSs, which can either be of the monopartite or bipartite type (A), importin IMPβ1 recognizes arg-rich NLSs (B), whereas IMPβ2 recognizes PY-NLSs. Proteins carrying the various types of NLS are listed in the boxes, with prototypical NLSs being shown in black, and herpesviral DNA replicase NLSs indicated in red.



3. Regulation of Nuclear Transport

Regulation of protein transport across the NPC, and the kinetics thereof, plays a central role in controlling cellular processes such as cell growth and differentiation, and response to pathogens [45]. In the case of transcription factors, this can determine whether a critical intranuclear concentration is achieved or not, thus ultimately determining its effect on cellular transcription [46]. Mechanisms regulating nuclear import include alteration of the expression levels and/or function of IMPs, or components of the NPC [47], and/or modulation of the affinity of IMP recognition of the cargo by post-translational modification, such as by phosphorylation [46]. The latter strategy offers several advantages, including the ability of modulating intranuclear protein concentration without requiring *ex-novo* protein synthesis or degradation, and affords rapid and reversible control of nuclear/cytoplasmic localization [48]. Indeed the localization of several crucial cellular key molecules is controlled by this mechanism, including the *Drosophila* morphogen Dorsal, where protein kinase A (PKA) mediated phosphorylation 22 residues upstream of the NLS are associated with a c. seven-fold increase in IMP α binding affinity, and a marked increase in transport efficiency [49]. Another extremely well characterized example of phosphorylation-regulated nuclear import of cargo is the SV40 T-ag, where two synergistically acting kinases (protein kinase CK2 and dsDNA-dependent kinase) phosphorylate residues located 10–12 aa upstream of the NLS, thus increasing the affinity of NLS recognition by IMP α / β 1 by about 100-fold to achieve high efficiency of T-ag nuclear targeting [50]. T-ag nuclear import is also controlled by a phosphorylation site for the cyclin-dependent kinase cdk1 adjacent to the NLS, where phosphorylation confers binding to the cytoplasmic retention factor BRCA-1 binding protein 2 (BRAP2), a general negative regulator of nuclear import [51].

4. Nuclear Transport of Herpesviral DNA Replicating Enzymes

In order to mediate viral DNA replication, the proteins forming the herpesviral DNA replication core need to reach the cell nucleus. Besides ssDNA binding protein, the two main complexes are a multimeric DNA polymerase holoenzyme and helicase/primase complexes. Given the high molecular weight of such proteins, their nuclear transport has been shown to be an active process mediated by a number of different IMPs, which recognize specific NLSs on the cargo proteins [52–59]. In some cases, individual subunits are capable of localizing to the cell nucleus independently of the others, like in the case of HSV and HCMV DNA polymerase holoenzyme subunits [52–56]. However in the case of KSHV and EBV DNA polymerase holoenzymes [59,60] and most of their helicase/primase complex subunits, formation of the holoenzyme appears to be a pre-requisite for nuclear transport [61–64] (see Table 1).

Table 1. Identified NLSs and nuclear import pathways of Herpesviral DNA replicating enzymes. The NLSs identified so far for each herpesviral DNA replication proteins are indicated, as well as the IMPs binding them. The subcellular localization of each protein when individually expressed is also indicated (*loc. alone*). Additional details are reported in respect to the nuclear import pathways of DNA polymerase and helicase/primase holoenzymes. Pol, DNA polymerase catalytic subunit; PAP, DNA polymerase processivity factor; hNLS, hydrophobic NLS; cNLS, classical NLS; bip, bipartite NLS; N, nucleus; C, cytoplasm; IMPs, importins.

HSV-1					HCMV			
<i>protein</i>	<i>name</i>	<i>NLS</i>	<i>IMPs</i>	<i>loc. alone</i>	<i>name</i>	<i>NLS</i>	<i>IMPs</i>	<i>loc. alone</i>
processivity factor	UL42	1 cNLS bip [55]	α/β	N [55]	UL44	1 cNLS [52]	α/β	N [52]
DNA polymerase	UL30	1 cNLS bip [53] 1 NLS hyd [53,56]	α/β	N [53,56,87]	UL54	1 cNLS [54] 1 NLS hyd [56]	α/β	N [56]
primase	UL5			C [63,64]	UL70			C [52]
helicase	UL52			C [63,64]	UL105			N in infected cells [110]
primase-associated factor	UL8	putative PY-NLS (see text for details)		C [63,64]	UL102			
ssDNA binding protein	UL29			?	UL57			
DNA polymerase holoenzyme nuclear import	Individual subunits and holoenzyme [55]				Individual subunits and holoenzyme [54]			
PAP status	Monomer [76]				Dimer [77,106]; transported to the nucleus as dimer [105]			
Helicase primase complex nuclear import	Transported as a trimeric complex C [63,64]				Not known, but UL70 can localize to the nucleus independently of the other subunits if the cellular chaperone DNAJB6-a is overexpressed [109]			
EBV					KHSV			
<i>protein</i>	<i>name</i>	<i>NLS</i>	<i>IMPs</i>	<i>loc. alone</i>	<i>name</i>	<i>NLS</i>	<i>IMPs</i>	<i>loc. alone</i>
processivity factor	BMRF1	1 cNLS [84]		N [84]	ORF59	1 cNLS [59]		N [59]
DNA polymerase	BALF5			C [60]	ORF9			C [59]
primase	BSLF1			C [61]	ORF44	putative PY-NLS (see text for details)		C [59]
helicase	BBLF4			C [61]	ORF56			C [59]
primase-associated factor	BBLF2/3			C [61]	ORF40/41			C [59]
ssDNA binding protein	BALF2				ORF6			
DNA polymerase holoenzyme nuclear import	Piggy back as mediated by PAP [60]				Piggy back as mediated by PAP [59]			
PAP status	Multimer [78,114]				Dimer [79,115]; transported to the nucleus as dimer [116]			
Helicase primase complex nuclear import	Imported as a complex; BBLF4 can be imported if expressed with the ZTA transactivator [61]				pol/PAP and all the helicase primase subunits need to be simultaneously expressed to achieve nuclear targeting [62]			

5. DNA Polymerase Holoenzyme Nuclear Transport

All human herpesviruses encode a two subunit DNA polymerase. The pol subunit is known to perform the synthesis of leading and lagging strands, while the PAP tethers the holoenzyme to the DNA, thus conferring processivity [65–67]. Since the replication of all known herpesviruses takes place in the host cell nucleus [68], these high molecular weight proteins need to be actively transported through the NPC to reach the site of viral DNA replication. Different herpesviruses have evolved different strategies to ensure correct subcellular localization of their DNA polymerase holoenzyme. Indeed different proteins can be translocated independently, or only as fully assembled holoenzymes, wherein a NLS present on the PAP mediates import of the catalytic subunit via a piggy-back mechanism.

In all known herpesviruses, a functional NLS has been characterized on the PAP [52,55,57–59], which plays a central role in nuclear targeting of the holoenzyme (Figure 2 and Table 2). In addition, in the case of HCMV a complex phosphorylation-dependent regulation system finely tunes nuclear import of the PAP [51,52,69,70]. Herpesviral PAPs can be generally divided into two functional domains: a *N*-terminal catalytic domain and a *C*-terminal nuclear targeting domain. The *N*-terminal domain is capable of performing all known biochemical activities, including binding to pol and to dsDNA in the absence of ATP and clamp loaders, and stimulation of the holoenzyme activity [71–75]. Structural studies showed that the *N*-terminal domains of HSV-1 (UL42), HCMV (UL44), EBV (BMRF1) and KSHV (PF-8) PAPs share a very similar three dimensional architecture, which closely resembles that of monomers of the eukaryotic DNA polymerase processivity factor PCNA, the main difference being that UL42 is a monomer, while UL44, BMRF1 and PF/8 form head to head dimers [76–80]. On the other hand, the *C*-terminal domains appear to be largely unstructured, do not possess any known biochemical property, and carry an NLS. Functional NLSs have been identified at the *C*-terminus of HSV-1, HCMV, HHV-7, EBV and KSHV PAPs, all of which are known to be able to localize independently of pol into the cell nucleus [52,55,57–59], probably reflecting the fact that they are also capable of performing additional functions during the viral life cycle. The latter hypothesis is also suggested by several other lines of evidence, including their higher and more temporally delayed expression kinetics when compared to pol, and the fact that they have been shown to possess transcriptional regulation properties [81–85]. In the case of HCMV, HHV-7, EBV and KSHV PAPs, the cNLS is monopartite, closely resembling that identified on SV40 T-ag [52,57–59]. In the case of HSV-1, the NLS is also located at the *C*-terminus, but appears to be bipartite, with two basic clusters of aas being required for optimal nuclear targeting [55]. Biochemical and functional studies revealed that nuclear import of such cargoes is energy and Ran-dependent, and mediated by the IMP α / β 1 heterodimer [55,57] (see Table 2). It is not clear why HSV-1 has evolved a bipartite NLS on its PAP UL42, but this might reflect the ability of HSV-1 to infect neuronal cells *in vivo*. Indeed different IMP α members exhibit unique temporal expression profiles in neurons [86], and immunoprecipitation experiments have shown that UL42's bipartite NLS can bind to both IMP α -S and α -Q members, while the HCMV PAP UL44 monopartite NLS only binds efficiently to IMP α -S members [55]. Regardless of these differences, all known PAPs bear a functional NLS. In contrast, not all known pol subunits are able to localize to the cell nucleus independently of their PAP. Indeed, while HSV-1 and HCMV pols [87] or portions thereof [54] localize to the cell nucleus when expressed in the absence of other viral proteins in cell culture, this is not the case for pols from EBV and KSHV, wherein NLSs have not been identified, and nuclear targeting entirely relies on the presence of the respective PAP via a piggy-back mechanism [59,60]. The reason why these viruses evolved such different strategies to ensure nuclear targeting of their DNA polymerase catalytic subunits is not clear, but it is possible that the ability of HSV and HCMV DNA pols to be independently transported into the nucleus is critical to their infectious cycles. Indeed, both viruses evolved two distinct but functional NLSs on their pol enzymes. The first to be discovered were atypical hydrophobic NLSs (hNLSs), which lie within the PAP binding domain, and thus appear to be active only when pol is not associated with the PAP [56]; these have not been fully characterized. cNLSs, at the very *C*-terminus of the proteins outside the PAP binding domains similar to those identified in the respective PAPs (see below) were subsequently described that confer energy- and Ran-dependent nuclear targeting to reporter proteins, as well as recognition by the

IMP α/β heterodimer through direct interaction with IMP α [53,54]. In addition, in contrast to the hNLSs, which are masked upon PAP binding [54,55], these signals are functional in the context of the DNA polymerase holoenzymes. Due to the presence of these additional NLSs, HSV and HCMV pols are capable of translocating their respective PAPs to the nucleus via a piggy-back mechanism once the NLS on the latter are inactivated by mutagenesis [54,55]. Thus, it appears that HCMV and HSV have evolved several mechanisms to maximize the nuclear import possibilities of their DNA polymerase subunits, so that they are capable of both being imported individually or as complexes in holoenzymes. In contrast, functional NLSs are not present on the pols from EBV and KSHV-8 and their nuclear translocation entirely relies on their interaction with their PAP. Although the IMPs responsible for nuclear targeting of the EBV and KSHV-1 PAPs have not been identified experimentally, the process is likely mediated by the IMP α/β heterodimer, since EBV and KSHV PAPs possess highly basic sequences resembling the T-ag cNLS (see Table 2). This apparently obligates the cotranslocation into the nucleus of DNA polymerase holoenzyme subunits in the case of KSHV and EBV, and likely provides a means for coordinating and facilitating the processes of viral DNA recognition and processive DNA synthesis.

The fact that several herpesviral DNA polymerase subunits are imported by the IMP α/β heterodimer has important implications for antiviral therapy. Indeed Ivermectin, a broad-spectrum anti-parasite medication has been recently identified as a specific inhibitor of IMP α/β -mediated nuclear import [88]. Treatment of cells expressing proteins translocated to the nucleus by the IMP α/β heterodimer such as HCMV UL44 and 54, as well as Dengue virus NS5 and HIV-1 integrase (IN), resulted in a strong inhibition of nuclear localization of such cargoes [89]. More importantly, treatment of cells infected with HIV-1 and DENV resulted in a strong reduction of virus production [89,90]. It is therefore not unreasonable to suggest that inhibition of IMP α/β nuclear transport could also have a broad anti herpetic activity.

Figure 2. Nuclear transport of DNA polymerase holoenzymes. Different herpesviruses have evolved distinct mechanisms to mediate nuclear targeting of their DNA polymerase holoenzymes. In the case of HSV-1 and HCMV both pol and PAP enzymes possess intrinsic, NLSs and can be imported to the nucleus either alone or as a fully assembled enzyme. In contrast, EBV and KSHV pols are incapable of nuclear targeting in the absence of the respective PAPs, bearing a C-terminally located NLS. A requirement of Hsp90 activity has also been reported for the nuclear import of HSV-1 and EBV pol. Black vertical bars indicate direct protein-protein interactions; red vertical bars indicate direct protein-protein interactions depending on Hsp90 activity.




Table 2. NLSs identified on Herpesviral DNA polymerase subunits and nuclear import molecular mechanisms. Pol, DNA polymerase catalytic subunit; PAP, DNA polymerase processivity factor; hNLS, hydrophobic NLS; cNLS, classical NLS. The single letter amino acid code is used; hydrophobic residues in hNLS are *underlined*, whereas basic residues in cNLSs are in *boldface*. s indicates the ability of the NLS to function when the subunit is uncomplexed; h indicates the ability of the signal to function once the holoenzyme is assembled. In the column “Ran” the sensitivity of nuclear transport to overexpression of the transdominant negative RanQ69L is reported; in the column “ATP”, sensitivity to depletion of intracellular ATP levels; X, sensitivity; ?, unknown.

Virus	Protein	Name	NLS type	Sequence	IMPs	Ran	ATP	Nuclear import	Notes	Reference
HSV-1	pol	UL30	hNLS	RR <u>ML</u> HR-1129	?	?	?	s	overlapping PAP binding domain	[56]
			cNLS bip	PA KR PRETPSPADPPGGASK PRK -1136	α/β	X	X	s + h		[53]
	PAP	UL42	cNLS bip	PT TKRGR SGGEDARADAL KKPK -413	α/β	X	X	s + h		[55]
HCMV	pol	UL54	hNLS	PR <u>RLHL</u>	?	?	?	s	overlapping PAP binding domain	[56]
			cNLS	PA KKRAR -1159	α/β	X	X	s + h		[54]
	PAP	UL44	cNLS	PNT KKQK -431	α/β	X	X	s + h	regulated by phosphorylation	[52]
HHV-7	pol	U38	/	/	/	/	/	/		
	PAP	U27	cNLS	PNS KRQR -361	?	X	X	s		[57]
KHSV-1	pol	pol-8	no NLS					h	requires PF-8 for import	[59]
	PAP	PF-8	cNLS	KRPHKRRSD -377	?	?	?	s + h	mediates translocation of holoenzyme	[59]
EBV	pol	BALF1	no NLS					h	requires BMRF1 for import	[60]
	PAP	BMRF1	cNLS	KHPKK -396	?	?	?	s + h	mediates translocation of holoenzyme	[58]

6. MCV UL44 by IMPs


Beside inhibition of the $\text{IMP}\alpha/\beta$ dependent nuclear import generally, with associated problems of host cell toxicity, a potentially more specific approach to inhibit viral replication would be to inhibit the recognition between specific NLS bearing viral cargoes and the transporter [91], as recently described for the HIV-1 matrix protein [92] and for the HCMV viral phosphoprotein UL84, which plays a crucial, still not yet completely characterized role during viral infection [93]. Indeed, the above mentioned redundancy of NLS on DNA polymerase holoenzymes of HSV-1 and HCMV but not on EBV and KSHV implies that the NLSs located on the PAPs from the latter represent potential targets for the development of antiviral drugs, since their function is not redundant for viral replication in cell culture [59,60]. On the other hand, in the case of both HSV and HCMV, simply targeting the PAPs NLSs might not be sufficient to block viral replication, since inhibition of DNA polymerase holoenzymes nuclear transport would require the simultaneous inhibition of the cNLSs located on the PAPs and on the catalytic subunits. Consistent with this hypothesis, HSV-1 UL42 C-terminal domain, containing the functional cNLS and absolutely required for nuclear import of UL42 when expressed individually [55], is completely dispensable for virus growth in cell culture [94]. Surprisingly, this is in stark contrast to reports for HCMV, where deletion of the HCMV UL44 C-terminal domain, including the cNLS [52], results in the inability of HCMV to replicate in cell culture [95,96]. This is surprising, given the fact that the UL44 C-terminal deletion mutant can still localize to the nucleus through piggy-back by the UL54 pol catalytic subunit, and addition of the T-ag cNLS does not restore viral replication [52,95,96]. The clear implication is that the UL44 C-terminal domain plays other roles in the HCMV life cycle, in addition to conferring nuclear localization to the protein. The UL44 C-terminus (residues 409–433) is in fact the target of extensive post-translational modifications. These include multiple phosphorylation sites (S413, S415, S418 and T42T) and a sumoylation site (K410), which regulate UL44 subcellular localization by controlling its ability to interact with host cell proteins [52,69,70,97,98], although the exact role of UL44 sumoylation is still unknown [97]. Depending on the target residues, phosphorylation can either negatively or positively regulate UL44 nuclear import. Indeed, *in vitro* data showed that the viral kinase UL97 is capable of phosphorylating T427. This residue lies within UL44 cNLS sequence (PNTKKQK431) and mediates interaction with the cellular cytoplasmic retention factor BRAP2 [51,99]: negative charge on T427 enables BRPA2 binding and reduces UL44 nuclear accumulation more than 10-fold [51,69]. Thus, phosphorylation of T427 can prevent UL44 interaction with the $\text{IMP}\alpha/\beta$ heterodimer by two mechanisms: first of all, the negative charge in position +1 of the NLS can directly decrease binding to the $\text{IMP}\alpha$ NLS-binding pocket due to electrostatic repulsion between the phosphate group of the T residue and the NLS binding groove [70,100]. Secondly, binding of BRAP2 to UL44 competes directly with $\text{IMP}\alpha/\beta$ for binding to the NLS [51,69]. In contrast, phosphorylation of S413, S415 and S418 strongly increases UL44 nuclear accumulation in transient transfection assays [52]. Intriguingly, the three residues are part of a phosphorylation cascade triggered by protein kinase CK2 (CK2) that phosphorylates S413, that in turn promotes protein kinase CK1 (CK1) phosphorylation of S418, which then enables CK2 to phosphorylate S415 [69]. Analysis of UL44 subcellular localization of UL44 point mutant derivatives demonstrates that each phosphorylation event increases UL44 nuclear accumulation [69], with pharmacological inhibition of CK2 able to reduce nuclear accumulation [52,70]. The molecular basis

for improved nuclear import through phosphorylation of S413, S415 and S418 appears to reside in altered binding affinity for the IMP α/β heterodimer, in similar fashion to SV40 T-ag and the key role of the CK2 site upstream of its NLS [50]. Consistent with this idea, a GFP-UL44(410–433) fusion protein containing sequences upstream of the NLS binds to IMP α/β with much stronger affinity than a GFP-UL44(425–433) fusion protein containing only the NLS, as shown using native gel electrophoresis [52]. Strikingly, the key phosphorylation sites in HCMV UL44 are conserved in the HHV-6 and seven homologues of UL44, suggesting that β -herpesviruses utilize a common strategy to enhance nuclear import of their PAPs [52]. These regulatory mechanisms, which enable the extent of nuclear targeting of a cargo to be controlled by modulating the affinity of recognition between the IMPs and the cNLS within the cargo itself, are analogous to those originally described for the SV40 T-ag [101]. In the latter case, CK2- and dsDNA-dependent kinase cooperatively phosphorylate T-ag upstream of its NLS to enhance its nuclear localization and cdk1 phosphorylates in close proximity to the NLS itself to prevent nuclear targeting; comparable regulatory mechanisms apply to other viral (Human papillomavirus 11-E1; see [102]) and cellular proteins (for example p53; see [48]). HCMV has thus evolved to be able to finely tune the nuclear transport of its PAP by modulating its interaction with the specific IMPs responsible for PAP nuclear targeting. Therefore UL44 can be translocated to the nucleus at three different rates (Figure 3), according to its phosphorylation status, which is of importance in the context of the viral life cycle. Indeed, when UL44 is phosphorylated on residues S413, S415 and S418, its NLS is recognized with very high affinity by IMP α/β , resulting in highly efficient/rapid nuclear accumulation. This may prevent UL44 from associating with other viral and cellular proteins in the cytoplasm, thus supplying a source of uncomplexed nuclear protein able to perform DNA replication-related (for example by associating with UL54) or unrelated (for example by regulating transcription [83]) functions. Similarly, phosphorylation of T427 by UL97 or cdk1/2 [69,103] would prevent interaction with IMP α/β by promoting interaction with BRAP2, causing UL44 to be retained in the cytoplasm, to perform other as yet unknown functions. Finally, a lack of phosphorylation on UL44 C-terminal residues would result in low affinity interaction with IMP α/β and accompanying lower efficiency nuclear import, to enable UL44 to interact with several proteins before being translocated into the nucleus. These include the catalytic subunit of the DNA polymerase holoenzyme, UL54 [54], the uracil DNA glycosylase UL114 [104] and UL44 itself [105,106]. Thus, it seems likely that the essential role of the UL44 C-terminus in HCMV replication depends on its nuclear transport regulation properties, and the regulation thereof by phosphorylation in particular. Of importance in this context is firstly that all nuclear transport enhancing sites (S413, S415 and S418) have been found to be phosphorylated to high levels in cells infected with HCMV *in vitro* [103], while the introduction of phosphonull mutations at these sites abolishes both UL44 nuclear accumulation and viral replication [103]. Secondly, introduction of phosphonull or phosphomimetic mutations on T427 decreases HCMV DNA replication in *OriLyt* transcomplementation assays [69]. Thirdly, UL44 is known to transport to the nucleus other viral factors, including the uracil-DNA glycosylase UL114 [104] and the UL54 pol catalytic subunit [54], so that regulation of its nuclear transport activity would also influence the subcellular localization of these factors (Figure 3). Finally, the key phosphorylation sites are all conserved in other β -Herpesviruses, including HHV-6A and 6B and HHV-7 [52].

However, it is also possible that the role of the UL44 C-terminus in HCMV replication does not relate exclusively to its ability to confer phosphorylation-dependent nuclear accumulation, since the

HCMV C-terminus is capable of conferring binding to factors other than BRAP2 and IMPs, including the viral phosphoprotein UL112-113 p84 [96] and the SUMO E2 conjugase Ubc9 [97]; the exact role these interactions play in the HCMV life cycle, however, is unclear. What is clear is that phosphorylation through a CK2-triggered phosphorylation cascade [69] of UL44 residues S413, S415 and S418 is absolutely required for nuclear targeting and viral replication [103], implying that pharmacological inhibition of CK2 activity would also result in inhibition of viral replication. Whether recently developed, highly selective CK2 inhibitors, [107] may have application in anti-HCMV therapy remains to be seen, and cell toxicity associated to the use of such compounds will have to be considered [108].

Figure 3. Phosphorylation-regulated nuclear transport of HCMV PAP. HCMV PAP UL44 nuclear import depends on the phosphorylation of its C-terminus. (A) Unphosphorylated UL44 C-terminus can be recognized by the IMP α / β heterodimer with medium affinity. This results in a low rate nuclear import of UL44, possibly enabling interaction with other cytoplasmic factors before import occurs. X indicates either another UL44 molecule, UL54 or the viral uracil DNA glycosylase; (B) Phosphorylation of T427, as mediated by UL97 or cdk1 enables binding to the cytoplasmic retention factor BRAP2, thereby preventing IMP α / β recognition and nuclear targeting. Phosphorylation of S413, S415 and S418 by CK2 and CK1 enhances IMP α / β binding, thereby promoting rapid nuclear import, possibly preventing UL44 to interact with other partners before nuclear import (C).



7. Nuclear Import of DNA Primase Helicase Complexes and the Formation of Composite NLSs

Despite the fact that several subunits of the polymerase holoenzyme possess functional NLSs, such signals have still not been identified on any herpesviral primase/helicase subunit. Indeed, ectopic expression of any of these three subunits in isolation has always been reported to be cytoplasmic [61,63,64,109], despite the fact that all three clearly need to be translocated to the nucleus

in order to perform their role in viral replication, and can be readily detected in the nucleus of infected cells [109,110]. In all studies to date, the simultaneous expression of at least all three subunits appears to be necessary for nuclear targeting; in the case of HSV-1 and EBV, coexpression of the three subunits appears to be sufficient for nuclear targeting of the complexes [61,63,64]. In the case of KSHV, the DNA polymerase holoenzyme and the ssDNA binding protein also need to be coexpressed with the helicase/primase holoenzyme to ensure nuclear localization [62]. Thus, herpesviruses appear to have evolved mechanisms to prevent nuclear entry of individual, unassembled helicase/primase subunits, presumably to avoid their potential interaction with cellular rather than viral DNA [61–64].

Based on these observations, it is currently believed that assembly of the DNA helicase/primase holoenzymes generates an as yet unidentified but functional NLS, which may depend on conformational changes induced by interaction of the subunits either to unmask an otherwise masked nuclear targeting peptide, or to create a composite NLS which is formed by amino acids belonging to each individual subunits. Consistent with the first possibility, bioinformatics analysis has identified putative hPY-NLSs on HSV-1 PAF UL8 and KSHV primase ORF52 (respectively PGALAPVFAFLGPEFEVRGGPVPY₃₅₈ and TFQSQVAWLRTKFTALRKLYKMTPSPY₁₆₀), although it remains to be elucidated whether these signals are functional. In the case of KSHV, this scenario is completed by the evidence that PAF (ORF40/41) can also be partially translocated to the nucleus when expressed in the presence of other viral proteins, such as K8 or MTA, implying that piggy-back transport through individual viral factors may also contribute to nuclear accumulation [62]. The IMPs responsible have not been identified, but a recent report has implicated the cellular chaperone DNAJB6 in the nuclear transport of HCMV primase (UL70) [109]. Significantly, DNAB6 is not the only cellular chaperone playing a role in the nuclear targeting of herpesviral replicating proteins (see below).

8. An Emerging Role for Cellular Chaperones in Nuclear Import

There is growing evidence that chaperones such as heat shock protein (Hsp) 90 and Hsp40 play an important role in nuclear targeting of herpesviral DNA replication proteins. The first example is represented by HSV-1 UL30, which bears two independently acting NLSs (see Table 2 and Figure 2) and which can localize to the nucleus independently of its cognate PAP [53,55,56]. Once inside the cell nucleus, UL30 co-localizes in replication compartments with Hsp90. Drug-mediated inhibition of Hsp90 activity resulted in cytoplasmic retention of UL30 and its degradation via the proteasome [87]. The inability of UL30 to reach the nuclear compartment is probably the reason for the simultaneous drop in viral DNA replication and infectious particle production observed upon inhibition of Hsp90 in HSV-1 infected cells *in vitro* [87]. It is likely that Hsp90 activity is required for correct folding of UL30, thus exposing the NLSs and promoting their interaction with IMPs for nuclear targeting. Importantly, a recent report showed that either pharmacological or genetic ablation of Hsp90 causes a drop in EBV DNA replication during lytic infection [60]. As in the case of HSV-1, this inhibition is most likely dependent on the observed mislocalization of the EBV pol (BALF1) to the cytosol. Immunoprecipitation data suggested that Hsp90 is required to enable BALF1 to interact with BMRF1. Therefore, while in the case of HSV-1, Hsp90 activity is likely required for correct folding of UL30 to enable interaction with IMPs/subsequent nuclear transport, Hsp90 probably assists the folding of EBV BALF1 to enable its interaction with BMRF1, which then drives nuclear targeting of the DNA

polymerase holoenzyme through its C-terminally located cNLS (see Table 2 and Figure 2) [58]. Another herpesvirus whose life cycle can be targeted by Hsp90 inhibitors is VZV, where nuclear localization of the VZV ssDNA-binding protein ORF29p relies on Hsp90 activity. ORF29p localizes to the nucleus of infected cells during lytic infection, and when individually expressed in the absence of other viral proteins, through an IMP α/β recognized, N-terminally located atypical NLS of 144 aas [111]; inhibition of Hsp90 activity by geldanamycin results in ORF29p mislocalization to the cytoplasm, where it is rapidly subjected to proteasomal degradation [112]. Moreover in the case of VZV-infected cells, Hsp90 inhibition causes a dramatic reduction of both viral genome replication and viral progeny production. Interestingly, although colocalizing with Hsp90, ORF29 does not appear to bind Hsp90 directly, but potentially indirectly through the host cochaperone/adaptor protein BAG3 [112]. Consistent with this idea, siRNA-mediated inhibition of BAG3 expression results in delayed/ineffective VZV spread and replication [112].

Taken together these data indicated that various herpesviruses can exploit the activity of Hsp90 in order to enable viral lytic cycle progression, implying that Hsp90 may represent a common target for approaches to prevent herpesvirus replication. Indeed pharmacological ablation of HSP90 inhibits the HSV-1, EBV and VSZ life cycles [60,87,112]. Consistent with the idea that chaperone proteins may play a role in infection by all herpesviruses is the fact that HCMV primase UL70 has recently been shown to interact with the Hsp40 family member DNAJB6. Two different isoforms of DNAJB6 protein, generated by alternative splicing, have been thus far identified: DNAJB6-a and DNAJB6-b, which exhibit distinct subcellular localization. While DNAJB6-a is nuclear, DNAJB6-b localizes exclusively in the cytoplasm, possibly through the fact that DNAJB6-b lacks the C-terminal domain, which contains a functional NLS [113]. Both isoforms interact with UL70, as demonstrated in the yeast two hybrid system and coimmunoprecipitations, suggesting that the interaction does not require the C-terminal domain of DNAJB6 [109]. The relative levels of DNAJB6 isoforms appear to play a crucial role in regulating the subcellular localization of HCMV UL70 protein. As reported for several herpesviral helicase/primase subunits, individual expression of UL70 in the absence of other viral proteins results in cytoplasmic localization [109]. Upon coexpression with DNAJB6-a, but not DNAJB6-b, UL70 accumulates strongly in the nucleus, suggesting that DNAJB6-a plays a crucial role in mediating UL70 nuclear accumulation, presumably via a piggy-back mechanism. In contrast, cells overexpressing HCMV primase and DNAJB6-b showed marked cytoplasmic localization of both proteins. Similar results have been obtained in HCMV-infected cells, where siRNA-mediated knockdown of DNAJB6-a or overexpression of DNAJB6-b prior to infection resulted in dramatically reduced HCMV replication and virus production, while silencing of DNAJB6-b or overexpression of DNAJB6-a causes an increase in HCMV replication and virus production [109]; levels of HCMV replication and virus production correlate well with nuclear accumulation of myc-tagged cotransfected UL70. Thus, it appears that HCMV has evolved mechanisms to exploit variations in the levels of DNAJB6 isoforms to regulate the subcellular localization of its primase UL70 and possibly the whole DNA primase/helicase complex.

9. Conclusions

Nuclear targeting of herpesvirus DNA replication factors remains largely uncharacterized, despite the fact that it represents a potential target for therapeutic intervention. Key unanswered questions include why HSV-1 and HCMV DNA polymerase holoenzymes subunits can be imported as individual proteins or in a complex, while this is not the case for γ -herpesviruses [52–56,59,60]; why HCMV evolved such a sophisticated mechanism of regulation by multiple phosphorylation sites of nuclear targeting of its PAP [51,52,69,95], and whether Hsp90 activity is required for other herpesviral proteins in addition to EBV BALF5, HSV-1 UL30 and VZV protein ORF29p [60,87,112]. In addition, the exact mechanism enabling nuclear transport of helicase/primase complexes, as well as the cellular factors (including IMPs) responsible are still unknown [61–64,109].

Answering these questions is not only important for fundamental understanding of herpesviral biology, but also for the development of urgently needed antiviral drugs to fight herpesvirus infections. Inhibition of IMP α/β mediated nuclear import has been recently shown to impair both HIV-1 and DENV replication [89,90]; since several crucial herpesviral proteins are dependent on IMP α/β for transport to the nucleus, it seems highly likely that pharmacological targeting of IMP α/β may impact on herpesvirus production. Along the same lines, it is possible that inhibition of kinases phosphorylating the C-terminus of HCMV PAP, including CK2, may result in impaired HCMV replication, by impairing PAP nuclear import. Finally, Hsp90 inhibitors may represent a novel class of anti-herpetic drugs able to block nuclear import of several DNA replicating machinery proteins. Further studies will be required to verify all of these exciting possibilities.

Acknowledgments

This study was founded by PRIN 2010-2011 (grant no. 2010Z9FLM8) to G.P. and Fondi attività di ricerca istituzionali to G.A.

Conflicts of Interest

The authors declare no conflict of interest.

References and Notes

1. Roizman, B.; Pellett, P.E. The family Herpesviridae: A brief introduction. In *Fields Virology*, 4th ed.; Knipe, D.M., Howley, P.M., Griffin, D.E., Lamb, R.A., Martin, M.A., Roizman, B., Straus, S.E., Eds.; Lippincott Williams & Wilkins: Philadelphia, PA, USA, 2001; Volume 2.
2. Wilkinson, D.E.; Weller, S.K. The role of DNA recombination in herpes simplex virus DNA replication. *IUBMB Life* **2003**, *55*, 451–458.
3. Strang, B.L.; Boulant, S.; Coen, D.M. Nucleolin associates with the human cytomegalovirus DNA polymerase accessory subunit UL44 and is necessary for efficient viral replication. *J. Virol.* **2010**, *84*, 1771–1784.
4. Anders, D.G.; Kacica, M.A.; Pari, G.; Punturieri, S.M. Boundaries and structure of human cytomegalovirus oriLyt, a complex origin for lytic-phase DNA replication. *J. Virol.* **1992**, *66*, 3373–3384.

5. Pari, G.S.; Anders, D.G. Eleven loci encoding trans-acting factors are required for transient complementation of human cytomegalovirus oriLyt-dependent DNA replication. *J. Virol.* **1993**, *67*, 6979–6988.
6. Pari, G.S.; Kacica, M.A.; Anders, D.G. Open reading frames UL44, IRS1/TRS1, and UL36-38 are required for transient complementation of human cytomegalovirus oriLyt-dependent DNA synthesis. *J. Virol.* **1993**, *67*, 2575–2582.
7. Anders, D.G.; McCue, L.A. The human cytomegalovirus genes and proteins required for DNA synthesis. *Intervirology* **1996**, *39*, 378–388.
8. Tran, E.J.; Wentz, S.R. Dynamic nuclear pore complexes: Life on the edge. *Cell* **2006**, *125*, 1041–1053.
9. Cardarelli, F.; Serresi, M.; Bizzarri, R.; Giacca, M.; Beltram, F. *In vivo* study of HIV-1 Tat arginine-rich motif unveils its transport properties. *Mol. Ther.* **2007**, *15*, 1313–1322.
10. Fahrenkrog, B.; Aebi, U. The nuclear pore complex: Nucleocytoplasmic transport and beyond. *Nat. Rev. Mol. Cell Biol.* **2003**, *4*, 757–766.
11. Katahira, J.; Yoneda, Y. Nucleocytoplasmic transport of microRNAs and related small RNAs. *Traffic* **2011**, *12*, 1468–1474.
12. Weis, K. Importins and exportins: How to get in and out of the nucleus. *Trends Biochem. Sci.* **1998**, *23*, 185–189.
13. Czernilofsky, A.P.; Levinson, A.D.; Varmus, H.E.; Bishop, J.M.; Tischer, E.; Goodman, H.M. Nucleotide sequence of an avian sarcoma virus oncogene (src) and proposed amino acid sequence for gene product. *Nature* **1980**, *287*, 198–203.
14. Balch, W.E.; McCaffery, J.M.; Plutner, H.; Farquhar, M.G. Vesicular stomatitis virus glycoprotein is sorted and concentrated during export from the endoplasmic reticulum. *Cell* **1994**, *76*, 841–852.
15. Kalderon, D.; Richardson, W.D.; Markham, A.F.; Smith, A.E. Sequence requirements for nuclear location of simian virus 40 large-T antigen. *Nature* **1984**, *311*, 33–38.
16. Kalderon, D.; Roberts, B.L.; Richardson, W.D.; Smith, A.E. A short amino acid sequence able to specify nuclear location. *Cell* **1984**, *39*, 499–509.
17. Lange, A.; Mills, R.E.; Lange, C.J.; Stewart, M.; Devine, S.E.; Corbett, A.H. Classical nuclear localization signals: Definition, function, and interaction with importin alpha. *J. Biol. Chem.* **2007**, *282*, 5101–5105.
18. Conti, E.; Uy, M.; Leighton, L.; Blobel, G.; Kuriyan, J. Crystallographic analysis of the recognition of a nuclear localization signal by the nuclear import factor karyopherin alpha. *Cell* **1998**, *94*, 193–204.
19. Kobe, B. Autoinhibition by an internal nuclear localization signal revealed by the crystal structure of mammalian importin alpha. *Nat. Struct. Biol.* **1999**, *6*, 388–397.
20. Lee, S.J.; Matsuura, Y.; Liu, S.M.; Stewart, M. Structural basis for nuclear import complex dissociation by RanGTP. *Nature* **2005**, *435*, 693–696.
21. Scheffzek, K.; Klebe, C.; Fritz-Wolf, K.; Kabsch, W.; Wittinghofer, A. Crystal structure of the nuclear Ras-related protein Ran in its GDP-bound form. *Nature* **1995**, *374*, 378–381.

22. Vetter, I.R.; Nowak, C.; Nishimoto, T.; Kuhlmann, J.; Wittinghofer, A. Structure of a Ran-binding domain complexed with Ran bound to a GTP analogue: Implications for nuclear transport. *Nature* **1999**, *398*, 39–46.
23. Izaurralde, E.; Kutay, U.; von Kobbe, C.; Mattaj, I.W.; Gorlich, D. The asymmetric distribution of the constituents of the Ran system is essential for transport into and out of the nucleus. *EMBO J.* **1997**, *16*, 6535–6547.
24. Bischoff, F.R.; Klebe, C.; Kretschmer, J.; Wittinghofer, A.; Ponstingl, H. RanGAP1 induces GTPase activity of nuclear Ras-related Ran. *Proc. Natl. Acad. Sci. USA* **1994**, *91*, 2587–2591.
25. Bischoff, F.R.; Ponstingl, H. Catalysis of guanine nucleotide exchange on Ran by the mitotic regulator RCC1. *Nature* **1991**, *354*, 80–82.
26. Ribbeck, K.; Lipowsky, G.; Kent, H.M.; Stewart, M.; Gorlich, D. NTF2 mediates nuclear import of Ran. *EMBO J.* **1998**, *17*, 6587–6598.
27. Kutay, U.; Bischoff, F.R.; Kostka, S.; Kraft, R.; Gorlich, D. Export of importin alpha from the nucleus is mediated by a specific nuclear transport factor. *Cell* **1997**, *90*, 1061–1071.
28. Hogarth, C.A.; Calanni, S.; Jans, D.A.; Loveland, K.L. Importin alpha mRNAs have distinct expression profiles during spermatogenesis. *Dev. Dyn.* **2006**, *235*, 253–262.
29. Huber, J.; Cronshagen, U.; Kadokura, M.; Marshallsay, C.; Wada, T.; Sekine, M.; Luhrmann, R. Snurportin1, an m3G-cap-specific nuclear import receptor with a novel domain structure. *EMBO J.* **1998**, *17*, 4114–4126.
30. Alvisi, G.; Poon, I.K.; Jans, D.A. Tumor-specific nuclear targeting: Promises for anti-cancer therapy? *Drug Resist. Updates* **2006**, *9*, 40–50.
31. Kuusisto, H.V.; Wagstaff, K.M.; Alvisi, G.; Jans, D.A. The C-terminus of apoptin represents a unique tumor cell-enhanced nuclear targeting module. *Int. J. Cancer* **2008**, *123*, 2965–2969.
32. Kuusisto, H.V.; Wagstaff, K.M.; Alvisi, G.; Roth, D.M.; Jans, D.A. Global enhancement of nuclear localization-dependent nuclear transport in transformed cells. *FASEB J.* **2012**, *26*, 1181–1193.
33. Truant, R.; Cullen, B.R. The arginine-rich domains present in human immunodeficiency virus type 1 Tat and Rev function as direct importin beta-dependent nuclear localization signals. *Mol. Cell. Biol.* **1999**, *19*, 1210–1217.
34. Cherezova, L.; Burnside, K.L.; Rose, T.M. Conservation of complex nuclear localization signals utilizing classical and non-classical nuclear import pathways in LANA homologs of KSHV and RFHV. *PLoS One* **2011**, *6*, e18920.
35. Fontes, M.R.; Teh, T.; Kobe, B. Structural basis of recognition of monopartite and bipartite nuclear localization sequences by mammalian importin-alpha. *J. Mol. Biol.* **2000**, *297*, 1183–1194.
36. Chook, Y.M.; Suel, K.E. Nuclear import by karyopherin-betas: Recognition and inhibition. *Biochim. Biophys. Acta* **2011**, *1813*, 1593–1606.
37. Pollard, V.W.; Michael, W.M.; Nakielny, S.; Siomi, M.C.; Wang, F.; Dreyfuss, G. A novel receptor-mediated nuclear protein import pathway. *Cell* **1996**, *86*, 985–994.
38. Kawamura, H.; Tomozoe, Y.; Akagi, T.; Kamei, D.; Ochiai, M.; Yamada, M. Identification of the nucleocytoplasmic shuttling sequence of heterogeneous nuclear ribonucleoprotein D-like protein JKTBP and its interaction with mRNA. *J. Biol. Chem.* **2002**, *277*, 2732–2739.

39. Rebane, A.; Aab, A.; Steitz, J.A. Transportins 1 and 2 are redundant nuclear import factors for hnRNP A1 and HuR. *RNA* **2004**, *10*, 590–599.
40. Siomi, M.C.; Fromont, M.; Rain, J.C.; Wan, L.; Wang, F.; Legrain, P.; Dreyfuss, G. Functional conservation of the transportin nuclear import pathway in divergent organisms. *Mol. Cell. Biol.* **1998**, *18*, 4141–4148.
41. Truant, R.; Kang, Y.; Cullen, B.R. The human tap nuclear RNA export factor contains a novel transportin-dependent nuclear localization signal that lacks nuclear export signal function. *J. Biol. Chem.* **1999**, *274*, 32167–32171.
42. Lee, B.J.; Cansizoglu, A.E.; Suel, K.E.; Louis, T.H.; Zhang, Z.; Chook, Y.M. Rules for nuclear localization sequence recognition by karyopherin beta 2. *Cell* **2006**, *126*, 543–558.
43. Imasaki, T.; Shimizu, T.; Hashimoto, H.; Hidaka, Y.; Kose, S.; Imamoto, N.; Yamada, M.; Sato, M. Structural basis for substrate recognition and dissociation by human transportin 1. *Mol. Cell* **2007**, *28*, 57–67.
44. Cansizoglu, A.E.; Lee, B.J.; Zhang, Z.C.; Fontoura, B.M.; Chook, Y.M. Structure-based design of a pathway-specific nuclear import inhibitor. *Nat. Struct. Mol. Biol.* **2007**, *14*, 452–454.
45. Okada, N.; Ishigami, Y.; Suzuki, T.; Kaneko, A.; Yasui, K.; Fukutomi, R.; Isemura, M. Importins and exportins in cellular differentiation. *J. Cell. Mol. Med.* **2008**, *12*, 1863–1871.
46. Jans, D.A.; Xiao, C.Y.; Lam, M.H. Nuclear targeting signal recognition: A key control point in nuclear transport? *Bioessays* **2000**, *22*, 532–544.
47. Terry, L.J.; Shows, E.B.; Wentz, S.R. Crossing the nuclear envelope: Hierarchical regulation of nucleocytoplasmic transport. *Science* **2007**, *318*, 1412–1416.
48. Poon, I.K.; Jans, D.A. Regulation of nuclear transport: Central role in development and transformation? *Traffic* **2005**, *6*, 173–186.
49. Briggs, L.J.; Stein, D.; Goltz, J.; Corrigan, V.C.; Efthymiadis, A.; Hubner, S.; Jans, D.A. The cAMP-dependent protein kinase site (Ser312) enhances dorsal nuclear import through facilitating nuclear localization sequence/importin interaction. *J. Biol. Chem.* **1998**, *273*, 22745–22752.
50. Hubner, S.; Xiao, C.Y.; Jans, D.A. The protein kinase CK2 site (Ser111/112) enhances recognition of the simian virus 40 large T-antigen nuclear localization sequence by importin. *J. Biol. Chem.* **1997**, *272*, 17191–17195.
51. Fulcher, A.J.; Roth, D.M.; Fatima, S.; Alvisi, G.; Jans, D.A. The BRCA-1 binding protein BRAP2 is a novel, negative regulator of nuclear import of viral proteins, dependent on phosphorylation flanking the nuclear localization signal. *FASEB J.* **2010**, *24*, 1454–1466.
52. Alvisi, G.; Jans, D.A.; Guo, J.; Pinna, L.A.; Ripalti, A. A protein kinase CK2 site flanking the nuclear targeting signal enhances nuclear transport of human cytomegalovirus ppUL44. *Traffic* **2005**, *6*, 1002–1013.
53. Alvisi, G.; Musiani, D.; Jans, D.A.; Ripalti, A. An importin alpha/beta-recognized bipartite nuclear localization signal mediates targeting of the human herpes simplex virus type 1 DNA polymerase catalytic subunit pUL30 to the nucleus. *Biochemistry* **2007**, *46*, 9155–9163.
54. Alvisi, G.; Ripalti, A.; Ngankeu, A.; Giannandrea, M.; Caraffi, S.G.; Dias, M.M.; Jans, D.A. Human cytomegalovirus DNA polymerase catalytic subunit pUL54 possesses independently acting nuclear localization and ppUL44 binding motifs. *Traffic* **2006**, *7*, 1322–1332.

55. Alvisi, G.; Avanzi, S.; Musiani, D.; Camozzi, D.; Leoni, V.; Ly-Huynh, J.D.; Ripalti, A. Nuclear import of HSV-1 DNA polymerase processivity factor UL42 is mediated by a C-terminally located bipartite nuclear localization signal. *Biochemistry* **2008**, *47*, 13764–13777.
56. Loregian, A.; Piaia, E.; Cancellotti, E.; Papini, E.; Marsden, H.S.; Palu, G. The catalytic subunit of herpes simplex virus type 1 DNA polymerase contains a nuclear localization signal in the UL42-binding region. *Virology* **2000**, *273*, 139–148.
57. Takeda, K.; Haque, M.; Nagoshi, E.; Takemoto, M.; Shimamoto, T.; Yoneda, Y.; Yamanishi, K. Characterization of human herpesvirus 7 U27 gene product and identification of its nuclear localization signal. *Virology* **2000**, *272*, 394–401.
58. Zhang, Q.; Holley-Guthrie, E.; Dorsky, D.; Kenney, S. Identification of transactivator and nuclear localization domains in the Epstein-Barr virus DNA polymerase accessory protein, BMRF1. *J. Gen. Virol.* **1999**, *80*, 69–74.
59. Chen, Y.; Ciustea, M.; Ricciardi, R.P. Processivity factor of KSHV contains a nuclear localization signal and binding domains for transporting viral DNA polymerase into the nucleus. *Virology* **2005**, *340*, 183–191.
60. Kawashima, D.; Kanda, T.; Murata, T.; Saito, S.; Sugimoto, A.; Narita, Y.; Tsurumi, T. Nuclear transport of epstein-barr virus DNA polymerase is dependent on the BMRF1 polymerase processivity factor and molecular chaperone Hsp90. *J. Virol.* **2013**, *87*, 6482–6491.
61. Gao, Z.; Krithivas, A.; Finan, J.E.; Semmes, O.J.; Zhou, S.; Wang, Y.; Hayward, S.D. The Epstein-Barr virus lytic transactivator Zta interacts with the helicase-primase replication proteins. *J. Virol.* **1998**, *72*, 8559–8567.
62. Wu, F.Y.; Ahn, J.H.; Alcendor, D.J.; Jang, W.J.; Xiao, J.; Hayward, S.D.; Hayward, G.S. Origin-independent assembly of Kaposi's sarcoma-associated herpesvirus DNA replication compartments in transient cotransfection assays and association with the ORF-K8 protein and cellular PML. *J. Virol.* **2001**, *75*, 1487–1506.
63. Calder, J.M.; Stow, E.C.; Stow, N.D. On the cellular localization of the components of the herpes simplex virus type 1 helicase-primase complex and the viral origin-binding protein. *J. Gen. Virol.* **1992**, *73*, 531–538.
64. Barnard, E.C.; Brown, G.; Stow, N.D. Deletion mutants of the herpes simplex virus type 1 UL8 protein: Effect on DNA synthesis and ability to interact with and influence the intracellular localization of the UL5 and UL52 proteins. *Virology* **1997**, *237*, 97–106.
65. Ertl, P.F.; Powell, K.L. Physical and functional interaction of human cytomegalovirus DNA polymerase and its accessory protein (ICP36) expressed in insect cells. *J. Virol.* **1992**, *66*, 4126–4133.
66. Gottlieb, J.; Marcy, A.I.; Coen, D.M.; Challberg, M.D. The herpes simplex virus type 1 UL42 gene product: A subunit of DNA polymerase that functions to increase processivity. *J. Virol.* **1990**, *64*, 5976–5987.
67. Loregian, A.; Case, A.; Cancellotti, E.; Valente, C.; Marsden, H.S.; Palu, G. Cloning, expression, and functional characterization of the equine herpesvirus 1 DNA polymerase and its accessory subunit. *J. Virol.* **2006**, *80*, 6247–6258.
68. Stinski, M.F. Herpesviruses. In *Herpesviruses*; Roizman, B., Ed.; Plenum Press: New York, NY, USA, 1983; Volume 2, pp. 67–113.

69. Alvisi, G.; Marin, O.; Pari, G.; Mancini, M.; Avanzi, S.; Loregian, A.; Jans, D.A.; Ripalti, A. Multiple phosphorylation sites at the C-terminus regulate nuclear import of HCMV DNA polymerase processivity factor ppUL44. *Virology* **2011**, *417*, 259–267.
70. Alvisi, G.; Rawlinson, S.M.; Ghildyal, R.; Ripalti, A.; Jans, D.A. Regulated nucleocytoplasmic trafficking of viral gene products: A therapeutic target? *Biochim. Biophys. Acta* **2008**, *1784*, 213–227.
71. Alvisi, G.; Roth, D.M.; Camozzi, D.; Pari, G.S.; Loregian, A.; Ripalti, A.; Jans, D.A. The flexible loop of the human cytomegalovirus DNA polymerase processivity factor ppUL44 is required for efficient DNA binding and replication in cells. *J. Virol.* **2009**, *83*, 9567–9576.
72. Weiland, K.L.; Oien, N.L.; Homa, F.; Wathen, M.W. Functional analysis of human cytomegalovirus polymerase accessory protein. *Virus Res.* **1994**, *34*, 191–206.
73. Digard, P.; Chow, C.S.; Pirrit, L.; Coen, D.M. Functional analysis of the herpes simplex virus UL42 protein. *J. Virol.* **1993**, *67*, 1159–1168.
74. Gallo, M.L.; Dorsky, D.I.; Crumpacker, C.S.; Parris, D.S. The essential 65-kilodalton DNA-binding protein of herpes simplex virus stimulates the virus-encoded DNA polymerase. *J. Virol.* **1989**, *63*, 5023–5029.
75. Gallo, M.L.; Jackwood, D.H.; Murphy, M.; Marsden, H.S.; Parris, D.S. Purification of the herpes simplex virus type 1 65-kilodalton DNA-binding protein: Properties of the protein and evidence of its association with the virus-encoded DNA polymerase. *J. Virol.* **1988**, *62*, 2874–2883.
76. Zuccola, H.J.; Filman, D.J.; Coen, D.M.; Hogle, J.M. The crystal structure of an unusual processivity factor, herpes simplex virus UL42, bound to the C terminus of its cognate polymerase. *Mol. Cell* **2000**, *5*, 267–278.
77. Appleton, B.A.; Loregian, A.; Filman, D.J.; Coen, D.M.; Hogle, J.M. The cytomegalovirus DNA polymerase subunit UL44 forms a C clamp-shaped dimer. *Mol. Cell* **2004**, *15*, 233–244.
78. Murayama, K.; Nakayama, S.; Kato-Murayama, M.; Akasaka, R.; Ohbayashi, N.; Kamewari-Hayami, Y.; Terada, T.; Shirouzu, M.; Tsurumi, T.; Yokoyama, S. Crystal structure of Epstein-Barr virus DNA polymerase processivity factor BMRF1. *J. Biol. Chem.* **2009**, *284*, 35896–35905.
79. Baltz, J.L.; Filman, D.J.; Ciustea, M.; Silverman, J.E.; Lautenschlager, C.L.; Coen, D.M.; Ricciardi, R.P.; Hogle, J.M. The crystal structure of PF-8, the DNA polymerase accessory subunit from Kaposi's sarcoma-associated herpesvirus. *J. Virol.* **2009**, *83*, 12215–12228.
80. Krishna, T.S.; Kong, X.P.; Gary, S.; Burgers, P.M.; Kuriyan, J. Crystal structure of the eukaryotic DNA polymerase processivity factor PCNA. *Cell* **1994**, *79*, 1233–1243.
81. Stinski, M.F. Sequence of protein synthesis in cells infected by human cytomegalovirus: Early and late virus-induced polypeptides. *J. Virol.* **1978**, *26*, 686–701.
82. Geballe, A.P.; Leach, F.S.; Mocarski, E.S. Regulation of cytomegalovirus late gene expression: Gamma genes are controlled by posttranscriptional events. *J. Virol.* **1986**, *57*, 864–874.
83. Bocconi, M.C.; Campanini, F.; Battista, M.C.; Bergamini, G.; Dal Monte, P.; Ripalti, A.; Landini, M.P. Human cytomegalovirus product UL44 downregulates the transactivation of HIV-1 long terminal repeat. *AIDS* **1998**, *12*, 365–372.

84. Zhang, Q.; Hong, Y.; Dorsky, D.; Holley-Guthrie, E.; Zalani, S.; Elshiekh, N.A.; Kiehl, A.; Le, T.; Kenney, S. Functional and physical interactions between the Epstein-Barr virus (EBV) proteins BZLF1 and BMRF1: Effects on EBV transcription and lytic replication. *J. Virol.* **1996**, *70*, 5131–5142.
85. Advani, S.J.; Weichselbaum, R.R.; Roizman, B. Herpes simplex virus 1 activates cdc2 to recruit topoisomerase II alpha for post-DNA synthesis expression of late genes. *Proc. Natl. Acad. Sci. USA* **2003**, *100*, 4825–4830.
86. Perry, R.B.; Fainzilber, M. Nuclear transport factors in neuronal function. *Semin. Cell Dev. Biol.* **2009**, *20*, 600–606.
87. Burch, A.D.; Weller, S.K. Herpes simplex virus type 1 DNA polymerase requires the mammalian chaperone hsp90 for proper localization to the nucleus. *J. Virol.* **2005**, *79*, 10740–10749.
88. Wagstaff, K.M.; Rawlinson, S.M.; Hearps, A.C.; Jans, D.A. An AlphaScreen(R)-based assay for high-throughput screening for specific inhibitors of nuclear import. *J. Biomol. Screen* **2011**, *16*, 192–200.
89. Wagstaff, K.M.; Sivakumaran, H.; Heaton, S.M.; Harrich, D.; Jans, D.A. Ivermectin is a specific inhibitor of importin alpha/beta-mediated nuclear import able to inhibit replication of HIV-1 and dengue virus. *Biochem. J.* **2012**, *443*, 851–856.
90. Tay, M.Y.; Fraser, J.E.; Chan, K.K.; Moreland, N.J.; Rathore, A.P.; Wang, C.; Vasudevan, S.G.; Jans, D.A. Nuclear localization of dengue virus (DENV) 1–4 non-structural protein 5; protection against all 4 DENV serotypes by the inhibitor Ivermectin. *Antivir. Res.* **2013**, *99*, 301–306.
91. Caly, L.; Wagstaff, K.M.; Jans, D.A. Nuclear trafficking of proteins from RNA viruses: Potential target for antivirals? *Antivir. Res.* **2012**, *95*, 202–206.
92. Haffar, O.; Dubrovsky, L.; Lowe, R.; Berro, R.; Kashanchi, F.; Godden, J.; Vanpouille, C.; Bajorath, J.; Bukrinsky, M. Oxadiazols: A new class of rationally designed anti-human immunodeficiency virus compounds targeting the nuclear localization signal of the viral matrix protein. *J. Virol.* **2005**, *79*, 13028–13036.
93. Kaiser, N.; Lischka, P.; Wagenknecht, N.; Stamminger, T. Inhibition of human cytomegalovirus replication via peptide aptamers directed against the nonconventional nuclear localization signal of the essential viral replication factor pUL84. *J. Virol.* **2009**, *83*, 11902–11913.
94. Gao, M.; DiTusa, S.F.; Cordingley, M.G. The C-terminal third of UL42, a HSV-1 DNA replication protein, is dispensable for viral growth. *Virology* **1993**, *194*, 647–653.
95. Silva, L.A.; Loregian, A.; Pari, G.S.; Strang, B.L.; Coen, D.M. The carboxy-terminal segment of the human cytomegalovirus DNA polymerase accessory subunit UL44 is crucial for viral replication. *J. Virol.* **2010**, *84*, 11563–11568.
96. Kim, Y.E.; Ahn, J.H. Role of the specific interaction of UL112-113 p84 with UL44 DNA polymerase processivity factor in promoting DNA replication of human cytomegalovirus. *J. Virol.* **2010**, *84*, 8409–8421.
97. Sinigalia, E.; Alvisi, G.; Segre, C.V.; Mercorelli, B.; Muratore, G.; Winkler, M.; Hsiao, H.H.; Urlaub, H.; Ripalti, A.; Chiocca, S.; *et al.* The human cytomegalovirus DNA polymerase processivity factor UL44 is modified by SUMO in a DNA-dependent manner. *PLoS One* **2012**, *7*, e49630.

98. Mattosco, D.; Segré, C.V.; Chiocca, S. Viral manipulation of cellular protein conjugation pathways: The SUMO lesson. *World J. Virol.* **2013**, *2*, 79–90.
99. Davies, R.G.; Wagstaff, K.M.; McLaughlin, E.; Loveland, K.A.; Jans, D.A. The BRCA1-binding protein BRAP2 can act as a cytoplasmic retention factor for nuclear and nuclear envelope-localizing testicular proteins. *Biochim. Biophys. Acta* **2013**, doi:10.1016/j.bbamcr.2013.05.015.
100. Hodel, M.R.; Corbett, A.H.; Hodel, A.E. Dissection of a nuclear localization signal. *J. Biol. Chem.* **2001**, *276*, 1317–1325.
101. Rihs, H.P.; Peters, R. Nuclear transport kinetics depend on phosphorylation-site-containing sequences flanking the karyophilic signal of the Simian virus 40 T-antigen. *EMBO J.* **1989**, *8*, 1479–1484.
102. Fulcher, A.J.; Jans, D.A. Regulation of nucleocytoplasmic trafficking of viral proteins: An integral role in pathogenesis? *Biochim. Biophys. Acta* **2011**, *1813*, 2176–2190.
103. Silva, L.A.; Strang, B.L.; Lin, E.W.; Kamil, J.P.; Coen, D.M. Sites and roles of phosphorylation of the human cytomegalovirus DNA polymerase subunit UL44. *Virology* **2011**, *417*, 268–280.
104. Prichard, M.N.; Lawlor, H.; Duke, G.M.; Mo, C.; Wang, Z.; Dixon, M.; Kemble, G.; Kern, E.R. Human cytomegalovirus uracil DNA glycosylase associates with ppUL44 and accelerates the accumulation of viral DNA. *Virol. J.* **2005**, *2*, doi:10.1186/1743-422X-2-55.
105. Alvisi, G.; Jans, D.A.; Ripalti, A. Human cytomegalovirus (HCMV) DNA polymerase processivity factor ppUL44 dimerizes in the cytosol before translocation to the nucleus. *Biochemistry* **2006**, *45*, 6866–6872.
106. Sinigalia, E.; Alvisi, G.; Mercorelli, B.; Coen, D.M.; Pari, G.S.; Jans, D.A.; Ripalti, A.; Palu, G.; Loregian, A. Role of homodimerization of human cytomegalovirus DNA polymerase accessory protein UL44 in origin-dependent DNA replication in cells. *J. Virol.* **2008**, *82*, 12574–12579.
107. Cozza, G.; Pinna, L.A.; Moro, S. Kinase CK2 inhibition: An update. *Curr. Med. Chem.* **2013**, *20*, 671–693.
108. Duncan, J.S.; Gyenis, L.; Lenehan, J.; Bretner, M.; Graves, L.M.; Haystead, T.A.; Litchfield, D.W. An unbiased evaluation of CK2 inhibitors by chemoproteomics: Characterization of inhibitor effects on CK2 and identification of novel inhibitor targets. *Mol. Cell. Proteomics* **2008**, *7*, 1077–1088.
109. Pei, Y.; Fu, W.; Yang, E.; Shen, A.; Chen, Y.C.; Gong, H.; Chen, J.; Huang, J.; Xiao, G.; Liu, F. A Hsp40 chaperone protein interacts with and modulates the cellular distribution of the primase protein of human cytomegalovirus. *PLoS Pathog.* **2012**, *8*, e1002968.
110. Smith, J.A.; Jairath, S.; Crute, J.J.; Pari, G.S. Characterization of the human cytomegalovirus UL105 gene and identification of the putative helicase protein. *Virology* **1996**, *220*, 251–255.
111. Stallings, C.L.; Silverstein, S. Dissection of a novel nuclear localization signal in open reading frame 29 of varicella-zoster virus. *J. Virol.* **2005**, *79*, 13070–13081.
112. Kyratsous, C.A.; Silverstein, S.J. BAG3, a host cochaperone, facilitates varicella-zoster virus replication. *J. Virol.* **2007**, *81*, 7491–7503.
113. Hanai, R.; Mashima, K. Characterization of two isoforms of a human DnaJ homologue, HsJ2. *Mol. Biol. Rep.* **2003**, *30*, 149–153.

114. Makhov, A.M.; Subramanian, D.; Holley-Guthrie, E.; Kenney, S.C.; Griffith, J.D. The Epstein-Barr virus polymerase accessory factor BMRF1 adopts a ring-shaped structure as visualized by electron microscopy. *J. Biol. Chem.* **2004**, *279*, 40358–40361.
115. Chen, X.; Lin, K.; Ricciardi, R.P. Human Kaposi's sarcoma herpesvirus processivity factor-8 functions as a dimer in DNA synthesis. *J. Biol. Chem.* **2004**, *279*, 28375–28386.
116. Zhou, X.; Liao, Q.; Ricciardi, R.P.; Peng, C.; Chen, X. Kaposi's sarcoma-associated herpesvirus processivity factor-8 dimerizes in cytoplasm before being translocated to nucleus. *Biochem. Biophys. Res. Commun.* **2010**, *397*, 520–525.

© 2013 by the authors; licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).