

## RESEARCH PAPER

# Cholecystokinin CCK<sub>2</sub> receptors mediate the peptide's inhibitory actions on the contractile activity of human distal colon via the nitric oxide pathway

M Fornai<sup>1</sup>, R Colucci<sup>1</sup>, L Antonioli<sup>1</sup>, F Crema<sup>2</sup>, P Buccianti<sup>3</sup>, M Chiarugi<sup>3</sup>, F Baschiera<sup>1</sup>, N Ghisu<sup>1</sup>, M Tuccori<sup>1</sup>, C Blandizzi<sup>1</sup> and M Del Tacca<sup>1</sup>

<sup>1</sup>Division of Pharmacology and Chemotherapy, Department of Internal Medicine, University of Pisa, Pisa, Italy; <sup>2</sup>Department of Internal Medicine and Therapeutics, University of Pavia, Pavia, Italy and <sup>3</sup>Department of Surgery, University of Pisa, Pisa, Italy

**Background and purpose:** Cholecystokinin is known to exert stimulant actions on intestinal motility via activation of type 1 cholecystokinin receptors (CCK<sub>1</sub>). However, the role played by cholecystokinin 2 (CCK<sub>2</sub>) receptors in the regulation of gut motility remains undetermined. This study was designed to examine the influence of CCK<sub>2</sub> receptors on the contractile activity of human distal colon.

**Experimental approach:** The effects of compounds acting on CCK<sub>2</sub> receptors were assessed *in vitro* on motor activity of longitudinal smooth muscle, under basal conditions as well as in the presence of KCl-induced contractions or transmural electrical stimulation.

**Key results:** Cholecystokinin octapeptide sulphate induced concentration-dependent contractions which were enhanced by GV150013 (CCK<sub>2</sub> receptor antagonist; +57% at 0.01 μM). These effects were unaffected by tetrodotoxin. The enhancing actions of GV150013 on contractions evoked by cholecystokinin octapeptide sulphate were unaffected by N<sup>ω</sup>-propyl-L-arginine (NPA, neuronal nitric oxide synthase inhibitor), while they were prevented by N<sup>ω</sup>-nitro-L-arginine methylester (L-NAME, non-selective nitric oxide synthase inhibitor). In the presence of KCl-induced contractions, cholecystokinin octapeptide sulphate elicited concentration-dependent relaxations (-36%), which were unaffected by NPA, but were counteracted by GV150013 or L-NAME. The application of electrical stimuli evoked phasic contractions which were enhanced by GV150013 (+41 % at 0.01 μM).

**Conclusions and implications:** CCK<sub>2</sub> receptors mediate inhibitory actions of cholecystokinin on motor activity of human distal colon. It is suggested that CCK<sub>2</sub> receptors exert their modulating actions through a nitric oxide pathway, independent of the activity of the neuronal nitric oxide synthase isoform.

*British Journal of Pharmacology* (2007) **151**, 1246–1253; doi:10.1038/sj.bjp.0707339; published online 18 June 2007

**Keywords:** cholecystokinin; CCK<sub>2</sub> receptors; human colon; intestinal motility; nitric oxide

**Abbreviations:** CCK, cholecystokinin; CCK<sub>1</sub>, cholecystokinin 1 receptor subtype; CCK<sub>2</sub>, cholecystokinin 2 receptor subtype; CNS, central nervous system; GV150013, N-(+)-[1-adamant-1-ylmethyl]-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro-1H-1,5-benzodiazepin-3-yl]-N phenylurea; ICC, interstitial cells of Cajal; L-NAME, N<sup>ω</sup>-nitro-L-arginine methylester; nNOS, neuronal nitric oxide synthase; NOS, nitric oxide synthase; NPA, N<sup>ω</sup>-propyl-L-arginine

## Introduction

Cholecystokinin (CCK) regulates a variety of physiological functions acting as a neuropeptide in the central nervous system (CNS) and as both hormone and neurotransmitter in

the gastrointestinal tract (Crawley and Corwin, 1994; Miyasaka and Funakoshi, 2003). Several studies, based on functional, pharmacological and molecular approaches, have indicated that the regulatory actions of CCK are mediated by two receptor subtypes, designated as CCK<sub>1</sub> and CCK<sub>2</sub>. CCK<sub>1</sub> receptors are mainly located in the gut and in few areas of the CNS. CCK<sub>2</sub> receptors are widely expressed in the CNS as well as in the digestive tract, where they are mainly located in the stomach, but can also be found throughout the small and large intestines (Noble *et al.*, 1999; Miyasaka and Funakoshi, 2003).

Correspondence: Professor C Blandizzi, Divisione di Farmacologia e Chemioterapia, Dipartimento di Medicina Interna, Università di Pisa, Via Roma, 55, 56126 Pisa, Italy.  
E-mail: c.blandizzi@virgilio.it  
Received 22 March 2007; accepted 15 May 2007; published online 18 June 2007

CCK and its related peptides have been implicated in the pathophysiology of functional digestive diseases, such as irritable bowel syndrome and functional dyspepsia. Previous studies suggested that either an altered release of CCK or abnormal responses to this peptide could contribute to symptoms of irritable bowel syndrome (Sjolund *et al.*, 1996; Chey *et al.*, 2001). In particular, CCK may take part in the development of digestive symptoms, such as abdominal discomfort, pain and altered sensations to luminal stress (Chua and Keeling, 2006). On this basis, the pharmacological modulation of CCK pathways might be useful for treatment of symptoms associated with functional digestive disorders (Herranz, 2003).

In the gastrointestinal tract, CCK<sub>1</sub> receptors have been localized in the smooth muscle and myenteric neurons as well as abdominal afferent nerves (Sternini *et al.*, 1999), and they appear to be mainly involved in the control of gut motility and visceral sensation (Noble *et al.*, 1999; Varga *et al.*, 2004). The main functions regulated by CCK<sub>2</sub> receptors in the digestive system include the stimulation of acid secretion from parietal cells (Kulaksiz *et al.*, 2000; Ochi *et al.*, 2005) and the release of histamine from enterochromaffin-like cells in the stomach (Waldum *et al.*, 2002), but it has been suggested that these receptors might also contribute to the control of gut motor functions (Dal Forno *et al.*, 1992; Giralt and Vergara, 1999).

CCK actions on intestinal motility can vary in nature, depending on the species and gut region examined. In general, direct effects of CCK on smooth muscle are thought to result in contractile responses, whereas neurally mediated actions can evoke either contractile or relaxant activity, depending on the transmitter being released (Varga *et al.*, 2004). Studies in guinea pigs showed that CCK can stimulate intestinal motor activity through recruitment of CCK<sub>1</sub> receptors located on smooth muscle or myenteric nerve pathways (Dal Forno *et al.*, 1992; Fornai *et al.*, 2007). In addition, following the development of non-peptidic CCK<sub>2</sub> receptor antagonists (Herranz, 2003), *in vivo* studies with canine or rat models suggested the involvement of these receptors in the inhibitory control of gut motor functions (Vergara *et al.*, 1996; Giralt and Vergara, 1999) and, more recently, Fornai *et al.* (2007) observed that CCK inhibits the motility of isolated guinea-pig colon via activation of CCK<sub>2</sub> receptors leading to nitric oxide (NO) release from myenteric nerves.

When considering the effects of CCK on human gut, current evidence indicates that CCK induces contractions of colon and that these stimulant actions are mediated by CCK<sub>1</sub> receptors located on smooth muscle (D'Amato *et al.*, 1991; Morton *et al.*, 2002a). However, the possible involvement of CCK<sub>2</sub> receptors in the control of human bowel motility has not been previously demonstrated. Accordingly, the present study was designed to investigate the *in vitro* effects of compounds acting at CCK<sub>2</sub> receptors on both spontaneous and stimulated human colonic motility.

## Methods

### *Tissue excision and preparation*

Colonic specimens were obtained from patients undergoing surgery for uncomplicated neoplastic conditions. Patients

not older than 60 years, without history of gastrointestinal inflammatory diseases, digestive functional disorders, abdominal surgery or intestinal obstruction, were selected. Samples consisted of whole wall sections of distal colon from a macroscopically normal region taken at a distance of at least 10 cm from any visible lesion. Care was taken to verify the absence of alterations by histological examination. Portions of tissue were fixed in cold 4% paraformaldehyde, diluted in phosphate-buffered saline, for routine histology. The remaining parts of colonic tissues were placed into pre-oxygenated Krebs solution and transported on ice to laboratory. Longitudinal muscle strips of approximately 3 mm width and 20 mm length were prepared as described previously (Fornai *et al.*, 2005). Informed patient consent was obtained before surgery, and the experimental protocol was approved by the Ethics Committee of our University Hospital.

### *Recording of longitudinal muscle contractile activity*

The contractile activity of colonic longitudinal smooth muscle was recorded as described previously by Fornai *et al.* (2005). Preparations were set up in 10-ml organ baths containing Krebs solution at 37°C, bubbled with 95% O<sub>2</sub> + 5% CO<sub>2</sub>. Preparations were connected to isotonic transducers (Basile, Comerio, Italy), under a constant load of 1 g, and allowed to equilibrate for at least 30 min. Krebs solution had the following composition (mM): NaCl 113, KCl 4.7, CaCl<sub>2</sub> 2.5, KH<sub>2</sub>PO<sub>4</sub> 1.2, MgSO<sub>4</sub> 1.2, NaHCO<sub>3</sub> 25, glucose 11.5 (pH 7.4 ± 0.1). Longitudinal muscle activity was recorded by polygraphs (Basile, Comerio, Italy). Transmural electrical stimulation was delivered by a BM-ST6 stimulator (Biomedica Mangoni, Pisa, Italy). Stimuli were applied as 10-s single trains of square wave pulses (0.5 ms, 30 mA) at 10 Hz. The interval between successive periods of electrical stimulation was about 30 min. Each preparation was repeatedly challenged with electrical stimulations, and experiments started when reproducible responses were obtained (usually after 2–3 stimulations).

In the first set of experiments, the effects of CCK octapeptide sulphate (CCK-8S) were evaluated on spontaneous motor activity of colonic preparations. For this purpose, non-cumulative concentration–response curves were constructed for CCK-8S (0.001–1 μM) on basal contractility of longitudinal muscle. In the second set, the effect of CCK-8S (0.1 μM) was examined on spontaneous colonic activity after incubation with increasing concentrations of the selective CCK<sub>2</sub> receptor antagonist GV150013 (0.001–0.1 μM).

To investigate whether CCK<sub>2</sub> receptors regulate colonic motility at neural and/or muscular level, the effects of CCK-8S and its interaction with GV150013 (0.01 μM) were also assessed after blockade of myenteric nerves. For this purpose, colonic tissues were maintained in Krebs solution containing tetrodotoxin (1 μM). In addition, to evaluate the involvement of NO in the effects mediated by CCK<sub>2</sub> receptors, CCK-8S was assayed on colonic preparations maintained in Krebs solution containing the non-selective NO synthase (NOS) inhibitor N<sup>ω</sup>-nitro-L-arginine methylester (L-NAME, 100 μM) or the selective neuronal NOS (nNOS) inhibitor N<sup>ω</sup>-propyl-

L-arginine (NPA, 0.1  $\mu\text{M}$ ). In a separate group of experiments, colonic preparations were pre-contracted with KCl (60 mM). Under these conditions, the effect of CCK-8S was assessed either alone or in the presence of GV150013 (0.01  $\mu\text{M}$ ), NPA (0.1  $\mu\text{M}$ ) or L-NAME (100  $\mu\text{M}$ ). In the final set of experiments, the effects of GV150013 were assayed on contractile responses induced by electrical stimulation, applied as reported above.

In all experiments, CCK-8S was added to the organ bath solution at 20-min intervals, and the exposure time never exceeded 2 min, to avoid desensitization. The CCK<sub>2</sub> antagonist was added to the bathing fluid 25 min before the application of CCK-8S or electrical stimulation. Concentration–response curves were obtained from distinct preparations obtained from the same patient.

#### Statistical analysis

Results are given as mean  $\pm$  standard error of mean (s.e.m.). The significance of differences was evaluated on raw data, before percentage normalization, by analysis of variance for repeated measures or unpaired data, followed by *post hoc* analysis with Dunnett or Student–Newman–Keuls test, as appropriate.  $P < 0.05$  was considered significant. Colonic preparations included in each test group were obtained from distinct patients and therefore the number of experiments refers also to the number of patients assigned to each group. EC<sub>50</sub> values were interpolated from concentration–response curves and variability was expressed as 95% confidence intervals (95% CI). Calculations were performed using commercial software (GraphPad Prism, version 3.0 from GraphPad Software Inc., San Diego, CA, USA).

#### Drugs and reagents

Human CCK octapeptide sulphate, L-NAME (Sigma Chemical, St Louis, MO, USA); tetrodotoxin, NPA (Tocris Cookson, Bristol, UK); devazepide (kindly provided by Merck Sharp and Dohme, UK); GV150013 [*N*-(+)-[1-adamant-1-ylmethyl)-2,4-dioxo-5-phenyl-2,3,4,5-tetrahydro-1*H*-1,5-benzodiazepin-3-yl]-*N* phenylurea] (kindly provided by Glaxo-

Wellcome, Verona, Italy). CCK-8S was dissolved in physiological saline. GV150013 was dissolved in dimethyl sulphoxide and further dilutions were made with saline solution. Dimethyl sulphoxide concentration in organ bath never exceeded 0.5%.


## Results

Before performing experiments on colonic contractions, care was taken to verify the presence of CCK receptors in the experimental model. By means of the reverse transcription-polymerase chain reaction, evidence was obtained that mRNAs coding for both CCK<sub>1</sub> and CCK<sub>2</sub> receptors were expressed in preparations of human colon longitudinal muscle (data not shown).


#### Studies on CCK-8S-induced motor activity

During the equilibration period, colonic preparations developed a spontaneous motor activity which appeared low in amplitude and, in most cases, remained stable throughout the experimental period. The application of CCK-8S (0.001–1  $\mu\text{M}$ ) to the bathing fluid induced contractions of longitudinal smooth muscle (Figure 1a). The construction of concentration–response curves for the contractile responses elicited by CCK-8S provided an estimate of the maximal effect, equivalent to  $78.4 \pm 4.3\%$  of electrically-evoked contraction, at the concentration of 1  $\mu\text{M}$ , with a mean EC<sub>50</sub> value of 11.7 nM (95% CI 3–49 nM) (Figure 1b). The stimulant actions of CCK-8S on colonic contractions were not modified by tetrodotoxin (1  $\mu\text{M}$ ) (Figure 2c), but they were significantly reduced by the CCK<sub>1</sub> receptor antagonist devazepide (0.01–1  $\mu\text{M}$ ) (data not shown), consistently with the notion that CCK<sub>1</sub> receptors mediate direct effects of CCK on human colonic smooth muscle (D'Amato *et al.*, 1991).

The incubation of colonic tissues with GV150013 (0.001–0.1  $\mu\text{M}$ ) did not affect the spontaneous motor activity (data not shown). However, in the presence of GV150013, CCK-8S-evoked contractions were enhanced in a concentration-dependent fashion (Figures 2a and b). In addition, the



**Figure 1** (a) Representative tracing showing the effect of CCK-8S (0.1  $\mu\text{M}$ ) on the contractile activity of longitudinal smooth muscle isolated from distal colon. (b) Effects of increasing concentrations of CCK-8S (0.001–1  $\mu\text{M}$ ) on the motor activity of longitudinal smooth muscle. CCK-induced contractions were expressed as % of the contraction induced by transmural electrical stimulation (TES). Each point represents the mean of 6–8 experiments  $\pm$  s.e.m. (vertical bars). CCK-8S, CCK octapeptide sulphate.




**Figure 2** (a) Representative tracings showing the effects of GV150013 (0.01 μM) on contractions evoked by CCK-8S (0.1 μM). (b) Effects of increasing concentrations of GV150013 (0.001–0.1 μM) on the contractile responses elicited by CCK-8S (0.1 μM). (c) Representative trace recordings showing the effects of CCK-8S (0.1 μM), in the absence or in the presence of tetrodotoxin (TTX, 1 μM), on colonic longitudinal smooth muscle, and the effects of GV150013 (0.01 μM) on the contractile responses induced by CCK-8S in the presence of TTX. (d) Effects of TTX or TTX plus GV150013 on CCK-8S-induced contractions. Each column represents the mean of 6–8 experiments ± s.e.m. (vertical bars). \**P* < 0.05, significant difference vs CCK-8S alone. CCK-8S, CCK octapeptide sulphate; W, wash.

enhancing effect of GV150013 (0.01 μM) on the motor responses evoked by CCK-8S (0.1 μM) was unaffected when colonic preparations were incubated with tetrodotoxin (1 μM) (Figures 2c and d). In colonic tissues maintained in Krebs solution containing the non-selective NOS inhibitor L-NAME (100 μM), contractions induced by CCK-8S (0.1 μM) were significantly increased. Under these conditions, the enhancing effects of GV150013 (0.01 μM) no longer occurred. By contrast, the selective nNOS inhibitor NPA (0.1 μM) did not influence the CCK-8S-evoked contractions and, in the presence of NPA, GV150013 was still able to enhance the motor responses elicited by CCK-8S (Figure 3).

#### Studies on tissues precontracted with KCl

The exposure of colonic preparations to KCl (60 mM) induced a fast phasic contraction followed by a period of sustained contraction which, in most cases, remained stable for at least 5 min (Figure 4a). The amplitude of the sustained contractions was equivalent to 85.5 ± 6.4% of contractile responses evoked by electrical stimulation. Under these conditions, CCK-8S (0.001–1 μM) elicited concentration-dependent relaxations of smooth muscle (Figures 4a and b). Preincubation of colonic tissues with GV150013 (0.01 μM) did not alter contractions evoked by KCl (data not shown). However, the relaxant responses evoked by CCK-8S were completely prevented by GV150013 (Figure 4c). The relaxant



**Figure 3** Effects of L-NAME (100  $\mu\text{M}$ ), GV150013 (0.01  $\mu\text{M}$ ), L-NAME plus GV150013, NPA (0.1  $\mu\text{M}$ ) or NPA plus GV150013 on the contractile responses evoked by CCK-8S (0.1  $\mu\text{M}$ ) in longitudinal smooth muscle preparations isolated from human distal colon. Each column represents the mean of seven to eight experiments  $\pm$  s.e.m. (vertical bars). \* $P < 0.05$ , significant difference vs CCK-8S alone. CCK-8S, CCK octapeptide sulphate; L-NAME,  $N^G$ -nitro-L-arginine methylester; NPA,  $N^G$ -propyl-L-arginine.


effects induced by CCK-8S were counteracted by L-NAME (100  $\mu\text{M}$ ), but not by NPA (0.1  $\mu\text{M}$ ) (Figure 4c). Moreover, in the presence of NPA, the blunting effect of GV150013 on CCK-8S-induced relaxations was unaffected (Figure 4c).

#### Studies on electrically-evoked contractions

The application of electrical stimulation to colonic preparations evoked reproducible phasic contractions (Figure 5a), which were prevented by tetrodotoxin (1  $\mu\text{M}$ ) (data not shown). The contractile responses elicited by electrical stimulation were significantly enhanced by GV150013 (0.01  $\mu\text{M}$ ) (Figures 5a and b).

#### Discussion

In recent years, a number of studies have evaluated the effects of CCK receptor ligands on gastrointestinal motor functions, as these agents might represent novel pharmacological tools for the management of symptoms associated with functional digestive disorders (Varga *et al.*, 2004; Galligan and Vanner, 2005). In view of these premises, the present study was performed to characterize the role played by CCK<sub>2</sub> receptors in the control of motor activity in human distal colon. Before addressing this issue, we confirmed the findings of previous reports indicating that both CCK<sub>1</sub> and CCK<sub>2</sub> receptors are expressed in the muscular compartment of human colon (Rettenbacher and Reubi, 2001; Morton *et al.*, 2002b), and that CCK analogues are able to induce



**Figure 4** (a) Representative tracing showing the effect of CCK-8S (0.1  $\mu\text{M}$ ) on the motor activity of human colonic longitudinal smooth muscle pre-contracted with KCl (60 mm). (b) Effects of increasing concentrations of CCK-8S (0.001–1  $\mu\text{M}$ ) on KCl-induced contractions. (c) Effects of GV150013 (0.01  $\mu\text{M}$ ), L-NAME (100  $\mu\text{M}$ ), NPA (0.1  $\mu\text{M}$ ) or NPA plus GV150013 on the relaxant effects of CCK-8S in colonic preparations pre-contracted with KCl. Each column represents the mean of seven to eight experiments  $\pm$  s.e.m. (vertical bars). \* $P < 0.05$ , significant difference vs control. # $P < 0.05$ , significant difference vs CCK-8S. CCK-8S, CCK octapeptide sulphate; L-NAME,  $N^G$ -nitro-L-arginine methylester; NPA,  $N^G$ -propyl-L-arginine; W, wash.


tetrodotoxin-insensitive colonic contractions, which can be counteracted by selective CCK<sub>1</sub> receptor antagonists (D'Amato *et al.*, 1991; Morton *et al.*, 2002a).

After verification of previous available information, in the present study a series of functional experiments was


designed to assay the effects of compounds acting at CCK<sub>2</sub> receptors on the contractile activity of human colon. The results showed that CCK-8S elicited contractile responses that were enhanced by GV150013, either in the absence or in the presence of tetrodotoxin, thus suggesting that CCK<sub>2</sub> receptors mediate inhibitory effects that do not seem to involve the recruitment of neural pathways. To the best of our knowledge, this is the first evidence demonstrating inhibitory effects mediated by CCK<sub>2</sub> receptors on human colonic longitudinal smooth muscle, as previous studies have reported excitatory effects mediated by CCK<sub>1</sub> receptors in this tissue. Morton *et al.* (2002a) examined the effects of a CCK<sub>2</sub> receptor antagonist, JB93182, on human colon *in vitro*, and observed that this compound did not influence CCK-8S-induced contractions. However, Morton *et al.* (2002a) performed their experiments on preparations of circular smooth muscle, which were isolated from the ascending colon and connected to isometric force transducers. These experimental variables are not directly comparable with those adopted in our study with longitudinal smooth muscle isolated from descending colon and connected to isotonic transducers.

The present experiments showed that the enhancing effect of GV150013 on motor responses evoked by CCK-8S was still evident in the presence of NPA, but not L-NAME. These findings allow us to hypothesize that CCK<sub>2</sub> receptors mediated their inhibitory actions in the distal colon through the release of NO, which did not appear to be generated by the nNOS isoform. To clarify this mechanism further, CCK-8S was tested on colonic preparations maintained under sustained contraction with KCl. In this setting, CCK-8S elicited concentration-dependent relaxations, which were antagonized by GV150013 or L-NAME, but not by NPA, thus providing further evidence that CCK<sub>2</sub> receptors can modulate human colonic motility via NO release, independent of nNOS activity.

The existence of NO pathways regulated by CCK<sub>2</sub> receptors in the muscular compartment of human colon has not been previously demonstrated and represents a major novel finding of the present study. Consistent with this proposal, previous reports indicated that NOS is expressed in digestive



**Figure 5** (a) Representative tracings showing the effects of transmurial electrical stimulation (TES, 1 ms, 30 mA, 10 Hz, 100 pulses) on the motor activity of human colonic longitudinal smooth muscle, either alone or in the presence of GV150013 (0.01 μM). (b) Effects of GV150013 (0.01 μM) on the contractile responses induced by TES. Each column represents the mean of eight experiments ± s.e.m. (vertical bars). \**P* < 0.05, significant difference vs control.



**Figure 6** Schematic diagram summarizing the proposed modulating actions exerted by the CCK<sub>2</sub> receptor subtype on the motor activity of human colonic longitudinal smooth muscle. CCK<sub>2</sub> receptors are localized on ICC and are activated by neurally released CCK. CCK<sub>2</sub> receptors then stimulate NOS activity, with subsequent NO release and relaxation of smooth muscle cells. CCK, cholecystokinin; ICC, interstitial cell of Cajal; LSMC, longitudinal smooth muscle cell; MN, myenteric neuron; NO, nitric oxide; NOS, nitric oxide synthase.

smooth muscle cells of rabbits and humans and that smooth muscle represents a major source of NO at enteric level (Jin *et al.*, 1996; Teng *et al.*, 1998). Other investigations demonstrated that NOS is expressed in interstitial cells of Cajal (ICC) of mouse colon (Vannucchi *et al.*, 2002), and there is evidence to suggest that ICC contribute significantly to the NO-dependent modulation of canine colonic motility (Publicover *et al.*, 1993). A more recent study indicated also that the administration of L-NAME to normal rats caused an increase in colonic motility, whereas in transgenic rats lacking ICC this effect no longer occurred, suggesting that NO is involved in the regulatory actions exerted by ICC on colonic motility (Takahashi *et al.*, 2005). These observations, together with the fact that CCK<sub>2</sub> receptors have been found expressed mostly in ICC of human colon (Rettenbacher and Reubi, 2001), support the view that the inhibitory actions mediated by CCK<sub>2</sub> receptors on colonic motility, as shown in the present study, might be driven by NO release from enteric cells, which are likely to correspond to ICC (Figure 6).

An important point to consider is the cellular source(s) of CCK-related peptides responsible for the recruitment of CCK receptor pathways. Indeed, at gastrointestinal level CCK can be produced and released by endocrine I cells in the duodenal wall or by intramural peptidergic neurons located within myenteric plexus of small intestine and colon (Liddle, 1997; Rehfeld, 2004; Varga *et al.*, 2004). In the present study, an attempt was made to verify whether CCK<sub>2</sub> receptors of distal colon can be activated by CCK of neuronal origin. For this purpose, a subset of experiments was performed on colonic preparations subjected to electrical stimulation to elicit contractile events mediated by the release of neural transmitters. Since, in our hands, electrical stimulation evoked tetrodotoxin-sensitive contractile responses, which were enhanced by GV150013, it can be suggested that CCK<sub>2</sub> receptors involved in the inhibitory control of colonic motility are likely to interact mainly with CCK-like peptides released from enteric nerves (Figure 6).

In conclusion, the present results indicate that CCK<sub>2</sub> receptors mediate inhibitory actions of CCK on motor functions of human distal colon. In particular, CCK seems to act via CCK<sub>2</sub> receptors to induce relaxant responses mediated by recruitment of NO-dependent pathways. In view of these findings, the isolated human distal colon can be regarded as a suitable model to assay the intestinal motor effects of novel drugs designed to interact with CCK<sub>2</sub> receptors.

## Acknowledgements

We thank Dr Massimo D'Amato (Rotta Research Laboratorio S.p.A., Monza, Italy) for his expert and critical review of the manuscript.

## Conflict of interest

The authors state no conflict of interest

## References

- Chey WY, Jin HO, Lee MH, Sun SW, Lee KY (2001). Colonic motility abnormalities in patients with irritable bowel syndrome exhibiting abdominal pain and diarrhea. *Am J Gastroenterol* **96**: 1499–1506.
- Chua ASB, Keeling PWN (2006). Cholecystokinin hyperresponsiveness in functional dyspepsia. *World J Gastroenterol* **12**: 2688–2693.
- Crawley JN, Corwin RL (1994). Biological actions of cholecystokinin. *Peptides* **15**: 731–755.
- Dal Forno G, Pietra C, Urciuoli M, Van Amsterdam FT, Toson G, Gaviraghi G *et al.* (1992). Evidence for two cholecystokinin receptors mediating the contraction of the guinea pig isolated ileum longitudinal muscle myenteric plexus. *J Pharmacol Exp Ther* **261**: 1056–1063.
- D'Amato M, Stanford IF, Bennett A (1991). Studies of three non-peptide cholecystokinin antagonists (devazepide, lorglumide and loxiglumide) in human isolated alimentary muscle and guinea-pig ileum. *Br J Pharmacol* **102**: 391–395.
- Fornai M, Blandizzi C, Colucci R, Antonioli L, Bernardini N, Segnani C *et al.* (2005). Role of cyclooxygenase 1 and 2 in the modulation of neuromuscular functions in the distal colon of humans and mice. *Gut* **54**: 608–616.
- Fornai M, Colucci R, Antonioli L, Baschiera F, Ghisu N, Tuccori M *et al.* (2007). CCK<sub>2</sub> receptors mediate inhibitory effects of cholecystokinin on the motor activity of guinea-pig distal colon. *Eur J Pharmacol* **557**: 212–220.
- Galligan JJ, Vanner S (2005). Basic and clinical pharmacology of new motility promoting agents. *Neurogastroenterol Motil* **17**: 643–653.
- Giralt M, Vergara P (1999). Both afferent and efferent nerves are implicated in cholecystokinin motor actions in the small intestine of the rat. *Regul Pept* **81**: 73–80.
- Herranz R (2003). Cholecystokinin antagonists: pharmacological and therapeutic potential. *Med Res Rev* **23**: 559–605.
- Jin JG, Murthy KS, Grider JR, Makhlof GM (1996). Stoichiometry of neurally induced VIP release, NO formation, and relaxation in rabbit gastric muscle. *Am J Physiol* **34**: G357–G369.
- Kulaksiz H, Arnold R, Goke B, Maronde E, Meyer M, Fahrenholz F *et al.* (2000). Expression and cell-specific localization of the cholecystokinin B/gastrin receptor in the human stomach. *Cell Tissue Res* **299**: 289–298.
- Liddle RA (1997). Cholecystokinin cells. *Annu Rev Physiol* **59**: 221–242.
- Miyasaka K, Funakoshi A (2003). Cholecystokinin and cholecystokinin receptors. *J Gastroenterol* **38**: 1–13.
- Morton MF, Harper EA, Tavares IA, Shankley NP (2002b). Pharmacological evidence for putative CCK(1) receptor heterogeneity in human colon smooth muscle. *Br J Pharmacol* **136**: 873–882.
- Morton MF, Welsh NJ, Tavares IA, Shankley NP (2002a). Pharmacological characterization of cholecystokinin receptors mediating contraction of human gallbladder and ascending colon. *Regul Pept* **105**: 59–64.
- Noble F, Wank SA, Crawley JN, Bradwejn J, Seroogy KB, Hamon M *et al.* (1999). International Union of Pharmacology. XXI. Structure, distribution, and functions of cholecystokinin receptors. *Pharmacol Rev* **51**: 745–781.
- Ochi Y, Horie S, Maruyama T, Watanabe K, Yano S (2005). Necessity of intracellular cyclic AMP in inducing gastric acid secretion via muscarinic M3 and cholecystokinin2 receptors on parietal cells in isolated mouse stomach. *Life Sci* **77**: 2040–2050.
- Publicover NG, Hammond EM, Sanders KM (1993). Amplification of nitric oxide signalling by interstitial cells isolated from canine colon. *Proc Natl Acad Sci USA* **90**: 2087–2091.
- Rehfeld JF (2004). Clinical endocrinology and metabolism. Cholecystokinin. *Best Pract Res Clin Endocrinol Metab* **18**: 569–586.
- Rettenbacher M, Reubi JC (2001). Localization and characterization of neuropeptide receptors in human colon. *Naunyn Schmiedeberg Arch Pharmacol* **364**: 291–304.
- Sjolund K, Ekman R, Lindgren S, Eehfeld JF (1996). Disturbed motilin and cholecystokinin release in irritable bowel syndrome. *Scand J Gastroenterol* **31**: 1110–1114.
- Sternini C, Wong H, Pham T, De Giorgio R, Miller IJ, Kunts SM *et al.* (1999). Expression of cholecystokinin A receptors in neurons

- innervating the rat stomach and intestine. *Gastroenterology* **117**: 1136–1146.
- Takahashi A, Takeshi T, Kaneko H, Hatori R, Ishige T, Suzuki M *et al.* (2005). *In vivo* recording of colonic motility in conscious rats with deficiency of interstitial cells of Cajal, with special reference to the effects of nitric oxide on colonic motility. *J Gastroenterol* **40**: 1043–1048.
- Teng BQ, Murthy KS, Kuemmerle JF, Grider JR, Sase K, Michel T *et al.* (1998). Expression of endothelial nitric oxide synthase in human and rabbit gastrointestinal smooth muscle cells. *Am J Physiol* **38**: G342–G351.
- Vannucchi MG, Corsani L, Bani D, Fausone-Pellegrini MS (2002). Myenteric neurons and interstitial cells of Cajal of mouse colon express several nitric oxide synthase isoforms. *Neurosci Lett* **326**: 191–195.
- Varga G, Balint A, Burghardt B, D'Amato M (2004). Involvement of endogenous CCK and CCK-1 receptors in colonic motor function. *Br J Pharmacol* **141**: 1275–1284.
- Vergara P, Woskowska Z, Cipris S, Fox-Threlkeld JE, Daniel EE (1996). Mechanisms of action of cholecystokinin in the canine gastrointestinal tract: role of vasoactive intestinal peptide and nitric oxide. *J Pharmacol Exp Ther* **279**: 306–316.
- Waldum HL, Kleveland PM, Sandvik AK, Brenna E, Syversen U, Bakke I *et al.* (2002). The cellular localization of the cholecystokinin 2 (gastrin) receptor in the stomach. *Pharmacol Toxicol* **91**: 359–362.