

SHORT REPORT

Mapping of a new autosomal dominant non-syndromic hearing loss locus (DFNA43) to chromosome 2p12

E Flex, M Mangino, M Mazzoli, A Martini, V Migliosi, A Colosimo, R Mingarelli, A Pizzuti, B Dallapiccola

J Med Genet 2003;**40**:278–282

Hearing impairment (HI) is the most frequent sensory defect with wide genetic heterogeneity. Approximately 80% of genetic hearing loss is non-syndromic and 15-25% of exhibit autosomal dominant inheritance. We analysed an Italian three generation family in which non-syndromic hearing impairment is transmitted as an autosomal dominant trait. Onset of HI in all affected subjects occurred in the second decade of life, with subsequent gradual progression from moderate to profound loss. HI was bilateral and symmetrical, involving all frequencies. After exclusion of the known DFNA loci with markers from the Hereditary Hearing Loss Homepage (URL: <http://dnalab-www.uia.ac.be/dnalab/hhh>), a genome wide scan was carried out using 358 highly informative microsatellite markers. Significant linkage ($Z_{\max}=4.21$, $\theta=0$) was obtained with chromosome 2p12 markers. The results were confirmed by multipoint analysis ($Z_{\max}=4.51$), using the location score method. Haplotype analysis defined a 9.6 cM disease gene interval on chromosome 2 without overlap with the other identified loci. Fine mapping and identification of candidate genes are in progress.

Hearing loss is a common form of sensory impairment, affecting millions of people world wide.¹ Deafness can be the result of genetic or environmental causes or a combination of them. Approximately 1 out 1000 infants is affected by either severe or profound deafness at birth and during early childhood, that is, in the prelingual period. About 60% of these cases are considered to have a genetic basis, with deafness mostly occurring as a non-syndromic defect.² Non-syndromic hearing loss is characterised by extensive genetic heterogeneity. More than 70 loci have been mapped³ and 29 genes identified so far.

We have performed genetic linkage studies in a three generation Italian family segregating an autosomal dominant form of non-syndromic hearing impairment (NSHI). A genome wide scan localised a novel NSHI locus (DFNA 43) to chromosome 2p12.

MATERIALS AND METHODS

Subjects

We examined a three generation Italian family with an autosomal dominant form of hearing loss (fig 1). Informed consent was obtained from all the enrolled patients and from




Figure 1 Haplotypes of nine DNA markers on chromosome 2p12 in the branch of the family examined. The blackened bars indicate the haplotypic region in affected subjects that define the critical DFNA43 region between flanking markers D2S2114 and D2S2333.




Figure 2 Audiograms of affected subjects from three generations (■ III.7, 24 years old; ◆ III.1, 35 years old; ▲ II.7, 54 years old; ▼ II.14, 74 years old; ◇ I.2, 96 years old), left ear (A) and right ear (B).

parents of patients younger than 18 years. Physical examination was performed on all the family members. An additional clinical history interview was conducted in order to identify potential environmental causes of hearing loss, such as infections, trauma, and exposure to ototoxic drugs. Tests to detect the presence and type of hearing loss were conducted. Air conduction thresholds were measured at 250 Hz, 500 Hz, 1 kHz, 2 kHz, 4 kHz, 6 kHz, and 8 kHz. Bone conduction thresholds were determined to ascertain whether there was any evidence for a conductive component in patients with hearing loss. Oto-emittance measurements were obtained for all subjects that were otoscopically examined.

Genotyping

Genome scanning was performed using 358 microsatellite markers at a distance of ~10 cM (ABI PRISM™ Linkage Mapping Set). PCR was performed using 50 ng of DNA in a 15 µl reaction mixture containing 1.5 µl buffer (100 mmol/l Tris HCl, pH 8.3, 500 mmol/l KCl), 1.5 µl MgCl₂ (25 mmol/l), 1.5 µl dNTPs mix (2.5 mmol/l), 1 µl primer mix (5 µmol/l), and 0.6 U of AmpliTaq Gold™. PCR products were analysed on a model 3100 automated fluorescent DNA sequencer, a four colour detection system. One µl of PCR reaction was combined with 10 µl of formamide and 0.5 µl of a fluorescent size marker (ROX GS-500). Each sample was run for 30 minutes. During the electrophoresis, fluorescence detected in the laser scanning region was collected and stored using Genescan Collection software (version 1.0.1). The fluorescent data collected during the run were analysed by the Genescan Analysis program (version 3.5). Each marker was examined by the Genotyper program (version 3.7) in order to analyse inheritance patterns and prepare the allele labels for export to linkage applications.

Linkage analysis

Linkage analysis was performed by the LINKAGE 5.1 program package.⁴ Two point lod scores between the disease gene and each marker were calculated by means of the MLINK program. The phenotype was coded as a fully penetrant autosomal dominant trait, with a disease allele frequency of 0.0001. Equal recombination frequencies for male and females were assumed. The order of the marker loci and the recombination distances used for multipoint linkage analysis were based on the Généthon linkage map.⁵⁻⁷ Multipoint analysis was performed with the VITESSE computer program.⁸

Candidate gene analysis

Sequence analysis of the whole cadherin associated protein (α -catenin) (*CTNNA2*, MIM 114025) coding region was performed. PCR primers and annealing conditions for *CTNNA2* exon PCRs are available on request. PCR products were purified using the Microcon 100 system (Millipore) and sequenced using a 3100 automated fluorescent DNA sequencer and the ABI Prism BigDye Terminator cycle sequencing kit. Electropherograms were analysed by the Sequencing Analysis software (version 3.7). We also analysed the intragenic marker D2S219. One primer of the microsatellite was labelled with the 6-FAM fluorochrome, which was analysed using standard conditions on a 3100 automated fluorescent DNA sequencer.


RESULTS

Subjects

We analysed an Italian three generation family in which NSHI was transmitted as an autosomal dominant trait (fig 1), characterised by bilateral symmetrical sensorineural hearing impairment starting during the second/third decade of life as

Table 1 Lod scores

Markers	Lod score at $\theta =$						
	0.0	0.01	0.05	0.1	0.2	0.3	0.4
D2S109	−∞	−3.83	−1.23	−0.29	0.36	0.45	0.30
D2S2114	−∞	2.15	2.60	2.57	2.16	1.52	0.74
D2S2116	2.41	2.37	2.21	1.99	1.53	1.01	0.42
D2S139	4.21	4.45	4.16	3.78	2.96	2.04	0.98
D2S2180	2.71	2.67	2.49	2.25	1.74	1.16	0.50
D2S2333	−∞	2.15	2.60	2.57	2.16	1.52	0.7
D2S2216	−∞	2.15	2.60	2.57	2.16	1.52	0.74
D2S160	−∞	−4.13	−1.51	−0.54	0.16	0.31	0.22

**Figure 3** Multipoint lod score analysis for the region 2p12. The maximum multipoint lod score within the interval of ~9.6 cM between markers D2S2114 and D11S2333 was 4.52.

a dip at 4000 Hz, similar to a noise induced hearing loss. The hearing impairment was slowly progressive, involving at first the high frequencies and extending to all frequencies by the fifth to sixth decade of life. The level of impairment became severe at later ages. Tympanometry and stapedia reflexes were normal. The hearing impairment was cochleopathic, with presence of recruitment and with tinnitus as a variable manifestation. In comparison with other genetic non-syndromic conditions, there was little intrafamilial variability and differences could be mainly attributed to the age related progression (fig 2).

Linkage analysis

Known loci for autosomal dominant hearing loss were excluded using the reference markers described on the Hereditary Hearing Loss Homepage (URL: <http://dnalab-www.uia.ac.be/dnalab/hhh>). A genome search was then undertaken in 19 members of the family using microsatellite markers at intervals of ~10 cM. A maximum lod score of 4.21 ($\theta=0$) for marker D2S139 (table 1) located the disease gene on the short arm of chromosome 2. Haplotype analysis disclosed a key recombination event between D2S2114 and D2S2116 in II.4, defining the telomeric boundary of the disease interval (fig 1). The centromeric limit is determined by a crossover between D2S2180 and D2S2333, observed in subject II.13 (fig 1). These recombinations place the disease candidate region between markers D2S2114 and D2S2333 (multipoint lod score of 4.52, fig 3). The interval spans 9.6 cM on the Génethon genetic map, without overlap with other NSHI loci previously identified.

Candidate gene analysis

A genome database survey of the candidate region showed the presence of several genes. Among them is the cadherin associated protein (α -catenin) gene (*CTNNA2*, MIM 114025), which is expressed by the cochlea cells (Morton Fetal Cochlea, dbEST Library: ID.371).⁹ We first analysed the intragenic marker D2S2191. As shown in fig 1, three affected subjects were heterozygous. The results excluded the presence of large dele-

tions in *CTNNA2*. Then, the gene coding regions were sequenced without finding any variation.

DISCUSSION

We performed a genome search in a three generation Italian family, in which a non-syndromic hearing impairment was transmitted as an autosomal dominant trait. We found linkage with markers located on chromosome 2p12. The DFNA43 critical region does not overlap with any other locus for non-syndromic hearing loss previously mapped on chromosome 2 (DFNA16, DFNB9, DFNB27).¹⁰⁻¹² In an effort to identify the DFNA43 gene, we tested the cadherin associated protein (α -catenin) gene (*CTNNA2*). Several pieces of evidence pointed to *CTNNA2* as a DFNA43 candidate gene. Nonsense and frameshift mutations in the cadherin-like protein CDH23 have been identified in Usher syndrome 1D patients (USH1D, MIM 601067),^{13 14} while missense mutations can cause severe to profound hearing loss (DFNB12).¹⁵ Moreover, mutations in two cadherin related genes were reported in the deaf mouse mutants *waltzer*¹⁶ and *Ames-waltzer*.¹⁷ Cadherins are thought to be involved in cell-cell interactions within the inner ear, but the presence of disorganised stereocilia in *waltzer* and *Ames-waltzer* mutants also suggests involvement in the lateral or tip links joining adjacent stereocilia.¹⁶ Proteins such as vezatin or harmonin could also have a role in these linkage complexes.³ The activity of cadherins in mediating homophilic cell-cell Ca^{2+} dependent association would rely on their anchorage to the cytoskeleton via the catenins.^{18 19} However, mutation screening of *CTNNA2* gene coding regions in DFNA43 patients was negative, apparently excluding it as a candidate. To date, no other gene mapping in the region appears a suitable candidate for this disease. The analysis of new families linked to the DFNA43 locus, as well as a more complete expression map of the region will help the task of isolating the DFNA43 gene.

ACKNOWLEDGEMENTS

The first two authors contributed equally to this work.

Authors' affiliations

E Flex, M Mangino, V Migliosi, A Colosimo, R Mingarelli, A Pizzuti, B Dallapiccola, IRCCS, S Giovanni Rotondo CCS-Mendel Institute, Rome, Italy

E Flex, A Pizzuti, B Dallapiccola, Department of Experimental Medicine and Pathology, University of Rome La Sapienza, Rome, Italy

M Mazzoli, UOA Otorinolaringoiatria, Ospedaliera di Padova, Padova, Italy

A Martini, Department of Audiology, University of Ferrara, Ferrara, Italy
A Colosimo, Department of Biomedical Science, University of Chieti, Chieti, Italy

Correspondence to: Dr M Mangino, IRCCS, CCS-Mendel Institute, Rome, Italy; m.mangino@css-mendel.it

Revised version received 19 December 2002
Accepted for publication 23 December 2002

REFERENCES

- Cohen MM**, Gorlin RJ. Epidemiology, etiology and genetic patterns. In: Gorlin RJ, Toriello HV, Cohen MM, eds. *Hereditary hearing loss and its syndromes*. Oxford: Oxford University Press, 1995:9-21.
- Van Camp G**, Willems PJ, Smith RJ. Nonsyndromic hearing impairment: unparalleled heterogeneity. *Am J Hum Genet* 1997;**60**:758-64.
- Steel KP**, Kros CJ. A genetic approach to understanding auditory function. *Nat Genet* 2001;**27**:143-9.
- Lathrop GM**, Lalouel JM. Easy calculations of lod scores and genetic risks on small computers. *Am J Hum Genet* 1984;**36**:460-5.
- Dib C**, Faure S, Fizames C, Samson D, Drouot N, Vignal A, Millasseau P, Marc S, Hazan J, Seboun E, Lathrop M, Gyapay G, Morissette J, Weissenbach J. A comprehensive genetic map of the human genome based on 5,264 microsatellites. *Nature* 1996;**380**:152-4.
- Gyapay G**, Morissette J, Vignal A, Dib C, Fizames C, Millasseau P, Marc S, Bernardi G, Lathrop M, Weissenbach J. The 1993-94 Genethon human genetic linkage map. *Nat Genet* 1994;**7**:246-339.
- Weissenbach J**, Gyapay G, Dib C, Vignal A, Morissette J, Millasseau P, Vaysseix G, Lathrop M. A second-generation linkage map of the human genome. *Nature* 1992;**359**:794-801.
- O'Connell JR**, Weeks DE. The VITESSE algorithm for rapid exact multilocus linkage analysis via genotype set-recoding and fuzzy inheritance. *Nat Genet* 1995;**11**:402-8.
- Robertson NG**, Khetarpal U, Gutierrez-Espeleta GA, Bieber FR, Morton CC. Isolation of novel and known genes from a human fetal cochlear cDNA library using subtractive hybridization and differential screening. *Genomics* 1994;**23**:42-50.
- Pulley LJ**, Jackson AP, Roberts E, Carridice A, Muxworthy C, Houseman M, Al-Gazali L, Lench NJ, Markham AF, Mueller RF. A new locus for autosomal recessive non-syndromal sensorineural hearing impairment (DFNB27) on chromosome 2q23-q31. *Eur J Hum Genet* 2000;**8**:991-3.
- Fukushima K**, Kasai N, Ueki Y, Nishizaki K, Sugata K, Hirakawa S, Masuda A, Gunduz M, Ninomiya Y, Masuda Y, Sato M, McGuirt WT, Coucke P, Van Camp G, Smith RJ. A gene for fluctuating, progressive autosomal dominant nonsyndromic hearing loss, DFNA16, maps to chromosome 2q23-24.3. *Am J Hum Genet* 1999;**65**:141-50.
- Chaib H**, Place C, Salem N, Chardenoux S, Vincent C, Weissenbach J, El-Zir E, Loiselet J, Petit C. A gene responsible for a sensorineural nonsyndromic recessive deafness maps to chromosome 2p22-23. *Hum Mol Genet* 1996;**5**:155-8.
- Bolz H**, von Brederlow B, Ramirez A, Bryda EC, Kutsche K, Nothwang HG, Seeliger M, del CSCM, Vila MC, Molina OP, Gal A, Kubisch C. Mutation of CDH23, encoding a new member of the cadherin gene family, causes Usher syndrome type 1D. *Nat Genet* 2001;**27**:108-12.
- Petit C**. Usher syndrome: from genetics to pathogenesis. *Annu Rev Genomics Hum Genet* 2001;**2**:271-97.
- Bork JM**, Peters LM, Riazuddin S, Bernstein SL, Ahmed ZM, Ness SL, Polomeno R, Ramesh A, Schloss M, Srisailpathy CR, Wayne S, Bellman S, Desmukh D, Ahmed Z, Khan SN, Kaloustian VM, Li XC, Lalwani A, Riazuddin S, Bitner-Glindzicz M, Nance WE, Liu XZ, Wistow G, Smith RJ, Griffith AJ, Wilcox ER, Friedman TB, Morell RJ. Usher syndrome 1D and nonsyndromic autosomal recessive deafness DFNB12 are caused by allelic mutations of the novel cadherin-like gene CDH23. *Am J Hum Genet* 2001;**68**:26-37.
- Di Palma F**, Holme RH, Bryda EC, Belyantseva IA, Pellegrino R, Kachar B, Steel KP, Noben-Trauth K. Mutations in Cdh23, encoding a new type of cadherin, cause stereocilia disorganization in waltzer, the mouse model for Usher syndrome type 1D. *Nat Genet* 2001;**27**:103-7.
- Alagramam KN**, Murcia CL, Kwon HY, Pawlowski KS, Wright CG, Woychik RP. The mouse Ames waltzer hearing-loss mutant is caused by mutation of Pcdh15, a novel protocadherin gene. *Nat Genet* 2001;**27**:99-102.
- Kemler R**, Ozawa M. Uvomorulin-catenin complex: cytoplasmic anchorage of a Ca²⁺-dependent cell adhesion molecule. *Bioessays* 1989;**11**:88-91.
- Ozawa M**, Baribault H, Kemler R. The cytoplasmic domain of the cell adhesion molecule uvomorulin associates with three independent proteins structurally related in different species. *EMBO J* 1989;**8**:1711-17.



Want to know more?

Data supplements

Limited space in printed journals means that interesting data and other material are often edited out of articles; however, limitless cyberspace means that we can include this information online. Look out for additional tables, references, illustrations.

www.jmedgenet.com



Mapping of a new autosomal dominant non-syndromic hearing loss locus (DFNA43) to chromosome 2p12

E Flex, M Mangino, M Mazzoli, A Martini, V Migliosi, A Colosimo, R Mingarelli, A Pizzuti and B Dallapiccola

J Med Genet 2003 40: 278-281
doi: 10.1136/jmg.40.4.278

Updated information and services can be found at:
<http://jmg.bmj.com/content/40/4/278>

	<i>These include:</i>
References	This article cites 18 articles, 1 of which you can access for free at: http://jmg.bmj.com/content/40/4/278#BIBL
Email alerting service	Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.
Topic Collections	Articles on similar topics can be found in the following collections Molecular genetics (1244)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>