

Short Communication

Expression of genes encoding protein kinases during flower opening in two cut rose cultivars with different longevity

Hanife Hajizadeh^{1,2,3}, Khadije Razavi^{1*}, Younes Mostofi^{2*}, Giovanni Cacco³, Amir Mousavi¹, Zabihollah Zamani², Piergiorgio Stevanato³

¹National Institute of Genetic Engineering and Biotechnology, Tehran-Karaj Highway, P.O. Box 14965/161, Tehran, I.R. Iran ²Department of Horticultural Sciences, University of Tehran, P.O. Box 4111, Karaj, I.R. Iran ³Department of Agricultural Biotechnology, University of Padova, Viale dell'Università 16, 35020 Legnaro (PD), Italy

Abstract

Ethylene plays an important role in wide-ranging aspects of plant growth and development, including fruit ripening, leaf and flower senescence. In this study, the expression patterns of two genes involved in the ethylene signal transduction pathway (*RhCTR1* and *RhCTR2*) were investigated during the flower opening stages in two *Rosa hybrida* cultivars, 'Black magic' and 'Maroussia', which are characterized by short and long vase lives, respectively. *RhCTR1* expression increased significantly during flower opening in both cultivars, but its expression level in cv. Maroussia was significantly higher than that in cv. Black magic. No variation in gene expression was detected for *RhCTR2* in both cultivars. Therefore, this study showed that the vase life of the two cultivars correlated with the expression of *RhCTR1*, but not with that of *RhCTR2*, the behavior of which is typical of a constitutive gene.

Keywords: Cut rose (*Rosa hybrida*); Ethylene sensitivity; Gene expression; *RhCTR1* and *RhCTR2*

Rose is one of the most important cut-flowers in ornamental plants. For the purpose of commercial produc-

tion, cut roses are usually harvested at a specific bud stage. Depending on the cultivar and the handling conditions, the flowers often do not open (Tan *et al.*, 2006). The gaseous hormone ethylene plays an important role in diverse aspects of plant growth and development, ranging from seed germination (Abeles *et al.*, 1992) to senescence and abscission of flowers, fruits and leaves (Giovannoni, 2001; Johnson and Ecker 1998). Also, in ornamental plants, ethylene influences flower opening (Reid *et al.*, 1989), petal senescence (Nukui *et al.*, 2004; Shibuya *et al.*, 2000) and abscission (Kuroda *et al.*, 2003). Current evidence indicates that ethylene is sensed by a small family of receptors, the Ethylene Responsive Elements (ETRs), which are able to form a complex with a kinase, namely *CTR1*. In the absence of ethylene, this complex actively represses the ethylene response pathways. The receptor/*CTR1* complex appears to operate by negatively regulating *EIN2* which is thought to activate response pathways through the *EIN3* family of transcriptional regulators (Bleeker, 2001). The loss-of-function *CTR1* mutants result in a constitutive ethylene-response phenotype, indicating that *CTR1* negatively regulates the ethylene response pathway (Chang and Stadler, 2001; Kieber *et al.*, 1993). Therefore, *CTR1* is a key gene in the ethylene signal transduction pathway. In tomato, unlike *Arabidopsis thaliana*, which only contains one constitutively expressed *AtCTR1* gene, a family of four

*Correspondence to: Khadije Razavi and Younes Mostofi, Ph.D.
Tel/Fax: +98 261 2248721
E-mail: razavi@nigeb.ac.ir; ymostofi@ut.ac.ir

CTR1-like genes (*LeCTR1-LeCTR4*) has been identified (Lin *et al.*, 1998), and expression analysis has revealed that the expression of *LeCTR1* increases during fruit ripening and flower senescence (Leclercq *et al.*, 2002; Zegzouti *et al.*, 1999), whereas *LeCTR2* is expressed constitutively (Alexander and Grierson, 2002). A homologue of *CTR1* has been cloned and characterized in *Cucurbita pepo*, and during flower development, *Cup-CTR1* is up-regulated in male flowers but not in female flowers (Manzano *et al.*, 2008).

The induction of petal senescence or abscission by ethylene or pollination is associated with transcriptional regulation of the *ACS* and *ACO* (Fernández-Otero *et al.*, 2006; Jones, 2003) and ethylene receptor genes (such as *DC-ERS1*, *DC-ERS2* and *DC-ETR1* in *Dianthus caryophyllus* L. and *Dl-ERS1* type-1 and *Dl-ERS1* type-2 in *Delphinium*) (Kuroda *et al.*, 2003; Shibuya *et al.*, 2002). Besides, this induction is also accompanied with an increase in the *CTR* genes of some ornamental plant species (Kuroda *et al.*, 2004; Müller *et al.*, 2002). Flower opening and senescence in roses is sensitive to ethylene, although the degree of this sensitivity varies in different cultivars (Müller *et al.*, 2001; Reid *et al.*, 1989). In miniature potted roses, this difference could be due to the different expression levels of receptor genes rather than the ethylene biosynthetic genes (Müller *et al.*, 2000 a,b). The aim of this study was to evaluate the expression pattern of two critical genes (*RhCTR1* and *RhCTR2*) in the ethylene signal transduction pathway for a better understanding of the principles of gene regulation during rose flower senescence.

Rosa hybrida cv. Maroussia (long vase life) and cv. Black magic (short vase life) were grown in a hydroponic greenhouse. Petal samples were collected at 2 stages of rose flower opening (stage 2: partially opened bud; stage 8: fully-opened flower), as described by Wang *et al.* (2004). Petal samples were immediately detached and frozen in liquid nitrogen and stored at -80°C for eventual total RNA isolation. Rose petals from each stage (Fig. 1) were grounded by means of the high-speed mixer mill (MM301, Retsch, Haan, Germany). High-quality total RNA was success-

fully isolated from 100 µg of petal tissue using the RNeasy Plant Mini Kit (Qiagen, Valencia, CA, USA), according to the manufacturer's protocol. Total RNA was quantified by spectrophotometric analysis using a Biophotometer (Eppendorf, Germany).

First-strand cDNA was synthesized from 5 µg of total RNA using the ProtoScript® First Strand cDNA Synthesis Kit (New England Biolabs, UK) following the manufacturer's instructions. cDNA was quantified using the Biophotometer.

Real-time PCR was carried out by the Applied Biosystems 7500 Fast Real-Time PCR system using the DYNAmo HS SYBR Green qPCR kit (Finnzymes, Finland), in accordance with the recommendations of the manufacturer. Reaction mixtures (30 µl) contained 15 µl of 2 × SYBR-Green mixed with 0.5 µl of Rox solution, 1 µl each of 10 µM forward and 10 µM reverse primers, 5 µl of cDNA and 7.5 µl of distilled water. Primers' sequences, as reported by Müller *et al.* (2000 a,b), are presented in Table 1. The thermal profile used consisted of 2 min at 50°C, 10 min at 95°C and 40 repeats of denaturation at 95°C for 15 s and annealing-extension-fluorescence data acquisition at 60°C for 1 min. In a 96-well plate, each reaction was performed in triplicate. An endogenous 18S rRNA was used as an internal standard. Relative expression levels were calculated using the delta (Δ) threshold cycle (C_t) method (Applied Biosystems). Data from quantitative PCR were subjected to ANOVA using Statistica8.0 package (StatSoft Inc., Tulsa, OK, USA). A randomized complete block design with three independent biological replications was used. 40- delta ΔC_t was taken as an dependent variable. Means were separated using the least significant difference test (LSD) at the 0.01 level of probability.

A significant increase ($P < 0.01$) was observed for the expression level of *RhCTR1* between the two flower opening stages, in both rose cultivars (Fig. 1 and Table 2). Analogous results were observed in tomato during fruit ripening and flower senescence (Leclercq *et al.*, 2002; Zegzouti *et al.*, 1999), and in the miniature potted roses (Müller *et al.*, 2002) and two cut rose cultivars (Tan *et al.*, 2006) during flower

Table 1. Primer sequences of *CTR1*, *CTR2* and 18S rRNA.

Gene	Forward primer	Reverse primer
<i>CTR1</i>	5'- GAT GGC GCC AGA AGT CC-3'	5'- GCC CAG CAA GCC TCA AT-3'
<i>CTR2</i>	5'- GTC GCG CTT GAA ACA TAA CA -3'	5'- AAC AGG GGG ATC AAC TTC TTT-3'
18S rRNA	5'- CGG GGA GGT AGT GAC AAT AAA TAA CA-3'	5'- CCA CCA CCC ATA GAA TCA AGA AAG AG-3'




Figure 1. Expression level of *RhCTR* genes in two cut roses cv. Black magic and cv. Maroussia between two stages of flower opening. The vertical bar is the least significant difference (LSD; $P < 0.01$), for measuring significant difference between cultivars.

senescence. The expression level of *RhCTR1* was higher in cv. Maroussia when compared to that in cv. Black magic, but the increase in expression of this gene, through stages 2 and 8, was greater in cv. Black magic in comparison to cv. Maroussia (Fig. 1 and Table 2). This suggests that the vase life might be reduced by increasing the level of expression of *RhCTR1* during the different flower opening stages in spite of its expression level. Further studies will be needed to confirm this hypothesis.

Results also suggest that cv. Maroussia may not be sensitive to ethylene according to previous experiments (unpublished data), since the standard model for the ethylene signal transduction pathway confirms that a high level of *CTR* expression should result in decreased ethylene sensitivity (Tan *et al.*, 2006). These findings are consistent with other studies in *Arabidopsis thaliana* (Hua and Meyerowitz, 1998 a), tomato (Tieman *et al.*, 2000), and carnation (Shibuya *et al.*, 2002), supporting the role of ethylene receptors in the negative regulation model (Chen *et al.*, 2005; Hua *et al.*, 1998 b).

The comparison of the expression level of *RhCTR2* between the two cultivars revealed no significant difference, as shown in Figure 1 and Table 2. No significant difference was also found for *RhCTR2* expression between the two stages of the flower opening stages. Furthermore, results showed that expression levels of

Table 2. Regression slope and percentage variation of *RhCTR* gene expression between two stages of flower opening in two cut roses cv. Black magic and cv. Maroussia.

Gene	Cultivar	Slope	Percentage	P-value
<i>RhCTR1</i>	Black magic	4.8	17.3%	$P < 0.01$
	Maroussia	2.0	5.7%	$P < 0.05$
<i>RhCTR2</i>	Black magic	-0.4	-1.3%	ns
	Maroussia	-0.2	-0.4%	ns

RhCTR2 were lower than that of *RhCTR1*. Therefore, it can be hypothesized that the behavior of this gene is similar to that of a constitutive gene as previously observed in other studies on two cut rose cultivars (Tan *et al.*, 2006) and miniature potted roses (Müller *et al.*, 2002).

Acknowledgments

This study was supported by the National Institute of Genetic Engineering and Biotechnology (NIGEB) grant No. 314 and a grant from the University of Tehran and the Ministry of Science, Research and Technology of Iran. The authors wish to thank Dr. Alessandro Paparella for admitting us to use the real-time RT-PCR assays and Prof. Livio Trainotti for supporting us to clone the ESTs in Padua, Italy.

References

- Abeles FB, Morgan PW, Saltveit, Jr ME (1992). Ethylene in plant biology. 2 ed. Academic Press, New York.
- Alexander L, Grierson D (2002). Ethylene biosynthesis and action in tomato: a model for climacteric fruit ripening. *J Exp Bot.* 53: 2039-2055.
- Bleeker A (2001). Ethylene. *Current Biol.* 11: 23- R952.
- Chang C, Stadler R (2001). Ethylene hormone receptor action in *Arabidopsis*. *Bioessays* 23: 619-627.
- Chen YF, Etheridge N, Schaller GE (2005). Ethylene signal transduction. *Ann Bot.* 95: 901-915.
- Fernández-Otero C, Matilla AJ, Rasori A, Ramina A, Bonghi C (2006). Regulation of ethylene biosynthesis in reproductive organs of damson plum (*Prunus domestica* L. subsp. Syriaca). *Plant Sci.* 171: 74-83.
- Giovannoni JJ (2001). Molecular biology of fruit maturation and ripening. *Ann Rev Plant Physiol Plant Mol Biol.* 52: 725-749.
- Hua J, Meyerowitz EM (1998 a). Ethylene responses are negatively regulated by a receptor gene family in *Arabidopsis thaliana*. *Cell* 94: 261-271.
- Hua J, Sakai H, Nourizadeh S, Chen QG, Bleeker AB, Ecker JR, Meyerowitz EM (1998 b). EIN4 and ERS2 are members of the putative ethylene receptor gene family in *Arabidopsis*. *Plant Cell.* 10: 1321-1332.
- Johnson PR, Ecker JR (1998). The ethylene gas signal transduction

- pathway: A molecular perspective. *Ann Rev Genet.* 32: 32227-32254.
- Jones ML (2003). Ethylene biosynthetic genes are differentially regulated by ethylene and ACC in carnation styles. *Plant Growth Regul.* 40: 129-138.
- Kieber JJ, Rothenburg M, Roman G, Feldmann KA, Ecker JK (1993). CTR1 a negative regulator of the ethylene response pathway in Arabidopsis, encodes a member of the Raf family of protein kinases. *Cell* 72: 427-441.
- Kuroda S, Hakata M, Hirose Y, Shiraishi M, Abe S (2003). Ethylene production and enhanced transcription of an ethylene receptor gene, ERS1, in Delphinium during abscission of florets. *Plant Physiol Biochem.* 41: 812-820.
- Kuroda S, Hirose Y, Shiraishi M, Davies E, Abe S (2004). Co-expression of an ethylene receptor gene, ERS1, and ethylene signaling regulator gene, CTR1, in Delphinium during abscission of florets. *Plant Physiol Biochem.* 42: 745-751.
- Leclercq J, Adams-Phillips LC, Zegzouti H, Jones B, Latche' A, Giovannoni JJ, Pech JC, Bouzayen M (2002). LeCTR1, a tomato CTR1-like gene, demonstrates ethylene signaling ability in Arabidopsis and novel expression patterns in tomato. *Plant Physiol.* 130: 1132-1142.
- Lin Z, Hackett R, Payton S, Gerierson D (1998). A tomato sequence, TCTR2 (accession number AJ005077), encoding an Arabidopsis CTR1 homologue. *Plant Physiol.* 117: 1126.
- Manzano S, Gómez P, Garrido D, Jamilena M (2008). Cloning and characterization of a CTR1-like gene in *Cucurbita pepo*. In: Pitrat, M. (Ed), Proc. IX EUCARPIA Meeting, Avignon. PP. 575-580.
- Müller R, Lind-iversen S, Stummann BM, Serek M (2000 a). Expression of genes for ethylene biosynthetic enzymes and an ethylene receptor in senescing flowers of miniature roses. *J Hort Sci Biotech.* 75: 12-18.
- Müller R, Owen CA, Xue ZT, Welander M, Stummann BM (2002). Characterization of two CTR-like protein kinases in *Rosa hybrida* and their expression during flower senescence and in response to ethylene. *J Exp Bot.* 53: 1223-1225.
- Müller R, Stummann BM, Serek M (2000 b). Characterization of an ethylene receptor family with differential expression in rose (*Rosa hybrida* L.) flowers. *Plant Cell Rep.* 19: 1232-1239.
- Müller R, Stummann BM, Sisler EC, Serek M (2001). Cultivar differences in regulation of ethylene production in miniature rose flowers (*Rosa hybrida* L.). *Gartenbauwissenschaft* 1: 34-38.
- Nukui H, Kudo S, Yamashita A, Satoh S (2004). Repressed ethylene production in the gynoecium of long-lasting flowers of the carnation "White Candle": role of the gynoecium in carnation flower senescence. *J Exp Bot.* 55: 641-650.
- Reid MS, Evans RY, Dodge LL, Mor Y (1989). Ethylene and silver thiosulphate influence opening of cut rose flowers. *J Am Soc Hort.* 114: 436-440.
- Shibuya K, Nagata M, Tanikawa N, Yoshioka T, Hashiba T, Satoh S (2002). Comparison of mRNA levels of three ethylene receptors in senescing flowers of carnation (*Dianthus caryophyllus* L.). *J Exp Bot.* 53: 399-406.
- Shibuya K, Yoshioka T, Hashiba T, Satoh S (2000). Role of the gynoecium in natural senescence of carnation (*Dianthus caryophyllus* L.) flowers. *J Exp Bot.* 51: 2067-2073.
- Tan H, Liu X, Ma N, Xue J, Lu W, Bai J, Gao J (2006). Ethylene-influenced flower opening and expression of genes encoding *Etrs*, *Ctrs*, and *Ein3s* in two cut rose cultivars. *Postharvest Biol Technol.* 40: 97-105.
- Tieman DM, Taylor MG, Ciardi JA, Klee HJ (2000). The tomato ethylene receptors NR and LeETR4 are negative regulators of ethylene response and exhibit functional compensation within a multigene family. *PNAS* 97: 5663-5668.
- Wang D, Fan J, Ranu RS (2004). Cloning and expression of 1-aminocyclopropane-1-carboxylate synthase cDNA from rose (*Rosa x hybrida*). *Plant Cell Rep.* 22: 422-429.
- Zegzouti H, Jones B, Frasse P, Marty C, Maitre B, Latche A, Pech JC, Bouzayen M (1999). Ethylene-regulated gene expression in tomato fruit: characterization of novel ethylene-response and ripening-related genes isolated by differential-display. *Plant J.* 18: 589-600.