

## Accuracy of the Semi-Classical Approximation: the Pullen-Edmonds Hamiltonian (\*).

S. GRAFFI<sup>(1)</sup>, V. R. MANFREDI<sup>(2)</sup>(\*\*)(\*\*\*) and L. SALASNICH<sup>(3)</sup>

<sup>(1)</sup> *Dipartimento di Matematica dell'Università*

*Piazza di Porta S. Donato 5, I 40127 Bologna, Italy*

<sup>(2)</sup> *Dipartimento di Fisica «G. Galilei» dell'Università - Padova, Italy*

*Interdisciplinary Laboratory, SISSA - Trieste, Italy*

<sup>(3)</sup> *Dipartimento di Fisica dell'Università - Firenze, Italia*

*INFN, Sezione di Firenze - Largo E. Fermi 2, I 50125 Firenze, Italy*

(ricevuto il 5 Aprile 1994; approvato il 5 Settembre 1994)

**Summary.** — A test on the numerical accuracy of the semi-classical approximation as a function of the principal quantum number has been performed for the Pullen-Edmonds model, a two-dimensional, non-integrable, scaling-invariant perturbation of the resonant harmonic oscillator. A perturbative interpretation is obtained of the recently observed phenomenon of the accuracy decrease on the approximation of individual energy levels at the increase of the principal quantum number. Moreover, the accuracy provided by the semi-classical approximation formula is on average the same as that provided by quantum perturbation theory.

PACS 03.65 – Quantum theory; quantum mechanics.

PACS 05.45 – Theory and models of chaotic systems.

Recently, there has been considerable renewed interest in the various aspects of the semi-classical approximation (SCA), a powerful motivation behind that being the problem of the so-called quantum chaos (see, for example, ref.[1-5]). An important aspect is represented by the effectiveness of the semi-classical quantization formula approximating the quantum energy levels, and in this connection one recent work [6] shows that the predictions of individual levels by SCA (by this we mean the Bohr-Sommerfeld formula, or one of its generalizations to the non-integrable case, such as EBK; see, *e.g.*, [3,4]), worsen as the quantum number increases, contrary to the naive expectation. We argue that this result can be interpreted as follows: if  $\hbar$ , no

---

(\*) This work has been partially supported by the Ministero dell'Università e della Ricerca Scientifica e Tecnologica (MURST).

(\*\*) Permanent address: INFN, Sezione di Padova, Via Marzolo 8, I 35131 Padova, Italy.

(\*\*\*) Author to whom all correspondence and reprint requests should be addressed.

E-mail: vxafpd::manfredi, manfredi@padova.infn.it.

matter how small, is kept fixed, the SCA on the individual levels has the meaning of a perturbation theory (PT) in  $\hbar$ . Therefore, the accuracy of the approximation decreases for higher levels (to get good agreement it is necessary, as is well known, to implement the classical limit  $\hbar \rightarrow 0$ ,  $n \rightarrow \infty$ ,  $n\hbar = I$  classical action; see, *e.g.*, [7]).

The aim of this paper is to clarify this point, from the theoretical standpoint and from the computational one as well, considering a scaling-invariant potential, which makes ordinary quantum PT strictly equivalent to a power expansion in  $\hbar$ . We do actually observe that, for  $\hbar$  fixed, the perturbation strength has to be decreased to keep the accuracy at a constant value as the quantum number increases; however, we also observe that the algorithm provided by the appropriate SCA is always comparable to the algorithm provided by ordinary quantum PT. A very good agreement between the «exact» eigenvalues, obtained by numerical diagonalization of the Schrödinger operator, and the semi-classical ones, is indeed observed in the presence of high unperturbed degeneracy.

The most significant examples to carry out this comparison are represented by non-separable two-dimensional systems exhibiting both regular and irregular spectrum [8], *i.e.*, in particular, non-uniform behaviour of the level spacing, and among these the simplest one is the Pullen-Edmonds model [9]. Its quantum Hamiltonian is

$$(1) \quad H = -\frac{\hbar^2}{2} \left[ \frac{\partial^2}{\partial q_1^2} + \frac{\partial^2}{\partial q_2^2} \right] + \frac{1}{2}(q_1^2 + q_2^2) + \chi q_1^2 q_2^2,$$

where  $m$  has been put equal to 1. For  $\chi = 0$ , (1) reduces to a resonant two-dimensional harmonic oscillator of levels  $(n_1 + n_2 + 1)\hbar \equiv m_1\hbar$ ,  $m_1 = 1, 2, \dots$ , of multiplicity  $m_1$ .

The scaling transformation  $q_1 \rightarrow \sqrt{\hbar} q_1$ ,  $q_2 \rightarrow \sqrt{\hbar} q_2$  yields the unitarily equivalent operator

$$(2) \quad \tilde{H} = -\hbar \left[ \frac{1}{2} \left( \frac{\partial^2}{\partial q_1^2} + \frac{\partial^2}{\partial q_2^2} \right) + \frac{1}{2}(q_1^2 + q_2^2) + \chi \hbar q_1^2 q_2^2 \right].$$

The coupling constant has become  $\chi\hbar$ , which clearly shows equivalence between expansions in powers of  $\chi$  or  $\hbar$ . (An analogous result holds for any other polynomial perturbation.) Moreover, the symmetry of the potential enables us to split the Hamiltonian matrix, computed on the harmonic-oscillator basis, into submatrices reducing the computer storage required. The matrix elements of (1) can be written:

$$(3) \quad \langle n'_1 n'_2 | H | n_1 n_2 \rangle = \hbar(n_1 + n_2 + 1) \delta_{n'_1 n_1} \delta_{n'_2 n_2} + \\ + \chi \frac{\hbar^2}{4} \left[ \sqrt{n_1(n_1 - 1)} \delta_{n'_1 n_1 - 2} + \sqrt{(n_1 + 1)(n_1 + 2)} \delta_{n'_1 n_1 + 2} + (2n_1 + 1) \delta_{n'_1 n_1} \right] \cdot \\ \cdot \left[ \sqrt{n_2(n_2 - 1)} \delta_{n'_2 n_2 - 2} + \sqrt{(n_2 + 1)(n_2 + 2)} \delta_{n'_2 n_2 + 2} + (2n_2 + 1) \delta_{n'_2 n_2} \right]$$

and each submatrix can be labelled by the parity of the occupation numbers  $n_1, n_2$ . We restrict from now on to the invariant subspace spanned by  $m_1$  even, *i.e.*  $n_1$  and  $n_2$  of opposite parity. The eigenvalues of  $H$  in this subspace have constant multiplicity 2 [9]. Therefore, the level  $m_1\hbar = 2s\hbar$  splits into  $s$  levels for  $\chi > 0$ .

The appropriate SCA is here provided by the Bohr-Sommerfeld quantization of

the resonant (or secular) canonical perturbation theory[10], also known, in this particular case, as the Birkhoff-Gustafson normal form[11,12], which we now construct at first order. Starting from the classical Pullen-Edmonds Hamiltonian

$$(4) \quad H_{cl} = \frac{1}{2}(p_1^2 + p_2^2) + \frac{1}{2}(q_1^2 + q_2^2) + \chi q_1^2 q_2^2,$$

we introduce the standard action-angle variables  $(I, \theta)$  by the canonical transformation:

$$(5) \quad \begin{cases} q_i = \sqrt{2I_i} \cos \theta_i, \\ p_i = \sqrt{2I_i} \sin \theta_i, \end{cases} \quad i = 1, 2.$$

Then (4) becomes

$$(6) \quad H_{cl} = I_1 + I_2 + 4\chi I_1^2 I_2^2 \cos^2 \theta_1 \cos^2 \theta_2.$$

The second canonical transformation into the well-known «slow» and «fast» variables,

$$(7) \quad \begin{cases} A_1 = I_1 + I_2, \\ A_2 = I_1 - I_2, \end{cases} \quad \begin{cases} \theta_1 = \phi_1 + \phi_2, \\ \theta_2 = \phi_1 - \phi_2, \end{cases}$$

eliminates the dependence on the «slow action»  $A_2$  in the unperturbed part, so that the Hamiltonian becomes

$$(8) \quad H_{cl} = A_1 + \chi(A_1^2 - A_2^2) \cos^2(\phi_1 + \phi_2) \cos^2(\phi_1 - \phi_2).$$

We now eliminate the dependence on the angles up to terms of order  $\chi^2$  by resonant (or secular) canonical perturbation theory[10]. To eliminate the dependence on the «fast angle»  $\phi_1$ , it is enough to average the perturbation on this variable. This yields

$$(9) \quad \frac{1}{2\pi} \int_0^{2\pi} d\phi_1 \cos^2(\phi_1 + \phi_2) \cos^2(\phi_1 - \phi_2) = \frac{1}{8}(2 + \cos 4\phi_2),$$

and thus

$$(10) \quad \tilde{H}_{cl} = A_1 + \frac{\chi}{8}(A_1^2 - A_2^2)(2 + \cos 4\phi_2).$$

The dependence on  $\phi_2$  o the perturbation part can now be eliminated by af further canonical transformation. The Hamiltonian-Jacobi equation for the perturbation part is in fact

$$(11) \quad \left[ A_1^2 - \left( \frac{\partial S}{\partial \phi_2} \right)^2 \right] (2 + \cos 4\phi_2) = K,$$

$$(12) \quad \frac{\partial S}{\partial \phi_2} = \pm \sqrt{\frac{A_1^2(2 + \cos 4\phi_2) - K}{2 + \cos 4\phi_2}},$$

and thus the Hamiltonian (9) becomes

$$(13) \quad \tilde{H}_{cl} = B_1 + \frac{\chi}{8}K(B_1, B_2),$$

where

$$(14) \quad B_1 = A_1, \quad B_2 = \frac{1}{2\pi} \oint d\phi_2 \frac{\partial S}{\partial \phi_2}.$$

It appears from the structure of eq. (12) that the motions generated by the perturbation part of our system have the following qualitative behaviour:

$$(15) \quad \begin{cases} 0 < K < B_1^2 & \text{rotational motion,} \\ K = B_1^2 & \text{separatrix,} \\ B_1^2 < K < 3B_1^2 & \text{librational motion.} \end{cases}$$

The appearance of a separatrix (which is not immediately obvious in the  $(p, q)$  coordinates) accounts, as is well known (see, e.g., [3]), for the stochastic layers originating near it. This corresponds to local irregular behaviour of the quantum spectrum; one of its manifestations is (see Delande in [4]) the local shrinking of the level spacing and the presence of avoided crossings. This shrinking of the level spacing is best accounted for by the SCA, as we will discuss below.

On the separatrix we have

$$(16) \quad B_1^2 (2 + \cos 4\phi_2) = K,$$

while, in general,

$$(17) \quad B_2 = \pm \frac{2}{\pi} \int_a^b dx \sqrt{\frac{B_1^2 (2 + \cos 4x) - K}{2 + \cos 4x}},$$

where

$$(18) \quad \begin{cases} a = 0, & b = \frac{\pi}{2} & \text{rotational motion,} \\ a = \phi_-(K, B_1), & b = \phi_+(K, B_1) & \text{librational motion,} \end{cases}$$

with

$$(19) \quad \phi_{\pm}(K, B_1) = \pm \frac{1}{4} \arccos \left( \frac{K}{B_1^2} - 2 \right).$$

Now the approximate Hamiltonian (13) depends only on the actions, so that a semi-classical quantization formula for the  $m_1$  even part of spectrum of the operator (1) can be obtained by a straightforward application of the Bohr-Sommerfeld quantization rules [10]. Set therefore

$$(20) \quad \begin{cases} I_1 = (n_1 + 1/2) \hbar, \\ I_2 = (n_2 + 1/2) \hbar, \end{cases}$$

whence, from (6),

$$(21) \quad \begin{cases} A_1 = (n_1 + n_2 + 1) \hbar, \\ A_2 = (n_1 - n_2) \hbar. \end{cases}$$

Set

$$(22) \quad \begin{cases} A_1 = m_1 \hbar, \\ A_2 = m_2 \hbar. \end{cases}$$

By comparison of (19) and (20) we obtain

$$(23) \quad \begin{cases} m_1 = n_1 + n_2 + 1, \\ m_2 = n_1 - n_2, \end{cases}$$

where  $m_1 = 2, 4, \dots$  and  $m_2 = \pm(m_1 - 1), \pm(m_1 - 3), \pm(m_1 - 5), \dots$

Finally,

$$(24) \quad B_1 = m_1 \hbar, \quad B_2 = m \hbar;$$

then the semi-classical approximation to the quantum spectrum is

$$(25) \quad E_{m_1, m_2} = m_1 \hbar + \frac{\chi}{8} K(m_1 \hbar, m_2 \hbar),$$

where  $K$  is implicitly defined by the relation

$$(26) \quad m_2 \hbar = \pm \frac{2}{\pi} \int_a^b dx \sqrt{\frac{(m_1 \hbar)^2 (2 + \cos 4x) - K}{2 + \cos 4x}}$$

and

$$(27) \quad \begin{cases} a = 0, & b = \pi/2, & 0 < K < (m_1 \hbar)^2, \\ a = \phi_-(K, m_1 \hbar), & b = \phi_+(K, m_1 \hbar) & (m_1 \hbar)^2 < K < 3(m_1 \hbar)^2. \end{cases}$$

Remark that for  $|m_2| < [x m_1]$  we obtain the quantization of the rotational motions, while for  $|m_2| > [x m_1]$  ( $[x]$  = integer part of  $x$ ) we have the quantization of




Fig. 1. – Comparison between the «exact» levels (a) the semi-classical ones (b) and the levels obtained by first-order perturbation theory (c), for  $\chi = 10^{-3}$ ,  $\hbar = 0.1$ ,  $m_1 = 30$ .




Fig. 2. - Comparison between the «exact» levels (a), the semi-classical ones (b), and the levels obtained by first-order perturbation theory (c), for  $m_1 = 60$ ,  $\chi = 10^{-5}$ .

the librational ones. Here, by (17)

$$(28) \quad \alpha = \frac{2}{\pi} \int_0^{\pi/2} dx \sqrt{\frac{1 + \cos 4x}{2 + \cos 4x}} \approx 0.602.$$

Moreover, it is easy to see that for  $m_1$  fixed, the function  $K$ , and hence the semi-classical energy  $E_{m_1, m_2}$ , is a decreasing function of the secondary quantum number  $m_2$ . It is furthermore proved in [13] that (25) coincides with the exact quantum spectrum up to terms of order  $\hbar$  and  $\chi^2$ . The numerical computations (see fig. 4 below) show that at order 1 in  $\chi$  the corrections of order  $\hbar$  affect at most the eighth decimal figure.

The «exact» levels have been computed, and compared with the semi-classical ones as well as with the levels computed by degenerate first-order quantum perturbation theory [14], for  $m_1 = 1, \dots, 60$  at  $\hbar = 0.1$  and for different values of  $\chi$  (given the degeneracy, this is equivalent to compute 1800 different levels). The




Fig. 3. - The difference  $\Delta$  between the «exact» levels and the semi-classical ones vs.  $m_1$ , with  $m_2 = (m_1 - 1)$  and  $m_1 \hbar = 1$ .

results obtained for  $m_1 = 30, \hbar = 0.1, \chi = 10^{-3}$  and  $m_1 = 60, \hbar = 0.1, \chi = 10^{-5}$  are shown in fig. 1 and 2, respectively. The local shrinking of the spacing, reproduced by both methods, can be immediately noticed; remark that the corresponding semi-classical levels are those near the separatrix (by (28),  $m_2 \sim 18$  and  $m_2 \sim 36$ , respectively).

In fig. 3 the function

$$(29) \quad \Delta = |E^{\text{Ex}} - E^{\text{SCA}}|$$

vs.  $m_1$  is plotted for  $m_1 \chi = 1$ ; this shows that, if the coupling constant is decreased in inverse proportion to the principal quantum number the accuracy of SCA not only




Fig. 4. - a) The difference between the «exact» levels and the semi-classical ones; b) the difference between the «exact» levels and the first-order quantum PT ones; ( $m_1 = 60, \hbar = 0.1, \chi = 10^{-5}$ ).




Fig. 5. - The difference between the semi-classical levels and the first-order quantum PT ones; ( $m_1 = 60, \hbar = 0.1, \chi = 10^{-5}$ ).

remains constant but actually improves, as anticipated because the scaling invariance makes the limit  $m_1 \rightarrow \infty$ ,  $\chi \rightarrow 0$ ,  $m_1 \chi \rightarrow \text{const}$  equivalent to the classical limit  $m_1 \rightarrow \infty$ ,  $\hbar \rightarrow 0$ ,  $m_1 \hbar \rightarrow \text{const}$ .

In fig. 4 the accuracies obtained through semi-classical and quantum first-order perturbation theories are compared for  $m_1 = 60$ ,  $\chi = 10^{-5}$ ,  $\hbar = 0.1$ , and in fig. 5 the difference between the two perturbation theories is plotted (remark that the energy decreases as  $m_2$  increases). As can be seen, the agreement with the «exact» levels is very good and the accuracy is on average the same. Remark however that, as it should be expected (the Bohr-Sommerfeld rules take no account of tunnelling [15]), the lowest accuracy of the SCA is reached near  $m_2 = 36$  which corresponds to the levels near the separatrix: for these levels the quantum PT is therefore better than SCA.

\* \* \*

The authors are greatly indebted to Dr. Stefano Isola for many useful discussions and to Mr. G. Salmaso for his valuable computational assistance.

## REFERENCES

- [1] V. P. MASLOV and M. V. FEDORIUK: *Semi-Classical Approximation in Quantum Mechanics* (Reidel Publishing Company, 1981).
- [2] A. B. MIGDAL: *Qualitative Methods in Quantum Theory* (Benjamin, 1977).
- [3] M. C. GUTZWILLER: *Chaos in Classical and Quantum Physics* (Springer-Verlag, 1991); A. R. P. RAU: *Rev. Mod. Phys.*, **64**, 623 (1992).
- [4] D. DELANDE: *Chaos and Quantum Physics*, in *Les Houches Summer School, Course LII, 1989*, edited by M. J. GIANNONI, A. VOROS and J. ZINN-JUSTIN (Elsevier Science Publishing, 1989).
- [5] P. A. BRAUN: *Rev. Mod. Phys.*, **65**, 115 (1993).
- [6] T. PROSEN and M. ROBNIK: *J. Phys. A*, **26**, L37 (1993).
- [7] G. ALVAREZ, S. GRAFFI and H. J. SILVERSTONE: *Phys. Rev. A*, **38**, 1697 (1988).
- [8] I. C. PERCIVAL: *J. Phys. B*, **6**, L229 (1973).
- [9] R. A. PULLEN and A. R. EDMONDS: *J. Phys. A*, **14**, L477 (1981).
- [10] M. BORN: *Mechanics of the Atom* (Bell, London, 1960).
- [11] W. P. REINHARDT: in *The Mathematical Analysis of Physical Systems*, edited by R. E. MICKENS and R. VAN NOSTRAND (1984).
- [12] T. UZER, D. W. NOID and R. A. MARCUS: *J. Chem. Phys.*, **79**, 4412 (1983).
- [13] S. GRAFFI: in *Probabilistic Methods in Mathematical Physics*, edited by F. GUERRA, M. I. LOFFREDO and C. MARCHIORO (World Scientific, 1992).
- [14] L. LANDAU and E. LIFSHITZ: *Quantum Mechanics: Non Relativistic Theory* (Pergamon Press, Oxford, 1978).
- [15] A. M. OZORIO DE ALMEIDA: *J. Phys. Chem.*, **88**, 6139 (1984).