

*Title Page*

**When, where and how? Focus on neuronal calcium dysfunctions in Alzheimer's Disease**

*Authors*

**Mario Agostini<sup>a</sup> and Cristina Fasolato<sup>a\*</sup>**

*Affiliations:*

<sup>a</sup>Department of Biomedical Sciences, University of Padua, Italy

\*To whom correspondence should be addressed:

Cristina Fasolato  
Department of Biomedical Sciences  
University of Padua  
Via Ugo Bassi 58B  
35131 Padua, Italy  
Tel: +39 049 8276065  
FAX: +39 049 8276049

*E-mail addresses:*

Cristina Fasolato: [cristina.fasolato@unipd.it](mailto:cristina.fasolato@unipd.it)

Mario Agostini: [mario.agostini@phd.unipd.it](mailto:mario.agostini@phd.unipd.it)

## **ABSTRACT**

Alzheimer's disease (AD), since its characterization as a precise form of dementia with its own pathological hallmarks, has captured scientists' attention because of its complexity. The last 30 years have been filled with discoveries regarding the elusive aetiology of this disease and, thanks to advances in molecular biology and live imaging techniques, we now know that an important role is played by calcium ( $\text{Ca}^{2+}$ ).  $\text{Ca}^{2+}$ , as ubiquitous second messenger, regulates a vast variety of cellular processes, from neuronal excitation and communication, to muscle fibre contraction and hormone secretion, with its action spanning a temporal scale that goes from microseconds to hours. It is therefore very challenging to conceive a single hypothesis that can integrate the numerous findings on this issue with those coming from the classical fields of AD research such as amyloid-beta ( $\text{A}\beta$ ) and tau pathology. In this contribution, we will focus our attention on the  $\text{Ca}^{2+}$  hypothesis of AD, dissecting it, as much as possible, in its subcellular localization, where the  $\text{Ca}^{2+}$  signal meets its specificity. We will also follow the temporal evolution of the  $\text{Ca}^{2+}$  hypothesis, providing some of the most updated discoveries. Whenever possible, we will link the findings regarding  $\text{Ca}^{2+}$  dysfunction to the other players involved in AD pathogenesis, hoping to provide a crossover body of evidence, useful to amplify the knowledge that will lead towards the discovery of an effective therapy.

## **Keywords:**

Alzheimer's disease, calcium homeostasis, amyloid-beta, VOCCs, nAChRs, glutamate receptors, CALHM1, SOCCs, endoplasmic reticulum, mitochondria, lysosomes.

## INTRODUCTION

When the  $\text{Ca}^{2+}$  hypothesis of AD arose during the late '80s and early '90s, it was somehow disconnected from the amyloid cascade hypothesis that was already moving its first steps [1] [2]. When, later in the decade, studies linking  $\text{A}\beta$  accumulation and deposition to  $\text{Ca}^{2+}$  dysregulation began to appear the amyloid and the  $\text{Ca}^{2+}$  hypotheses became linked [3]. It was soon clear that not only amyloid plaques but also soluble  $\text{A}\beta$  aggregates, and specifically  $\text{A}\beta$  oligomers ( $\text{A}\beta\text{o}$ ), were exerting significant toxicity, either endogenously produced or exogenously administered to cells and primary neurons in culture [4][5] [6] (**Figure 1**). The  $\text{Ca}^{2+}$  hypothesis of AD, by positing intracellular  $\text{Ca}^{2+}$  dysregulation as the direct consequence of  $\text{A}\beta$  toxicity, offered one of the richest grounds for researchers in the field. It obscured the fact, however, that  $\text{Ca}^{2+}$  dysregulation at the onset of AD can occur independently of  $\text{A}\beta$  accumulation and toxicity [7]. AD is now considered as a multiple hits disease where amyloid and tau dysfunctions are aggravated by oxidative stress, intracellular  $\text{Ca}^{2+}$  imbalance and metabolic disturbance. A large body of evidence strongly points towards  $\text{Ca}^{2+}$  as a possible unifying element underlying AD pathophysiology. According to the amyloid cascade hypothesis,  $\text{A}\beta$  toxicity precedes tau pathology but exactly how  $\text{A}\beta$  and tau cooperate in synaptic dysfunction is still a controversial issue [8]. Studies with AD patients as well as with amyloid precursor protein (APP) and/or tau mouse models have shown that tau defects affect the lateral entorhinal cortex at an early stage but require  $\text{A}\beta$  to reach other cortical sites [9].

In this contribution, we will mainly focus on the relationship between cellular  $\text{Ca}^{2+}$  handling and AD, paying special attention to the crosstalk with  $\text{A}\beta$ . Given the large amount of data collected, the often contrasting hypotheses and the vast array of cellular players more directly involved in  $\text{Ca}^{2+}$  homeostasis, the review will be divided in thematic paragraphs, based on the “anatomy” of the  $\text{Ca}^{2+}$  signal (**Figure 2**).

## 2. PLASMA MEMBRANE CHANNELS

It was initially suggested that  $\text{A}\beta$  peptides are capable of forming plasma membrane channels [10]; indeed, remarkable increases in the cytosolic  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_{\text{cyt}}$ ) were reported upon challenging cultured cells with either  $\text{A}\beta\text{40}$  or  $\text{A}\beta\text{42}$  [11] [12] [13]. The “ion channel hypothesis” for  $\text{A}\beta$  toxicity, as proposed by Arispe’s group, was mainly supported by *in vitro* studies demonstrating the formation of cation selective channels, whose permeation is inhibited by metals such as  $\text{Zn}^{2+}$  and  $\text{Cu}^{2+}$  and by particular peptides designed to line the pore-forming region [14] [15]. The ion channel hypothesis got further support from Demuro’s studies showing that, in oocytes,  $\text{A}\beta$  injection causes cytosolic  $\text{Ca}^{2+}$  rises due to  $\text{Ca}^{2+}$  entry across  $\text{A}\beta$  channels formed in the plasma membrane as well as  $\text{Ca}^{2+}$  release via stimulation of the Gq/PLC pathway [16]. This hypothesis contrasts with *in vivo* data showing that exogenously applied  $\text{A}\beta$  enhance the overall plasma membrane permeability to anions and cations rather than forming a  $\text{Ca}^{2+}$ -selective pathway [17]. Since the first observations describing the increase in the  $[\text{Ca}^{2+}]_{\text{cyt}}$  of cells directly harvested from AD patients [18], many groups had focused their attention on this phenomenon changing the paradigm from the channel-forming to the channel-modulating hypothesis. This intense effort led to a plethora of hypothetical candidates capable of mediating cellular  $\text{Ca}^{2+}$  overload by  $\text{A}\beta$  since neuronal cells have a vast array of  $\text{Ca}^{2+}$ -permeable plasma

membrane channels at their disposal. As suggested players implicated in the pathogenesis of AD, and possibly mediating A $\beta$  toxicity, we will here consider the voltage-operated Ca<sup>2+</sup> channels (VOCCs), the nicotinic acetylcholine receptors (nAChRs), the ionotropic glutamate receptors [N-Methyl-D-aspartate receptors (NMDARs) and  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors (AMPA)], the Ca<sup>2+</sup> homeostasis modulator 1 (CALHM1), as well as the store-operated Ca<sup>2+</sup> channels (SOCCs). For a thorough investigation on this issue, also including the plethora of A $\beta$  binding receptors, the readers should refer to excellent more extensive reviews of which only a few are cited here.

## 2.1. VOCCs

VOCCs open following membrane depolarization: the firing of an action potential by the neuron, that is the integrated answer of the afferent presynaptic stimuli, causes the plasma membrane to depolarize, allowing Ca<sup>2+</sup> to enter from the outside along its huge electrochemical gradient. This Ca<sup>2+</sup> signal causes the opening of a subtype of potassium channels that helps membrane repolarization. Ca<sup>2+</sup> entry is coupled with synaptic vesicle release and intracellular signalling cascades that lead to gene transcription of several players involved in synapses' maintenance and plasticity.

VOCCs, especially those of the L-type subfamily, have long been implicated in aging and AD. Several studies have reported an elevated activity of these channels, with a subsequently larger Ca<sup>2+</sup>-dependent hyperpolarization, as a possible explanation for the deterioration of synaptic efficiency [19][20] [21] [22]. Given that the Ca<sup>2+</sup> signal is spatially and temporally well defined, it is conceivable that even a little delay in its clearance from the cytosol after depolarization can have dramatic effects; for instance, the differential activation of the Ca<sup>2+</sup>-Calmodulin-dependent kinase II (CaMKII) and the phosphatase calcineurin (CN) is based on their different reaction times to cytosolic Ca<sup>2+</sup> elevations [23] [24].

An impaired intracellular Ca<sup>2+</sup> handling, that derives either from an increased and sustained entry through the membrane channels or from an inefficient clearance, can lead to an imbalance between long-term potentiation (LTP) and long-term depression (LTD), the electrophysiological correlates that support learning and memory, thus eventually resulting in disbanding memory storage. L-type ion channels are required for LTP at synapses between mossy fibres and CA3 pyramidal neurons. This type of LTP does not depend on NMDARs and involves postsynaptic Ca<sup>2+</sup> influx carried by L-type VOCCs [25]. Patients with AD show higher L-type VOCC expression in the hippocampus compared with healthy individuals [26]. Forette and co-workers found that the blockage of L-type VOCCs with dihydropyridines, a class of drugs mainly used to treat systolic hypertension, was able to prevent dementia in some AD patients [27]. Consistently, in CA1 neurons of the hippocampus, Ca<sup>2+</sup> currents through L-type VOCCs increase during aging in rats, together with a marked decline in learning and memory faculties. Chronic nimodipine treatment ameliorates memory loss, suggesting that excessive Ca<sup>2+</sup> influx, through L-type Ca<sup>2+</sup> channels, impairs learning and memory [28]. However, it has been demonstrated that in cultured neurons from APP<sub>sw,Lo</sub> mice, A $\beta$  suppress gene transcription that is mediated by Ca<sup>2+</sup> influx (and cAMP surge) and agonists of L-type VOCCs efficiently reverse these transcriptional deficits, implicating a defect in Ca<sup>2+</sup> influx rather than an augmentation [29]. Furthermore, in cell lines and primary cerebellar

granules from the Tg2576 AD mice, dihydropyridines, and especially the most used nimodipine, increase A $\beta$ 42 secretion through a mechanism that is independent of Ca<sup>2+</sup> entry-blockage thus raising doubts to the possible therapeutic use of these compounds [30]. The effect of A $\beta$  on L-type VOCCs, whenever present, is likely dose- and time-dependent, since a 1-hour exposure of mouse cortical wild-type neurons to A $\beta$ 42o at submicromolar concentrations does not affect Ca<sup>2+</sup> peaks and long lasting plateaus due to membrane depolarization [31]. Interestingly, A $\beta$  globulomers (the 48-kDa component of A $\beta$ o) suppress spontaneous synaptic activity of both GABAergic and glutamatergic synapses by acting selectively on presynaptic P/Q type Ca<sup>2+</sup> currents [32]. At the presynaptic level, A $\beta$ o also affect Ca<sup>2+</sup> influx through VOCCs, possibly explaining fast axonal transport defects [33].

## 2.2. nAChRs

A $\beta$  were reported to affect cholinergic neurons precociously. In the central nervous system, their cell bodies are packed in dense nuclei in the basal forebrain whilst their axonal projections spread throughout the whole cortex, exerting a modulatory effect rather than pure excitation. Acetylcholine binds to two families of receptors, the ionotropic nicotinic receptor (nAChR) and the metabotropic muscarinic one (mAChR). The first one is a Ca<sup>2+</sup>-permeable non-selective cation channel with a pentameric structure being formed by alpha or beta subunits, or a mix of the two. The subunit composition influences the pharmacological and kinetic properties of the channel, varying its permeability to K<sup>+</sup>, Na<sup>+</sup> and Ca<sup>2+</sup> [34].

Ionotropic nAChR exist in various forms, among which the homomeric ( $\alpha$ 7) and the heteromeric  $\alpha$ 4 $\beta$ 2-nAChR were the first to be investigated in AD and they still represent the most promising targets [35] [36]. Recently an heteromeric  $\alpha$ 7 $\beta$ 2 nAChR has been added to the list of A $\beta$ -binding nAChRs [37]. They differ in their Ca<sup>2+</sup> permeability and sensitivity to  $\alpha$ -bungarotoxin, and, functionally speaking, they are thought to take part in memory formation and attention and they have been long known to interact with A $\beta$ . In particular, several studies demonstrated the different affinity and exerted effects upon binding to A $\beta$ 40 or A $\beta$ 42 by the  $\alpha$ 7-nAChR and the  $\alpha$ 4 $\beta$ 2-nAChR [38]. Summarizing, we can say that both receptor subtypes bind A $\beta$ , preferentially the 42 amino-acid long peptides, which activate  $\alpha$ 7-nAChR in the picomolar range and inhibit it in the nanomolar one, whilst  $\alpha$ 4 $\beta$ 2-nAChR binds A $\beta$  only in the nanomolar range with an inhibitory effect. Given that amyloid deposition is a spatially defined phenomenon, with certain brain areas exposed to premature deposition, it is conceivable that widely different interstitial A $\beta$  concentrations can be achieved, causing diverse effects on nAChRs. Albeit contrasting with regard of the A $\beta$  concentrations, species (oligomers vs monomers) and models used, almost all the evidence points towards a disruption of the physiologically nAChRs-driven LTP, which predicts an impairment in memory formation [39]. This was tested further by generating *in vivo* models bearing AD-related APP mutations in a  $\alpha$ 7-nAChR knock-out (KO) context. However, depending on the APP mutations employed, different results were obtained. The KO of  $\alpha$ 7-nAChR in a APP<sub>swe,Indiana</sub> mouse model was protected against synaptic loss and memory deficits, when tested at 13 and 16 months of age [40], while the same KO in a APP<sub>swe</sub> mouse model worsened both memory and A $\beta$  pathology, with increased amount of

soluble high molecular weight (HMW) oligomers when tested at 5 months of age [41]. To our knowledge, no studies were conducted to shed light on the intracellular pathway involved in these effects and, therefore,  $\text{Ca}^{2+}$  contribution is inferred only on the basis of the high  $\text{Ca}^{2+}$  permeability of neuronal nAChRs. Indeed, a likely recruitment by A $\beta$  of astrocytic  $\alpha$ 7-nAChR, altering both  $\text{Ca}^{2+}$  homeostasis and gliotransmission, further complicates the picture [42] [43].

### 2.3. Ionotropic Glutamate Receptors

A vast majority of excitatory neurons rely on glutamate as neurotransmitter and are therefore defined glutamatergic. Glutamate exerts its excitatory effect interacting with a diverse array of ionotropic receptors (NMDARs, AMPARs, KainateRs) and metabotropic receptors (mGluRs). In the AD/aging field, the best characterized is the NMDAR since it is pivotal in LTP. It has been proposed that the interaction of A $\beta$  with NMDARs is an initiating/propagating event in the pathogenesis of AD. A high percentage of the reports concerning the physical and functional interactions between A $\beta$  and NMDARs relies on synthetically prepared, poorly characterized oligomers, thus leading to contrasting and misleading results. Co-immunoprecipitation with the NMDAR subunit NR1, together with genetic knockdown of the latter and pharmacological blockade of the whole channel had proven that A $\beta$  preferentially target the synapses due to the presence of NMDARs [44] [45]. Recently, it was shown that soluble A $\beta$  assemblies derived from the brains of individuals with AD interact with cellular prion protein (PrP<sup>C</sup>) to activate Fyn [46] [47]. A $\beta$  engagement of PrP<sup>C</sup>-Fyn signalling requires mGluR5 as co-receptor, which leads to phosphorylation of the GluN2B subunit, culminating in loss of surface NMDARs and dendritic spines [48]. Furthermore, PrP<sup>C</sup> KO prevents the toxic effects on LTP ascribed to A $\beta$  [48]. Nonetheless it was reported that PrP<sup>C</sup> exerts a modulatory effect on the glutamatergic signalling, also in physiological conditions, with neurons from a PrP<sup>C</sup> null mice exhibiting sustained and aberrant NMDAR mediated currents [49]; a competition between PrP<sup>C</sup> and A $\beta$  for the modulation of the NMDAR activity could be present since they exert opposite effects on glycine binding at the co-agonist site, with PrP<sup>C</sup> negatively modulating it [50].

It has recently been suggested that A $\beta$ , by triggering astrocytic release of glutamate to the extrasynaptic space, preferentially affect extra-synaptic GluN2B-NMDARs that, at variance with synaptic GluN2A-NMDARs, further promote the amyloidogenic pathway and A $\beta$  production [51]. The increased amyloid burden can eventually lead to neuronal toxicity by inhibition of excitatory amino acid transporters (EAATs). The subsequent increase in the extracellular concentration of glutamate in turn activates its receptors in an aberrant way; albeit never demonstrated, the interaction between A $\beta$  and EAATs is extremely likely and there are studies reporting an altered expression pattern of EAATs in the AD brain [52]. Taken together, these findings suggest a neurotoxic action mediated by A $\beta$  through the disbandment of the excitatory glutamatergic transmission. However, the only strong evidence at the *in vivo* level is the positive effect of the weak NMDAR antagonist memantine, which has long been used in AD treatment, and its successor nitro-memantine. Both are efficient at blocking or reversing the deleterious actions of A $\beta$  largely because of their selectivity for extrasynaptic NMDARs [53] [51]. The hypothesis of A $\beta$  toxicity involving extra-synaptic NMDARs got only

partial validation given that the GluN2B-NMDAR antagonist Ro 25-6981 is protective against excitotoxic neuronal death and A $\beta$ -induced synapse loss in some AD mouse models [54][55], and was not confirmed in others [56].

Among other intracellular events prompted by A $\beta$ , the activation of calcineurin and the dephosphorylation of NMDARs and AMPARs are primary mechanisms prompting receptor internalization [57][58][59]. Of note, the synaptic loss induced by A $\beta$  through NMDAR activation seems independent of Ca<sup>2+</sup> influx, relying mainly on p38-MAPK activation [60].

An issue of extreme importance, that has unfortunately gone almost unseen, is the fact that a substantial amount of data on the interaction between A $\beta$  and glutamate receptors come from studies employing synthetic A $\beta$  that have been prepared by dissolving and aging the monomers in DMEM/F12 medium [61]. The latter type of medium specifically causes cytosolic Ca<sup>2+</sup> increases on its own, likely because of its high content in glutamate and, especially of glutamine that undergoes degradation during the aging procedure. It has been estimated that a 100  $\mu$ M stock solution of A $\beta$  aged in DMEM-F12 medium might contain up to 100  $\mu$ M glutamate [62]. Using A $\beta$  at concentrations equal or above 1  $\mu$ M thus means stimulating cells with glutamate at concentrations sufficient to elicit Ca<sup>2+</sup> rises through NMDA- and mGlu-Rs [62]. With this caveat, data interpretation is extremely complicated. Moreover, A $\beta$  were often administered to neuronal cultures or slices, at concentrations considerably higher than those found in the AD brain (5-10  $\mu$ M). Neuronal Ca<sup>2+</sup> rises, due to addition of different culture media used for preparing A $\beta$ , were recently described also by Caballero et al. [63]. The same authors show that culture media devoid of glutamate and glutamine but containing sub-micromolar amounts of Zn<sup>2+</sup>, Cu<sup>2+</sup>, Fe<sup>2+</sup> allow production of low molecular weight (LMW) oligomers that are responsible for cytosolic and mitochondria Ca<sup>2+</sup> overload [63]. We have recently demonstrated that neither synthetic A $\beta$  monomers nor oligomers (ranging from low to high MW), when aged in distilled water, can elicit neuronal Ca<sup>2+</sup> increases when acutely applied to neuronal cultures and PC12 cells at concentrations ranging from 0.5 to 5  $\mu$ M (**Figs 3,4**) [31]. Conversely, a 1-hour exposure of cortical neurons to A $\beta$  profoundly disturbs neuronal Ca<sup>2+</sup> stores through a pathway that involves mGluR5 but not NMDARs [31]. Our findings point to a more subtle but insidious effect of A $\beta$ , rather than to a massive Ca<sup>2+</sup> load. Indeed, the idea that a long-lasting disease, such as AD, might act simply through the disruption of a neurotransmission as ubiquitous as the glutamatergic one, shed shadows on a generalized, massive involvement of neuronal glutamate receptors, and forces to investigate AD dysfunctions also at different levels such as the glia and the neurovascular unit [64][65].

#### **2.4. CALHM1**

Another relevant player in the crosstalk between Ca<sup>2+</sup> and A $\beta$  is the Ca<sup>2+</sup> homeostasis modulator 1 (CALHM1), whose polymorphism P86L was shown to segregate with early-onset forms of sporadic AD [66]. Initially thought to be a Ca<sup>2+</sup>-selective entry pathway and later proposed as an ATP-exiting pore majorly involved in taste transmission [67], CALHM1 is also more generally implicated in neuronal excitability [68]. Interestingly, the channel shares homology with connexins, pannexins and innexins [69]. It was also described as a relevant

component of neuronal ER  $\text{Ca}^{2+}$  leak, which is blocked by a class of compounds similar to benzodiazepines [70][71]. In *Calhm1* knockout mice, A $\beta$  accumulation is increased by 50% because of reduced clearance by insulin degrading enzyme; evidence was provided showing that channel activation and  $\text{Ca}^{2+}$  influx are required for the antiamyloidogenic effect of CALHM1 [72][73].

## 2.5. SOCCs

All the major components (Orai and STIM proteins) of the Store-Operated  $\text{Ca}^{2+}$  Channels (SOCCs) have been described at the neuronal level and a significant amount of studies have underlined the relevance of Capacitative  $\text{Ca}^{2+}$  Entry (CCE) on the pathophysiology of the CNS [74][75][76] and AD [77][78]. Notwithstanding, very little is known on the effects of A $\beta$  at this level. As originally reported by Verkhratsky and coll. [79], in central but not peripheral rat neurons, caffeine-sensitive  $\text{Ca}^{2+}$  stores are almost empty at rest, likely because of a highly dynamic ER  $\text{Ca}^{2+}$  efflux, caused by the ongoing neuronal spiking activity and neurotransmitter release. Despite the fact that these neurons can efficiently refill their store  $\text{Ca}^{2+}$  content during KCl-evoked depolarization, they lose the releasable  $\text{Ca}^{2+}$  very quickly [79]. Similarly, in mouse cortical neurons  $\text{IP}_3$ -sensitive  $\text{Ca}^{2+}$  stores are very labile, and can be maximally refilled by prolonged depolarization [80]. On these premises it is expected that CCE, to name all the  $\text{Ca}^{2+}$  entry pathways linked to store depletion, is constitutively active at rest [81]. We have recently shown that acute treatment of mouse cortical neurons with A $\beta$  before the refilling process does not significantly modify the store  $\text{Ca}^{2+}$  content, suggesting that neuronal CCE is not acutely blocked by soluble A $\beta$  [31]. We can also exclude that a 1-hour exposure to A $\beta$  exerts an indirect effect on CCE given that the total store  $\text{Ca}^{2+}$  content is not affected, as assayed by ionomycin treatment [31]. Taken together, we do not envisage a direct effect of A $\beta$  on CCE. Indeed, the opposite is likely true, with CCE having a direct effect on A $\beta$  secretion: a high  $\text{Ca}^{2+}$  influx through CCE reduces A $\beta$  secretion, while a low influx increases it [82]. Given the fact that FAD-linked mutant PSs reduce CCE, their effect on A $\beta$  accumulation is twofold: favouring A $\beta$  production as well as secretion [83]. Recently, it has been shown that the STIM2/neuronal SOC/CaMKII pathway is down-regulated in PS1KI mice, a condition that leads to loss of mushroom spines, the “stable memory spines” that are directly involved in long-term memory by making functionally stronger synapses [84]. The same group similarly demonstrated that, in hippocampal neurons of APPKI mice, mushroom spine loss results as accumulation of extracellular A $\beta$ 42 [85]. The effect is due to overstimulation of mGluR5 by A $\beta$ 42, causing ER  $\text{Ca}^{2+}$  overload and consequently STIM2 down-regulation [85]. Unfortunately, the A $\beta$  employed in that study were aged in F12 medium raising doubts about the effective toxic agent. Of note, the ER  $\text{Ca}^{2+}$  overload induced by A $\beta$ , described in APPKI neurons [85], is a chronic and indirect effect, since different FAD-linked APP mutations modify neither the cytosolic nor the store  $\text{Ca}^{2+}$  level ([86]) but see also [87] for a reduced ER  $\text{Ca}^{2+}$  loading capacity in APP<sub>swe</sub>-expressing cells.

It is worth noting that a major role for CCE is expected in astrocytes and microglia cells, whose signalling is largely based on ER stores. Interestingly, a reduction in  $\text{Ca}^{2+}$  release and sustained  $\text{Ca}^{2+}$  influx upon agonist stimulation was reported in microglia of AD patients compared to non-demented individuals [88] whereas chronic incubation of astrocytes with A $\beta$  at nanomolar concentrations potentiates  $\text{Ca}^{2+}$  influx caused by



activation of mGluRs [89].

By using plasma membrane-targeted aequorin, it was recently shown that in mouse cerebellar granules, PrP<sup>C</sup> exerts a tonic inhibition on SOC Entry channels by reducing the level of STIM1 phosphorylation [90]. It was suggested that A $\beta$  target mGluRs through the interaction with PrP<sup>C</sup> [91][48] one of the major A $\beta$  binding sites. Of interest, potentiation of SOCE by A $\beta$  is lost in PrP<sup>C</sup>-KO mice [90].

### 3. ENDOPLASMIC RETICULUM CHANNELS

Amidst the complex scenario of ER Ca<sup>2+</sup> dysregulation, another set of findings has shaken the scientific community over the past 15 years. At the beginning of the 21st century, several groups reported that presenilins (PS1 and PS2), the catalytic subunits of the  $\gamma$ -secretase, are able to modify the ER Ca<sup>2+</sup> handling when overexpressed in cell lines (see for reviews [83][92]). Among the most successful sets of studies was that produced by Bezprozvanny and co-workers, who defined the well-known paradigm of the “Ca<sup>2+</sup> overload” hypothesis of AD [93]. According to this model, endogenous wild-type (wt) PSs work as the elusive ER Ca<sup>2+</sup> leak channel that has puzzled scientists for decades, a function that is impaired when one PS allele bears a FAD-linked mutation. This situation leads to accumulation of Ca<sup>2+</sup> inside the ER and consequently exaggerated, toxic release when the IP3Rs and RyRs open. This fascinating hypothesis has been questioned both by ourselves and other groups [94]. In fact, by using organelle-targeted Ca<sup>2+</sup> probes, we first discovered that different types of PS2 mutants reduce the ER and Golgi apparatus Ca<sup>2+</sup> content, with PS1 mutants exerting no effect or even partially mimicking the PS2 mutants [80][95] [7][96]. These latter exert their effects through a reduction of the SERCA pump with minor effects on ER Ca<sup>2+</sup> leak [97]. By using both cytosolic and ER Ca<sup>2+</sup> probes, we provided evidence that not only cell lines but also neurons in cultures and slices from PS2-N141I transgenic mice display a reduced store Ca<sup>2+</sup> content and responsiveness to IP3-generating agonists, albeit in the presence of larger responses to ryanodine receptor (RyR) activation by caffeine [98] [99]. No clear differences in ER Ca<sup>2+</sup> levels and dynamics were found between wt and PS double knockout (DKO) MEFs [97]. Furthermore, Foskett and coworkers similarly found no clear evidence of ER Ca<sup>2+</sup> overload due to the expression of PS1 or PS2 mutants [94], and partly solved the issue demonstrating that the “apparent” Ca<sup>2+</sup> overload is due to exaggerated Ca<sup>2+</sup> release as a consequence of IP3R-gating modulation by PS mutants [100]. Moreover, they demonstrated that in 3xTg mice, reduction of IP3R1 expression levels attenuates A $\beta$  accumulation and tau hyper-phosphorylation and rescues hippocampal LTP and memory deficits [101]. In spite of the conflicting findings related to ER Ca<sup>2+</sup> content and IP3R release, it might be possible to envisage a unifying hypothesis that sees the RyR over-expression/activation as a common pathway that brings together PS1 and PS2 mutants as well as A $\beta$  [102][103]. Notably, dantrolene, a RyR1 blocker used to treat malignant hyperthermia, has been reported to soften the symptoms and to reduce the amyloid burden in AD mouse models [87][104]. Since RyR is present in three different isoforms, all of them represented at the brain level, and due to the lack of commercially available specific antibodies, it is difficult to prove which specific isoform is involved. In the previous work, using quantitative PCR (qPCR), the potentiated RyR response was attributed to upregulation of RyR2, the isoform also linked to synaptic loss and neuronal death [105][106]. Increased

levels of RyR2 also characterize the mouse brain of transgenic mice based on mutant PS2 [99]. In contrast, the RyR3 isoform, which exerts a pro-survival effect, was not involved [107].

#### 4. THE MITOCHONDRIA CHANNELS

Mitochondria are currently considered among the primary targets and players in neurodegenerative diseases, including AD, with mitochondrial dysfunctions being considered a hallmark of A $\beta$ -induced neuronal toxicity. It has been shown that soluble A $\beta$  accumulate precociously in neuronal mitochondria of Tg2576 mice [108]. Additionally, APP has been suggested to enter the mitochondrial protein import machinery (TOM/TIM) with its C-terminal domain, leaving the huge N-terminal domain outside [109]. The APP intracellular domain (AICD) is then cleaved by mitochondrial  $\gamma$ -secretases and dysfunctions might thus arise from P3 and A $\beta$  generated inside the mitochondrial matrix [110]. *In vitro* studies have shown that incubation of A $\beta$  with cells or purified mitochondria leads to decreased mitochondrial function [111]. Others have described deleterious effects of A $\beta$  on the outer mitochondrial membrane (OMM) where, by activating the apoptotic BAK pore, they induce cytochrome *c* release and cell death [112]. A $\beta$  interaction with BAK on the OMM is difficult to conceive giving that A $\beta$  do not display a cytosolic localization, being found extracellularly, following both production at the plasma membrane (less than 10%) and active release, and intracellularly inside vesicles - endosomes and multivesicular bodies - where they are produced upon APP internalization and from which they cannot easily escape. Thus, under pathophysiological conditions, direct effects of A $\beta$  on intracellular channels (MCU, PTP, VDAC, RyR, IP3R, ER leak channels, to mention a few) are not expected, unless they are translocated across membranes. Whereas A $\beta$  mitochondrial localization and uptake were confirmed in brain human samples and in *in vitro* studies [108][109], mitochondrial localization of both APP and  $\gamma$ -secretase components is still waiting, leaving the topology conundrum of A $\beta$  transfer across membranes yet an open question. Furthermore, it remains to be explained what physiological role APP is playing at the mitochondrial level.

Among the dysfunctions through which mitochondria are likely responsible for neuronal demise in AD are those occurring at the ER-mitochondria network level. We discovered that upon agonist stimulation, mitochondria from cell lines and neurons carrying FAD-linked human PS2 mutations experience much higher Ca<sup>2+</sup> transients despite a lower ER Ca<sup>2+</sup> content. The phenomenon is due to an increased number of ER-mitochondria close contact sites occurring at the level of mitochondria-associated ER membranes (MAMs), a finding reported also with PS2 but not PS1 mutants [99][113] [98][114][115]. Interestingly, A $\beta$  can be produced at the MAM level and knocking down mitofusin-2 increases ER-mitochondria contact sites [116] and inhibits A $\beta$  production [117].

Indirect effects on the expression level of VDAC1 and IP3R3, the corresponding ER facing IP3Rs, have also been reported for A $\beta$  at nanomolar concentrations [113]. Indeed, the effect is rapid since a 1-hour incubation with A $\beta$  is sufficient to significantly increase the IP3R3 mRNA level in cultured mouse cortical neurons (**Figure 5**).

How an increased ER-mitochondria coupling will ultimately cause pathological alterations at different levels

such as energy metabolism, lipid synthesis, mitochondria remodelling and apoptotic signalling has yet to be fully investigated.

## 5. THE LYSOSOMAL CHANNELS

A growing body of evidence has linked failure of the autophagic pathway to multiple outcomes in neurodegenerative disease. In AD, lysosomal  $\text{Ca}^{2+}$  storage/release defects accelerate amyloidogenesis, neuritic dystrophy and apoptosis. It is well known that vesicular  $\text{Ca}^{2+}$  is required for efficient vesicle fusion including lysosomal vesicles [118][119]. Lysosomal  $\text{Ca}^{2+}$  defects have been reported in  $\text{PSEN}^{-/-}$  cells and neurons, thus suggesting an additional route through which PS dysfunctions impair cellular proteostasis and autophagic fluxes. Nixon and coworkers have shown that PS1 deficiency causes a primary defect in V-ATPase that determines lysosomal alkalization [120]. Furthermore, the same authors have suggested that anomalous alkalization causes hyperactivation of the transient receptor potential mucolipin-1 (TRPML1) channels located in the lysosomal membrane, thus inducing both organelle  $\text{Ca}^{2+}$  depletion and cytosolic  $\text{Ca}^{2+}$  overload [121]. Whereas in  $\text{PSEN-DKO}$  MEFs and  $\text{PSEN1}^{-/-}$  hippocampal neurons defects in lysosomal  $\text{Ca}^{2+}$  content and fusion were fully confirmed, defective V-ATPase and lysosomal alkalization were not reported [122]. How lysosomal  $\text{Ca}^{2+}$  depletion contributes to AD neuronal dysfunction and death is yet to be fully elucidated. Unfortunately, obtaining reliable measurements of  $\text{Ca}^{2+}$  dynamics inside acidic vesicles is still an unsolved challenge. Regarding AD mouse models based on PS2-N141I, we did not observe any significant effects of mutant PS2 on acidic  $\text{Ca}^{2+}$  pools: the combined use of ionomycin to release neutral stores, followed by monensin to collapse the proton gradient of acidic vesicles and allow ionomycin to release their  $\text{Ca}^{2+}$  content, demonstrated that the acidic pools of PS2-N141I expressing neurons have the same  $\text{Ca}^{2+}$  content of neurons from control mice [99]. Notwithstanding, small pH differences in acidic pools cannot be detected by this indirect approach.

## 6. CONCLUDING REMARKS

The brain consists of a vast array of interconnected neuronal and astrocytic cells. As such, the aforementioned intracellular  $\text{Ca}^{2+}$  dysregulations, when spread across different cell types and brain areas, can have a prominent effect on overall brain performance. Recent technical advances in  $\text{Ca}^{2+}$  imaging at the *in vivo* level, together with a deeper knowledge about the role of the different cell populations, have enabled us to look at the brain as a whole, spotting the effects of the subtle intracellular defects at the network level [123]. We also reported an increased susceptibility to a pro-epileptogenic protocol employing picrotoxin in acute slices of two-week old AD transgenic mice based on mutant PS2 [99], therefore highlighting the possibility that hyper-excitability and hyper-synchronicity are early network deficits that precede amyloid deposition [124]. Other works, focusing their attention on the astrocytic network of the APP<sup>swe</sup>/PS1 $\Delta$ E9 AD mice and imaging  $\text{Ca}^{2+}$  waves *in vivo*, have shown that at 4-5 months of age astrocytes display an elevated resting  $\text{Ca}^{2+}$  level independent of plaque proximity. Nonetheless, spontaneous intracellular  $\text{Ca}^{2+}$  waves were observed in astrocytes close to

plaques, while they were able to spread over a distance of at least 200  $\mu\text{m}$  [125]. The same authors found that neurons in close vicinity of  $\text{A}\beta$  plaques are characterized by an increased activity driven by  $\text{Ca}^{2+}$  dyshomeostasis and affecting both structure and function that leads to a final disruption of the neuronal network [126]. In the frontal cortex of APP23xPS45 mice, hyperactive neurons were found in proximity of plaques but without a direct contact with them, suggesting a halo effect of soluble intermediates [127]. Indeed, the same authors demonstrated a massive increase in overall hippocampal activity and an altered ratio between hyperactive and hypoactive hippocampal neurons in young APP23xPS45 mice in the absence of plaques, thus concluding that neuronal hyperactivity is an early neuronal dysfunction of AD mouse models [128]. The noxious stimulus is likely represented by  $\text{A}\beta_0$  such as dimers, trimers and higher order oligomers, species that exist before plaque formation and to whom plaques, once formed, represent a reservoir. In fact, local application of soluble  $\text{A}\beta_0$  in the form of synthetic dimers mimicked hyperactivity of CA1 neurons in wild-type mice *in vivo* [128]. However, these works do not distinguish the role of excitatory pyramidal neurons and inhibitory interneurons, an issue of profound importance when it comes to specific circuitry functionality. It is worth noting that AD patients show a 5- to 10-fold increased probability of experiencing epileptic seizures, which correlate with the progression of the cognitive decline [129][130]. The degree of neuronal activity might influence the level of  $\text{A}\beta$  plaques deposition: it has recently been reported that cortical areas associated with elevated basal activity are more prone to develop amyloid deposits, and that the active unilateral vibrissae hyper-stimulation causes an increased plaque burden in the controlateral barrel cortex [131].

Increased neuronal hyper-excitability was observed in hippocampal slices not only from PS2APP mice but also from single mutant PS2 mice, *i.e.*, with an extremely low level of  $\text{A}\beta$ -load. Our data suggest that either a few pg/mg of wet tissue of  $\text{A}\beta_{42}$  are sufficient to eliciting neuronal hyperactivity or that the ER  $\text{Ca}^{2+}$  defects themselves, due to the mutant PS2, are the initiating cause of altered network activity [99]. Curiously ER  $\text{Ca}^{2+}$  defects also characterize human fibroblasts obtained from FAD PS2-T122R twin patients, one of the two in the total absence of cognitive decline [7]. Taken together, these findings indicate that  $\text{Ca}^{2+}$  dysregulation, particularly at the store level, occurs precociously, likely accelerating the disease progression and could be instrumental to find early biomarkers of the disease onset.

At the brain network level, the concerted activity of excitatory and inhibitory neurons originates signals oscillating in different frequency domains. The correct storage and retrieval of memories in the hippocampus strongly relies on this rhythmic activity, which is most prominently featured by the so-called theta (4-12 Hz) and gamma (25-100 Hz) oscillations. Theta and gamma rhythms can interact in a way that is known as cross-frequency coupling, where the phase of the theta oscillation influences the amplitude of gamma wave. This synchronicity mechanism underlies the formation of the working memory, a functionality that is severely impaired in the AD patient [132][133]. These aspects of brain activity can be measured using a variety of methods, even on human subjects, but, in spite of their relevance, very little work has been carried out on this topic. It has recently been reported that, prior to plaque deposition, APP23 mice display a deeply altered rhythmic pattern, with an overall increased burst activity, met by a decreased theta rhythm power, an increased gamma component and an impaired cross-frequency coupling [134]. Both the balance between the two rhythms

and their coupling were influenced by riluzole, that blocks voltage-gated sodium channels, and MK-801, a NMDA receptor blocker, thus highlighting the importance of the activity-evoked  $\text{Ca}^{2+}$  signal [134]. Taken together, these results suggest how intracellular events linked to  $\text{Ca}^{2+}$  dysregulation, which affect neurons as well as other cell types in the brain, can ultimately lead to a diffuse dyshomeostasis on different activities, causing structural and functional damages that eventually result in brain dysfunction and cognitive decline.

#### **Disclosure statement**

The authors declare no conflict of interest.

#### **Acknowledgements**

We are grateful to P. Magalhães, R. Fontana and T. Pozzan for support and critical discussion and to C. Lazzari and V. Gobbo for executing the qPCR and TEM experiments respectively. This work was supported by the Italian Ministry of University and Scientific Research and the University of Padua (FASO-PRAT16-01 to CF); the Strategic Projects of the University of Padua and the Italian National Research Council, CNR (Ageing project to TP). M.A. is supported by a Ph.D. fellowship from the University of Padua.

## FIGURE LEGENDS:

**FIGURE 1. A $\beta$  aggregation pathways.** A $\beta$ 42 can aggregate following different pathways that lead to different, albeit coexisting, products [121]. The left side shows a trimer-based aggregation process resulting in soluble aggregates of Low and High Molecular Weight (A $\beta$ o) that are responsible for a vast array of cellular effects [122]. The right side shows the pathway leading to the formation of insoluble fibrils. Both pathway eventually can lead to plaques. Modified from [135].

**FIGURE 2. Synaptic A $\beta$ o targets.** A $\beta$ o impair the functionality of several targets localized at the synaptic level. We have here reviewed some of the major pathways involving Ca<sup>2+</sup> dynamics, among which: metabotropic glutamate receptor (mGluR), store-operated Ca<sup>2+</sup> channel (SOCC), cellular prion protein (PrP<sup>C</sup>), N-methyl-D-aspartate ionotropic glutamate receptor (NMDAR), nicotinic acetylcholine receptor ( $\alpha$ 7-nAChR), voltage-operated Ca<sup>2+</sup> channel (VOCC). Other targets less directly involved on Ca<sup>2+</sup> homeostasis such as the neuroligin-1 (NL-1) adhesion molecule and the low affinity Nerve Growth Factor receptor (p75), have not been included but see refs [123]. Only receptors located at the postsynaptic site are here highlighted; however, also targets at the presynaptic site should be considered to fully explain A $\beta$ o toxicity. Modified from [136].

**FIGURE 3. Preparation of synthetic A $\beta$ (1-42) monomers and oligomers.** Human synthetic dry powdered A $\beta$ (1-42) (Anaspec, Fremont, CA, USA) is dissolved in 1,1,1,3,3,3-hexafluoro-2-isopropanol (HFIP) at room temperature (RT) for 1 hour to final concentration of 200  $\mu$ M and stored as 10-20  $\mu$ l aliquots at -80°C until use. A $\beta$ 42o and A $\beta$ 42m are prepared according to the following procedure: aliquots are spun for 10 minutes (13,000 rpm, 5 min, 4°C) and then the HFIP is carefully removed by air fluxing with a pipette for several times and allowed to dry for about 10 minutes. The dried HFIP film is dissolved in DMSO (500  $\mu$ M) and then diluted (50  $\mu$ M) in double distilled water. When used immediately, the stock solution contained mainly monomers and low molecular weight (LMW) oligomers (6-17 kDa) whereas when left at RT for 48-72 hours it also contained high molecular weight (HMW) oligomers (50-100 kDa) that disappeared upon boiling [124] as shown from the Western blot on the right side. Since no quantification of HMW oligomers is carried out, the level of oligomers in the mixture is expressed as the equivalent concentration of monomers. Cells are routinely exposed to either A $\beta$ 42m or A $\beta$ 42o (0.5  $\mu$ M) by adding 5  $\mu$ L of the stock solution (50  $\mu$ M, 10% DMSO) to 0.5 mL of the bathing medium. Before cell treatment, the A $\beta$ 42 stock solution was spun (13,000 rpm, 10 min) and immediately used. Preparations are tested by transmission electron microscopy (TEM). Images are acquired on a Tecnai 12 microscope (FEI, Hillsboro, Oregon, USA) operating at 100kV. The stock A $\beta$ 42 solution (50  $\mu$ M) of either oligomers or monomers is diluted 5 times in bi-distilled water, and 20  $\mu$ l samples are placed on Formvar/carbon-coated nickel grids and blotted off after 2 minutes. Samples are negatively stained with uranyl acetate and observed at 90,000X magnifications. A $\beta$ 42o particle diameter mainly ranges from 5 to 15 nm, consistent with a HMW oligomer population; A $\beta$ 42m and LMW aggregates are not revealed by TEM [125].

**FIGURE 4. Characterization of A $\beta$ o by Western blot.** Oligomeric preparations are separated by gel electrophoresis using a tris-tricine 4-10-16% polyacrilamide gel. A $\beta$ 42o are diluted in loading sample buffer

(LSB; in mM: 50 Tris, 50 tricine, 2% SDS, 12% glycerol, 80 dithiothreitol, pH 8.25 at RT) at the final concentration of 20 ng/μl. Samples (200-400 ng) were loaded in LSB on a 4-10-16% (w/v) Tris-Tricine discontinuous gel in Tris-Tricine buffer with or without a boiling step (95°C, 10 min). Nitrocellulose membranes are saturated 1 hour at RT with a blocking solution containing 5% (w/v) non-fat dry milk (BioRad) in PBS plus Tween-20 [PBS-T; in mM: 136.89 NaCl, 6.7 KCl, 20 Na<sub>2</sub>HPO<sub>4</sub>, 3.67 KH<sub>2</sub>PO<sub>4</sub>, 0.1% (w/v) Tween-20, pH 7.4], washed twice with PBS-T (10 min) and incubated 24 hours at 4°C with the primary antibody (6E10, Covance). The sample composition at 96 hours does not differ from that at 48 hours and it comprises monomers, dimers, trimers, tetramers and HMW oligomers, which are disrupted, alongside with the majority of tetramers and trimers, upon a 10-minute 95°C boiling procedure. See the enlarged monomer and dimer immuno-band in the boiled sample.

**FIGURE 5. Quantitative reverse-transcription polymerase chain reaction (qPCR).** Cortical neuronal cultures at DIV 11-13, seeded in 12-well culture vessels are washed twice with mKRB and then incubated for 1 h at 37°C with Aβ<sub>42</sub>o (0.5 μM) or vehicle (0.1% DMSO) [30]. Total RNA was extracted using the NucleoSpin RNA purification kit (Macherey-Nagel) and then quantified with NanoDrop 2000 (Thermo Scientific). For each sample, 500 ng of total RNA were reverse-transcribed into cDNA using the SuperScript II Reverse Transcriptase (Invitrogen) following the manufacturer's instructions. PCR amplification was performed in an iQ5 Thermal Cycler (BioRad), using SYBR Green Supermix (BioRad) as reagent. For each cDNA, the efficiency of the reaction was estimated with a standard curve by using the cDNA reverse transcript from mouse brain. The transcript of Hypoxanthine phosphoribosyl-transferase 1 (HPRT1) was used as internal control. The following primers (Invitrogen) were used:

RyR2: 5'-GGCTTGCTCCAGATGAAACT-3' and 5'-CTTCGATCCGGTGCCTAAC-3';

RyR3: 5'-CAAACCTGCCCTCCCTAAACA-3' and 5'-AAACACGGTCCAAGAACAGC-3';

IP3R isoform 1: FW: 5'-CTCATGGACTGATTATGGACAGGAC-3'

and 5'-GCAGGTCAGCAAAGAACTTATAGCC-3';

IP3R- 3: 5'-ATAAGAAGGAGAGACCCTCGG-3' and 5'-CCACACATCTTGTTTCAGCCT-3';

HPRT1: 5'-CTACAGAGTGCCTGACCT-3' and 5'-TCGTAGATCCCTCCGCATAC-3'.

Quantification followed the method of ΔCt comparison. Data are expressed as percentage ratio change relative to the HPRT1 transcript (mean ± SEM, n = number of wells).

FIGURE 1




FIGURE 2






FIGURE 3




FIGURE 4





FIGURE 5



## REFERENCES

- [1] J. Hardy, D. Allsop, Amyloid deposition as the central event in the aetiology of Alzheimer's disease, *Trends Pharmacol. Sci.*, 12 (1991) 383-388.
- [2] Z.S. Khachaturian, Hypothesis on the regulation of cytosol calcium concentration and the aging brain, *Neurobiol. Aging*, 8 (1987) 345-346.
- [3] J.A. Hardy, G.A. Higgins, Alzheimer's disease: the amyloid cascade hypothesis, *Science*, 256 (1992) 184-185.
- [4] C. Haass, D.J. Selkoe, Soluble protein oligomers in neurodegeneration: lessons from the Alzheimer's amyloid  $\beta$ -peptide, *Nat. Rev. Mol. Cell Biol.*, 8 (2007) 101-112.
- [5] J. Hardy, D.J. Selkoe, The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics, *Science*, 297 (2002) 353-356.
- [6] S. Lesne, M.T. Koh, L. Kotilinek, R. Kaye, C.G. Glabe, A. Yang, M. Gallagher, K.H. Ashe, A specific amyloid-beta protein assembly in the brain impairs memory, *Nature*, 440 (2006) 352-357.
- [7] M. Giacomello, L. Barbiero, G. Zatti, R. Squitti, G. Binetti, T. Pozzan, C. Fasolato, R. Ghidoni, P. Pizzo, Reduction of  $\text{Ca}^{2+}$  stores and capacitative  $\text{Ca}^{2+}$  entry is associated with the familial Alzheimer's disease presenilin-2 T122R mutation and anticipates the onset of dementia, *Neurobiol. Dis.*, 18 (2005) 638-648.
- [8] D. Liao, E.C. Miller, P.J. Teravskis, Tau acts as a mediator for Alzheimer's disease-related synaptic deficits, *Eur. J. Neurosci.*, 39 (2014) 1202-1213.
- [9] U.A. Khan, L. Liu, F.A. Provenzano, D.E. Berman, C.P. Profaci, R. Sloan, R. Mayeux, K.E. Duff, S.A. Small, Molecular drivers and cortical spread of lateral entorhinal cortex dysfunction in preclinical Alzheimer's disease, *Nat. Neurosci.*, 17 (2014) 304-311.
- [10] N. Arispe, J.C. Diaz, O. Simakova,  $\text{A}\beta$  ion channels. Prospects for treating Alzheimer's disease with  $\text{A}\beta$  channel blockers, *Biochim. Biophys. Acta*, 1768 (2007) 1952-1965.
- [11] E. Alberdi, M.V. Sanchez-Gomez, F. Cavaliere, A. Perez-Samartin, J.L. Zugaza, R. Trullas, M. Domercq, C. Matute, Amyloid beta oligomers induce  $\text{Ca}^{2+}$  dysregulation and neuronal death through activation of ionotropic glutamate receptors, *Cell Calcium*, 47 (2010) 264-272.
- [12] A. Demuro, E. Mina, R. Kaye, S.C. Milton, I. Parker, C.G. Glabe, Calcium dysregulation and membrane disruption as a ubiquitous neurotoxic mechanism of soluble amyloid oligomers, *J. Biol. Chem.*, 280 (2005) 17294-17300.
- [13] R. Resende, E. Ferreira, C. Pereira, C. Resende de Oliveira, Neurotoxic effect of oligomeric and fibrillar species of amyloid-beta peptide 1-42: involvement of endoplasmic reticulum calcium release in oligomer-induced cell death, *Neuroscience*, 155 (2008) 725-737.
- [14] N. Arispe, J. Diaz, S.R. Durell, Y. Shafir, H.R. Guy, Polyhistidine peptide inhibitor of the  $\text{A}\beta$  calcium channel potently blocks the  $\text{A}\beta$ -induced calcium response in cells. Theoretical modeling suggests a cooperative binding process, *Biochemistry*, 49 (2010) 7847-7853.
- [15] O. Simakova, N.J. Arispe, Early and late cytotoxic effects of external application of the Alzheimer's A $\beta$  result from the initial formation and function of A $\beta$  ion channels, *Biochemistry*, 45 (2006) 5907-5915.
- [16] A. Demuro, I. Parker, Cytotoxicity of intracellular  $\text{A}\beta$ 42 amyloid oligomers involves  $\text{Ca}^{2+}$  release from the endoplasmic reticulum by stimulated production of inositol trisphosphate, *J. Neurosci.*, 33 (2013) 3824-3833.
- [17] Y. Sokolov, J.A. Kozak, R. Kaye, A. Chanturiya, C. Glabe, J.E. Hall, Soluble amyloid oligomers increase bilayer conductance by altering dielectric structure, *J. Gen. Physiol.*, 128 (2006) 637-647.
- [18] A. Adunsky, D. Baram, M. Hershkowitz, Y.A. Mekori, Increased cytosolic free calcium in lymphocytes of Alzheimer patients, *J. Neuroimmunol.*, 33 (1991) 167-172.
- [19] J.C. Gant, M.M. Sama, P.W. Landfield, O. Thibault, Early and simultaneous emergence of multiple hippocampal biomarkers of aging is mediated by  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release, *Journal of Neuroscience*, 26 (2006) 3482-3490.

- [20] P.W. Landfield, Increased hippocampal  $\text{Ca}^{2+}$  channel activity in brain aging and dementia. Hormonal and pharmacologic modulation, *Ann. N.Y. Acad. Sci.*, 747 (1994) 351-364.
- [21] O. Thibault, R. Hadley, P.W. Landfield, Elevated postsynaptic  $[\text{Ca}^{2+}]_i$  and L-type calcium channel activity in aged hippocampal neurons: relationship to impaired synaptic plasticity, *Journal of Neuroscience*, 21 (2001) 9744-9756.
- [22] G.E. Stutzmann, I. Smith, A. Caccamo, S. Oddo, F.M. Laferla, I. Parker, Enhanced ryanodine receptor recruitment contributes to  $\text{Ca}^{2+}$  disruptions in young, adult, and aged Alzheimer's disease mice, *J. Neurosci.*, 26 (2006) 5180-5189.
- [23] M.J. Berridge, Calcium hypothesis of Alzheimer's disease, *Pflugers Arch.*, 459 (2010) 441-449.
- [24] L.C. Reese, G. Tagliavola, A role for calcineurin in Alzheimer's disease, *Curr. Neuropharmacol.*, 9 (2011) 685-692.
- [25] A. Kapur, M.F. Yeckel, R. Gray, D. Johnston, L-Type calcium channels are required for one form of hippocampal mossy fiber LTP, *J. Neurophysiol.*, 79 (1998) 2181-2190.
- [26] A.L. Coon, D.R. Wallace, C.F. Mactutus, R.M. Booze, L-type calcium channels in the hippocampus and cerebellum of Alzheimer's disease brain tissue, *Neurobiol. Aging*, 20 (1999) 597-603.
- [27] F. Forette, M.L. Seux, J.A. Staessen, L. Thijs, M.R. Babarskiene, S. Babeanu, A. Bossini, R. Fagard, B. Gil-Extremera, T. Laks, Z. Kobalava, C. Sarti, J. Tuomilehto, H. Vanhanen, J. Webster, Y. Yodfat, W.H. Birkenhager, S.-E. Investigators, The prevention of dementia with antihypertensive treatment, *Arch. Intern. Med.*, 162 (2002) 2046-2052.
- [28] L.M. Veng, M.H. Mesches, M.D. Browning, Age-related working memory impairment is correlated with increases in the L-type calcium channel protein  $\alpha_{1D}$  ( $\text{Ca}_v1.3$ ) in area CA1 of the hippocampus and both are ameliorated by chronic nimodipine treatment, *Brain Res. Mol. Brain Res.*, 110 (2003) 193-202.
- [29] J. Espana, J. Valero, A.J. Minano-Molina, R. Masgrau, E. Martin, C. Guardia-Laguarta, A. Lleo, L. Gimenez-Llort, J. Rodriguez-Alvarez, C.A. Saura,  $\beta$ -Amyloid Disrupts Activity-Dependent Gene Transcription Required for Memory through the CREB Coactivator CRTCL1, *Journal of Neuroscience*, 30 (2010) 9402-9410.
- [30] F. Facchinetti, C. Fasolato, E. Del Giudice, A. Burgo, S. Furegato, M. Fusco, E. Basso, R. Seraglia, A. D'Arrigo, A. Leon, Nimodipine selectively stimulates  $\beta$ -amyloid 1-42 secretion by a mechanism independent of calcium influx blockage, *Neurobiol. Aging*, 27 (2006) 218-227.
- [31] C. Lazzari, M.J. Kipanyula, M. Agostini, T. Pozzan, C. Fasolato,  $\text{A}\beta_{42}$  oligomers selectively disrupt neuronal calcium release, *Neurobiol. Aging*, 36 (2015) 877-885.
- [32] V. Nimmrich, C. Grimm, A. Draguhn, S. Barghorn, A. Lehmann, H. Schoemaker, H. Hillen, G. Gross, U. Ebert, C. Bruehl, Amyloid beta oligomers (A  $\beta_{1-42}$ ) globulomer suppress spontaneous synaptic activity by inhibition of P/Q-type calcium currents, *J. Neurosci.*, 28 (2008) 788-797.
- [33] K.J. Gan, M.A. Silverman, Dendritic and axonal mechanisms of  $\text{Ca}^{2+}$  elevation impair BDNF transport in  $\text{A}\beta$  oligomer-treated hippocampal neurons, *Mol. Biol. Cell.*, 26 (2015) 1058-1071.
- [34] E.X. Albuquerque, E.F. Pereira, M. Alkondon, S.W. Rogers, Mammalian nicotinic acetylcholine receptors: from structure to function, *Physiol. Rev.*, 89 (2009) 73-120.
- [35] D. Puzzo, W. Gulisano, O. Arancio, A. Palmeri, The Keystone of Alzheimer Pathogenesis Might Be Sought in  $\text{A}\beta$  Physiology, *Neuroscience*, 307 (2015) 26-36.
- [36] J. Shen, J. Wu, Nicotinic Cholinergic Mechanisms in Alzheimer's Disease, *Int. Rev. Neurobiol.*, 124 (2015) 275-292.
- [37] J. Wu, Q. Liu, P. Tang, J.D. Mikkelsen, J. Shen, P. Whiteaker, J.L. Yakel, Heteromeric  $\alpha 7\beta 2$  Nicotinic Acetylcholine Receptors in the Brain, *Trends Pharmacol. Sci.*, 37 (2016) 562-574.

- [38] H.Y. Wang, D.H. Lee, M.R. D'Andrea, P.A. Peterson, R.P. Shank, A.B. Reitz,  $\beta$ -Amyloid<sub>1-42</sub> binds to  $\alpha_7$  nicotinic acetylcholine receptor with high affinity. Implications for Alzheimer's disease pathology, *J. Biol. Chem.*, 275 (2000) 5626-5632.
- [39] S. Lombardo, U. Maskos, Role of the nicotinic acetylcholine receptor in Alzheimer's disease pathology and treatment, *Neuropharmacology*, 96 (2015) 255-262.
- [40] G. Dziewczapolski, C.M. Glogowski, E. Masliah, S.F. Heinemann, Deletion of the  $\alpha_7$  nicotinic acetylcholine receptor gene improves cognitive deficits and synaptic pathology in a mouse model of Alzheimer's disease, *Journal of Neuroscience*, 29 (2009) 8805-8815.
- [41] C.M. Hernandez, R. Kaye, H. Zheng, J.D. Sweatt, K.T. Dineley, Loss of  $\alpha_7$  nicotinic receptors enhances  $\beta$ -amyloid oligomer accumulation, exacerbating early-stage cognitive decline and septohippocampal pathology in a mouse model of Alzheimer's disease, *Journal of Neuroscience*, 30 (2010) 2442-2453.
- [42] L. Lee, P. Kosuri, O. Arancio, Picomolar amyloid- $\beta$  peptides enhance spontaneous astrocyte calcium transients, *J. Alzheimers Dis.*, 38 (2014) 49-62.
- [43] T.M. Pirttimaki, N.K. Codadu, A. Awani, P. Pratik, D.A. Nagel, E.J. Hill, K.T. Dineley, H.R. Parri,  $\alpha_7$  Nicotinic receptor-mediated astrocytic gliotransmitter release: A $\beta$  effects in a preclinical Alzheimer's mouse model, *PLoS One*, 8 (2013) e81828.
- [44] F.G. De Felice, P.T. Velasco, M.P. Lambert, K. Viola, S.J. Fernandez, S.T. Ferreira, W.L. Klein, Abeta oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine, *J. Biol. Chem.*, 282 (2007) 11590-11601.
- [45] H. Decker, K.Y. Lo, S.M. Unger, S.T. Ferreira, M.A. Silverman, Amyloid- $\beta$  peptide oligomers disrupt axonal transport through an NMDA receptor-dependent mechanism that is mediated by glycogen synthase kinase 3 $\beta$  in primary cultured hippocampal neurons, *Journal of Neuroscience*, 30 (2010) 9166-9171.
- [46] J. Lauren, D.A. Gimbel, H.B. Nygaard, J.W. Gilbert, S.M. Strittmatter, Cellular prion protein mediates impairment of synaptic plasticity by amyloid- $\beta$  oligomers, *Nature*, 457 (2009) 1128-1134.
- [47] J.W. Um, H.B. Nygaard, J.K. Heiss, M.A. Kostylev, M. Stagi, A. Vortmeyer, T. Wisniewski, E.C. Gunther, S.M. Strittmatter, Alzheimer amyloid-beta oligomer bound to postsynaptic prion protein activates Fyn to impair neurons, *Nat. Neurosci.*, 15 (2012) 1227-1235.
- [48] J.W. Um, A.C. Kaufman, M. Kostylev, J.K. Heiss, M. Stagi, H. Takahashi, M.E. Kerrisk, A. Vortmeyer, T. Wisniewski, A.J. Koleske, E.C. Gunther, H.B. Nygaard, S.M. Strittmatter, Metabotropic glutamate receptor 5 is a coreceptor for Alzheimer  $\alpha\beta$  oligomer bound to cellular prion protein, *Neuron*, 79 (2013) 887-902.
- [49] H. Khosravani, Y.F. Zhang, S. Tsutsui, S. Hameed, C. Altier, J. Hamid, L.N. Chen, M. Villemain, Z. Ali, F.R. Jirik, G.W. Zamponi, Prion protein attenuates excitotoxicity by inhibiting NMDA receptors, *Journal of Cell Biology*, 181 (2008) 551-565.
- [50] H.T. You, S. Tsutsui, S. Hameed, T.J. Kannanayakal, L.N. Chen, P. Xia, J.D.T. Engbers, S.A. Lipton, P.K. Stys, G.W. Zamponi, A $\beta$  neurotoxicity depends on interactions between copper ions, prion protein, and N-methyl-D-aspartate receptors, *Proc. Natl. Acad. Sci. U.S.A.*, 109 (2012) 1737-1742.
- [51] T. Rush, A. Buisson, Reciprocal disruption of neuronal signaling and Abeta production mediated by extrasynaptic NMDA receptors: a downward spiral, *Cell Tissue Res.*, 356 (2014) 279-286.
- [52] H.L. Scott, D.V. Pow, A.E.G. Tannenberg, P.R. Dodd, Aberrant expression of the glutamate transporter excitatory amino acid transporter 1 (EAAT1) in Alzheimer's disease, *Journal of Neuroscience*, 22 (2002) Vol. 22 RC206.
- [53] A.C. Paula-Lima, J. Brito-Moreira, S.T. Ferreira, Deregulation of excitatory neurotransmission underlying synapse failure in Alzheimer's disease, *J. Neurochem.*, 126 (2013) 191-202.

- [54] Y.T. Liu, T.P. Wong, M. Aarts, A. Rooyackers, L.D. Liu, T.W. Lai, D.C. Wu, J. Lu, M. Tymianski, A.M. Craig, Y.T. Wang, NMDA receptor subunits have differential roles in mediating excitotoxic neuronal death both in vitro and in vivo, *Journal of Neuroscience*, 27 (2007) 2846-2857.
- [55] R. Ronicke, M. Mikhaylova, S. Ronicke, J. Meinhardt, U.H. Schroder, M. Fandrich, G. Reiser, M.R. Kreutz, K.G. Reymann, Early neuronal dysfunction by amyloid  $\beta$  oligomers depends on activation of NR2B-containing NMDA receptors, *Neurobiol. Aging*, 32 (2011) 2219-2228.
- [56] J.E. Hanson, J.F. Pare, L. Deng, Y. Smith, Q. Zhou, Altered GluN2B NMDA receptor function and synaptic plasticity during early pathology in the PS2APP mouse model of Alzheimer's disease, *Neurobiol. Dis.*, 74 (2015) 254-262.
- [57] S. Jurgensen, L.L. Antonio, G.E. Mussi, J. Brito-Moreira, T.R. Bomfim, F.G. De Felice, E.R. Garrido-Sanabria, E.A. Cavaleiro, S.T. Ferreira, Activation of D1/D5 dopamine receptors protects neurons from synapse dysfunction induced by amyloid- $\beta$  oligomers, *J. Biol. Chem.*, 286 (2011) 3270-3276.
- [58] P. Kurup, Y. Zhang, J. Xu, D.V. Venkitaramani, V. Haroutunian, P. Greengard, A.C. Nairn, P.J. Lombroso, A $\beta$ -mediated NMDA receptor endocytosis in Alzheimer's disease involves ubiquitination of the tyrosine phosphatase STEP61, *Journal of Neuroscience*, 30 (2010) 5948-5957.
- [59] A.J. Minano-Molina, J. Espana, E. Martin, B. Barneda-Zahonero, R. Fado, M. Sole, R. Trullas, C.A. Saura, J. Rodriguez-Alvarez, Soluble oligomers of amyloid- $\beta$  peptide disrupt membrane trafficking of  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor contributing to early synapse dysfunction, *J. Biol. Chem.*, 286 (2011) 27311-27321.
- [60] J.H. Birnbaum, J. Bali, L. Rajendran, R.M. Nitsch, C. Tackenberg, Calcium flux-independent NMDA receptor activity is required for A $\beta$  oligomer-induced synaptic loss, *Cell Death Dis.*, 6 (2015).
- [61] M.P. Lambert, A.K. Barlow, B.A. Chromy, C. Edwards, R. Freed, M. Liosatos, T.E. Morgan, I. Rozovsky, B. Trommer, K.L. Viola, P. Wals, C. Zhang, C.E. Finch, G.A. Krafft, W.L. Klein, Diffusible, nonfibrillar ligands derived from A $\beta$ 1-42 are potent central nervous system neurotoxins, *Proc. Natl. Acad. Sci. U.S.A.*, 95 (1998) 6448-6453.
- [62] H. Zempel, E. Thies, E. Mandelkow, E.M. Mandelkow, A $\beta$  oligomers cause localized Ca<sup>2+</sup> elevation, missorting of endogenous Tau into dendrites, Tau phosphorylation, and destruction of microtubules and spines, *J. Neurosci.*, 30 (2010) 11938-11950.
- [63] E. Caballero, M. Calvo-Rodriguez, A. Gonzalo-Ruiz, C. Villalobos, L. Nunez, A new procedure for amyloid beta oligomers preparation enables the unambiguous testing of their effects on cytosolic and mitochondrial Ca<sup>2+</sup> entry and cell death in primary neurons, *Neurosci. Lett.*, 612 (2016) 66-73.
- [64] D. Lim, J.J. Rodriguez-Arellano, V. Parpura, R. Zorec, F. Zeidan-Chulia, A.A. Genazzani, A. Verkhratsky, Calcium signalling toolkits in astrocytes and spatio-temporal progression of Alzheimer's disease, *Curr. Alzheimer Res.*, 13 (2016) 359-369.
- [65] B. De Strooper, E. Karran, The Cellular Phase of Alzheimer's Disease, *Cell*, 164 (2016) 603-615.
- [66] U. Dreses-Werringloer, J.C. Lambert, V. Vingtdeux, H.T. Zhao, H. Vais, A. Siebert, A. Jain, J. Koppel, A. Rovelet-Lecrux, D. Hannequin, F. Pasquier, D. Galimberti, E. Scarpini, D. Mann, C. Lendon, D. Campion, P. Amouyel, P. Davies, J.K. Foskett, F. Campagne, P. Marambaud, A polymorphism in CALHM1 influences Ca<sup>2+</sup> homeostasis, A $\beta$  levels, and Alzheimer's disease risk, *Cell*, 133 (2008) 1149-1161.
- [67] A. Taruno, V. Vingtdeux, M. Ohmoto, Z.M. Ma, G. Dvoryanchikov, A. Li, L. Adrien, H.T. Zhao, S. Leung, M. Abernethy, J. Koppel, P. Davies, M.M. Civan, N. Chaudhari, I. Matsumoto, G. Hellekant, M.G. Tordoff, P. Marambaud, J.K. Foskett, CALHM1 ion channel mediates purinergic neurotransmission of sweet, bitter and umami tastes, *Nature*, 495 (2013) 223-226.
- [68] Z. Ma, A.P. Siebert, K.H. Cheung, R.J. Lee, B. Johnson, A.S. Cohen, V. Vingtdeux, P. Marambaud, J.K. Foskett, Calcium homeostasis modulator 1 (CALHM1) is the pore-forming

- subunit of an ion channel that mediates extracellular  $\text{Ca}^{2+}$  regulation of neuronal excitability, *Proceedings of the National Academy of Sciences of the United States of America* 109 (2012) E1963-1971.
- [69] Z. Ma, J.E. Tanis, A. Taruno, J.K. Foskett, Calcium homeostasis modulator (CALHM) ion channels, *Pflugers Arch.*, 468 (2016) 395-403.
- [70] S. Gallego-Sandin, M.T. Alonso, J. Garcia-Sancho, Calcium homeostasis modulator 1 (CALHM1) reduces the calcium content of the endoplasmic reticulum (ER) and triggers ER stress, *Biochem. J.*, 437 (2011) 469-475.
- [71] A.J. Moreno-Ortega, F.J. Martinez-Sanz, R. Lajarin-Cuesta, C. de los Rios, M.F. Cano-Abad, Benzothiazepine CGP37157 and its 2'-isopropyl analogue modulate  $\text{Ca}^{2+}$  entry through CALHM1, *Neuropharmacology*, 95 (2015) 503-510.
- [72] J. Koppel, F. Campagne, V. Vingtdeux, U. Dreses-Werringloer, M. Ewers, D. Rujescu, H. Hampel, M.L. Gordon, E. Christen, J. Chapuis, B.S. Greenwald, P. Davies, P. Marambaud, CALHM1 P861 Polymorphism Modulates CSF A $\beta$  Levels in Cognitively Healthy Individuals at Risk for Alzheimer's Disease, *Mol. Med.*, 17 (2011) 974-979.
- [73] V. Vingtdeux, P. Chandakkar, H.T. Zhao, L. Blanc, S. Ruiz, P. Marambaud, CALHM1 ion channel elicits amyloid- $\beta$  clearance by insulin-degrading enzyme in cell lines and in vivo in the mouse brain, *J. Cell Sci.*, 128 (2015) 2330-2338.
- [74] F. Moccia, E. Zuccolo, T. Soda, F. Tanzi, G. Guerra, L. Mapelli, F. Lodola, E. D'Angelo, Stim and Orai proteins in neuronal  $\text{Ca}^{2+}$  signaling and excitability, *Front. Cell. Neurosci.*, 9 (2015) 153.
- [75] R. Hooper, B.S. Rothberg, J. Soboloff, Neuronal STIMulation at rest, *Sci. Signal.*, 7 (2014) pe18.
- [76] R. Kraft, STIM and ORAI proteins in the nervous system, *Channels*, (2015) 235-243.
- [77] J.W. Putney, Jr., C.M. Ribeiro, Signaling pathways between the plasma membrane and endoplasmic reticulum calcium stores, *Cell. Mol. Life Sci.*, 57 (2000) 1272-1286.
- [78] M.J. Berridge, Calcium regulation of neural rhythms, memory and Alzheimer's disease, *J. Physiol.*, 592 (2014) 281-293.
- [79] A. Shmigol, S. Kirischuk, P. Kostyuk, A. Verkhratsky, Different properties of caffeine-sensitive  $\text{Ca}^{2+}$  stores in peripheral and central mammalian neurones, *Pflugers Arch.*, 426 (1994) 174-176.
- [80] G. Zatti, A. Burgo, M. Giacomello, L. Barbiero, R. Ghidoni, G. Sinigaglia, C. Florean, S. Bagnoli, G. Binetti, S. Sorbi, P. Pizzo, C. Fasolato, Presenilin mutations linked to familial Alzheimer's disease reduce endoplasmic reticulum and Golgi apparatus calcium levels, *Cell Calcium*, 39 (2006) 539-550.
- [81] J. Lalonde, G. Saia, G. Gill, Store-operated calcium entry promotes the degradation of the transcription factor Sp4 in resting neurons, *Sci. Signal.*, 7 (2014) ra51.
- [82] W. Zeiger, K.S. Vetrivel, V. Buggia-Prevot, P.D. Nguyen, S.L. Wagner, M.L. Villereal, G. Thinakaran,  $\text{Ca}^{2+}$  Influx through Store-operated  $\text{Ca}^{2+}$  Channels Reduces Alzheimer Disease  $\beta$ -Amyloid Peptide Secretion, *Journal of Biological Chemistry*, 288 (2013) 26955-26966.
- [83] K. Honarnejad, J. Herms, Presenilins: role in calcium homeostasis, *Int. J. Biochem. Cell Biol.*, 44 (2012) 1983-1986.
- [84] S.Y. Sun, H. Zhang, J. Liu, E. Popugaeva, N.J. Xu, S. Feske, C.L. White, I. Bezprozvanny, Reduced Synaptic STIM2 Expression and Impaired Store-Operated Calcium Entry Cause Destabilization of Mature Spines in Mutant Presenilin Mice, *Neuron*, 82 (2014) 79-93.
- [85] H. Zhang, L. Wu, E. Pchitskaya, O. Zakharova, T. Saito, T. Saido, I. Bezprozvanny, Neuronal Store-Operated Calcium Entry and Mushroom Spine Loss in Amyloid Precursor Protein Knock-In Mouse Model of Alzheimer's Disease, *Journal of Neuroscience*, 35 (2015) 13275-13286.
- [86] E. Stieren, W.P. Werchan, A. El Ayadi, F. Li, D. Boehning, FAD mutations in amyloid precursor protein do not directly perturb intracellular calcium homeostasis, *PLoS One*, 5 (2010) e11992.

- [87] B. Oules, D. Del Prete, B. Greco, X. Zhang, I. Lauritzen, J. Sevalle, S. Moreno, P. Paterlini-Brechot, M. Trebak, F. Checler, F. Benfenati, M. Chami, Ryanodine receptor blockade reduces amyloid- $\beta$  load and memory impairments in Tg2576 mouse model of Alzheimer disease, *J. Neurosci.*, 32 (2012) 11820-11834.
- [88] J.G. McLarnon, H.B. Choi, L.F. Lue, D.G. Walker, S.U. Kim, Perturbations in calcium-mediated signal transduction in microglia from Alzheimer's disease patients, *J. Neurosci. Res.*, 81 (2005) 426-435.
- [89] E. Alberdi, A. Wyssenbach, M. Alberdi, M.V. Sanchez-Gomez, F. Cavaliere, J.J. Rodriguez, A. Verkhratsky, C. Matute,  $\text{Ca}^{2+}$ -dependent endoplasmic reticulum stress correlates with astrogliosis in oligomeric amyloid  $\beta$ -treated astrocytes and in a model of Alzheimer's disease, *Aging Cell*, 12 (2013) 292-302.
- [90] A. De Mario, A. Castellani, C. Peggion, M.L. Massimino, D. Lim, A.F. Hill, M.C. Sorgato, A. Bertoli, The prion protein constitutively controls neuronal store-operated  $\text{Ca}^{2+}$  entry through Fyn kinase, *Front. Cell. Neurosci.*, 9 (2015) 416.
- [91] L.T. Haas, S.V. Salazar, M.A. Kostylev, J.W. Um, A.C. Kaufman, S.M. Strittmatter, Metabotropic glutamate receptor 5 couples cellular prion protein to intracellular signalling in Alzheimer's disease, *Brain*, 139 (2016) 526-546.
- [92] J. Shen, Function and Dysfunction of Presenilin, *Neurodegenerative Diseases*, 13 (2014) 61-63.
- [93] H. Tu, O. Nelson, A. Bezprozvanny, Z. Wang, S.F. Lee, Y.H. Hao, L. Serneels, B. De Strooper, G. Yu, I. Bezprozvanny, Presenilins form ER  $\text{Ca}^{2+}$  leak channels, a function disrupted by familial Alzheimer's disease-linked mutations, *Cell*, 126 (2006) 981-993.
- [94] D. Shilling, D.O. Mak, D.E. Kang, J.K. Foskett, Lack of evidence for presenilins as endoplasmic reticulum  $\text{Ca}^{2+}$  leak channels, *J. Biol. Chem.*, (2012).
- [95] G. Zatti, R. Ghidoni, L. Barbiero, G. Binetti, T. Pozzan, C. Fasolato, P. Pizzo, The presenilin 2 M239I mutation associated with familial Alzheimer's disease reduces  $\text{Ca}^{2+}$  release from intracellular stores, *Neurobiol. Dis.*, 15 (2004) 269-278.
- [96] L. Fedrizzi, D. Lim, E. Carafoli, M. Brini, Interplay of the  $\text{Ca}^{2+}$ -binding protein dream with presenilin in neuronal  $\text{Ca}^{2+}$  signaling, *J. Biol. Chem.*, (2008).
- [97] L. Brunello, E. Zampese, C. Florean, T. Pozzan, P. Pizzo, C. Fasolato, Presenilin-2 dampens intracellular  $\text{Ca}^{2+}$  stores by increasing  $\text{Ca}^{2+}$  leakage and reducing  $\text{Ca}^{2+}$  uptake, *J. Cell. Mol. Med.*, 13 (2009) 3358-3369.
- [98] E. Zampese, C. Fasolato, M.J. Kipanyula, M. Bortolozzi, T. Pozzan, P. Pizzo, Presenilin 2 modulates endoplasmic reticulum (ER)-mitochondria interactions and  $\text{Ca}^{2+}$  cross-talk, *Proc. Natl. Acad. Sci. U.S.A.*, 108 (2011) 2777-2782.
- [99] M.J. Kipanyula, L. Contreras, E. Zampese, C. Lazzari, A.K.C. Wong, P. Pizzo, C. Fasolato, T. Pozzan,  $\text{Ca}^{2+}$  dysregulation in neurons from transgenic mice expressing mutant presenilin 2, *Aging Cell*, 11 (2012) 885-893.
- [100] K. Cheung, L. Mei, Daniel Mak D-O, I. Hayashi, T. Iwatsubo, D. Kang, J. Foskett, "Gain-of-Function Enhancement of IP3 Receptor Modal Gating by Familial Alzheimer's Disease Linked Presenilin Mutants in Human Cells and Mouse Neurons, *Sci. Signal.*, 3 (2010).
- [101] D. Shilling, M. Muller, H. Takano, D.O. Mak, T. Abel, D.A. Coulter, J.K. Foskett, Suppression of  $\text{InsP}_3$  receptor-mediated  $\text{Ca}^{2+}$  signaling alleviates mutant presenilin-linked familial Alzheimer's disease pathogenesis, *Journal of Neuroscience*, 34 (2014) 6910-6923.
- [102] D. Del Prete, F. Checler, M. Chami, Ryanodine receptors: physiological function and deregulation in Alzheimer disease, *Mol. Neurodegener.*, 9 (2014) 21.
- [103] M.P. Mattson, ER calcium and Alzheimer's disease: in a state of flux, *Sci. Signal.*, 3 (2010) pe10.
- [104] S. Chakroborty, C. Briggs, M.B. Miller, I. Goussakov, C. Schneider, J. Kim, J. Wicks, J.C. Richardson, V. Conklin, B.G. Cameransi, G.E. Stutzmann, Stabilizing ER  $\text{Ca}^{2+}$  channel function as an early preventative strategy for Alzheimer's disease, *PLoS One*, 7 (2012) e52056.

- [105] E. Ferreira, R. Resende, R. Costa, C.R. Oliveira, C.M. Pereira, An endoplasmic-reticulum-specific apoptotic pathway is involved in prion and amyloid- $\beta$  peptides neurotoxicity, *Neurobiol. Dis.*, 23 (2006) 669-678.
- [106] O. Thibault, J.C. Gant, P.W. Landfield, Expansion of the calcium hypothesis of brain aging and Alzheimer's disease: minding the store, *Aging Cell*, 6 (2007) 307-317.
- [107] C. Supnet, C. Noonan, K. Richard, J. Bradley, M. Mayne, Up-regulation of the type 3 ryanodine receptor is neuroprotective in the TgCRND8 mouse model of Alzheimer's disease, *J. Neurochem.*, 112 (2010) 356-365.
- [108] M. Manczak, T.S. Anekonda, E. Henson, B.S. Park, J. Quinn, P.H. Reddy, Mitochondria are a direct site of A $\beta$  accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression, *Hum. Mol. Genet.*, 15 (2006) 1437-1449.
- [109] C.A. Hansson, S. Frykman, M.R. Farmery, L.O. Tjernberg, C. Nilsberth, S.E. Pursglove, A. Ito, B. Winblad, R.F. Cowburn, J. Thyberg, M. Ankarcróna, Nicastrin, presenilin, APH-1, and PEN-2 form active  $\gamma$ -secretase complexes in mitochondria, *J. Biol. Chem.*, 279 (2004) 51654-51660.
- [110] P.F. Pavlov, B. Wiehager, J. Sakai, S. Frykman, H. Behbahani, B. Winblad, M. Ankarcróna, Mitochondrial  $\gamma$ -secretase participates in the metabolism of mitochondria-associated amyloid precursor protein, *Faseb Journal*, 25 (2011) 78-88.
- [111] P.J. Crouch, R. Blake, J.A. Duce, G.D. Ciccotosto, Q.X. Li, K.J. Barnham, C.C. Curtain, R.A. Cherny, R. Cappai, T. Dyrks, C.L. Masters, I.A. Trounce, Copper-dependent inhibition of human cytochrome c oxidase by a dimeric conformer of amyloid- $\beta_{1-42}$ , *Journal of Neuroscience*, 25 (2005) 672-679.
- [112] J. Kim, Y. Yang, S.S. Song, J.H. Na, K.J. Oh, C. Jeong, Y.G. Yu, Y.K. Shin, Beta-amyloid oligomers activate apoptotic BAK pore for cytochrome c release, *Biophys. J.*, 107 (2014) 1601-1608.
- [113] L. Hedskog, C.M. Pinho, R. Filadi, A. Ronnback, L. Hertwig, B. Wiehager, P. Larssen, S. Gellhaar, A. Sandebring, M. Westerlund, C. Graff, B. Winblad, D. Galter, H. Behbahani, P. Pizzo, E. Glaser, M. Ankarcróna, Modulation of the endoplasmic reticulum-mitochondria interface in Alzheimer's disease and related models, *Proc. Natl. Acad. Sci. U.S.A.*, 110 (2013) 7916-7921.
- [114] E. Zampese, C. Fasolato, T. Pozzan, P. Pizzo, Presenilin-2 modulation of ER-mitochondria interactions, *Communicative & Integrative Biology*, 4 (2011) 357-360.
- [115] R. Filadi, E. Greotti, G. Turacchio, A. Luini, T. Pozzan, P. Pizzo, Presenilin 2 Modulates Endoplasmic Reticulum-Mitochondria Coupling by Tuning the Antagonistic Effect of Mitofusin 2, *Cell Rep.*, 15 (2016) 2226-2238.
- [116] R. Filadi, E. Greotti, G. Turacchio, A. Luini, T. Pozzan, P. Pizzo, Mitofusin 2 ablation increases endoplasmic reticulum-mitochondria coupling, *Proc. Natl. Acad. Sci. U.S.A.*, 112 (2015) E2174-2181.
- [117] N.S. Leal, B. Schreiner, C.M. Pinho, R. Filadi, B. Wiehager, H. Karlstrom, P. Pizzo, M. Ankarcróna, Mitofusin-2 knockdown increases ER-mitochondria contact and decreases amyloid beta-peptide production, *J. Cell. Mol. Med.*, (2016).
- [118] P. Saftig, J. Klumperman, Lysosome biogenesis and lysosomal membrane proteins: trafficking meets function, *Nat. Rev. Mol. Cell Biol.*, 10 (2009) 623-635.
- [119] W.J.J.M. Scheenen, C.B. Wollheim, T. Pozzan, C. Fasolato, Ca<sup>2+</sup> depletion from granules inhibits exocytosis - A study with insulin-secreting cells, *Journal of Biological Chemistry*, 273 (1998) 19002-19008.
- [120] J.H. Lee, W.H. Yu, A. Kumar, S. Lee, P.S. Mohan, C.M. Peterhoff, D.M. Wolfe, M. Martinez-Vicente, A.C. Massey, G. Sovak, Y. Uchiyama, D. Westaway, A.M. Cuervo, R.A. Nixon, Lysosomal proteolysis and autophagy require presenilin 1 and are disrupted by Alzheimer-related PS1 mutations, *Cell*, 141 (2010) 1146-1158.
- [121] J.H. Lee, M.K. McBrayer, D.M. Wolfe, L.J. Haslett, A. Kumar, Y. Sato, P.P. Lie, P. Mohan, E.E. Coffey, U. Kompella, C.H. Mitchell, E. Lloyd-Evans, R.A. Nixon, Presenilin 1 Maintains



- Lysosomal Ca<sup>2+</sup> Homeostasis via TRPML1 by Regulating vATPase-Mediated Lysosome Acidification, *Cell Rep.*, 12 (2015) 1430-1444.
- [122] K. Coen, R.S. Flannagan, S. Baron, L.R. Carraro-Lacroix, D. Wang, W. Vermeire, C. Michiels, S. Munck, V. Baert, S. Sugita, F. Wuytack, P.R. Hiesinger, S. Grinstein, W. Annaert, Lysosomal calcium homeostasis defects, not proton pump defects, cause endo-lysosomal dysfunction in PSEN-deficient cells, *Journal of Cell Biology*, 198 (2012) 23-35.
- [123] M.A. Busche, A. Konnerth, Neuronal hyperactivity - A key defect in Alzheimer's disease?, *Bioessays*, 37 (2015) 624-632.
- [124] A. Stargardt, D.F. Swaab, K. Bossers, Storm before the quiet: neuronal hyperactivity and A $\beta$  in the presymptomatic stages of Alzheimer's disease, *Neurobiol. Aging*, 36 (2015) 1-11.
- [125] K.V. Kuchibhotla, C.R. Lattarulo, B.T. Hyman, B.J. Bacskai, Synchronous hyperactivity and intercellular calcium waves in astrocytes in Alzheimer mice, *Science*, 323 (2009) 1211-1215.
- [126] K.V. Kuchibhotla, S.T. Goldman, C.R. Lattarulo, H.Y. Wu, B.T. Hyman, B.J. Bacskai, Abeta plaques lead to aberrant regulation of calcium homeostasis in vivo resulting in structural and functional disruption of neuronal networks, *Neuron*, 59 (2008) 214-225.
- [127] M.A. Busche, G. Eichhoff, H. Adelsberger, D. Abramowski, K.H. Wiederhold, C. Haass, M. Staufenbiel, A. Konnerth, O. Garaschuk, Clusters of hyperactive neurons near amyloid plaques in a mouse model of Alzheimer's disease, *Science*, 321 (2008) 1686-1689.
- [128] M.A. Busche, X. Chen, H.A. Henning, J. Reichwald, M. Staufenbiel, B. Sakmann, A. Konnerth, Critical role of soluble amyloid-beta for early hippocampal hyperactivity in a mouse model of Alzheimer's disease, *Proc. Natl. Acad. Sci. U.S.A.*, 109 (2012) 8740-8745.
- [129] J.C. Amatniek, W.A. Hauser, C. DelCastillo-Castaneda, D.M. Jacobs, K. Marder, K. Bell, M. Albert, J. Brandt, Y. Stern, Incidence and predictors of seizures in patients with Alzheimer's disease, *Epilepsia*, 47 (2006) 867-872.
- [130] P. Imfeld, M. Bodmer, M. Schuerch, S.S. Jick, C.R. Meier, Seizures in patients with Alzheimer's disease or vascular dementia: a population-based nested case-control analysis, *Epilepsia*, 54 (2013) 700-707.
- [131] A.W. Bero, P. Yan, J.H. Roh, J.R. Cirrito, F.R. Stewart, M.E. Raichle, J.M. Lee, D.M. Holtzman, Neuronal activity regulates the regional vulnerability to amyloid- $\beta$  deposition, *Nat. Neurosci.*, 14 (2011) 750-756.
- [132] N. Axmacher, M.M. Henseler, O. Jensen, I. Weinreich, C.E. Elger, J. Fell, Cross-frequency coupling supports multi-item working memory in the human hippocampus, *Proc. Natl. Acad. Sci. U.S.A.*, 107 (2010) 3228-3233.
- [133] O. Jensen, L.L. Colgin, Cross-frequency coupling between neuronal oscillations, *Trends Cogn. Sci.*, 11 (2007) 267-269.
- [134] A.A. Ittner, A. Gladbach, J. Bertz, L.S. Suh, L.M. Ittner, p38 MAP kinase-mediated NMDA receptor-dependent suppression of hippocampal hypersynchronicity in a mouse model of Alzheimer's disease, *Acta Neuropathol. Commun.*, 2 (2014) 149.
- [135] S.E. Lesne, Toxic oligomer species of amyloid-beta in Alzheimer's disease, a timing issue, *Swiss medical weekly*, 144 (2014) w14021.
- [136] M.C. Dinamarca, J.A. Rios, N.C. Inestrosa, Postsynaptic Receptors for Amyloid- $\beta$  Oligomers as Mediators of Neuronal Damage in Alzheimer's Disease, *Front. Physiol.*, 3 (2012) 464.