

Neuronal Ca²⁺ dyshomeostasis in Huntington disease

Marta Giacomello,^{1,*} Juan C. Oliveros,² Jose R. Naranjo^{2,3} and Ernesto Carafoli^{4,*}

¹University of Geneva; Geneva, Switzerland; ²Centro Nacional de Biotecnología; Consejo Superior de Investigaciones Científicas (CSIC); Madrid, Spain; ³Centro Investigación Biomedica En Red de Enfermedades Neurodegenerativas (CIBERNED); Madrid, Spain; ⁴Venetian Institute of Molecular Medicine; Padova, Italy

Keywords: Huntington disease, calcium, mitochondria, transcription, Huntington

The expansion of the N-terminal poly-glutamine tract of the huntingtin (Htt) protein is responsible for Huntington disease (HD). A large number of studies have explored the neuronal phenotype of HD, but the molecular aetiology of the disease is still very poorly understood. This has hampered the development of an appropriate therapeutic strategy to at least alleviate its symptoms. In this short review, we have focused our attention on the alteration of a specific cellular mechanism common to all HD models, either genetic or induced by treatment with 3-NPA, i.e., the cellular dyshomeostasis of Ca²⁺. We have highlighted the direct and indirect (i.e., transcriptionally mediated) effects of mutated Htt on the maintenance of the intracellular Ca²⁺ balance, the correct modulation of which is fundamental to cell survival and the disturbance of which plays a key role in the death of the cell.

Introduction

All patients affected by Huntington disease (HD) have as a common genetic defect, the expansion in the number of CAG triplets located in the N-terminal region of the protein huntingtin (Htt). Normally, the wild type protein is characterized by 15–35 CAG repeats, while in the Htt forms associated with HD the repeats increase up to 40–180. One of the hallmarks of HD is the presence, within the cell body, of insoluble inclusions, composed of aggregates of Htt fragments, produced by the cleavage of the protein by caspases, characterized by the trapping of very diverse proteins, among them calmodulin, transcription factors, components of the ubiquitin-proteasome system and polyubiquitin binding proteins, e.g., p62 that is involved in autophagy.^{1,2}

Even if Htt is ubiquitously expressed, HD is characterized principally by specific motor and cognitive impairments, suggesting a precise role of Htt in certain tissues and brain domains. The exact function of Htt has not been clarified; however, many tasks for it have been proposed based on the results obtained in different models of HD that have highlighted impairment in organelle and vesicular trafficking, cholesterol biosynthesis and propensity to apoptosis.^{3,4} Numerous papers have also reported

an imbalance in the generation and scavenging of reactive oxygen species (ROS),^{5,6} as well as defects in the respiratory chain complexes^{7,8} and in mitochondrial functions and morphology.⁹ A detailed discussion of these specific alterations, which have been already summarized elsewhere,^{4,10} is beyond the scope of this review, which will focus on the changes of intracellular Ca²⁺ homeostasis induced by mutated Htt.

The concentration of Ca²⁺ ([Ca²⁺]) within the cell is finely tuned by a series of mechanisms, since Ca²⁺ is a messenger that modulates different signal transduction pathways that are vital to cells: their disturbance can eventually even lead to cell death. Among the main actors involved in [Ca²⁺] handling there are proteins that act as Ca²⁺ buffers, proteins that export Ca²⁺ from the cytosol toward the extracellular medium (the plasma membrane Ca²⁺ pumps and Na⁺/Ca²⁺ exchangers) or to the lumen of organelles and proteins that mediate Ca²⁺ entry in the cytoplasm: the influx of Ca²⁺ is mediated by a number of Ca²⁺ channels (among them those formed by the STIM/Orai proteins which are involved in the store operated Ca²⁺ entry (SOCE) process). Finally, there are proteins whose function is activated by binding to Ca²⁺, such as DREAM and calmodulin.^{11,12} The membrane of some organelles, e.g., that of mitochondria and the endoplasmic reticulum (ER), also contain systems that take up/extrude Ca²⁺, establishing a Ca²⁺-linked crosstalk with neighboring organelles.^{13,14}

Work on various animal models in which HD was either induced by the genetically produced presence of mutated Htt or by the treatment with 3-nitropropionic acid (3-NPA), an inhibitor of a component of the mitochondrial respiratory chain, complex II, that has been shown to induce a HD phenotype,¹⁵ has indeed suggested that one of the hallmarks of HD is the impairment of the intracellular [Ca²⁺] modulation. The presence of mutated Htt fragments has been associated to the altered expression of some genes involved in Ca²⁺ homeostasis both in human patients and in HD murine models.^{4,17} Direct binding of mutated Htt fragments to proteins involved in Ca²⁺ handling has also been reported.¹⁶

Both effects have been supported by convincing evidence. However, it is still unclear whether the transcriptional effects in HD neurons are a cell adaptation response to the variations of intracellular [Ca²⁺], which could be due to the direct interaction of mutated Htt (and/or Htt fragments) with Ca²⁺ binding/Ca²⁺ transport proteins, or whether the changes of Ca²⁺ levels are instead a secondary consequence of primary effects of mutated

*Correspondence to: Marta Giacomello and Ernesto Carafoli;
Email: marta.giacomello@unige.ch and ernesto.carafoli@unipd.it
Submitted: 11/29/12; Revised: 12/27/12; Accepted: 01/11/13
<http://dx.doi.org/10.4161/pri.23581>

Htt on the expression of regulatory proteins. It is of course still possible that transcriptional and non-transcriptional processes run in parallel and contribute jointly to the onset of HD.

Intracellular Ca²⁺ Dyshomeostasis in HD

Defects of intracellular Ca²⁺ homeostasis have been reported in the majority, if not all, the experimental models of HD. However, different alterations have been described depending on the model used. For instance, in medium spiny neurons from mice expressing exon 1 of the human HD gene (R6/2 mice) an increase in the basal cytosolic [Ca²⁺] has been detected that could induce an adaptation to excitotoxic stress.¹⁸ In clonal striatal progenitor cell lines obtained from knock-in mouse embryos carrying a 111 polyglutamine repeat, we have reported an increased release of Ca²⁺ from the ER induced by the blockade of the SERCA pump with cyclopiazonic acid with respect to the wild type controls.⁹ Defects in the morphology of mitochondria have also been observed in the same cell line. The organelles appeared more fragmented: the deregulation of cytosolic Ca²⁺ contributed to the upregulation of the activity of the Ca²⁺-dependent phosphatase calcineurin that could dephosphorylate (and thus activate) the pro-fission dynamin related protein 1 (Drp1) leading to the fragmentation of mitochondria and increased propensity to apoptosis.⁹

Interestingly, the majority of the papers on Ca²⁺ dyshomeostasis in HD describe principally defects on Ca²⁺ uptake/extrusion by mitochondria, which are known to act as central regulators of intracellular Ca²⁺ levels. When the concentration of cytosolic [Ca²⁺] in the vicinity of mitochondria increases significantly, the low-affinity mitochondrial Ca²⁺ uniporter (MCU) becomes activated and mediates Ca²⁺ entry into the matrix. Mitochondria release Ca²⁺ by means of a Na⁺/Ca²⁺ exchanger¹³ or, in case of Ca²⁺ overload, through the opening of the permeability transition pore (PTP),¹⁴ which results in collapse of the membrane potential, rupture of the outer membrane and release of cytochrome c.¹⁴ Alterations in the Ca²⁺ buffering/handling activities of mitochondria have been reported in both early and late stages of HD, indicating a key role of these defects in its pathogenesis. Mitochondria isolated from cells obtained both from HD patients and HD murine models (carrying 72 or 150 polyglutamine repeats) are more sensitive to Ca²⁺ loads, i.e., they have a lower [Ca²⁺] threshold for the depolarization of their inner membrane and higher propensity to PTP opening.^{19,20} Similar mitochondrial Ca²⁺ handling defects have been also reported in clonal striatal cells expressing an Htt form carrying 111Q.²¹

Results of this type could easily explain the higher susceptibility of HD cells to apoptosis; however, they have not been reproduced in all HD experimental models. For instance, striatal mitochondria from knock-in mouse models expressing different variants of Htt (with 20, 80, 92 or 111 glutamine stretches), from R6/2 mice (3 months old) and from 12 months YAC128 mice (carrying a mutated Htt with 128 CAG triplets) were equally or even less susceptible to Ca²⁺ loads than their wild type counterparts.^{22,23} In addition, the sensitivity to Ca²⁺ of mitochondria from several HD models decreased in parallel with age and polyQ length,²² suggesting the intervention of compensatory

mechanisms that would eventually protect these organelles from PTP opening.

In addition to interfering with mitochondrial Ca²⁺ handling, Htt also influences the release of Ca²⁺ from the ER and the function of SOCE channels. As to the ER, mutated Htt (but not the wild type protein) has been claimed to interact directly with the C-terminal portion of the inositol 1,4,5-trisphosphate (InsP₃) receptor type 1 (InsP₃R1),¹⁶ sensitizing it to InsP₃²⁴ and thus modulating the efflux of Ca²⁺ from the ER. As suggested for other neurological disorders,²⁵ this could in turn affect the ability of mitochondria to take up Ca²⁺, that is known to depend on the close proximity of mitochondria to the Ca²⁺ releasing ER. A recent finding has shown that InsP₃R silencing reduces the aggregation of mutated Htt,²⁶ underlining the importance of the interaction between the two proteins for the molecular pathogenesis of HD.

Interestingly, the association of Htt fragments with ER membranes has been recently confirmed and reported to be dependent on ER stress:²⁷ the translocation of Htt from the ER to the nucleus was induced by ER stress. Both the binding to the ER and the shuttling of the Htt fragments to the nucleus were affected by the length of the glutamine repeat.^{26,28,29}

The capacitative, SOCE channels mediated entry of Ca²⁺ from the extracellular medium also appears to be affected in HD. Convincing experimental evidence has indeed shown that this process is enhanced in HD cells.³⁰ Inhibitors of the capacitative Ca²⁺ entry slow down the progression/appearance of HD phenotypes in transgenic flies and exert protective effects on primary cultures of medium spiny neurons from YAC128 mice.³⁰ SOCE channels could thus be a possible novel target for the design of new drugs to alleviate the symptoms of HD.

Full length and N-terminal fragments of Htt have also been reported to bind the synaptic region of N-type voltage dependent Ca²⁺ channels,³¹ possibly modulating their activity. Finally, mutated Htt can directly interact with numerous Ca²⁺ binding proteins.³² This ability should not be underestimated, since, as mentioned above, mutated Htt could sequester these proteins, preventing them from fulfilling their normal physiological tasks and leading to the impairment of signal transduction pathways.³³ Notably, mutated Htt fragments have been found associated to Calmodulin in high molecular weight complexes,³² disruption of this interaction exerting neuroprotective effects.^{34,35} Another of the sequestered proteins is Calretinin, a calcium-binding protein with six EF-hand Ca²⁺ binding domains: the interaction between Calretinin and mutated Htt seems to have protective effects since Calretinin knockdown increases the susceptibility of HD cells to cell death, while its overexpression alleviates the [Ca²⁺] alterations associated to the presence of mutated Htt.³⁶

One of the hallmarks of mutated Htt is its ability to aggregate, thus the question raises on whether Ca²⁺ dyshomeostasis is playing a role in the aggregation process or, in turn, whether the presence of Htt inclusions correlate with the appearance of Ca²⁺ deregulations. Actually the latter has been investigated in detail in a very elegant paper in which the authors overexpress in PC12 cells a truncated form of Htt (Httex1p) containing the polyglutamine repeat.³⁷ Overexpression of Httex1p containing

Table 1. Transcriptomic experiments compared in the bioinformatics analysis

GEO ID	Organism (GeneChip)	Experimental (replicates)	Control (replicates)
GSE3583	<i>Mus musculus</i> (MG430v2.0)	Wild type Q7 cells 3NP treated (3)	Wild type Q7 cells untreated (3)
		Mutant Q111 cells (3)	Wild Type Q7 cells (3)
GSE18551	<i>Mus musculus</i> (MG430v2.0)	YAC128 mice 12 mo old (4)	Wild type mice 12 mo old (4)
		YAC128 mice 24 mo old (6)	Wild type mice 24 mo old (4)
GSE1751	<i>Homo sapiens</i> (HGU133A)	HD symptomatic (12)	Healthy controls (14)
		Presymptomatic (5)	Healthy controls (14)

The Gene Expression Omnibus (NCBI-GEO, www.ncbi.nlm.nih.gov/geo/) ID of each experiment is shown on the right. In each case, the species and the experimental vs. the appropriate control are indicated. YAC128 is a transgenic mouse model for HD expressing a yeast artificial chromosome containing the mutated Huntingtin gene with 128 CAG repeats. Q7 and Q111 cells are neuroblasts cells derived from striatal neurons obtained from wild type or mHtt knock-in mice, respectively.

97 glutamines caused both an alteration of the basal Ca²⁺ levels as well as an increase in the intracellular Ca²⁺ levels induced by glutamate. On the contrary, expression of the wild type Httex1p caused no variations in the resting Ca²⁺ levels, suggesting a key role for the expansion of the glutamine repeat on Ca²⁺ dyshomeostasis. As to the effect of aggregates per se on Ca²⁺ dyshomeostasis, it is worth mentioning that expression of mutants forms of Httex1p carrying either 97 or 103 glutamines deprived of the first 17 amino acids (thus causing a reduction of their ability to aggregate) was sufficient to alter the basal Ca²⁺ levels, but not the response to glutamate stimulation.

Transcriptional Deregulation of Ca²⁺ Handling Proteins in HD

Transcriptional deregulation has been described in almost every model of HD, suggesting a fundamental role for it in the molecular pathogenesis of the disease.^{38,39} The transcriptional effects have been ascribed to a variety of mechanisms: impairment of microRNA mediated gene expression,^{40,41} direct binding to the DNA and interaction with transcription factors both inside the nucleus and within the cytoplasm.^{39,42}

The preferred mechanism seems to be direct binding: Htt fragments could behave like a “kidnapper,” sequestering negative and positive regulators of transcription in the Htt inclusion bodies. Among the molecules sequestered a number are indeed factors that regulate gene expression, some of which also contain glutamine stretches. This is the case of the CREB-binding protein (CBP),⁴³ a coactivator of CREB (cAMP-responsive element binding protein). Interestingly, transient overexpression of CBP has protective role in cellular models expressing mutated Htt,⁴³ and disruption of CREB function leads to a striatal phenotype reminiscent of HD.⁴⁴ The impairment of CBP activity could have particularly detrimental consequences given its function as a master regulator of the peroxisome proliferator-activated receptor γ (PPAR γ) coactivator-1 α (PGC1 α). PGC1 α is primarily involved in mitochondria biogenesis: thus, impairment of its expression could contribute to the production of the mitochondria defects described above. The possible role of the impairment of PGC1 α expression on the onset and progression of HD is supported by experimental evidence; the majority of HD models present changes in the level of PGC1 α , and its overexpression has

protective effects on striatal neurons expressing mutated Htt.⁴⁴ Finally, PGC1 α Knock out mice present lesions in the striatum and phenotype similarities with HD murine models.^{45,46}

The genes that show transcriptional deregulation in HD are involved in numerous physiological processes. They range from those coding for molecules that act as neurotrophins [this is the case of the brain-derived neurotrophic factor (BDNF), whose mRNA levels are significantly lower in human and murine HD samples,^{47,48} for protein kinases (such as protein kinase C β II¹⁷)] or for cytoskeletal proteins.¹⁷ Importantly, many genome-wide studies performed on various HD models have shown profound differences in the mRNA levels of genes coding for Ca²⁺ binding proteins, i.e., parvalbumin, calbindin or hippocalcin or proteins involved in the regulation of intracellular Ca²⁺ handling, i.e., the ryanodine receptor type 1 or different voltage-dependent plasma membrane Ca²⁺ channels.^{17,38} To better illustrate this point, we have performed a simple bioinformatics analysis using results archived in public databases. Thus, several public gene expression studies performed with Affymetrix genechips (Table 1) were combined in one single file of logRatios for approximately 650 probes corresponding to genes related to Ca²⁺ homeostasis in neurons according to Affymetrix annotations to probesets included in their genechips. Four representative HD model systems were selected, the mouse Q7/Q111 knock-in striatal neurons, Q7 (wild type) neurons exposed to the mitochondrial toxin 3-NPA, the striatum from the YAC128 transgenic mice and lymphocytes from HD patients or pre-symptomatic mHtt carriers with their appropriate healthy controls (Table 1). A total of six individual genechip hybridization experiments were included in the analysis (Table 1).

Prior to performing the bioinformatic analysis, to allow the comparison between human and mouse genechips, all probeset IDs from the human genechip (HGU133A) were converted to the orthologous probesets for the mouse Affymetrix GeneChip (MG430v2.0), by using the conversion tables provided by Affymetrix (www.affymetrix.com). Normalized log₂Intensity values provided by GEO database (www.ncbi.nlm.nih.gov/geo) were averaged over all replicates (AvgLogIntensities). For each comparison LogRatios were calculated as: LogRatio = AvgLogIntensity (experiment) – AvgLogIntensity (control). Twenty-five clusters of similar logRatio patterns were generated by K-means algorithm (Fig. 1). Some of the most significant clusters from Figure 1 are commented below.




Figure 1. Representation of the 25 clusters of similar logRatio patterns generated by K-means algorithm. LogRatios were calculated as described, from the normalized intensities downloaded from GEO database for each experiment. They represent differential expression of the genes in the altered samples (disease or transgenic) vs. the controls (healthy or wild type). LogRatios (from -3 to +3) are represented in the Y-axis in each cluster, as shown for cluster 25 at the bottom right corner. LogRatio equal to 0 is represented by the horizontal middle line in each cluster. Positive logRatios (above the horizontal middle line) indicate overexpression whereas negative logRatios (below the horizontal middle line) indicates under-expression. Clusters 1 to 25 are numbered in the bottom right corner. For each cluster the six experiments are labeled (A–F) (as shown for cluster 21, bottom left corner). (A) Q7–3NP vs. Q7; (B) Q111 vs. Q7; (C) YAC128-12 min vs. WT-12 min; (D) YAC128-24 min vs. WT-24 min; (E) HD vs. Normal and (F) Pre-symptomatic HD vs. Normal. Each cluster represents the coordinated changes in gene expression of a subset of calcium-related genes for each of the six experimental conditions (A–F) compared in this study.

Genes related to calcium homeostasis that are downregulated in lymphocytes from HD patients, Yac128 mice and Q111 knock in cells (clusters 22 and 11) are shown in **Figure 2** and in supplemental **Figure 1A**, respectively, while genes that are downregulated only in HD patients (cluster 17) or in HD patients and YAC128 mice (cluster 16) are shown in **Figure S1B and C**, respectively. Similarly, cluster 19 includes the most representative collection of genes related to calcium homeostasis that are induced in all four model-systems of HD (**Fig. 3**), while others in clusters 21 and 1 are mostly induced in human HD lymphocytes (**Fig. S1A and B**). Moreover, we have found that another significant number of genes change in opposite directions in the different model systems compared in this analysis. Thus, several genes included in clusters 10 and 15 are downregulated in human samples but are induced in non-human models (**Fig. 4A and B**). In summary, this preliminary bioinformatics analysis shows coordinated or divergent changes in the expression rate of many calcium homeostasis-related genes, that involve a wide variety of cellular functions related to calcium signaling, in HD. Several factors, however, preclude a more detailed commentary for each set of genes at this stage. For instance, the number of experiments introduced in the comparison is reduced specially when compared with the diversity of the experimental specimens

that are compared. Thus, differences observed in the behavior of certain sets of genes could be related to the different tissues subjected to genome-wide analysis, human lymphocytes vs. neuronal tissue or cells or to the different approaches used to generate the HD model, i.e., the generally poor agreement between the transcriptional changes observed in striatal Q111 neurons and those present in Q7 cells after exposure to the 3-NPA toxin (**Figs. 2–4**). Further comparative bioinformatic analyses including more genome-wide studies would be required to complete the assessment of changes in calcium homeostasis in HD that are controlled at the transcriptional level.

The transcriptional defects in the expression of proteins involved in the maintenance of intracellular Ca^{2+} homeostasis has certainly a role in the Ca^{2+} dyshomeostasis observed in HD models. The transcriptional defects could be further amplified by Ca^{2+} dependent transcriptional control mechanisms. This could occur by changes in the function of gene expression regulatory proteins modulated by Ca^{2+} binding. This is the case of the transcriptional repressor DREAM [downstream responsive element (DRE) antagonist modulator], that translocates to the nucleus following the increase of cytosolic $[\text{Ca}^{2+}]$ levels,^{49,50} or of the cofactor LMO4 (LIM domain only 4), the activity of which is induced by Ca^{2+} entry through voltage dependent Ca^{2+} channels.⁵¹

-3.0 0 3.0

Q7-3NP vs O7 (GSE3583)
 Q111 vs O7 (GSE3583)
 YAC128-12m vs WT-12m (GSE18551)
 YAC128-24m vs WT-24m (GSE18551)
 HD vs N (GSE1751)
 P vs N (GSE1751)

Cacna1b	calcium channel, voltage-dependent, N type, alpha 1B subunit
Slc11a2	solute carrier family 11 (proton-coupled divalent metal ion transporters), member 2
Trim9	tripartite motif-containing 9
Gpr12	G-protein coupled receptor 12
E2f4	E2F transcription factor 4
Slc24a1	solute carrier family 24 (sodium/potassium/calcium exchanger), member 1
Cabp5	calcium binding protein 5
Slc24a2	solute carrier family 24 (sodium/potassium/calcium exchanger), member 2
Atp6v0d1	ATPase, H+ transporting, lysosomal V0 subunit D1
Trpv4	transient receptor potential cation channel, subfamily V, member 4
Mmp24	matrix metalloproteinase 24
Cdh15	cadherin 15
Tesc	tescalcin
Necab2	N-terminal EF-hand calcium binding protein 2
Cd7	CD7 antigen
P2rx7	purinergic receptor P2X, ligand-gated ion channel, 7
Cab39l	calcium binding protein 39-like
Chrn2	cholinergic receptor, nicotinic, beta polypeptide 2 (neuronal)
Stat3	signal transducer and activator of transcription 3
Madd	MAP-kinase activating death domain
Cacna1b	calcium channel, voltage-dependent, N type, alpha 1B subunit
Kcnab2	potassium voltage-gated channel, shaker-related subfamily, beta member 2
Cdh10	cadherin 10
Stim1	stromal interaction molecule 1
Mmp24	matrix metalloproteinase 24
Trpv1	transient receptor potential cation channel, subfamily V, member 1
Hcrtr2	hypocretin (orexin) receptor 2
Mmp24	matrix metalloproteinase 24
P2ry2	purinergic receptor P2Y, G-protein coupled 2
Pde6a	phosphodiesterase 6A, cGMP-specific, rod, alpha
Cdk5	cyclin-dependent kinase 5
Nfatc2ip	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2 interacting protein
Gnao1	guanine nucleotide binding protein, alpha 0
Mef2b	myocyte enhancer factor 2B
Cabp5	calcium binding protein 5
Med1	mediator complex subunit 1
Clstn3	calsyntenin 3
Cdh7	cadherin 7, type 2
Ncs1	neuronal calcium sensor 1
Cycl	cytochrome c-1
Tesc	tescalcin
App	amyloid beta (A4) precursor protein
Ccl1	chemokine (C-C motif) ligand 1
Drd4	dopamine receptor D4
Syn2	synapsin II
Sec24c	Sec24 related gene family, member C (S. cerevisiae)
Nup214	nucleoporin 214
Mef2d	myocyte enhancer factor 2D
Pcdhgc4	protocadherin gamma subfamily C, 4
Cabin1	calcineurin binding protein 1
Tacr3	tachykinin receptor 3
Cnn2	calponin 2
Calb2	calbindin 2
Shpk	sedoheptulokinase
Gpr12	G-protein coupled receptor 12
Fat4	FAT tumor suppressor homolog 4 (Drosophila)
Glr1	glycine receptor, alpha 1 subunit
Nos3	nitric oxide synthase 3, endothelial cell
Slc34a1	solute carrier family 34 (sodium phosphate), member 1
Sdh	succinate dehydrogenase complex, subunit D, integral membrane protein
Tacr3	tachykinin receptor 3
Kcnmb4	potassium large conductance calcium-activated channel, subfamily M, beta member 4
Nfatc1	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1
Letm1	leucine zipper-EF-hand containing transmembrane protein 1
Ttyh1	tweety homolog 1 (Drosophila)
Tacr3	tachykinin receptor 3
Adam22	a disintegrin and metalloproteinase domain 22

Figure 2. For figure legend, see page 81.

Figure 2 (See opposite page). List of the 67 Ca^{2+} -related genes in cluster 22 that are downregulated in the model systems of HD shown on the top. Color-coded LogRatio from -3 to +3 is shown on the top. For each gene in the cluster the acronym and the full name is shown. MultiExperiment Viewer software⁵⁴ was used for generating and plotting the clusters.



Figure 3. List of the 32 Ca^{2+} -related genes in cluster 19 that are upregulated in the model systems of HD shown on the top.

Conclusions

This short review has succinctly summarized the present knowledge on the intracellular Ca^{2+} alterations that occur in HD models. Mutated Htt can directly bind to the InsP_3R , thus modulating the ability of its channel to release Ca^{2+} and thus in turn the crosstalk between the ER and mitochondria and the active mitochondrial Ca^{2+} uptake.^{16,52,53} This could be a general mechanism in many neuropathological processes, since disturbances in the formation of peri-mitochondrial Ca^{2+} microdomains has been already reported to contribute to other disorders, such as for example Alzheimer disease.²⁵

In parallel, strong evidence underlines the importance of transcriptional dysregulation in HD and of the ability of mutated

Htt fragments to sequester or modulate proteins involved in gene expression. Interestingly, this is reflected in the changes of the expression of numerous genes related to Ca^{2+} homeostasis,^{17,38} raising the question of whether these two key processes that are affected in the disease run in parallel or in series.

A conciliatory hypothesis on the transcriptional and non-transcriptional effects of mutated Htt could be suggested by the interesting finding of Htt translocation from the ER to the nucleus.²⁷⁻²⁹ The expansion of the glutamine repeat in mutated Htt could affect the nuclear shuttling of Htt itself. At the same time it could cause the formation of inclusion bodies where transcription factors could be sequestered, in turn again influencing the expression of various genes, among them some coding for proteins of the Ca^{2+} homeostasis machinery.



©2012 Landes Bioscience. Do not distribute

Figure 4 (See opposite page). Lists of the 21 and 17 Ca²⁺-related genes in clusters 10 (A) and 15 (B), respectively, that are differentially regulated in the different model systems of HD shown on the top.

Disclosure of Potential Conflict of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

Work in J.R.N. laboratory is supported by grants from CAM (Neurodegenmodels); ERA-Net (SAF2008-03469); DGICYT (SAF2010-21784); FISS (CIBERNED); Fundación Reina Sofía (PI 006/09) and EU 6th Framework Program (NeuroNE, LSHG-CT-2006-037627).

Work in E.C. laboratory is supported by the same ERA-Net grant and by a grant of the CARIPARO Foundation (Projects of Excellence 2008–2009).

Supplemental Materials

Supplemental materials may be found here: www.landesbioscience.com/journals/prion/article/23581

References

- Arrasate M, Finkbeiner S. Protein aggregates in Huntington's disease. *Exp Neurol* 2012; 238:1-11; PMID:22200539.
- Nagaoka U, Kim K, Jana NR, Doi H, Maruyama M, Mitsui K, et al. Increased expression of p62 in expanded polyglutamine-expressing cells and its association with polyglutamine inclusions. *J Neurochem* 2004; 91:57-68; PMID:15379887.
- Valenza M, Rigamonti D, Goffredo D, Zuccato C, Fenu S, Jamot L, et al. Dysfunction of the cholesterol biosynthetic pathway in Huntington's disease. *J Neurosci* 2005; 25:9932-9; PMID:16251441.
- Zuccato C, Valenza M, Cattaneo E. Molecular mechanisms and potential therapeutical targets in Huntington's disease. *Physiol Rev* 2010; 90:905-81; PMID:20664076.
- Bertoni A, Giuliano P, Galgani M, Rotoli D, Ulianich L, Adornetto A, et al. Early and late events induced by polyQ-expanded proteins: identification of a common pathogenic property of polyQ-expanded proteins. *J Biol Chem* 2011; 286:4727-41; PMID:21115499.
- Hands S, Sajjad MU, Newton MJ, Wyttenbach A. In vitro and in vivo aggregation of a fragment of huntingtin protein directly causes free radical production. *J Biol Chem* 2011; 286:44512-20; PMID:21984825.
- Tabrizi SJ, Workman J, Hart PE, Mangiarini L, Mahal A, Bates G, et al. Mitochondrial dysfunction and free radical damage in the Huntington R6/2 transgenic mouse. *Ann Neurol* 2000; 47:80-6; PMID:10632104.
- Arenas J, Campos Y, Ribacoba R, Martín MA, Rubio JC, Ablanado P, et al. Complex I defect in muscle from patients with Huntington's disease. *Ann Neurol* 1998; 43:397-400; PMID:9506560.
- Costa V, Giacomello M, Hudec R, Lopreiato R, Ermak G, Lim D, et al. Mitochondrial fission and cristae disruption increase the response of cell models of Huntington's disease to apoptotic stimuli. *EMBO Mol Med* 2010; 2:490-503; PMID:21069748.
- Giacomello M, Hudec R, Lopreiato R. Huntington's disease, calcium, and mitochondria. *Biofactors* 2011; 37:206-18; PMID:21674644.
- Saris NE, Carafoli E. A historical review of cellular calcium handling, with emphasis on mitochondria. *Biochemistry (Mosc)* 2005; 70:187-94; PMID:15807658.
- Naranjo JR, Mellström B. Ca²⁺-dependent transcriptional control of Ca²⁺ homeostasis. *J Biol Chem* 2012; 287:31674-80; PMID:22822058.
- Carafoli E, Tiozzo R, Lugli G, Crovetti F, Kratzing C. The release of calcium from heart mitochondria by sodium. *J Mol Cell Cardiol* 1974; 6:361-71; PMID:4855051.
- Giacomello M, Drago I, Pizzo P, Pozzan T. Mitochondrial Ca²⁺ as a key regulator of cell life and death. *Cell Death Differ* 2007; 14:1267-74; PMID:17431419.
- Brouillet E, Hantraye P. Effects of chronic MPTP and 3-nitropropionic acid in nonhuman primates. *Curr Opin Neurol* 1995; 8:469-73; PMID:8845934.
- Tang TS, Tu H, Chan EY, Maximov A, Wang Z, Wellington CL, et al. Huntingtin and huntingtin-associated protein 1 influence neuronal calcium signaling mediated by inositol-(1,4,5) triphosphate receptor type 1. *Neuron* 2003; 39:227-39; PMID:12873381.
- Luthi-Carter R, Hanson SA, Strand AD, Bergstrom DA, Chun W, Peters NL, et al. Dysregulation of gene expression in the R6/2 model of polyglutamine disease: parallel changes in muscle and brain. *Hum Mol Genet* 2002; 11:1911-26; PMID:12165554.
- Hansson O, Guatteo E, Mercuri NB, Bernardi G, Li XJ, Castilho RF, et al. Resistance to NMDA toxicity correlates with appearance of nuclear inclusions, behavioural deficits and changes in calcium homeostasis in mice transgenic for exon 1 of the huntingtin gene. *Eur J Neurosci* 2001; 14:1492-504; PMID:11722611.
- Panov AV, Gutekunst CA, Leavitt BR, Hayden MR, Burke JR, Strittmatter WJ, et al. Early mitochondrial calcium defects in Huntington's disease are a direct effect of polyglutamines. *Nat Neurosci* 2002; 5:731-6; PMID:12089530.
- Choo YS, Johnson GV, MacDonald M, Detloff PJ, Lesort M. Mutant huntingtin directly increases susceptibility of mitochondria to the calcium-induced permeability transition and cytochrome c release. *Hum Mol Genet* 2004; 13:1407-20; PMID:15163634.
- Milakovic T, Quintanilla RA, Johnson GV. Mutant huntingtin expression induces mitochondrial calcium handling defects in clonal striatal cells: functional consequences. *J Biol Chem* 2006; 281:34785-95; PMID:16973623.
- Brustovetsky N, LaFrance R, Purl KJ, Brustovetsky T, Keene CD, Low WC, et al. Age-dependent changes in the calcium sensitivity of striatal mitochondria in mouse models of Huntington's Disease. *J Neurochem* 2005; 93:1361-70; PMID:15935052.
- Oliveira JM, Jekabsons MB, Chen S, Lin A, Rego AC, Gonçalves J, et al. Mitochondrial dysfunction in Huntington's disease: the bioenergetics of isolated and in situ mitochondria from transgenic mice. *J Neurochem* 2007; 101:241-9; PMID:17394466.
- Zhang H, Das S, Li QZ, Dragatsis I, Repa J, Zeitlin S, et al. Elucidating a normal function of huntingtin by functional and microarray analysis of huntingtin-null mouse embryonic fibroblasts. *BMC Neurosci* 2008; 9:38; PMID:18412970.
- Kipanyula MJ, Contreras L, Zampese E, Lazzari C, Wong AK, Pizzo P, et al. Ca²⁺ dysregulation in neurons from transgenic mice expressing mutant presenilin 2. *Aging Cell* 2012; 11:885-93; PMID:22805202.
- Bauer PO, Hudec R, Ozaki S, Okuno M, Ebisui E, Mikoshiba K, et al. Genetic ablation and chemical inhibition of IP3R1 reduce mutant huntingtin aggregation. *Biochem Biophys Res Commun* 2011; 416:13-7; PMID:22056561.
- Atwal RS, Xia J, Pinchev D, Taylor J, Epand RM, Truant R. Huntingtin has a membrane association signal that can modulate huntingtin aggregation, nuclear entry and toxicity. *Hum Mol Genet* 2007; 16:2600-15; PMID:17704510.
- Benn CL, Landles C, Li H, Strand AD, Woodman B, Sathasivam K, et al. Contribution of nuclear and extra-nuclear polyQ to neurological phenotypes in mouse models of Huntington's disease. *Hum Mol Genet* 2005; 14:3065-78; PMID:16183657.
- Cornett J, Cao F, Wang CE, Ross CA, Bates GP, Li SH, et al. Polyglutamine expansion of huntingtin impairs its nuclear export. *Nat Genet* 2005; 37:198-204; PMID:15654337.
- Wu J, Shih HP, Vigont V, Hrdlicka L, Diggins L, Singh C, et al. Neuronal store-operated calcium entry pathway as a novel therapeutic target for Huntington's disease treatment. *Chem Biol* 2011; 18:777-93; PMID:21700213.
- Swayne LA, Chen L, Hameed S, Barr W, Charlesworth E, Colicos MA, et al. Crosstalk between huntingtin and syntaxin 1A regulates N-type calcium channels. *Mol Cell Neurosci* 2005; 30:339-51; PMID:16162412.
- Bao J, Sharp AH, Wagster MV, Becher M, Schilling G, Ross CA, et al. Expansion of polyglutamine repeat in huntingtin leads to abnormal protein interactions involving calmodulin. *Proc Natl Acad Sci U S A* 1996; 93:5037-42; PMID:8643525.
- Kreutz MR, Naranjo JR, Koch KW, Schwaller B. The Neuronal Functions of EF-Hand Ca²⁺-Binding Proteins. [j]. *Front Mol Neurosci* 2012; 5:92; PMID:22973191.
- Dudek NL, Dai Y, Muma NA. Neuroprotective effects of calmodulin peptide 76-121aa: disruption of calmodulin binding to mutant huntingtin. *Brain Pathol* 2010; 20:176-89; PMID:19338577.
- Dudek NL, Dai Y, Muma NA. Protective effects of interrupting the binding of calmodulin to mutant huntingtin. *J Neuropathol Exp Neurol* 2008; 67:355-65; PMID:18379433.
- Dong G, Gross K, Qiao F, Ferguson J, Callegari EA, Rezvani K, et al. Calretinin interacts with huntingtin and reduces mutant huntingtin-caused cytotoxicity. *J Neurochem* 2012; 123:437-46; PMID:22891683.
- Rockabrand E, Slepko N, Pantalone A, Nukala VN, Kazantsev A, Marsh JL, et al. The first 17 amino acids of Huntingtin modulate its sub-cellular localization, aggregation and effects on calcium homeostasis. *Hum Mol Genet* 2007; 16:61-77; PMID:17135277.
- Hodges A, Strand AD, Aragaki AK, Kuhn A, Sengstag T, Hughes G, et al. Regional and cellular gene expression changes in human Huntington's disease brain. *Hum Mol Genet* 2006; 15:965-77; PMID:16467349.
- Seredenina T, Luthi-Carter R. What have we learned from gene expression profiles in Huntington's disease? *Neurobiol Dis* 2012; 45:83-98; PMID:21820514.
- Savas JN, Makusky A, Ottosen S, Baillat D, Then F, Krainc D, et al. Huntington's disease protein contributes to RNA-mediated gene silencing through association with Argonaute and P bodies. *Proc Natl Acad Sci U S A* 2008; 105:10820-5; PMID:18669659.
- Lee ST, Chu K, Im WS, Yoon HJ, Im JY, Park JE, et al. Altered microRNA regulation in Huntington's disease models. *Exp Neurol* 2011; 227:172-9; PMID:21035445.

42. Hogel M, Laprairie RB, Denovan-Wright EM. Promoters are differentially sensitive to N-terminal mutant huntingtin-mediated transcriptional repression. *PLoS One* 2012; 7:e41152; PMID:22815947.
43. Nucifora FC Jr, Sasaki M, Peters MF, Huang H, Cooper JK, Yamada M, et al. Interference by huntingtin and atrophin-1 with cbp-mediated transcription leading to cellular toxicity. *Science* 2001; 291:2423-8; PMID:11264541.
44. Mantamadiotis T, Lemberger T, Bleckmann SC, Kern H, Kretz O, Martin Villalba A, et al. Disruption of CREB function in brain leads to neurodegeneration. *Nat Genet* 2002; 31:47-54; PMID:11967539.
45. Cui L, Jeong H, Borovecki F, Parkhurst CN, Tanese N, Krainc D. Transcriptional repression of PGC-1alpha by mutant huntingtin leads to mitochondrial dysfunction and neurodegeneration. *Cell* 2006; 127:59-69; PMID:17018277.
46. Lin J, Wu PH, Tarr PT, Lindenberg KS, St-Pierre J, Zhang CY, et al. Defects in adaptive energy metabolism with CNS-linked hyperactivity in PGC-1alpha null mice. *Cell* 2004; 119:121-35; PMID:15454086.
47. Gambazzi L, Gokce O, Seredenina T, Katsyuba E, Runne H, Markram H, et al. Diminished activity-dependent brain-derived neurotrophic factor expression underlies cortical neuron microcircuit hypoconnectivity resulting from exposure to mutant huntingtin fragments. *J Pharmacol Exp Ther* 2010; 335:13-22; PMID:20624994.
48. Zuccato C, Marullo M, Conforti P, MacDonald ME, Tartari M, Cattaneo E. Systematic assessment of BDNF and its receptor levels in human cortices affected by Huntington's disease. *Brain Pathol* 2008; 18:225-38; PMID:18093249.
49. Ledo F, Kremer L, Mellström B, Naranjo JR. Ca2+-dependent block of CREB-CBP transcription by repressor DREAM. *EMBO J* 2002; 21:4583-92; PMID:12198160.
50. Savignac M, Pintado B, Gutierrez-Adan A, Palczewska M, Mellström B, Naranjo JR. Transcriptional repressor DREAM regulates T-lymphocyte proliferation and cytokine gene expression. *EMBO J* 2005; 24:3555-64; PMID:16177826.
51. Kashani AH, Qiu Z, Jurata L, Lee SK, Pfaff S, Goebbels S, et al. Calcium activation of the LMO4 transcription complex and its role in the patterning of thalamocortical connections. *J Neurosci* 2006; 26:8398-408; PMID:16899735.
52. Giacomello M, Drago I, Bortolozzi M, Scorzeto M, Gianelle A, Pizzo P, et al. Ca2+ hot spots on the mitochondrial surface are generated by Ca2+ mobilization from stores, but not by activation of store-operated Ca2+ channels. *Mol Cell* 2010; 38:280-90; PMID:20417605.
53. Csordás G, Várnai P, Golenár T, Roy S, Purkins G, Schneider TG, et al. Imaging interorganelle contacts and local calcium dynamics at the ER-mitochondrial interface. *Mol Cell* 2010; 39:121-32; PMID:20603080.
54. Saeed AI, Bhagabati NK, Braisted JC, Liang W, Sharov V, Howe EA, et al. TM4 microarray software suite. *Methods Enzymol* 2006; 411:134-93; PMID:16939790.