

MEMOIRS

of the
American Mathematical Society

ISSN 0065-9266 (print)
ISSN 1947-6221 (online)

Number 1218

Perihelia Reduction and Global Kolmogorov Tori in the Planetary Problem

Gabriella Pinzari

September 2018 • Volume 255 • Number 1218 (first of 7 numbers)

MEMOIRS

ISSN 0065-9266 (print)
ISSN 1947-6221 (online)

of the
American Mathematical Society

Number 1218

Perihelia Reduction and Global Kolmogorov Tori in the Planetary Problem

Gabriella Pinzari

September 2018 • Volume 255 • Number 1218 (first of 7 numbers)

Library of Congress Cataloging-in-Publication Data

Names: Pinzari, Gabriella, 1966– author.

Title: Perihelia reduction and global Kolmogorov tori in the planetary problem / Gabriella Pinzari.

Description: Providence, RI : American Mathematical Society, 2018. | Series: Memoirs of the American Mathematical Society, ISSN 0065-9266 ; number 1218 | "September 2018. Volume 255. Number 1218 (first of 7 numbers)." | Includes bibliographical references.

Identifiers: LCCN 2018040533 | ISBN 9781470441029 (alk. paper)

Subjects: LCSH: Celestial mechanics. | Planetary theory. | Differential equations, Partial.

Classification: LCC QB351 .P527 2018 | DDC 521–dc23

LC record available at <https://lcn.loc.gov/2018040533>

DOI: <https://doi.org/10.1090/memo/1218>

Memoirs of the American Mathematical Society

This journal is devoted entirely to research in pure and applied mathematics.

Subscription information. Beginning with the January 2010 issue, *Memoirs* is accessible from www.ams.org/journals. The 2018 subscription begins with volume 251 and consists of six mailings, each containing one or more numbers. Subscription prices for 2018 are as follows: for paper delivery, US\$999 list, US\$799.20 institutional member; for electronic delivery, US\$879 list, US\$703.20 institutional member. Upon request, subscribers to paper delivery of this journal are also entitled to receive electronic delivery. If ordering the paper version, add US\$20 for delivery within the United States; US\$80 for outside the United States. Subscription renewals are subject to late fees. See www.ams.org/help-faq for more journal subscription information. Each number may be ordered separately; *please specify number* when ordering an individual number.

Back number information. For back issues see www.ams.org/backvols.

Subscriptions and orders should be addressed to the American Mathematical Society, P. O. Box 845904, Boston, MA 02284-5904 USA. *All orders must be accompanied by payment.* Other correspondence should be addressed to 201 Charles Street, Providence, RI 02904-2213 USA.

Copying and reprinting. Individual readers of this publication, and nonprofit libraries acting for them, are permitted to make fair use of the material, such as to copy select pages for use in teaching or research. Permission is granted to quote brief passages from this publication in reviews, provided the customary acknowledgment of the source is given.

Republication, systematic copying, or multiple reproduction of any material in this publication is permitted only under license from the American Mathematical Society. Requests for permission to reuse portions of AMS publication content are handled by the Copyright Clearance Center. For more information, please visit www.ams.org/publications/pubpermissions.

Send requests for translation rights and licensed reprints to reprint-permission@ams.org.

Excluded from these provisions is material for which the author holds copyright. In such cases, requests for permission to reuse or reprint material should be addressed directly to the author(s). Copyright ownership is indicated on the copyright page, or on the lower right-hand corner of the first page of each article within proceedings volumes.

Memoirs of the American Mathematical Society (ISSN 0065-9266 (print); 1947-6221 (online)) is published bimonthly (each volume consisting usually of more than one number) by the American Mathematical Society at 201 Charles Street, Providence, RI 02904-2213 USA. Periodicals postage paid at Providence, RI. Postmaster: Send address changes to *Memoirs*, American Mathematical Society, 201 Charles Street, Providence, RI 02904-2213 USA.

© 2018 by the American Mathematical Society. All rights reserved.

This publication is indexed in *Mathematical Reviews*[®], *Zentralblatt MATH*, *Science Citation Index*[®], *Science Citation IndexTM-Expanded*, *ISI Alerting ServicesSM*, *SciSearch*[®], *Research Alert*[®], *CompuMath Citation Index*[®], *Current Contents*[®]/*Physical, Chemical & Earth Sciences*.

This publication is archived in *Portico* and *CLOCKSS*.

Printed in the United States of America.

∞ The paper used in this book is acid-free and falls within the guidelines established to ensure permanence and durability.

Visit the AMS home page at <https://www.ams.org/>

10 9 8 7 6 5 4 3 2 1 23 22 21 20 19 18

Contents

Chapter 1. Background and results	1
Acknowledgments	7
Chapter 2. Kepler maps and the Perihelia reduction	9
2.1. The \mathcal{P} -map vs rotations and reflections	20
Chapter 3. The \mathcal{P} -map and the planetary problem	23
3.1. A general property of Kepler maps	23
3.2. The case of the \mathcal{P} -map	25
Chapter 4. Global Kolmogorov tori in the planetary problem	29
4.1. A domain of holomorphy	29
4.2. A normal form for the planetary problem	31
4.3. A “multi-scale” KAM Theorem and proof of Theorem A	33
Chapter 5. Proofs	37
5.1. Normalization of fast angles	37
5.2. Secular normalizations	41
Appendix A. Computing the domain of holomorphy	63
A.1. On the analyticity of the solution of Kepler equation	63
A.2. Proof of Proposition 4.2	65
Appendix B. Proof of Lemma 3.2	71
Appendix C. Checking the non-degeneracy condition	75
Appendix D. Some results from perturbation theory	77
D.1. A multi-scale normal form theorem	77
D.2. A slightly-perturbed integrable system	86
Appendix E. More on the geometrical structure of the \mathcal{P} -coordinates, compared to Deprit’s coordinates	87
Bibliography	91

Abstract

We prove the existence of an almost full measure set of $(3n - 2)$ -dimensional quasi-periodic motions in the planetary problem with $(1 + n)$ masses, with eccentricities arbitrarily close to the Levi–Civita limiting value and relatively high inclinations. This extends previous results, where smallness of eccentricities and inclinations was assumed. The question had been previously considered by V. I. Arnold (1963) in the 60s, for the particular case of the planar three-body problem, where, due to the limited number of degrees of freedom, it was enough to use the invariance of the system by the $SO(3)$ group.

The proof exploits nice parity properties of a new set of coordinates for the planetary problem, which reduces completely the number of degrees of freedom for the system (in particular, its degeneracy due to rotations) and, moreover, is well fitted to its reflection invariance. It allows the explicit construction of an associated close to be integrable system, replacing Birkhoff normal form, common tool of previous literature.

Received by the editor February 21, 2015 and, in revised form, February 9, 2016.

Article electronically published on June 25, 2018.

DOI: <https://doi.org/10.1090/memo/1218>

2010 *Mathematics Subject Classification*. Primary 34C20, 70F10, 37J10, 37J15, 37J40; Secondary 34D10, 70F07, 70F15, 37J25, 37J35.

Key words and phrases. Canonical coordinates, Jacobi’s reduction, Deprit’s reduction, Perihelia reduction, symmetries, quasi-periodic motions, Arnold’s theorem on the stability of planetary motions.

This research has been financially supported partly (up to February 28, 2016) by ERC Ideas-Project 306414 “Hamiltonian PDEs and small divisor problems: a dynamical systems approach” and partly (since March 1, 2016) by the ERC Project 677793 “Stable and Chaotic Motions in the Planetary Problem.”

The author is affiliated with the Dipartimento di Matematica “T. Levi-Civita”, via Trieste 63, 35131 Padova, Italy. Email: gabriella.pinzari@math.unipd.it.

©2018 American Mathematical Society

Background and results

In recent years, substantial progress on a statement by Vladimir Igorevich Arnold concerning the stability of the planetary system has been achieved [2, 9, 14, 20, 22, 23, 27, 33].

It sounds as follows.

“For the majority of initial conditions under which the instantaneous orbits of the planets are close to circles lying in a single plane, perturbation of the planets on one another produces, in the course of an infinite interval of time, little change on these orbits provided the masses of the planets are sufficiently small. [...] In particular [...] in the n -body problem there exists a set of initial conditions having a positive Lebesgue measure and such that, if the initial positions and velocities of the bodies belong to this set, the distances of the bodies from each other will remain perpetually bounded.” [2, Chapter III, p. 125].

Solving the differential equations of the motions of the planetary problem, i.e., n planets interacting among themselves and with a star via gravity is, for $n \geq 2$, a problem with ancient roots. This story goes back to Sir Isaac Newton – who brilliantly solved the case of two bodies and then, tackling the analogous one for three bodies, soon realized the necessity of turning to a “perturbative” study (except for naming it a “head ache problem”) – passed through investigations by eminent mathematicians like Delaunay, Lagrange, the prize publicly announced by King Oscar II of Sweden and Norway and awarded to Henri Poincaré, but its “solution” is nowadays open. Chaotic and stable regions may coexist [2, 11, 17].

The question received a new mathematical description, and a strong modern endorsement, after A. N. Kolmogorov announced, at the International Congress of Mathematicians of 1954 in Amsterdam, what is now almost unanimously considered the most important result of the last century for dynamical systems: the theorem of conservation of the invariant torus. This breakthrough result, next enriched by substantial contributions by J. Moser and V. I. Arnold himself [1, 22, 26], states that for a generic Hamiltonian system close to an integrable one, i.e., a system of the form

$$H(I, \varphi) = h(I) + \mu f(I, \varphi) \quad (I, \varphi) \in B \times \mathbb{T}^N \quad B \subset \mathbb{R}^N \quad \mathbb{T} := \mathbb{R}/(2\pi\mathbb{Z}) \quad \mu \ll 1,$$

the major part of unperturbed motions survives, after a small perturbation is switched on, provided suitable “non-degeneracy” conditions are verified by the “unperturbed part” h . Moreover, the theory provides precise arithmetic (“diophantine”) properties to be verified by the “unperturbed frequencies” $\omega_* = \partial h(I_*)$, so that they will be preserved in the full system.

In 1962, V. I. Arnold, extending Kolmogorov’s ideas, and looking for an application to the planetary problem, at the International Congress of Mathematicians of

Stockholm, announced the theorem of stability of planetary motions quoted above. In 1965 Kolmogorov and Arnold were awarded the Lenin Prize for their studies on the stability of the planetary problem – but the story was not finished there.

In order to introduce the results of this paper, we highlight basic facts of this story and its continuation, referring the reader to [5, 10, 16, 28, 29] for more notices.

The planetary problem is close to the integrable problem of n uncoupled two-body problems, where each planet interacts separately with the sun. The mutual interactions among planets are regarded as a perturbing function, the smallness of which is ruled by the planets' masses. However, as a perturbed system, the planetary problem has a limiting degeneracy. Its associated integrable system (the two-body problem) is “super-integrable”: it has more integrals than degrees of freedom. At a technical level, the limiting degeneracy is exhibited by the disappearance of degrees of freedom in the unperturbed part. Therefore, continuing the unperturbed motions to a *positive measure set* of quasi-periodic trajectories might, in general, be not possible, in absence of further informations on the perturbing function.

Arnold found, for the planetary problem, a brilliant solution to the problem of the limiting degeneracy. This led him to add to the assumptions and assertions that are proper of perturbation theories (e.g., “the masses of the planets are sufficiently small”, “set of initial conditions having a positive Lebesgue measure”, “the distances . . . will remain perpetually bounded”) a further requirement of smallness of eccentricities and inclinations of the unperturbed Keplerian ellipses (“the instantaneous orbits of the planets are close to circles lying in a single plane”). Let's summarize Arnold's ideas.

Choosing, as Arnold did, Poincaré coordinates [30] (see, also [2, Ch. III, §2], or, e.g., [8, 15]), the system takes the usual close to be integrable form

$$H_{\mathcal{P}oi} = h_{\text{Kep}} + \mu f_{\mathcal{P}oi},$$

where μ is a small parameter related to the planetary masses, but the unperturbed “Keplerian” part $h_{\text{Kep}}(\Lambda)$ depends on only n action variables $\Lambda = (\Lambda_1, \dots, \Lambda_n)$ (related to the semi-major axes of the instantaneous Keplerian ellipses), out of an overall of $3n$ degrees of freedom. The perturbing function, $f_{\mathcal{P}oi}$, on the other hand, depends on all the coordinates: the actions Λ , their conjugated angles $\ell = (\ell_1, \dots, \ell_n)$ (proportional to the areas of the elliptic sectors spanned by the planets), and, moreover, on some other coordinates $(\mathbf{p}, \mathbf{q}) = (p_1, \dots, p_{2n}, q_1, \dots, q_{2n})$, $4n$ -dimensional, related to those (“secular”) quantities (eccentricities, inclinations, nodes and perihelia of the ellipses) that in the unperturbed problem stay fixed, and for this reason do not appear in h_{Kep} .

It is of great help that the averaged perturbing function (with respect to the angles ℓ) $\overline{f_{\mathcal{P}oi}}(\Lambda, \mathbf{p}, \mathbf{q})$ enjoys several parities in the coordinates (\mathbf{p}, \mathbf{q}) , geometrically related to its invariance by rotations and reflections with respect to the coordinate planes. The “secular origin” $(\mathbf{p}, \mathbf{q}) = 0$, corresponding to all the planets moving on co-centric circles in the same plane, turns out to be an elliptic equilibrium point for the averaged perturbing function, for any value of Λ .

Arnold brilliantly argued to exploit this circumstance to his purpose. By Birkhoff theory, one might think to switch to another set of canonical coordinates $(\Lambda, \tilde{\ell}, \tilde{\mathbf{p}}, \tilde{\mathbf{q}})$, analogous to Poincaré's coordinates, possibly defined only for $(\tilde{\mathbf{p}}, \tilde{\mathbf{q}})$ in a small neighborhood of radius ε around the origin, such that the Hamiltonian of the system, or,

more precisely, its $\tilde{\ell}$ -averaged (“secular”) perturbing function $\overline{f_{\mathcal{B}ir}}$, takes a “normalized form” : it is a polynomial, $\overline{f_{\mathcal{B}ir, tr}}$, of some degree greater or equal than two in the combinations (“degenerate actions”) $\tau_i = \frac{\tilde{p}_i^2 + \tilde{q}_i^2}{2}$, $i = 1, \dots, 2n$, plus a remainder with a higher order. Roughly, Arnold projected to solve the limiting degeneracy by conjugating the planetary system to a new system, whose unperturbed part was just the truncated, normalized Hamiltonian

$$h_{\text{Kep}} + \mu \overline{f_{\mathcal{B}ir, tr}}$$

so as to recover the standard set up of KAM theory. With these ideas in mind, he proved the following impressive result and next applied it to the planar three-body problem. It states that stable trajectories occupy a positive measure set of the phase space, and are more and more dense closer to the elliptic equilibrium. Hence, the smaller eccentricities and inclinations are, the larger the number of stable motions is.

‘The Fundamental Theorem’ (V. I. Arnold, [2]) *If the Hessian matrix of h and the matrix of the coefficients of the second-order term in τ_i in $\overline{f_{\mathcal{B}ir}}$ (“torsion”, or “second-order Birkhoff invariants”) do not vanish identically, and if μ is suitably small with respect to ε , the system affords a positive measure set $\mathcal{K}_{\mu, \varepsilon}$ of quasi-periodic motions in phase space such that its density goes to 1 as $\varepsilon \rightarrow 0$.*

Arnold perfectly knew that, in order to apply the Fundamental Theorem to the problem in space, one should previously treat an unpleasant fact: one of the first order Birkhoff invariants vanishes identically. He was aware that the reason for this first-order degeneracy was to be sought into the existence of two non-commuting integrals, the two horizontal components of the total angular momentum of the systems. If, apparently, a vanishing eigenvalue strongly violates the construction of the normalized system (a deeper analysis of the symmetries of the perturbing function [8, 25], however, shows that the identically vanishing eigenvalue is not a real obstruction), a major problem definitely prevents the application of the Fundamental Theorem: an infinite number of coefficients of *any order* of the (formal) Birkhoff series vanishes identically, among which one entire row and a column in the torsion matrix, which so is identically singular, and the reason is again the invariance by rotations. The proof of this generalized degeneracy is in [8]. We recall here that even Herman had raised a question about the degeneracy of torsion [20, p. 24].

We do not know weather Arnold was aware of the infinite degeneracy of the normalized system (he did not even mention the vanishing of torsion in his paper). He however suggested two different strategies for the three- and the many-body case, of which he provided very few and somewhat controversial details. As for the three-body problem (his ideas for the many-body case will be recalled a few below), he proposed to reduce the integrals (hence, the number of degrees of freedom) of the system by switching to a system of canonical coordinates going back to the XIX century, worked out by Jacobi and Radau [21, 32], which in literature go under the name of *Jacobi reduction of the nodes*. The idea was later completely developed by P. Robutel [33], who, in a deeply quantitative study, checked the non-degeneracy assumptions required by the Fundamental Theorem.

Finding a system of canonical coordinates that do the job of Jacobi reduction of the nodes when the number of bodies is more than three has been a central difficulty for a long time [2, 25]. At this respect, Arnold sadly commented: “In the case of

more than three bodies there is no such elegant method [as Jacobi reduction of the nodes] of reducing the number of degrees of freedom.” [2, Ch. III, §5.5, p. 141].

Exactly twenty years later, F. Boigey and A. Deprit refuted this sentence [3, 12]. They indeed were able to extend Jacobi-Radau reduction to the four-body and general problem, respectively. It should be remarked, anyway, that, while the works by Jacobi, Radau and Boigey provide canonical coordinates on suitable submanifolds of the phase space, the one by Deprit is more general and clarifying, since it provides a set of canonical coordinates for the whole phase space and allows us to recover his predecessors by restriction.

The utility of Boigey-Deprit’s coordinates was not suddenly clear. Neither Boigey nor Deprit ever provided any motivation of their study, or foresaw applications. The only application that is known to the author up to 2008, concerning indeed Deprit’s coordinates, stands in a paper by Ferrer and Osácar, in the 90s, to the three body problem [18]. But this case is not really exhaustive, since for three bodies Deprit’s and Jacobi-Radau’s coordinates coincide. A reason why Boigey-Deprit’s coordinates have been forgotten so long might be that, for more than three bodies, they actually have a less natural aspect, compared to the classical case of Jacobi. A sort of “hierarchical” structure in the geometry of Deprit’s coordinates discouraged the author himself, who, at the end of his paper, declared: “Whether the new phase variables are practical in the general theory of perturbation is an open question. At least, for planetary theories, the answer is likely to be in the negative. But finding a natural system of coordinates for eliminating the nodes in a planetary cluster was not the intention of this note.” [12, p. 194].

In the meantime, in 2004, the first general proof of Arnold’s stability statement appeared. It was by Jacques Féjoz, who completed investigations by the late Michael Herman [14] – but the different procedure that Herman had in mind did not rely on the necessity of handling, explicitly, good coordinates. Indeed, Herman conceived a proof based, besides on a “twist-less” KAM theory going back to H. Russmann [34], on indirect arguments of Lagrangian intersections in order to bypass the so-called “secular resonances”. See [10] for more details.

In 2008, Boigey-Deprit’s coordinates were rediscovered by the author [27], in a slightly different, “planetary” form. The rediscovery was motivated by the purpose of realizing Arnold’s program (i.e., applying the Fundamental Theorem quoted above directly to the planetary Hamiltonian) in the general case, so as to obtain a detailed information about the tori frequencies, the measure of the invariant set and the symplectic structure of the phase space. The utility of Boigey-Deprit’s coordinates became suddenly clear: switching (in order to overcome certain singularities of the chart) to a regularized version, called “RPS” coordinates, (acronym standing for “Regular, Planetary and Symplectic”), allowed them to derive the Birkhoff normal form of the planetary problem, to prove its non-degeneracy, and hence to complete the application of the Fundamental Theorem to the general problem. These results have been published in [6, 7, 9].

Qualitatively, RPS coordinates are very different from JRBD (Jacobi-Radau-Boigey-Deprit); rather, they are more similar to Poincaré coordinates. The mentioned parities and the elliptic equilibrium of the averaged system are still present in the RPS-averaged system. But, as an advantage with respect to Poincaré coordinates,

the RPS perform¹ a “partial reduction” of the rotation symmetry – in contrast with JRBD coordinates, which reduce “fully”. This way, all the degeneracies of the Birkhoff series mentioned above are removed at once, and the non-degeneracy assumptions of the Fundamental Theorem may be checked.

We like to recall now Arnold’s strategy for the many-body case: more than forty years earlier, he foresaw to construct a system of coordinates analogous to RPS, via a Taylor series in Poincaré coordinates [2, Ch III, §5, n. 5, p. 141].

Indeed, both the reduction of the nodes and this latter reduction are available whatever the number of bodies is.

The possibility of switching from Delaunay-Poincaré to the more fruitful JRBD, or even RPS coordinates, is an effect of the limiting degeneracy. This gives in fact the opportunity of remixing coordinates related to secular quantities, and, simultaneously, keeping the Keplerian term h_{Kep} unvaried.

Following this idea, in this paper, we show that other systems of coordinates may be determined for the planetary problem which, as well as JRBD and RPS coordinates, are well adapted to overcome the degeneracy due to rotations, and, moreover, enjoy some different properties.

We present a full reduction, which we call \mathcal{P} -map, or perihelia reduction. It refines JRBD coordinates in two respects.

Firstly, the \mathcal{P} -map is well defined in the case of the planar problem, while JRBD coordinates are not. Everyone knows, in fact, that the starting point for the Radau-Jacobi reduction is the so-called “line of the nodes”, the straight line determined by the intersection between the planes of the two orbits. When the orbits of the two planets belong to the same plane, this is not defined. A similar circumstance arises for Boigey-Deprit’s coordinates, since their construction relies on certain straight lines in the space, which again lose their meaning in case of co-planarity.

The proof of Arnold’s theorem given in [9, 27] is not affected by such singularity, since, as said, it relies on RPS coordinates, which, at the expense of one more degree of freedom, are well defined for co-planar motions – in that case they reduce to the classical Poincaré coordinates.

It has its consequences when one wants to compare results for the fully reduced systems, in space or in the plane. The singularity of the chart does not allow one to state that motions in the spatial problem with minimum number of independent frequencies starting with very small inclinations stay close to the corresponding planar motions. Notwithstanding further studies appearing in [28], where this problem is partially overcome (via the construction of regular coordinates for co-planar motions defined locally), it would be nice, in principle, to handle a global system of action-angle coordinates which completely reduces rotations and is shared simultaneously by the planar and the spatial problem.

Secondly, the \mathcal{P} -map is well adapted to reflection symmetries of the problem, while JRBD coordinates are not, as discussed in [25, 29].

¹In the framework of the study of canonical coordinates for the planetary system, by “partial reduction”, we mean a system of canonical coordinates where a couple of conjugated coordinates consists of integrals (e.g., functions of the three components of the total angular momentum). By “full reduction”, we mean a partial reduction where also another integral appears among the coordinates. The terms “partial reduction”, “full reduction” have been coined in [25].

Reflection symmetries are parities of the Hamiltonian expressed in Cartesian coordinates. As known, this does not change under arbitrary changes of the signs of positions or momenta coordinates. They are not related to integrals. Therefore, it might be a nice fact, and in general useful for applications, to have a system of coordinates that, after integrals are reduced, parities associated to reflections are maintained. Quite often parities are associated to equilibria, and equilibria to stable motions; an example is provided a few lines below.

We shall apply the \mathcal{P} -map by proving a variant of Arnold's stability theorem. We shall face up to a question raised again by Arnold in his fantastic paper on the possibility of removing the constraint on eccentricities and inclinations. He indeed proved that, at least for the planar three-body problem, there is no need to assume their smallness. Rather, it is sufficient that the trajectories of the planets are away enough so as to avoid collisions. He obtained this stronger result by exploiting the convergence of the Birkhoff series associated to the averaged perturbation, a very particular and happy circumstance, due to the few degrees of freedom of the problem.

From the mathematical point of view, the question is whether strategies exist for finding stable motions other than the one of exploring the neighborhood of the elliptic equilibrium.

Concerning instead the physical relevance, asteroids or some trans-Neptunian objects have motions with relatively large eccentricities and inclinations and an almost continuous spectrum of frequencies.

Besides the mentioned stronger result by Arnold, some other statements in the same direction have been obtained for the case of the spatial three-body problem and the planar problem with any number of bodies [28]. Here, the measure of the invariant set has been estimated to be larger and larger as the planetary masses and the semi-axes ratios are small, no matter the smallness of the eccentricities and inclinations – the proof relies on an argument of convergence of a significant approximation of the Birkhoff series. Other results in this direction have been announced by J. Féjóz, since late 2013 [13].

Even though the arguments of [2, 28] do not apply to the general spatial problem, since no significant approximation of the Birkhoff series associated to the averaged perturbation is integrable, using the \mathcal{P} -map, we shall prove the following.

Theorem A. *Fix numbers $0 < \underline{e}_i < \bar{e}_i < 0.6627\dots$, $i = 1, \dots, n$. There exists a number N depending only on n and a number α_0 depending on \underline{e}_i , \bar{e}_i , and n such that, if $\alpha < \alpha_0$, $\mu \leq \alpha^N$, in a domain of planetary motions where the semi-major axes $a_1 < a_2 < \dots < a_n$ are spaced as follows*

$$(*) \quad a_i^- \leq a_i \leq a_i^+ \quad \text{with} \quad a_i^\pm := \frac{a_1^\pm}{\alpha^{\frac{1}{3}(2n+1-2n-i+2+1-i)}}$$

there exists a positive measure set $\mathcal{K}_{\mu,\alpha}$, the density of which in phase space can be bounded below as

$$\text{dens}(\mathcal{K}_{\mu,\alpha}) \geq 1 - (\log \alpha^{-1})^p \sqrt{\alpha},$$

consisting of quasi-periodic motions with $3n - 2$ frequencies where the planets' eccentricities e_i verify

$$\underline{e}_i \leq e_i \leq \bar{e}_i.$$

Before we switch to details, a few remarks.

Firstly, the claimed upper bound $0.6627\dots$ is classical. It is related to the fact that, as well as in [2, 28], the proof uses the machinery of real-analytic functions. We refer the reader to [24, 35] and references therein for general notices. A treatment of the argument, as needed in the present paper, is provided in Chapter A.1.

Secondly, as it may be seen to the choice of a_j^\pm , the distances among the planets' semi-axes are not of the same order but grow super-exponentially going towards the sun. This resembles a sort of belt arrangement, observed in nature for asteroids. It is possible to prove an analogous result, with increasing distances in the opposite direction.

Thirdly, the result in Theorem A (especially, the claimed growth of a_i^\pm) may be regarded as an alternative way of solving the problem of the limiting degeneracy – without Birkhoff normal form.

Acknowledgments

The author is indebted to Jacques Féjoz, who let the author know the work by Harrington [19], without which I would never have thought of this application of the \mathcal{P} -coordinates. Also, I am deeply grateful to A. Celletti and R. de la Llave, for their interest and for encouraging me with precious advices. Thanks finally to L. Biasco for his interest.

CHAPTER 2

Kepler maps and the Perihelia reduction

We introduce the *Perihelia reduction*, or \mathcal{P} -map, in the slightly general context of *Kepler maps*.

Fix a reference frame $\mathbf{G}_0 = (k^{(1)}, k^{(2)}, k^{(3)})$ in the Euclidean space E^3 . We identify the three chosen directions $k^{(1)}, k^{(2)}, k^{(3)}$ with the triples of coordinates with respect of the system of coordinates established by themselves:

$$k^{(1)} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} \quad k^{(2)} = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix} \quad k^{(3)} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

DEFINITION 2.1. An *ellipse* (with a focus in the origin and non-vanishing eccentricity) is a quadruplet $\mathfrak{E} = (a, e, N, P)$, where $a \in \mathbb{R}_+$ is the *semi-major axis*, $e \in (0, 1)$ is the *eccentricity*, $N \in \mathbb{R}^3 \cap S^2$ is the *normal direction* and $P \in N^\perp \cap S^2$ is the *perihelion direction*.

DEFINITION 2.2 (Kepler maps). Given $2n$ positive “mass parameters” m_1, \dots, m_n , $\mathfrak{M}_1, \dots, \mathfrak{M}_n$, a set $\mathfrak{X} \subset \mathbb{R}^{5n}$, we say that a map

$$\mathcal{K} : \mathbf{K} = (\mathbf{X}_{\mathcal{K}}, \ell) \in \mathcal{D} := \mathfrak{X} \times \mathbb{T}^n \rightarrow (y_{\mathcal{K}}, x_{\mathcal{K}}) \in \mathcal{C} := \mathcal{K}(\mathcal{D}) \subset (\mathbb{R}^3)^n \times (\mathbb{R}^3)^n$$

where

$$\begin{aligned} \ell &= (\ell_1, \dots, \ell_n), & (y_{\mathcal{K}}, x_{\mathcal{K}}) &= (y_{\mathcal{K}}^{(1)}, \dots, y_{\mathcal{K}}^{(n)}, x_{\mathcal{K}}^{(1)}, \dots, x_{\mathcal{K}}^{(n)}) \\ y_{\mathcal{K}}^{(j)} &= y_{\mathcal{K}}^{(j)}(\mathbf{X}_{\mathcal{K}}, \ell_j) & x_{\mathcal{K}}^{(j)} &= x_{\mathcal{K}}^{(j)}(\mathbf{X}_{\mathcal{K}}, \ell_j) \quad j = 1, \dots, n, \end{aligned}$$

is a *Kepler map* if there exists an injection

$$\tau_{\mathcal{K}} : \mathbf{X}_{\mathcal{K}} \in \mathfrak{X} \rightarrow \mathfrak{E}_{\mathcal{K}} = (\mathfrak{E}_{1,\mathcal{K}}, \dots, \mathfrak{E}_{n,\mathcal{K}})$$

which assigns to any $\mathbf{X}_{\mathcal{K}} \in \mathfrak{X}$ an n -plet $(\mathfrak{E}_{1,\mathcal{K}}, \dots, \mathfrak{E}_{n,\mathcal{K}})$ of (co-focal) ellipses

$$\mathfrak{E}_{j,\mathcal{K}} = (a_{j,\mathcal{K}}, e_{j,\mathcal{K}}, N_{\mathcal{K}}^{(j)}, P_{\mathcal{K}}^{(j)}), \quad j = 1, \dots, n$$

and \mathcal{K} acts in the following way. Letting $Q_{\mathcal{K}}^{(j)} := N_{\mathcal{K}}^{(j)} \times P_{\mathcal{K}}^{(j)}$, then

$$(2.1) \quad x_{\mathcal{K}}^{(j)} = a_{j,\mathcal{K}} P_{\mathcal{K}}^{(j)} + b_{j,\mathcal{K}} Q_{\mathcal{K}}^{(j)} \quad y_{\mathcal{K}}^{(j)} = a_{j,\mathcal{K}}^\circ P_{\mathcal{K}}^{(j)} + b_{j,\mathcal{K}}^\circ Q_{\mathcal{K}}^{(j)}$$

where, if $\zeta_{j,\mathcal{K}}$, the *eccentric anomaly*, is the solution of Kepler’s Equation

$$(2.2) \quad \zeta_{j,\mathcal{K}} - e_{j,\mathcal{K}} \sin \zeta_{j,\mathcal{K}} = \ell_j$$

then

(2.3)

$$\begin{aligned} a_{j,\mathcal{K}} &:= a_{j,\mathcal{K}}(\cos \zeta_{j,\mathcal{K}} - e_{j,\mathcal{K}}) & b_{j,\mathcal{K}} &:= a_{j,\mathcal{K}} \sqrt{1 - e_{j,\mathcal{K}}^2} \sin \zeta_{j,\mathcal{K}} \\ a_{j,\mathcal{K}}^\circ &:= -m_j \sqrt{\frac{\mathfrak{M}_j}{a_{j,\mathcal{K}}}} \frac{\sin \zeta_{j,\mathcal{K}}}{1 - e_{j,\mathcal{K}} \cos \zeta_{j,\mathcal{K}}} & b_{j,\mathcal{K}}^\circ &:= m_j \sqrt{\frac{\mathfrak{M}_j(1 - e_{j,\mathcal{K}}^2)}{a_{j,\mathcal{K}}}} \frac{\cos \zeta_{j,\mathcal{K}}}{1 - e_{j,\mathcal{K}} \cos \zeta_{j,\mathcal{K}}}. \end{aligned}$$

REMARK 2.1. The definition implies that

- (i) \mathcal{K} is a bijection of the sets \mathcal{D} and \mathcal{C} ;
- (ii) the *angular momenta* and the *energies*¹

$$(2.4) \quad C_{\mathcal{K}}^{(j)} := x_{\mathcal{K}}^{(j)} \times y_{\mathcal{K}}^{(j)}, \quad H_{\mathcal{K}}^{(j)} := \frac{\|y_{\mathcal{K}}^{(j)}\|^2}{2m_j} - \frac{m_j \mathfrak{M}_j}{\|x_{\mathcal{K}}^{(j)}\|}$$

do not depend on ℓ_j and are given by

$$(2.5) \quad C_{\mathcal{K}}^{(j)} = m_j \sqrt{\mathfrak{M}_j a_{j,\mathcal{K}} (1 - e_{j,\mathcal{K}}^2)} N_{\mathcal{K}}^{(j)}, \quad H_{\mathcal{K}}^{(j)} = -\frac{m_j \mathfrak{M}_j}{2a_{j,\mathcal{K}}};$$

- (iii) the couples $(y_{\mathcal{K}}^{(j)}, x_{\mathcal{K}}^{(j)})$ verify the system of ODEs

$$(2.6) \quad \begin{cases} m_j \sqrt{\frac{\mathfrak{M}_j}{a_{j,\mathcal{K}}^3}} \partial_{\ell_j} x_{\mathcal{K}}^{(j)} = y_{\mathcal{K}}^{(j)} \\ \sqrt{\frac{\mathfrak{M}_j}{a_{j,\mathcal{K}}^3}} \partial_{\ell_j} y_{\mathcal{K}}^{(j)} = -m_j \mathfrak{M}_j \frac{x_{\mathcal{K}}^{(j)}}{\|x_{\mathcal{K}}^{(j)}\|^3}. \end{cases}$$

- (iv) Even though *canonical* maps (with respect to the standard two-form) have a pre-eminent role in Hamiltonian Mechanics, Kepler maps are used also in different contexts in Astronomy, where being canonical is not required. For example, one can consider the Kepler map associated to the “elliptic elements” injection

$$\tau_{\mathcal{E}ell} : (a, e, P, i, \Omega) \rightarrow \mathfrak{E}_{\mathcal{E}ell}$$

where $a = (a_1, \dots, a_n)$ are the *semi-major axes*, $e = (e_1, \dots, e_n)$ are the *eccentricities*, $P = (P^{(1)}, \dots, P^{(n)})$ are the *perihelia*, $i = (i_1, \dots, i_n)$ are the *inclinations*, $\Omega = (\Omega_1, \dots, \Omega_n)$ are the *nodes’ longitudes*.

The only known examples up to now of *canonical Kepler maps* are the classical *Delaunay map* Del (its definition is recalled in the next Definition 2.5) and the map Dep [7, 27] related to Deprit’s coordinates [12], which is recalled in Appendix E. Below, we introduce a new canonical Kepler map.

DEFINITION 2.3 (perihelia reduction, or \mathcal{P} -map). We denote as \mathcal{P} , and call *perihelia reduction*, or \mathcal{P} -map, the Kepler map

$$(2.7) \quad \mathcal{P} : \mathbb{P} = (X_{\mathcal{P}}, \ell) \in \mathcal{D}_{\mathcal{P}} = \mathfrak{X}_{\mathcal{P}} \times \mathbb{T}^n \rightarrow (y, x) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n}$$

associated to the bijection

$$\tau_{\mathcal{P}} : X_{\mathcal{P}} = (\Theta, \chi, \Lambda, \vartheta, \kappa) \in \mathfrak{X}_{\mathcal{P}} \rightarrow (\mathfrak{E}_1, \dots, \mathfrak{E}_n) \in \mathbb{E}_{\mathcal{P}} = \tau_{\mathcal{P}}(\mathfrak{X}_{\mathcal{P}}) \subset E^{3n}$$

defined by means of Definition 2.4 and Proposition 2.1 below.

¹Here, $\|v\| := \sqrt{v_1^2 + v_2^2 + v_3^2}$ denotes the usual Euclidean norm of $v = (v_1, v_2, v_3) \in \mathbb{R}^3$.

DEFINITION 2.4. For a given $(\mathfrak{E}_1, \dots, \mathfrak{E}_n) \subset E^3 \times \dots \times E^3$, with

$$\mathfrak{E}_j = (a_j, e_j, N^{(j)}, P^{(j)}),$$

and masses $\mathfrak{m}_1, \dots, \mathfrak{m}_n, \mathfrak{M}_1, \dots, \mathfrak{M}_n$, define

$$(2.8) \quad C_{\mathcal{E}}^{(j)} := \mathfrak{m}_j \sqrt{\mathfrak{M}_j a_j (1 - e_j^2)} N^{(j)} \quad S_{\mathcal{E}}^{(j)} := \sum_{i=j}^n C_{\mathcal{E}}^{(i)} \quad 1 \leq j \leq n$$

be the *angular momenta* associated to \mathfrak{E}_j and the j^{th} *partial angular momenta*, so that

$$(2.9) \quad S_{\mathcal{E}}^{(1)} = \sum_{i=1}^n C_{\mathcal{E}}^{(i)} \quad S_{\mathcal{E}}^{(n)} = C_{\mathcal{E}}^{(n)}$$

are the *total angular momentum* and the angular momentum of the last ellipse, respectively. Define the \mathcal{P} -nodes

$$(2.10) \quad \nu_j := \begin{cases} k^{(3)} \times S_{\mathcal{E}}^{(1)} & j = 1 \\ P^{(j-1)} \times S_{\mathcal{E}}^{(j)} & j = 2, \dots, n \end{cases} \quad \mathfrak{n}_j := S_{\mathcal{E}}^{(j)} \times P^{(j)} \quad j = 1, \dots, n.$$

Finally, define

$$\mathcal{E}_{\mathcal{P}} := \{((\mathfrak{E}_1, \dots, \mathfrak{E}_n) \subset E^3 \times \dots \times E^3) : 0 < e_j < 1, \nu_j \neq 0, \mathfrak{n}_j \neq 0 \forall j = 1, \dots, n\},$$

and, on this set, the map

$$\tau_{\mathcal{P}}^{-1} : (\mathfrak{E}_1, \dots, \mathfrak{E}_n) \in \mathcal{E}_{\mathcal{P}} \rightarrow X_{\mathcal{P}} \in \mathfrak{X}_{\mathcal{P}} = \tau_{\mathcal{P}}^{-1}(\mathcal{E}_{\mathcal{P}})$$

where

$$X_{\mathcal{P}} = (\Theta, \chi, \Lambda, \vartheta, \kappa) \in \mathbb{R}^n \times \mathbb{R}_+^n \times \mathbb{R}_+^n \times \mathbb{T}^n \times \mathbb{T}^n$$

with

$$\Theta = (\Theta_0, \dots, \Theta_{n-1}), \quad \vartheta = (\vartheta_0, \dots, \vartheta_{n-1})$$

$$\chi = (\chi_0, \dots, \chi_{n-1}), \quad \kappa = (\kappa_0, \dots, \kappa_{n-1})$$

$$\Lambda = (\Lambda_1, \dots, \Lambda_n)$$

defined via the following formulae:

$$(2.11) \quad \Theta_{j-1} := \begin{cases} Z := S_{\mathcal{E}}^{(1)} \cdot k^{(3)} \\ S_{\mathcal{E}}^{(j)} \cdot P^{(j-1)} \end{cases} \quad \vartheta_{j-1} := \begin{cases} \zeta := \alpha_{k^{(3)}}(k^{(1)}, \nu_1) & j = 1 \\ \alpha_{P^{(j-1)}}(\mathfrak{n}_{j-1}, \nu_j) & 2 \leq j \leq n \end{cases}$$

$$\chi_{j-1} := \begin{cases} G := \|S_{\mathcal{E}}^{(1)}\| \\ \|S_{\mathcal{E}}^{(j)}\| \end{cases} \quad \kappa_{j-1} := \begin{cases} \mathfrak{g} := \alpha_{S_{\mathcal{E}}^{(1)}}(\nu_1, \mathfrak{n}_1) & j = 1 \\ \alpha_{S_{\mathcal{E}}^{(j)}}(\nu_j, \mathfrak{n}_j) & 2 \leq j \leq n \end{cases}$$

$$\Lambda_j := \mathfrak{M}_j \sqrt{\mathfrak{m}_j a_j}.$$

PROPOSITION 2.1. *Let $\mathfrak{X}_{\mathcal{P}}$ be the subset of $\mathbb{R}^n \times \mathbb{R}_+^n \times \mathbb{R}_+^n \times \mathbb{T}^n \times \mathbb{T}^n$ defined by the following inequalities*

$$(2.12) \quad \begin{aligned} & \sqrt{\chi_{i-1}^2 + \chi_i^2 - 2\Theta_i^2 + 2\sqrt{(\chi_i^2 - \Theta_i^2)(\chi_{i-1}^2 - \Theta_i^2)} \cos \vartheta_i} < \Lambda_i \\ & (\chi_{i-1} - \chi_i, \vartheta_i) \neq (0, \pi) \quad 0 < \chi_{n-1} < \Lambda_n \quad i = 1, \dots, n-1 \end{aligned}$$

and

$$(2.13) \quad |\Theta_0| < \chi_0 \quad |\Theta_i| < \min(\chi_{i-1}, \chi_i) \quad i = 1, \dots, n-1.$$

The map $\tau_{\mathcal{P}}^{-1}$ is a bijection of $\mathcal{E}_{\mathcal{P}}$ onto $\mathfrak{X}_{\mathcal{P}}$. The formulae of the inverse map

$$\tau_{\mathcal{P}} : \quad \mathbf{X}_{\mathcal{P}} = (\Theta, \chi, \Lambda, \vartheta, \kappa) \in \mathcal{D}_{\mathcal{P}} \rightarrow \mathfrak{E}_{\mathcal{P}} = (\mathfrak{E}_{1,\mathcal{P}}, \dots, \mathfrak{E}_{n,\mathcal{P}}) \in \mathcal{E}_{\mathcal{P}}$$

$$\mathfrak{E}_{j,\mathcal{P}} = (a_{j,\mathcal{P}}, e_{j,\mathcal{P}}, N_{\mathcal{P}}^{(j)}, P_{\mathcal{P}}^{(j)})$$

are as follows. Let $\iota_1, \dots, \iota_n, \mathfrak{i}_1, \dots, \mathfrak{i}_n \in (0, \pi)$ be defined via

$$(2.14) \quad \cos \iota_j = \frac{\Theta_{j-1}}{\chi_{j-1}}, \quad \cos \mathfrak{i}_j := \frac{\Theta_j}{\chi_{j-1}}, \quad 1 \leq j \leq n$$

(with $\Theta_n := 0$, so that $\mathfrak{i}_n = \frac{\pi}{2}$) and $\mathcal{T}_1, \dots, \mathcal{T}_n, \mathcal{S}_1, \dots, \mathcal{S}_n \in \text{SO}(3)$ via

$$(2.15) \quad \mathcal{T}_j := \mathcal{R}_3(\vartheta_j) \mathcal{R}_1(\iota_j) \quad \mathcal{S}_j := \mathcal{R}_3(\kappa_j) \mathcal{R}_1(\mathfrak{i}_j), \quad 1 \leq j \leq n$$

and let

$$(2.16) \quad \mathbf{C}_{\mathcal{P}}^{(j)} := \mathcal{T}_1 \mathcal{S}_1 \cdots \mathcal{T}_{j-1} \mathcal{S}_{j-1} \mathcal{T}_j (\chi_{j-1} k^{(3)} - \chi_j \mathcal{S}_j \mathcal{T}_{j+1} k^{(3)})$$

with $\chi_n := 0$, so that

$$(2.17) \quad \|\mathbf{C}_{\mathcal{P}}^{(j)}\| = \begin{cases} \sqrt{\chi_{j-1}^2 + \chi_j^2 - 2\Theta_j^2 + 2\sqrt{(\chi_j^2 - \Theta_j^2)(\chi_{j-1}^2 - \Theta_j^2)} \cos \vartheta_j} & j = 1, \dots, n-1 \\ \chi_{n-1} & j = n. \end{cases}$$

Then $\mathbf{C}_{\mathcal{P}}^{(j)} = \mathbf{C}_{\mathcal{E}}^{(j)} \circ \tau_{\mathcal{P}}$ and

$$(2.18) \quad \begin{aligned} a_{j,\mathcal{P}} &= \frac{1}{\mathfrak{M}_j} \left(\frac{\Lambda_j}{\mathfrak{m}_j} \right)^2 & e_{j,\mathcal{P}} &= \sqrt{1 - \frac{\|\mathbf{C}_{\mathcal{P}}^{(j)}\|^2}{\Lambda_j^2}} \\ N_{\mathcal{P}}^{(j)} &= \frac{\mathbf{C}_{\mathcal{P}}^{(j)}}{\|\mathbf{C}_{\mathcal{P}}^{(j)}\|} & P_{\mathcal{P}}^{(j)} &= \mathcal{T}_1 \mathcal{S}_1 \cdots \mathcal{T}_j \mathcal{S}_j k^{(3)}. \end{aligned}$$

REMARK 2.2.

(i) From $\mathbf{C}_{\mathcal{P}}^{(j)} = \mathbf{C}_{\mathcal{E}}^{(j)} \circ \tau_{\mathcal{P}}$, (2.4), (2.5) and (2.25), there follows that $\mathbf{C}_{\mathcal{P}}^{(j)} = x_{\mathcal{P}}^{(j)} \times y_{\mathcal{P}}^{(j)}$.

(ii) $P_{\mathcal{P}}^{(j)} \perp N_{\mathcal{P}}^{(j)}$. Indeed, using the definitions,

$$\begin{aligned} \mathbf{C}_{\mathcal{P}}^{(j)} \cdot P_{\mathcal{P}}^{(j)} &= \chi_{j-1} k^{(3)} \cdot (\mathcal{S}_j k^{(3)}) - \mathcal{T}_{j+1} \chi_j k^{(3)} \cdot (k^{(3)}) \\ &= \chi_{j-1} \cos \iota_j - \chi_j \cos \mathfrak{i}_{j+1} \\ &= 0. \end{aligned}$$

(iii) $\mathbf{S}_{\mathcal{P}}^{(j)} := \mathbf{S}_{\mathcal{E}}^{(j)} \circ \tau_{\mathcal{P}} = \sum_{i=j}^n \mathbf{C}_{\mathcal{P}}^{(i)} = \chi_{j-1} \mathcal{T}_1 \mathcal{S}_1 \cdots \mathcal{T}_{j-1} \mathcal{S}_{j-1} \mathcal{T}_j k^{(3)}$.

We shall prove that

THEOREM 2.1. *The \mathcal{P} -map preserves the standard 2-form*

$$\sum_{j=1}^n dy_{\mathcal{P}}^{(j)} \wedge dx_{\mathcal{P}}^{(j)} = \sum_{i=1}^n (d\Theta_{i-1} \wedge d\vartheta_{i-1} + d\chi_{i-1} \wedge d\kappa_{i-1} + d\Lambda_i \wedge d\ell_i).$$

REMARK 2.3. Actually, we shall prove a finer result: the change $\phi_{\mathcal{D}el}^{\mathcal{P}} := \mathcal{D}el^{-1} \circ \mathcal{P}$ which relates the \mathcal{P} -coordinates to the classical Delaunay coordinates (see the Definition 2.5) is homogeneous-canonical (compare Lemma 2.6).

PROOF OF PROPOSITION 2.1. The formula for $a_{j,\mathcal{P}}$ in (2.16) is immediate from the definition of Λ_j . Postponing to below that $C_{\mathcal{P}}^{(j)} := C_{\mathcal{E}}^{(j)} \circ \tau_{\mathcal{P}}$ has the expression in (2.16) (in turn this implies (2.17), the formula for $N^{(j)}$ and the one for $e_{j,\mathcal{P}}$ in (2.18)), we check that the image set $\tau_{\mathcal{P}}^{-1}(\mathcal{E}_{\mathcal{P}})$ is included in the domain $\mathfrak{X}_{\mathcal{P}}$ defined by inequalities (2.12), (2.13). From the formula for $e_{j,\mathcal{P}}$ in (2.18), we have that conditions $0 < e_{j,\mathcal{P}} < 1$ for all $j = 1, \dots, n$ correspond to relations in (2.12). Note that the first condition in the second line of (2.12) is equivalent to $e_{j,\mathcal{P}} \neq 1$, as one sees rewriting

$$(2.19) \quad \|C_{\mathcal{P}}^{(j)}\|^2 = (\sqrt{\chi_{j-1}^2 - \Theta_j^2} - \sqrt{\chi_j^2 - \Theta_j^2})^2 + 2\sqrt{(\chi_j^2 - \Theta_j^2)(\chi_{j-1}^2 - \Theta_j^2)}(1 + \cos \vartheta_j).$$

Next, recalling the definitions of Θ_0, χ_0 in (2.11), and noticing the relations

$$\Theta_j = S_{\mathcal{E}}^{(j+1)} \cdot P^{(j)} = (S_{\mathcal{E}}^{(j)} - C_{\mathcal{E}}^{(j)}) \cdot P^{(j)} = S_{\mathcal{E}}^{(j)} \cdot P^{(j)} \quad j = 1, \dots, n-1,$$

we immediately see that conditions $\nu_i \neq 0 \neq n_i$ imply (2.13). We have so checked what we wanted.

Now it remains to check the formula for $C_{\mathcal{P}}^{(j)}$ in (2.16) and the one for $P_{\mathcal{P}}^{(j)}$ in (2.18), for any $X_{\mathcal{P}} \in \mathfrak{X}_{\mathcal{P}}$. To this end, we consider the following chain of vectors

$$(2.20) \quad \begin{array}{cccccccc} k^{(3)} & \rightarrow & S_{\mathcal{E}}^{(1)} & \rightarrow & P^{(1)} & \rightarrow & \dots & \rightarrow & S_{\mathcal{E}}^{(j)} & \rightarrow & P^{(j)} & \rightarrow & \dots & \rightarrow & P^{(n)} \\ & & \Downarrow & & \Downarrow & & \vdots & & \Downarrow & & \Downarrow & & \vdots & & \Downarrow \\ & & \nu_1 & & n_1 & & \vdots & & \nu_j & & n_j & & \vdots & & n_n \end{array}$$

where $\nu_1, n_1, \dots, \nu_n, n_n$ are the \mathcal{P} -nodes in (2.10), given by the skew-product of the two consecutive vectors in the chain.

We associate to this chain of vectors the following chain of frames

$$(2.21) \quad G_0 \rightarrow F_1 \rightarrow G_1 \rightarrow \dots \rightarrow F_j \rightarrow G_j \rightarrow F_{j+1} \rightarrow \dots \rightarrow G_n$$

where $G_0 = (k^{(1)}, k^{(2)}, k^{(3)})$ is the initial prefixed frame and the frames, while F_i, G_i are frames defined via

$$(2.22) \quad F_j = (\nu_j, \cdot, S^{(j)}) \quad G_j = (n_j, \cdot, P^{(j)}) \quad j = 1, \dots, n.$$

By construction, each frame in the chain has its first axis coinciding with the intersection of horizontal plane with the horizontal plane of the previous frame (hence, in particular, $\nu_j \perp S^{(j)}$ and $n_j \perp P^{(j)}$). Denote as T_j the rotation matrix which describes the change of coordinates from G_{j-1} to F_j and as S_j the one from F_j to G_j . The matrices $\overline{T}_j, \overline{S}_j$ have just the expressions claimed in (2.14), (2.15).

This follows from the definitions of $(\Theta, \chi, \vartheta, \kappa)$ in (2.11). Then we have the following sequence of transformations

$$\begin{array}{ccccccccccc} T_1 & & S_1 & & \cdots & & S_j & & \cdots & & S_n \\ G_0 & \rightarrow & F_1 & \rightarrow & G_1 & \rightarrow & \cdots & \rightarrow & F_j & \rightarrow & G_j & \rightarrow & \cdots & \rightarrow & G_n \end{array}$$

connecting G_0 to any other frame in the chain. From this, and the definitions of the frames (2.22), the formulae for $P_{\mathcal{P}}^{(j)}$ in (2.18) and

$$S_{\mathcal{P}}^{(j)} = \chi_{j-1} T_1 S_1 \cdots T_{j-1} S_{j-1} T_j k^{(3)}$$

follow at once. Hence, also the formulae for $C_{\mathcal{P}}^{(j)}$, which is given by $C_{\mathcal{P}}^{(j)} = S_{\mathcal{P}}^{(j)} - S_{\mathcal{P}}^{(j+1)}$, with $S_{\mathcal{P}}^{(n+1)} \equiv 0$. \square

For the proof of Theorem 2.1, we shall use three auxiliary maps, that we shall denote as $\tilde{\mathcal{P}}$, $\widetilde{\mathcal{D}el}$ and $\mathcal{D}el$. The map $\tilde{\mathcal{P}}$ is very closely related to \mathcal{P} ; $\widetilde{\mathcal{D}el}$ and $\mathcal{D}el$ are well known: in the literature they are often referred to as (two variants of) *Delaunay* maps.

The map $\tilde{\mathcal{P}}$. Define the set

$$\mathcal{C}_{\tilde{\mathcal{P}}} := \left\{ (y, x) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n} : x^{(j)} \neq 0, \quad \tilde{n}_j \neq 0, \quad \tilde{\nu}_j \neq 0 \quad \forall j = 1, \dots, n \right\},$$

where, for $(y, x) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n}$, with $y = (y^{(1)}, \dots, y^{(n)})$, $x = (x^{(1)}, \dots, x^{(n)})$, $x^{(j)} \neq 0$, we let

$$\tilde{\nu}_j := \begin{cases} k^{(3)} \times S_{\mathcal{C}}^{(1)} & j = 1 \\ \frac{x^{(j-1)}}{\|x^{(j-1)}\|} \times S_{\mathcal{C}}^{(j)} & j = 2, \dots, n \end{cases} \quad \tilde{n}_j := S_{\mathcal{C}}^{(j)} \times \frac{x^{(j)}}{\|x^{(j)}\|}$$

with $j = 1, \dots, n$ and

$$(2.23) \quad C_{\mathcal{C}}^{(j)} := x^{(j)} \times y^{(j)}, \quad S_{\mathcal{C}}^{(j)} := \sum_{i=j}^n C^{(i)}.$$

Define a map

$$\tilde{\mathcal{P}}^{-1} : (y, x) \in \mathcal{C}_{\tilde{\mathcal{P}}} \rightarrow (\tilde{\Theta}, \tilde{\chi}, \tilde{\mathbf{R}}, \tilde{\vartheta}, \tilde{\kappa}, \tilde{\mathbf{r}}) \in \mathbb{R}^n \times \mathbb{R}_+^n \times \mathbb{R}^n \times \mathbb{T}^n \times \mathbb{T}^n \times \mathbb{R}_+^n$$

with

$$\begin{aligned} \tilde{\Theta} &= (\tilde{\Theta}_0, \dots, \tilde{\Theta}_{n-1}) & \tilde{\vartheta} &= (\tilde{\vartheta}_0, \dots, \tilde{\vartheta}_{n-1}) \\ \tilde{\chi} &= (\tilde{\chi}_0, \dots, \tilde{\chi}_{n-1}) & \tilde{\kappa} &= (\tilde{\kappa}_0, \dots, \tilde{\kappa}_{n-1}) \\ \tilde{\mathbf{R}} &= (\tilde{\mathbf{R}}_1, \dots, \tilde{\mathbf{R}}_n) & \tilde{\mathbf{r}} &= (\tilde{\mathbf{r}}_1, \dots, \tilde{\mathbf{r}}_n) \end{aligned}$$

via the following formulae:

$$\begin{aligned} \tilde{R}_j &= \frac{y^{(j)} \cdot x^{(j)}}{\|x^{(j)}\|} & \tilde{r}_j &= \|x^{(j)}\| & j &= 1, \dots, n \\ \tilde{\chi}_{j-1} &= \|S_C^{(j)}\| & \tilde{\kappa}_{j-1} &= \alpha_{S_C^{(j)}}(\tilde{\nu}_j, \tilde{n}_j) & j &= 1, \dots, n \\ \tilde{\Theta}_{j-1} &= \begin{cases} S_C^{(1)} \cdot k^{(3)} \\ S_C^{(j)} \cdot \frac{x^{(j-1)}}{\|x^{(j-1)}\|} \end{cases} & \tilde{\vartheta}_{j-1} &= \begin{cases} \alpha_{k^{(3)}}(k^{(1)}, \tilde{\nu}_1) & j = 1 \\ \alpha_{\frac{x^{(j-1)}}{\|x^{(j-1)}\|}}(\tilde{n}_{j-1}, \tilde{\nu}_j) & j = 2, \dots, n. \end{cases} \end{aligned}$$

LEMMA 2.1. *Let $\mathcal{D}_{\tilde{\mathcal{P}}}$ be the set of $(\tilde{\Theta}, \tilde{\chi}, \tilde{R}, \tilde{\vartheta}, \tilde{\kappa}, \tilde{r}) \in \mathbb{R}^n \times \mathbb{R}_+^n \times \mathbb{R}^n \times \mathbb{T}^n \times \mathbb{T}^n \times \mathbb{R}_+^n$ such that $(\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa})$ satisfies (2.13), and let \tilde{T}_j , \tilde{S}_j and $C_{\tilde{\mathcal{P}}}^{(j)}$ be the functions of $(\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa})$ defined in (2.14)-(2.16), with $(\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa})$ replacing $(\Theta, \chi, \vartheta, \kappa)$.*

The map $\tilde{\mathcal{P}}^{-1}$ is a bijection from $\mathcal{C}_{\tilde{\mathcal{P}}}$ onto the set $\mathcal{D}_{\tilde{\mathcal{P}}}$. Its inverse map

$$\tilde{\mathcal{P}} : (\tilde{\Theta}, \tilde{\chi}, \tilde{R}, \tilde{\vartheta}, \tilde{\kappa}, \tilde{r}) \in \mathcal{D}_{\tilde{\mathcal{P}}} \rightarrow (y_{\tilde{\mathcal{P}}}, x_{\tilde{\mathcal{P}}}) \in \mathbb{R}^n \times \mathbb{R}^n$$

has the following analytical expression:

$$(2.24) \quad \begin{cases} x_{\tilde{\mathcal{P}}}^{(j)} := \tilde{r}_j \tilde{T}_1 \tilde{S}_1 \cdots \tilde{T}_j \tilde{S}_j k^{(3)} \\ y_{\tilde{\mathcal{P}}}^{(j)} := \frac{\tilde{R}_j}{\tilde{r}_j} x_{\tilde{\mathcal{P}}}^{(j)} + \frac{1}{\tilde{r}_j^2} C_{\tilde{\mathcal{P}}}^{(j)} \times x_{\tilde{\mathcal{P}}}^{(j)} \quad 1 \leq j \leq n \end{cases}$$

Moreover, the following relation holds:

$$(2.25) \quad C_{\tilde{\mathcal{P}}}^{(j)} = C_C^{(j)} \circ \tilde{\mathcal{P}} = x_{\tilde{\mathcal{P}}}^{(j)} \times y_{\tilde{\mathcal{P}}}^{(j)}.$$

PROOF. With similar arguments as the ones of the proof of Proposition 2.1, but replacing, in the diagram (2.20), $S_{\mathcal{E}}^{(j)}$ with $S_C^{(j)}$, $P_{\tilde{\mathcal{P}}}^{(j)}$ with $\frac{x_{\tilde{\mathcal{P}}}^{(j)}}{\|x_{\tilde{\mathcal{P}}}^{(j)}\|}$ and the nodes ν_k , n_k with $\tilde{\nu}_k$, \tilde{n}_k , one finds the formula for $x_{\tilde{\mathcal{P}}}^{(j)}$ in (2.24), the formula for

$$S_{\tilde{\mathcal{P}}}^{(j)} := S_C^{(j)} \circ \tilde{\mathcal{P}} = \tilde{\chi}_{j-1} \tilde{T}_1 \tilde{S}_1 \cdots \tilde{T}_{j-1} \tilde{S}_{j-1} \tilde{T}_j k^{(3)}$$

and hence the formula for

$$C_C^{(j)} \circ \tilde{\mathcal{P}} = S_{\tilde{\mathcal{P}}}^{(j)} - S_{\tilde{\mathcal{P}}}^{(j+1)} = C_{\tilde{\mathcal{P}}}^{(j)}$$

being just the formula for $C_{\tilde{\mathcal{P}}}^{(j)}$ in (2.16), with $(\Theta, \chi, \vartheta, \kappa)$ replaced by $(\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa})$. With the same argument as in Remark 2.2(ii), we see that $x_{\tilde{\mathcal{P}}}^{(j)} \perp C_{\tilde{\mathcal{P}}}^{(j)}$. Finally, the formula for $y_{\tilde{\mathcal{P}}}^{(j)}$ is found taking for $y_{\tilde{\mathcal{P}}}^{(j)}$ the unique vector verifying

$$y_{\tilde{\mathcal{P}}}^{(j)} \cdot \frac{x_{\tilde{\mathcal{P}}}^{(j)}}{\|x_{\tilde{\mathcal{P}}}^{(j)}\|} = R_j \quad x_{\tilde{\mathcal{P}}}^{(j)} \times y_{\tilde{\mathcal{P}}}^{(j)} = C_{\tilde{\mathcal{P}}}^{(j)}.$$

□

LEMMA 2.2. *$\tilde{\mathcal{P}}$ preserves the standard Liouville 1-form:*

$$(2.26) \quad \sum_{j=1}^n y_{\tilde{\mathcal{P}}}^{(j)} \cdot dx_{\tilde{\mathcal{P}}}^{(j)} = \sum_{j=1}^n (\tilde{\Theta}_{j-1} d\tilde{\vartheta}_{j-1} + \tilde{\chi}_{j-1} d\tilde{\kappa}_{j-1} + \tilde{R}_j d\tilde{r}_j).$$

The proof of Lemma 2.2 uses the following easy lemma:

LEMMA 2.3 ([7]). *Let*

$$x = \mathcal{R}_3(\theta)\mathcal{R}_1(i)\bar{x}, \quad y = \mathcal{R}_3(\theta)\mathcal{R}_1(i)\bar{y}, \quad C := x \times y, \quad \bar{C} := \bar{x} \times \bar{y},$$

with $x, \bar{x}, y, \bar{y} \in \mathbb{R}^3$. Then,

$$y \cdot dx = C \cdot k^{(3)} d\theta + \bar{C} \cdot k^{(1)} di + \bar{y} \cdot d\bar{x}.$$

PROOF OF LEMMA 2.2. We may write

$$x_{\bar{\mathcal{P}}}^{(j)} = \tilde{\mathbb{T}}_1 \tilde{\mathbb{S}}_1 \cdots \tilde{\mathbb{T}}_j \tilde{\mathbb{S}}_j \tilde{x}^{(j)}, \quad y_{\bar{\mathcal{P}}}^{(j)} = \tilde{\mathbb{T}}_1 \tilde{\mathbb{S}}_1 \cdots \tilde{\mathbb{T}}_j \tilde{\mathbb{S}}_j \tilde{y}^{(j)}, \quad C_{\bar{\mathcal{P}}}^{(j)} = \tilde{\mathbb{T}}_1 \tilde{\mathbb{S}}_1 \cdots \tilde{\mathbb{T}}_j \tilde{\mathbb{S}}_j \tilde{C}^{(j)}$$

where

$$(2.27) \quad \begin{aligned} \tilde{x}^{(j)} &:= \tilde{\Gamma}_j k^{(3)} & j = 1, \dots, n-1 \\ \tilde{y}^{(j)} &:= \tilde{\mathbb{R}}_j k^{(3)} + \frac{1}{\tilde{\Gamma}_j} \tilde{C}^{(j)} \times k^{(3)} \\ \tilde{C}^{(j)} &:= \tilde{\chi}_{j-1} \tilde{\mathbb{S}}_j^{-1} k^{(3)} - \tilde{\chi}_j \tilde{\mathbb{T}}_{j+1} k^{(3)} = \tilde{x}^{(j)} \times \tilde{y}^{(j)} \end{aligned}$$

with $\tilde{\chi}_n := 0, \tilde{\mathbb{S}}_n := \text{id}$. We also let, for $1 \leq k \leq j \leq n$ and $1 \leq i \leq n-1$,

$$\begin{aligned} \hat{C}_k^{(j)} &= \tilde{\mathbb{S}}_k (\tilde{\mathbb{T}}_{k+1} \tilde{\mathbb{S}}_{k+1} \cdots \tilde{\mathbb{T}}_j \tilde{\mathbb{S}}_j) \tilde{C}^{(j)}, \quad \check{C}_k^{(j)} = \tilde{\mathbb{T}}_k \tilde{\mathbb{S}}_k \cdots \tilde{\mathbb{T}}_j \tilde{\mathbb{S}}_j \tilde{C}^{(j)}, \quad \check{C}_{j+1}^{(j)} := \tilde{C}^{(j)} \\ \hat{S}_k^{(j)} &:= \sum_{m=j}^n \hat{C}_k^{(m)}, \quad \check{S}_k^{(j)} := \sum_{m=j}^n \check{C}_k^{(m)}, \quad \check{S}_{i+1}^{(i)} := \tilde{C}^{(i)} + \check{S}_{i+1}^{(i+1)} \end{aligned}$$

where the product $\tilde{\mathbb{T}}_{k+1} \tilde{\mathbb{S}}_{k+1} \cdots \tilde{\mathbb{T}}_j \tilde{\mathbb{S}}_j$ is to be replaced with the identity when $k = j$. We have the following identities (implied by $S^{(j)} = \sum_{k=j}^n C^{(k)}$):

$$(2.28) \quad \check{S}_j^{(j)} = \sum_{k=j}^n \check{C}_j^{(k)} = \tilde{\chi}_{j-1} \tilde{\mathbb{T}}_j k^{(3)}, \quad \hat{S}_j^{(j)} = \sum_{k=j}^n \hat{C}_j^{(k)} = \tilde{\chi}_{j-1} k^{(3)}, \quad \check{S}_{i+1}^{(i)} = \tilde{\chi}_{j-1} \tilde{\mathbb{S}}_i^{-1} k^{(3)}.$$

Applying Lemma 2.3 repeatedly and using (as it follows from (2.27))

$$\tilde{y}^{(j)} \cdot d\tilde{x}^{(j)} = \tilde{\mathbb{R}}_j d\tilde{r}_j,$$

we have, for $1 \leq j \leq n$,

$$\begin{aligned} y_{\bar{\mathcal{P}}}^{(j)} \cdot x_{\bar{\mathcal{P}}}^{(j)} &= \sum_{k=1}^j (\check{C}_k^{(j)} \cdot k^{(3)} d\tilde{\vartheta}_{k-1} \\ &\quad + \hat{C}_k^{(j)} \cdot k^{(1)} d\tilde{u}_k + \hat{C}_k^{(j)} \cdot k^{(3)} d\tilde{\kappa}_{k-1} + \check{C}_{k+1}^{(j)} \cdot k^{(1)} d\tilde{i}_k) + \tilde{\mathbb{R}}_j d\tilde{r}_j \end{aligned}$$

where, as in the proof of Lemma 2.1, $\tilde{\iota}_j, \tilde{\mathbb{I}}_j$ denote the functions ι_j, \mathbb{I}_j in (2.14), with Θ_i, χ_i replaced by $\tilde{\Theta}_i, \tilde{\chi}_i$. Note that we have used $d\tilde{i}_n \equiv 0$, since, by definition, $\tilde{i}_n = \frac{\pi}{2}$. Taking the sum over $j = 1, \dots, n$,

$$\begin{aligned} \sum_{j=1}^n y_{\bar{\mathcal{P}}}^{(j)} \cdot dx_{\bar{\mathcal{P}}}^{(j)} &= \sum_{j=1}^n \check{S}_j^{(j)} \cdot k^{(3)} d\tilde{\vartheta}_{j-1} + \hat{S}_j^{(j)} \cdot k^{(1)} d\tilde{u}_j + \\ &\quad \hat{S}_j^{(j)} \cdot k^{(3)} d\tilde{\kappa}_{j-1} + \check{S}_{j+1}^{(j)} \cdot k^{(1)} d\tilde{i}_j + \sum_{j=1}^n \tilde{\mathbb{R}}_j d\tilde{r}_j. \end{aligned}$$

In view of (2.28) and of the definitions in (2.14)-(2.15), we then find (2.26). \square

The map $\widetilde{\mathcal{D}el}$. The map

$$\widetilde{\mathcal{D}el} : (\widetilde{\mathbf{H}}, \widetilde{\Gamma}, \widetilde{\mathbf{R}}, \widetilde{\mathbf{h}}, \widetilde{\mathbf{g}}, \widetilde{\mathbf{r}}) \in \mathcal{D}_{\widetilde{\mathcal{D}el}} \rightarrow (y_{\widetilde{\mathcal{D}el}}, x_{\widetilde{\mathcal{D}el}}) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n}$$

is defined on the set

$$\begin{aligned} \mathcal{D}_{\widetilde{\mathcal{D}el}} := & \left\{ (\widetilde{\mathbf{H}}, \widetilde{\Gamma}, \widetilde{\mathbf{R}}, \widetilde{\mathbf{h}}, \widetilde{\mathbf{g}}, \widetilde{\mathbf{r}}) = (\widetilde{\mathbf{H}}_1, \dots, \widetilde{\mathbf{H}}_n, \widetilde{\Gamma}_1, \dots, \widetilde{\Gamma}_n, \widetilde{\mathbf{R}}_1, \dots, \widetilde{\mathbf{R}}_n, \widetilde{\mathbf{h}}_1, \dots, \widetilde{\mathbf{h}}_n, \right. \\ & \widetilde{\mathbf{g}}_1, \dots, \widetilde{\mathbf{g}}_n, \widetilde{\mathbf{r}}_1, \dots, \widetilde{\mathbf{r}}_n) \in \mathbb{R}^{3n} \times \mathbb{T}^{2n} \times \mathbb{R}_+^n : \quad \widetilde{\mathbf{r}}_j > 0, \quad \widetilde{\Gamma}_j > 0, \quad \frac{|\widetilde{\mathbf{H}}_j|}{\widetilde{\Gamma}_j} < 1 \\ & \left. \forall j = 1, \dots, n \right\} \end{aligned}$$

via the following formulae:

$$x_{\widetilde{\mathcal{D}el}}^{(j)} := \mathbf{R}_3(\widetilde{\mathbf{h}}_j) \mathcal{R}_1(\widetilde{i}_j) \overline{x}_{\widetilde{\mathcal{D}el}}^{(j)}, \quad y_{\widetilde{\mathcal{D}el}}^{(j)} := \mathbf{R}_3(\widetilde{\mathbf{h}}_j) \mathcal{R}_1(\widetilde{i}_j) \overline{y}_{\widetilde{\mathcal{D}el}}^{(j)}$$

where

$$\begin{aligned} \widetilde{i}_j &:= \cos^{-1} \frac{\widetilde{\mathbf{H}}_j}{\widetilde{\Gamma}_j} \in (0, \pi) \\ \overline{x}_{\widetilde{\mathcal{D}el}}^{(j)} &:= \widetilde{\mathbf{r}}_j \cos \widetilde{\mathbf{g}}_j k^{(1)} + \widetilde{\mathbf{r}}_j \sin \widetilde{\mathbf{g}}_j k^{(2)} \\ \overline{y}_{\widetilde{\mathcal{D}el}}^{(j)} &:= \left(\widetilde{\mathbf{R}}_j \cos \widetilde{\mathbf{g}}_j - \frac{\widetilde{\Gamma}_j}{\widetilde{\mathbf{r}}_j} \sin \widetilde{\mathbf{g}}_j \right) k^{(1)} + \left(\widetilde{\mathbf{R}}_j \sin \widetilde{\mathbf{g}}_j + \frac{\widetilde{\Gamma}_j}{\widetilde{\mathbf{r}}_j} \cos \widetilde{\mathbf{g}}_j \right) k^{(2)}. \end{aligned}$$

LEMMA 2.4 (Delaunay). $\widetilde{\mathcal{D}el}$ is a bijection from the domain $\mathcal{D}_{\widetilde{\mathcal{D}el}}$ onto the set

$$\begin{aligned} \mathcal{C}_{\widetilde{\mathcal{D}el}} := & \left\{ (y, x) = (y^{(1)}, \dots, y^{(n)}, x^{(1)}, \dots, x^{(n)}) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n} : \right. \\ & \left. \widetilde{\mathbf{n}}_j := k^{(3)} \times \mathcal{C}_C^{(j)} \neq 0, \quad x^{(j)} \neq 0 \quad \forall j = 1, \dots, n \right\} \end{aligned}$$

where $\mathcal{C}_C^{(j)}$ is as in (2.23). The formulae for the inverse map

$$\widetilde{\mathcal{D}el}^{-1} : (y, x) \in \mathcal{C}_{\widetilde{\mathcal{D}el}} \rightarrow (\widetilde{\mathbf{H}}, \widetilde{\Gamma}, \widetilde{\mathbf{R}}, \widetilde{\mathbf{h}}, \widetilde{\mathbf{g}}, \widetilde{\mathbf{r}}) \in \mathcal{D}_{\widetilde{\mathcal{D}el}}$$

are

(2.29)

$$\begin{cases} \widetilde{\mathbf{H}}_j = \mathcal{C}_C^{(j)} \cdot k^{(3)} \\ \widetilde{\mathbf{h}}_j := \alpha_{k^{(3)}}(k^{(1)}, \widetilde{\mathbf{n}}_j) \end{cases} \quad \begin{cases} \widetilde{\Gamma}_j := \|\mathcal{C}_C^{(j)}\| \\ \widetilde{\mathbf{g}}_j := \alpha_{\mathcal{C}_C^{(j)}}(\widetilde{\mathbf{n}}_j, x^{(j)}) \end{cases} \quad \begin{cases} \widetilde{\mathbf{R}}_j = \frac{y^{(j)} \cdot x^{(j)}}{\|x^{(j)}\|} \\ \widetilde{\mathbf{r}}_j = \|x^{(j)}\| \end{cases}$$

Finally, $\widetilde{\mathcal{D}el}$ preserves the standard Liouville 1-form

$$\sum_{i=1}^n y_{\widetilde{\mathcal{D}el}}^{(i)} \cdot dx_{\widetilde{\mathcal{D}el}}^{(i)} = \sum_{i=1}^n (\widetilde{\mathbf{H}}_i d\widetilde{\mathbf{h}}_i + \widetilde{\Gamma}_i d\widetilde{\mathbf{g}}_i + \widetilde{\mathbf{R}}_i d\widetilde{\mathbf{r}}_i).$$

We omit the proof of Lemma 2.4, which may be found in classical textbooks.

The map $\mathcal{D}el$.

DEFINITION 2.5 (Delaunay map). Let

$$\begin{aligned} \mathfrak{X}_{\mathcal{D}el} := \left\{ X_{\mathcal{D}el} := (H, \Gamma, \Lambda, h, g) = (H_1, \dots, H_n, \Gamma_1, \dots, \Gamma_n, \Lambda_1, \dots, \Lambda_n, h_1, \dots, h_n, \right. \\ \left. g_1, \dots, g_n) \in \mathbb{R}^{3n} \times \mathbb{T}^{2n} : \quad \Gamma_j > 0, \quad \frac{|H_j|}{\Gamma_j} < 1, \quad \Lambda_j > 0 \right. \\ \left. \forall j = 1, \dots, n \right\} \end{aligned}$$

and let $\mathcal{E}_{\mathcal{D}el}$ be the set of n -plets $(\mathfrak{E}_1, \dots, \mathfrak{E}_n)$ where $\mathfrak{E}_j = (a_j, e_j, N^{(j)}, P^{(j)})$ satisfies

$$0 < e_j < 1, \quad \mathbf{n}_j := k^{(3)} \times N^{(j)} \neq 0, \quad \forall j = 1, \dots, n.$$

Fix positive numbers $\mathfrak{M}_1, \dots, \mathfrak{M}_n, \mathfrak{m}_1, \dots, \mathfrak{m}_n$. Define

$$\tau_{\mathcal{D}el} : X_{\mathcal{D}el} := (H, \Gamma, \Lambda, h, g) \in \mathfrak{X}_{\mathcal{D}el} \rightarrow \mathfrak{E}_{\mathcal{D}el} = (\mathfrak{E}_{1, \mathcal{D}el}, \dots, \mathfrak{E}_{n, \mathcal{D}el})$$

with $\mathfrak{E}_{j, \mathcal{D}el} = (a_{j, \mathcal{D}el}, e_{j, \mathcal{D}el}, N_{\mathcal{D}el}^{(j)}, P_{\mathcal{D}el}^{(j)})$ and

$$\begin{aligned} a_{j, \mathcal{D}el} &= \frac{1}{\mathfrak{M}_j} \left(\frac{\Lambda_j}{\mathfrak{m}_j} \right)^2, & e_{j, \mathcal{D}el} &= \sqrt{1 - \left(\frac{\Gamma_j}{\Lambda_j} \right)^2} \\ N_{\mathcal{D}el}^{(j)} &= R_3(h_j) \mathcal{R}_1(i_j) k^{(3)} & P_{\mathcal{D}el}^{(j)} &= R_3(h_j) \mathcal{R}_1(i_j) \mathcal{R}_3(g_j) k^{(1)} \end{aligned}$$

where $i_j := \cos^{-1} \frac{H_j}{\Gamma_j}$.

We call *Delaunay map* the map

$$(2.30) \mathcal{D}el : \text{Del} = (H, \Gamma, \Lambda, h, g, \ell) \in \mathcal{D}_{\mathcal{D}el} \rightarrow (y_{\mathcal{D}el}, x_{\mathcal{D}el}) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n}$$

which is defined on the domain

$$\mathcal{D}_{\mathcal{D}el} := \mathfrak{X}_{\mathcal{D}el} \times \mathbb{T}^n$$

as the Kepler map associated to $\tau_{\mathcal{D}el}$ via the following lemma (the proof of which may be found in classical textbooks).

LEMMA 2.5 (Delaunay). $\tau_{\mathcal{D}el}$ is a bijection of $\mathfrak{X}_{\mathcal{D}el}$ onto $\mathcal{E}_{\mathcal{D}el}$. Its inverse map

$$\tau_{\mathcal{D}el}^{-1} : \mathfrak{E}_{\mathcal{D}el} = (\mathfrak{E}_{1, \mathcal{D}el}, \dots, \mathfrak{E}_{n, \mathcal{D}el}) \in \mathcal{E}_{\mathcal{D}el} \rightarrow X_{\mathcal{D}el} \in \mathfrak{X}_{\mathcal{D}el}$$

is defined by equations

$$(2.31) \quad \begin{cases} H_j = C_{\mathcal{E}}^{(j)} \cdot k^{(3)} \\ h_j := \alpha_{k^{(3)}}(k^{(1)}, \mathbf{n}_j) \end{cases} \quad \begin{cases} \Gamma_j = \|C_{\mathcal{E}}^{(j)}\| \\ g_j := \alpha_{C_{\mathcal{E}}^{(j)}}(\mathbf{n}_j, P^{(j)}) \end{cases} \quad \Lambda_j = \mathfrak{m}_j \sqrt{\mathfrak{M}_j a_j},$$

where $C_{\mathcal{E}}^{(j)}$ is as in (2.9). Furthermore, Del preserves the standard 2-form

$$\sum_{j=1}^n dy_{\mathcal{D}el}^{(j)} \wedge dx_{\mathcal{D}el}^{(j)} = \sum_{j=1}^n (dH_j \wedge dh_j + d\Gamma_j \wedge dg_j + d\Lambda_j \wedge d\ell_j).$$

Now we are ready to complete the proof of Theorem 2.1.

PROOF OF THEOREM 2.1. Let

$$\mathcal{D}_{\mathcal{P}}^* := \left\{ P = (\Theta, \chi, \Lambda, \vartheta, \kappa, \ell) \in \mathcal{D}_{\mathcal{P}} : P(P) \in \mathcal{C}_{\mathcal{D}el} \right\}.$$

It is enough to prove Theorem 2.1 on $\mathcal{D}_{\mathcal{P}}^*$, since indeed the \mathcal{P} -map is regular on $\mathcal{D}_{\mathcal{P}} = \widehat{\mathcal{D}_{\mathcal{P}}^*}$. On $\mathcal{D}_{\mathcal{P}}^*$, we consider the map

$$\phi_{\mathcal{D}_{Del}}^{\mathcal{P}} := \mathcal{D}el^{-1} \circ \mathcal{P} :$$

$$P = (\Theta, \chi, \Lambda, \vartheta, \kappa, \ell) \in \mathcal{D}_{\mathcal{P}}^* \rightarrow \text{Del} = (H, \Gamma, \Lambda, h, g, \ell) \in \mathcal{D}_{\mathcal{D}el}^* := \phi_{\mathcal{D}_{Del}}^{\mathcal{P}}(\mathcal{D}_{\mathcal{P}}^*) \subset \mathcal{D}_{\mathcal{D}el}.$$

$\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}$ gives the Delaunay coordinates at left hand side in (2.30) in terms of the P-coordinates at left hand side of (2.7) in the subset $\mathcal{D}_{\mathcal{P}}^*$ of $\mathcal{D}_{\mathcal{P}}$ the \mathcal{P} -image of which lies in the $\mathcal{D}el$ -image of $\mathcal{D}_{\mathcal{D}el}$. Clearly, $\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}$ leaves the (Λ, ℓ) unvaried. More precisely, $\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}$ decouples into two disjoint maps: the identity on the (Λ, ℓ) , and a $4n$ -dimensional map

$$\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}} : (\Theta, \chi, \vartheta, \kappa) \in \widehat{\mathcal{D}_{\mathcal{P}}^*} \rightarrow (H, \Gamma, h, g) \in \widehat{\mathcal{D}_{\mathcal{D}el}^*} = \phi_{\mathcal{D}_{Del}}^{\mathcal{P}}(\widehat{\mathcal{D}_{\mathcal{P}}^*}) \subset \widehat{\mathcal{D}_{\mathcal{D}el}}$$

on the remaining coordinates, which turns out to be a bijection of the sets $\widehat{\mathcal{D}_{\mathcal{P}}^*}$ and $\widehat{\mathcal{D}_{\mathcal{D}el}^*}$. Here, the map $\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}}$ and the sets $\widehat{\mathcal{D}_{\mathcal{P}}^*}$ and $\widehat{\mathcal{D}_{\mathcal{D}el}^*}$ do not depend on (Λ, ℓ) . Indeed, the explicit expressions of $\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}}$, $\widehat{\mathcal{D}_{\mathcal{P}}^*}$ in terms of $P = (\Theta, \chi, \Lambda, \vartheta, \kappa, \ell)$; or of $\widehat{\mathcal{D}_{\mathcal{D}el}^*}$ in terms of $\text{Del} = (H, \Gamma, \Lambda, h, g, \ell)$ involve only the $C_{\mathcal{P}}^{(j)}$, $P_{\mathcal{P}}^{(j)}$; the $C_{\mathcal{D}el}^{(j)}$, $P_{\mathcal{D}el}^{(j)}$, that do not depend on (Λ, ℓ) : (2.31) (where one has to replace \mathcal{C} with \mathcal{P}), (2.15) and (2.18).

In view of the previous consideration and of Lemma 2.5, Theorem 2.1 is implied by

LEMMA 2.6. *The map $\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}}$ preserves that standard 1-form:*

$$\sum_{j=1}^n (H_j d h_j + \Gamma_j d g_j) = \sum_{j=1}^n (\Theta_{j-1} d \vartheta_{j-1} + \chi_{j-1} d \kappa_{j-1}).$$

PROOF. We look at the analogous map

$$\widehat{\phi_{\mathcal{D}_{Del}}^{\tilde{\mathcal{P}}}} : (\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa}) \in \widehat{\mathcal{D}_{\tilde{\mathcal{P}}}^*} \rightarrow (\tilde{H}, \tilde{\Gamma}, \tilde{h}, \tilde{g}) \in \widehat{\mathcal{D}_{\mathcal{D}el}^*} = \phi_{\mathcal{D}_{Del}}^{\tilde{\mathcal{P}}}(\widehat{\mathcal{D}_{\tilde{\mathcal{P}}}^*}) \subset \widehat{\mathcal{D}_{\mathcal{D}el}}.$$

The analytical expression of this map is identical to the one of $\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}}$. This follows from the fact that $\widehat{\phi_{\mathcal{D}_{Del}}^{\tilde{\mathcal{P}}}}$ depends on the coordinates $(\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa})$ only via $C_{\tilde{\mathcal{P}}}^{(j)}$ and $\frac{x_{\tilde{\mathcal{P}}}^{(j)}}{\|x_{\tilde{\mathcal{P}}}^{(j)}\|}$ exactly as $\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}}$ depends on $(\Theta, \chi, \vartheta, \kappa)$ only via $C_{\mathcal{P}}^{(j)}$ and $\Pi_{\mathcal{P}}^{(j)}$, that $C_{\tilde{\mathcal{P}}}^{(j)}$ and $\frac{x_{\tilde{\mathcal{P}}}^{(j)}}{\|x_{\tilde{\mathcal{P}}}^{(j)}\|}$ have exactly the same expressions of $C_{\mathcal{P}}^{(j)}$ and $P^{(j)}$, apart for replacing $(\Theta, \chi, \vartheta, \kappa)$ with $(\tilde{\Theta}, \tilde{\chi}, \tilde{\vartheta}, \tilde{\kappa})$. Compare (2.29) (where one has to replace $C_{\mathcal{E}}^{(j)}$ with $C_{\tilde{\mathcal{E}}}^{(j)}$), (2.31) (where one has to replace $C_{\mathcal{E}}^{(j)}$ with $C_{\tilde{\mathcal{E}}}^{(j)}$), (2.15), (2.18), (2.24) and (2.25). But Lemmata 2.2 and 2.4 imply that $\widehat{\phi_{\mathcal{D}_{Del}}^{\tilde{\mathcal{P}}}}$ preserves that standard 1-form:

$$\sum_{j=1}^n (\tilde{H}_j d \tilde{h}_j + \tilde{\Gamma}_j d \tilde{g}_j) = \sum_{j=1}^n (\tilde{\Theta}_{j-1} d \tilde{\vartheta}_{j-1} + \tilde{\chi}_{j-1} d \tilde{\kappa}_{j-1}).$$

Then $\widehat{\phi_{\mathcal{D}_{Del}}^{\mathcal{P}}}$ does. □

2.1. The \mathcal{P} -map vs rotations and reflections

Now we discuss how the \mathcal{P} -map behaves in presence of symmetries in the Hamiltonian due to rotations or reflections.

Let $H = H(y, x)$ be the Hamiltonian governing the motion of n particles, where such particles are expressed in the canonical coordinates $(y^{(1)}, x^{(1)}), \dots, (y^{(n)}, x^{(n)})$. Assume that H is left unvaried by rotations and reflections. Namely, if

$$\phi_{\mathcal{R}, \mathcal{S}} : (y^{(j)}, x^{(j)}) \rightarrow (\mathcal{R}y^{(j)}, \mathcal{S}x^{(j)}), \quad j = 1, \dots, n$$

where \mathcal{R}, \mathcal{S} are real 3×3 matrices, then rotation invariance is

$$H \circ \phi_{\mathcal{R}, \mathcal{R}} = H \quad \forall \mathcal{R} : \mathcal{R}\mathcal{R}^t = \text{id}$$

while reflection invariance is

$$H \circ \phi_{\mathcal{S}_\sigma, \mathcal{S}_\tau} = H \quad \text{for some} \quad \mathcal{S}_\sigma = \begin{pmatrix} \sigma_1 & 0 & 0 \\ 0 & \sigma_2 & 0 \\ 0 & 0 & \sigma_3 \end{pmatrix} \quad \mathcal{S}_\tau = \begin{pmatrix} \tau_1 & 0 & 0 \\ 0 & \tau_2 & 0 \\ 0 & 0 & \tau_3 \end{pmatrix}$$

$$\sigma_i, \tau_i = \pm 1.$$

Rotation invariance is associated to the conservation, through the motion, of the total angular momentum $S_C^{(1)}$ is (2.23). Reflection invariance is not associated to integrals.

The Hamiltonian H_{hel} in (3.33) is rotation and reflection invariant, and reflection invariance holds with any choice of σ, τ .

Let

$$H_{\mathcal{P}} := H \circ \mathcal{P}.$$

The fact that $S_C^{(1)}$ is preserved along the motions of H implies that the coordinates

$$\Theta_0 = Z, \quad \vartheta_0 = \zeta, \quad \kappa_0 = \mathfrak{g}$$

do not appear in $H_{\mathcal{P}}$. Indeed, Z and ζ are integrals, while \mathfrak{g} is conjugated to $G = \|S_{\mathcal{P}}^{(1)}\|$, which is an integral for $H_{\mathcal{P}}$. Thus, the number of degrees of freedom is naturally reduced by two units, once one regards G as a prefixed external parameter. Namely, for any fixed $\chi_0 = G$, $H_{\mathcal{K}}$ may be regarded as a function of the $2(3n - 1)$ dimensional coordinates

$$\bar{\mathcal{P}} := (\bar{\Theta}, \chi, \Lambda, \bar{\vartheta}, \kappa, \ell)$$

which does not depend on κ_0 . Here,

$$\bar{\Theta} = (\Theta_1, \dots, \Theta_{n-1}), \quad \bar{\vartheta} = (\vartheta_1, \dots, \vartheta_{n-1}).$$

An analogue property is also shared with the action-angle coordinates $(\Psi, \Gamma, \Lambda, \psi, \gamma, \ell)$ described in [7, 27], and related to a set of coordinates discovered by A. Deprit [12] in the 80s (compare also [36] or the Appendix E).

The main novelty introduced by the \mathcal{P} -coordinates (that does not hold for the coordinates of [7]) is how \mathcal{P} behaves relative to reflections.

We denote as

$$\mathcal{R}_2^- := \phi_{\mathcal{S}_{\sigma^{(2)}}, \mathcal{S}_{\sigma^{(2)}}} \quad \sigma^{(2)} = (1, -1, 1)$$

the reflection of the second coordinate both for the $y^{(j)}$'s and the $x^{(j)}$'s and we let

$$\mathcal{S}^-(\Theta, \chi, \Lambda, \vartheta, \kappa, \ell) := (-\Theta, \chi, \Lambda, -\vartheta, \kappa, \ell).$$

PROPOSITION 2.2.

$$(2.32) \quad \mathcal{R}_2^- \circ \mathcal{P} = \mathcal{P} \circ \mathcal{S}^-.$$

Therefore, if $\mathbf{H} = \mathbf{H}(y, x)$ satisfies

$$\mathbf{H} \circ \mathcal{R}_2^- = \mathbf{H}$$

then $\mathbf{H}_{\mathcal{P}} := \mathbf{H} \circ \mathcal{P}$ satisfies

$$\mathbf{H}_{\mathcal{P}} \circ \mathcal{S}^- = \mathbf{H}_{\mathcal{P}}.$$

Hence, any of the points

$$\Theta_0 = \cdots = \Theta_{n-1} = 0, \quad (\vartheta_0, \cdots, \vartheta_{n-1}) = (k_0, \cdots, k_{n-1})\pi \quad \text{mod } 2\pi\mathbb{Z}^n$$

is an equilibrium point for $\mathbf{H}_{\mathcal{P}}$, for any $(\chi, \Lambda, \kappa, \ell)$.

PROOF. Defining $\mathcal{R}^{(j)} := \mathbf{T}_j \mathbf{S}_j$, $s^{(j)} := \mathbf{T}_j k^{(3)}$, we write the vectors $P_{\mathcal{P}}^{(j)}$ and $\mathbf{S}_{\mathcal{P}}^{(j)}$ (compare Eq. (2.18) and Remark 2.2(iii)) as

$$P_{\mathcal{P}}^{(j)} = \mathcal{R}^{(1)} \cdots \mathcal{R}^{(j)} k^{(3)}, \quad \mathbf{S}_{\mathcal{P}}^{(j)} = \chi_{j-1} \mathcal{R}^{(1)} \cdots \mathcal{R}^{(j)} s^{(j)}.$$

The explicit expressions of $\mathcal{R}^{(j)}$ and $s^{(j)}$ are

$$\begin{aligned} \mathcal{R}_{11}^{(j)} &= \cos \kappa_{j-1} \cos \vartheta_{j-1} - \sin \kappa_{j-1} \cos \iota_j \sin \vartheta_{j-1} \\ \mathcal{R}_{21}^{(j)} &= \cos \kappa_{j-1} \sin \vartheta_{j-1} + \sin \kappa_{j-1} \cos \iota_j \cos \vartheta_{j-1} \\ \mathcal{R}_{31}^{(j)} &= \sin \kappa_{j-1} \sin \iota_j \\ \mathcal{R}_{12}^{(j)} &= -\cos i_j \sin \kappa_{j-1} \cos \vartheta_{j-1} + \sin \vartheta_{j-1} (-\cos i_j \cos \iota_j \cos \kappa_{j-1} + \sin \iota_j \sin i_j) \\ \mathcal{R}_{22}^{(j)} &= -\cos i_j \sin \kappa_{j-1} \sin \vartheta_{j-1} - \cos \vartheta_{j-1} (-\cos i_j \cos \iota_j \cos \kappa_{j-1} + \sin \iota_j \sin i_j) \\ \mathcal{R}_{32}^{(j)} &= \cos i_j \cos \kappa_{j-1} \sin \iota_j + \sin i_j \cos \iota_j \\ \mathcal{R}_{13}^{(j)} &= \sin i_j \sin \kappa_{j-1} \cos \vartheta_{j-1} + \sin \vartheta_{j-1} (\sin i_j \cos \iota_j \cos \kappa_{j-1} + \sin \iota_j \cos i_j) \\ \mathcal{R}_{23}^{(j)} &= \sin i_j \sin \kappa_{j-1} \sin \vartheta_{j-1} - \cos \vartheta_{j-1} (\sin i_j \cos \iota_j \cos \kappa_{j-1} + \sin \iota_j \cos i_j) \\ \mathcal{R}_{33}^{(j)} &= -\sin i_j \cos \kappa_{j-1} \sin \iota_j + \cos i_j \cos \iota_j \\ s_1^{(j)} &= \sin \iota_j \sin \vartheta_{j-1} \\ s_2^{(j)} &= -\sin \iota_j \cos \vartheta_{j-1} \\ s_3^{(j)} &= \cos \iota_j. \end{aligned}$$

Then \mathcal{S}^- lets $P_{\mathcal{P}}^{(j)}$ and $\mathbf{S}_{\mathcal{P}}^{(j)}$ respectively, into

$$(P_{\mathcal{P}}^{(j)})^- := \mathcal{R}_2^- P_{\mathcal{P}}^{(j)} \quad \text{and} \quad (\mathbf{S}_{\mathcal{P}}^{(j)})^- := -\mathcal{R}_2^- \mathbf{S}_{\mathcal{P}}^{(j)}.$$

Therefore, $C_{\mathcal{P}}^{(j)} = S_{\mathcal{P}}^{(j)} - S_{\mathcal{P}}^{(j+1)}$ (with $S_{\mathcal{P}}^{(n+1)} := 0$) and $Q_{\mathcal{P}}^{(j)} = \frac{C_{\mathcal{P}}^{(j)}}{\|C_{\mathcal{P}}^{(j)}\|} \times P_{\mathcal{P}}^{(j)}$ are transformed, respectively, into

$$(C_{\mathcal{P}}^{(j)})^- := -\mathcal{R}_2^- C_{\mathcal{P}}^{(j)}, \quad (Q_{\mathcal{P}}^{(j)})^- := \mathcal{R}_2^- Q_{\mathcal{P}}^{(j)}.$$

On the other hand, $a_{j,\mathcal{P}}$ and $e_{j,\mathcal{P}}$ are left unvaried by \mathcal{S}^- . In view of Definition 2.2 and Definition 2.3, the thesis (2.32) follows. \square

The \mathcal{P} -map and the planetary problem

After the reduction of the invariance by translations, a Hamiltonian governing the motions of n planets with masses $\mu m_1, \dots, \mu m_n$ interacting among themselves and with a star with mass m_0 can be taken to be the “heliocentric” one

$$(3.33) \quad H_{\text{hel}} := \sum_{1 \leq i \leq n} \left(\frac{\|y^{(i)}\|^2}{2\mathfrak{m}_i} - \frac{\mathfrak{m}_i \mathfrak{M}_i}{\|x^{(i)}\|} \right) + \mu \sum_{1 \leq i < j \leq n} \left(\frac{y^{(i)} \cdot y^{(j)}}{m_0} - \frac{m_i m_j}{\|x^{(i)} - x^{(j)}\|} \right)$$

where $(y, x) = (y^{(1)}, \dots, y^{(n)}, x^{(1)}, \dots, x^{(n)})$ are “Cartesian coordinates” taking values on the “collision-less” phase space $\mathbb{R}^{3n} \times \mathbb{R}^{3n} \setminus \Delta$, where

$$\Delta = \left\{ x = (x^{(1)}, \dots, x^{(n)}) \in \mathbb{R}^3 \times \dots \times \mathbb{R}^3 : 0 \neq x^{(i)} \neq x^{(j)} \quad \forall 1 \leq i < j \leq n \right\}$$

endowed with the standard 2- form

$$\Omega := dy \wedge dx := \sum_{i=1}^n \sum_{j=1}^3 dy_j^{(i)} \wedge dx_j^{(i)}$$

and with

$$(3.34) \quad \mathfrak{M}_i = m_0 + \mu m_i \quad \mathfrak{m}_i = \frac{m_0 m_i}{m_0 + \mu m_i}$$

being the so-called “reduced masses”.

In the following Chapter 3.1 we describe a general property of Kepler maps, in relation to their application to the Hamiltonian H_{hel} . Then (in Chapter 3.2) we shall specialize to the case of the \mathcal{P} -map.

3.1. A general property of Kepler maps

For a general Kepler map \mathcal{K} , we denote

$$H_{\mathcal{K}}(\mathbf{K}) := H_{\text{hel}} \circ \mathcal{K} = - \sum_{j=1}^n \frac{\mathfrak{m}_j \mathfrak{M}_j}{2a_{j,\mathcal{K}}(\mathbf{X}_{\mathcal{K}})} + \mu f_{\mathcal{K}}(\mathbf{K}),$$

where

$$f_{\mathcal{K}}(\mathbf{K}) := \sum_{1 \leq i < j \leq n} \left(\frac{y_{\mathcal{K}}^{(i)} \cdot y_{\mathcal{K}}^{(j)}}{m_0} - \frac{m_i m_j}{\|x_{\mathcal{K}}^{(i)} - x_{\mathcal{K}}^{(j)}\|} \right)$$

and $y_{\mathcal{K}}^{(j)}, x_{\mathcal{K}}^{(j)}$ are as in Definition 2.2.

We denote

$$(3.35) \quad \overline{f_{\mathcal{K}}}(\mathbf{X}_{\mathcal{K}}) := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} f_{\mathcal{K}}(\mathbf{X}_{\mathcal{K}}, \ell) d\ell,$$

so that

$$f_{\mathcal{K}} = \sum_{1 \leq i < j \leq n} f_{\mathcal{K}}^{ij}, \quad \overline{f_{\mathcal{K}}} = \sum_{1 \leq i < j \leq n} \overline{f_{\mathcal{K}}^{ij}}$$

$$f_{\mathcal{K}}^{ij} := \frac{y_{\mathcal{K}}^{(i)} \cdot y_{\mathcal{K}}^{(j)}}{m_0} - \frac{m_i m_j}{\|x_{\mathcal{K}}^{(i)} - x_{\mathcal{K}}^{(j)}\|}, \quad \overline{f_{\mathcal{K}}^{ij}} := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} f_{\mathcal{K}}^{ij} dl_1 \cdots dl_n.$$

For a general Kepler map, one always has, as a consequence of (2.6),

$$(3.36) \quad -\frac{1}{2\pi} \int_{\mathbb{T}} T_{\mathcal{K}}^{(j)} dl_j = \frac{1}{2\pi} \int_{\mathbb{T}} \frac{V_{\mathcal{K}}^{(j)}}{2} dl_j = T_{\mathcal{K}}^{(j)} + V_{\mathcal{K}}^{(j)} = -\frac{\mathbf{m}\mathfrak{M}}{2a_{j,\mathcal{K}}}$$

$$\frac{1}{2\pi} \int_{\mathbb{T}} y_{\mathcal{K}}^{(j)} dl_j = 0 \quad \frac{1}{2\pi} \int_{\mathbb{T}} \frac{x_{\mathcal{K}}^{(j)}}{\|x_{\mathcal{K}}^{(j)}\|^3} dl_j = 0,$$

where we have denoted as

$$T_{\mathcal{K}}^{(j)} := \frac{\|y_{\mathcal{K}}^{(j)}\|^2}{2\mathbf{m}_j} \quad V_{\mathcal{K}}^{(j)} := -\frac{\mathbf{m}_j \mathfrak{M}_j}{\|x_{\mathcal{K}}^{(j)}\|}$$

the kinetic, potential part of $H_{\mathcal{K}}^{(j)}$ in (2.4), respectively.

Consider the average $\overline{f_{\mathcal{K}}}(\mathbf{X}_{\mathcal{K}})$ in (3.35). Due to the fact that $y_{\mathcal{K}}^{(j)}$ has zero-average, one has that only the Newtonian part contributes to $\overline{f_{\mathcal{K}}}(\mathbf{X}_{\mathcal{K}})$:

$$\overline{f_{\mathcal{K}}} = - \sum_{1 \leq i < j \leq n} \frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{dl_i dl_j}{\|x_{\mathcal{K}}^{(i)} - x_{\mathcal{K}}^{(j)}\|}.$$

We now consider any of the contributions to this sum

$$\overline{f_{\mathcal{K}}^{ij}} = -\frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{dl_i dl_j}{\|x_{\mathcal{K}}^{(i)} - x_{\mathcal{K}}^{(j)}\|} \quad 1 \leq i < j \leq n$$

and expand any such terms

$$\overline{f_{\mathcal{K}}^{ij}} = \overline{f_{\mathcal{K}}^{ij}}^{(0)} + \overline{f_{\mathcal{K}}^{ij}}^{(1)} + \overline{f_{\mathcal{K}}^{ij}}^{(2)} + \cdots$$

where

$$\overline{f_{\mathcal{K}}^{ij}}^{(h)} := -\frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{1}{h!} \frac{d^h}{d\varepsilon^h} \frac{1}{\|\varepsilon x_{\mathcal{K}}^{(i)} - x_{\mathcal{K}}^{(j)}\|} \Big|_{\varepsilon=0} dl_i dl_j$$

is proportional to $\frac{1}{a_j} \left(\frac{a_i}{a_j}\right)^h$. Then the formulae in (3.36) imply that the two first terms of this expansion are given by

$$\overline{f_{\mathcal{K}}^{ij}}^{(0)} = -\frac{m_i m_j}{a_{j,\mathcal{K}}}, \quad \overline{f_{\mathcal{K}}^{ij}}^{(1)} = 0.$$

Namely, whatever is the Kepler map that is used, the first term that depends on the secular coordinates $\mathbf{X}_{\mathcal{K}}$ is the double average of the second order term

$$\overline{f_{\mathcal{K}}^{ij}}^{(2)}(\mathbf{X}_{\mathcal{K}}) = -\frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} \frac{3(x_{\mathcal{K}}^{(i)} \cdot x_{\mathcal{K}}^{(j)})^2 - \|x_{\mathcal{K}}^{(i)}\|^2 \|x_{\mathcal{K}}^{(j)}\|^2}{\|x_{\mathcal{K}}^{(j)}\|^5} dl_i dl_j.$$

Now we specialize to the case of the \mathcal{P} -map.

3.2. The case of the \mathcal{P} -map

We denote as

$$(3.37) \quad \mathcal{H}_{\mathcal{P}}(X_{\mathcal{P}}, \ell) = h_{\text{fast}}^0(\Lambda) + \mu f_{\mathcal{P}}(X_{\mathcal{P}}, \ell) \quad X_{\mathcal{P}} := (\Theta, \chi, \Lambda, \vartheta, \kappa)$$

where

$$(3.38) \quad h_{\text{fast}}^0(\Lambda) := - \sum_{j=1}^n \frac{m_j^3 \mathfrak{M}_j^2}{2\Lambda_j^2},$$

is the Hamiltonian (3.33) expressed in \mathcal{P} -coordinates.

Using the definitions, it not difficult to see that

LEMMA 3.1. $\overline{f_{\mathcal{P}}^{ij}}$, $f_{\mathcal{P}}^{ij}$ depend, respectively, only on the coordinates

$$\overline{X_{\mathcal{P}}^{ij}} := (\Theta_i, \dots, \Theta_{j \wedge (n-1)}, \chi_{i-1}, \dots, \chi_{j \wedge (n-1)}, \Lambda_i, \Lambda_j, \vartheta_i, \dots, \vartheta_{j \wedge (n-1)}, \kappa_i, \dots, \kappa_{j-1})$$

$$\overline{P^{ij}} := (\overline{X_{\mathcal{P}}^{ij}}, \ell_i, \ell_j)$$

with $a \wedge b$ denoting the minimum of a and b .

Accordingly to the previous lemma, the “nearest-neighbor” terms $\overline{f_{\mathcal{P}}^{i,i+1}}$, with $i = 1, \dots, n-1$, depend only on

$$(3.39) \quad \overline{X_{\mathcal{P}}^{i,i+1}} = \begin{cases} (\Theta_i, \Theta_{i+1}, \chi_{i-1}, \chi_i, \chi_{i+1}, \Lambda_i, \Lambda_{i+1}, \vartheta_i, \vartheta_{i+1}, \kappa_i) & 1 \leq i \leq n-2 \\ (\Theta_{n-1}, \chi_{n-2}, \chi_{n-1}, \Lambda_{n-1}, \Lambda_n, \vartheta_{n-1}, \kappa_{n-1}) & i = n-1. \end{cases}$$

However, for the functions $\overline{f_{\mathcal{P}}^{i,i+1}}^{(2)}$, we have a special rule. Indeed, for any Kepler map \mathcal{K} , the “exterior” angular momentum $\|C_{\mathcal{K}}^{(i+1)}\|$ is an integral for $\overline{f_{\mathcal{K}}^{i,i+1}}^{(2)}$. This readily implies that any $\overline{f_{\mathcal{K}}^{i,i+1}}^{(2)}$ is *integrable*, for having four degrees of freedom and four independent, commuting integrals ($\|C_{\mathcal{K}}^{(i)} + C_{\mathcal{K}}^{(i+1)}\|$, $(C_{\mathcal{K}}^{(i)} + C_{\mathcal{K}}^{(i+1)}) \cdot k^{(3)}$, $\|C_{\mathcal{K}}^{(i+1)}\|$ and $\overline{f_{\mathcal{K}}^{i,i+1}}^{(2)}$ itself). This fact has been firstly noticed, in the three-body case ($i = 1, n = 2$), by R. Harrington [19] who, using the Jacobi reduction of the nodes $\mathcal{J}ac$, where the coordinates are named

$$G_i, \quad g_i, \quad \Lambda_i, \quad \ell_i, \quad i = 1, 2$$

(with $G_i = \|C^{(i)}\|$, g_i related to the perihelia directions, and $G := \|C\|$, $C = C^{(1)} + C^{(2)}$ appearing as an external parameter), noticed that $\overline{f_{\mathcal{J}ac}^{12}}^{(2)}$ depends only on $(G, G_1, G_2, \gamma_1, \Lambda_1, \Lambda_2)$.

Let us now inspect how the integrability of $\overline{f_{\mathcal{P}}^{i,i+1}}^{(2)}$ is exhibited in terms of the \mathcal{P} -map. Since $\|C_{\mathcal{P}}^{(n)}\| = \chi_{n-1}$, one has that $\overline{f_{\mathcal{P}}^{n-1,n}}^{(2)}$ does not depend on κ_{n-1} , and hence, by (3.39), depends only on

$$\overline{X_{\mathcal{P}}^{n-1,n}} := (\Theta_{n-1}, \chi_{n-2}, \chi_{n-1}, \Lambda_{n-1}, \Lambda_n, \vartheta_{n-1}).$$

This fact, for $n \geq 3$, is no longer true for $i = 1, \dots, n-2$, because in that case $\chi_i \neq \|C_{\mathcal{P}}^{(i+1)}\|$ (indeed, $\chi_i = \|S_{\mathcal{P}}^{(i+1)}\|$). However, since, for $(\Theta_{i+1}, \vartheta_{i+1}) = (0, \pi)$,

$\|C_{\mathcal{P}}^{(i+1)}\|$ reduces to

$$\|C_{\mathcal{P}}^{(i+1)}\|_{(\Theta_{i+1}, \vartheta_{i+1})=(0, \pi)} = \chi_i - \chi_{i+1} \quad i = 1, \dots, n-2,$$

one has that the functions

$$\overline{\overline{f_{\mathcal{P}}^{i,i+1}}^{(2)}} := \overline{f_{\mathcal{P}}^{i,i+1}}^{(2)} \Big|_{(\Theta_{i+1}, \vartheta_{i+1})=(0, \pi)}, \quad i = 1, \dots, n-2$$

do not depend on κ_i and hence, by (3.39) depend only on

$$\overline{X_{\mathcal{P}}^{i,i+1}} := (\Theta_i, \chi_{i-1}, \chi_i, \chi_{i+1}, \Lambda_i, \Lambda_{i+1}, \vartheta_i), \quad i = 1, \dots, n-2.$$

In the following lemma we provide their explicit expressions.

LEMMA 3.2. *The function $\overline{f_{\mathcal{P}}^{n-1,n}}^{(2)}$ and, for $n \geq 3$ and $1 \leq i \leq n-2$, the functions $\overline{f_{\mathcal{P}}^{i,i+1}}^{(2)}$ have the following expressions*

$$\begin{aligned} \overline{f_{\mathcal{P}}^{n-1,n}}^{(2)} &= m_{n-1} m_n \frac{a_{n-1}^2}{4a_n^3} \frac{\Lambda_n^3}{\chi_{n-1}^5} \left[\frac{5}{2} (3\Theta_{n-1}^2 - \chi_{n-1}^2) \right. \\ &\quad - \frac{3}{2} \frac{4\Theta_{n-1}^2 - \chi_{n-1}^2}{\Lambda_{n-1}^2} \left(\chi_{n-2}^2 + \chi_{n-1}^2 - 2\Theta_{n-1}^2 \right. \\ &\quad \left. \left. + 2\sqrt{(\chi_{n-1}^2 - \Theta_{n-1}^2)(\chi_{n-2}^2 - \Theta_{n-1}^2)} \cos \vartheta_{n-1} \right) \right. \\ &\quad \left. + \frac{3}{2} \frac{(\chi_{n-1}^2 - \Theta_{n-1}^2)(\chi_{n-2}^2 - \Theta_{n-1}^2)}{\Lambda_{n-1}^2} \sin^2 \vartheta_{n-1} \right] \end{aligned} \quad (3.40)$$

and

$$\begin{aligned} \overline{f_{\mathcal{P}}^{i,i+1}}^{(2)} &= m_i m_{i+1} \frac{a_i^2}{4a_{i+1}^3} \frac{\Lambda_{i+1}^3}{\chi_i^2 (\chi_i - \chi_{i+1})^3} \left[\frac{5}{2} (3\Theta_i^2 - \chi_i^2) \right. \\ &\quad - \frac{3}{2} \frac{4\Theta_i^2 - \chi_i^2}{\Lambda_i^2} \left(\chi_{i-1}^2 + \chi_i^2 - 2\Theta_i^2 + 2\sqrt{(\chi_i^2 - \Theta_i^2)(\chi_{i-1}^2 - \Theta_i^2)} \cos \vartheta_i \right) \\ &\quad \left. + \frac{3}{2} \frac{(\chi_i^2 - \Theta_i^2)(\chi_{i-1}^2 - \Theta_i^2)}{\Lambda_i^2} \sin^2 \vartheta_i \right]. \end{aligned} \quad (3.41)$$

Lemma 3.2 is proved in Appendix B. Here, we limit to the following.

REMARK 3.3.

- (i) The formula in (3.41) holds also for complex values of the coordinates, provided that $\arg(\chi_i - \chi_{i+1}) \in (-\frac{\pi}{2}, \frac{\pi}{2}] \bmod 2\pi$.
- (ii) The importance of the formulae in (3.40) and (3.41), which is the main feature of the \mathcal{P} -map, is that, exploiting the equilibrium for $(\Theta_i, \vartheta_i) = (0, \pi)$, the integration of $\overline{f_{\mathcal{P}}^{n-1,n}}^{(2)}$ and of $\overline{f_{\mathcal{P}}^{i,i+1}}^{(2)}$ can be performed *explicitly*, switching to a suitable associated *convergent* Birkhoff series, as Lemma 3.4 below states. Direct integrations of $\overline{f_{\mathcal{K}}^{n-1,n}}^{(2)}$, for example, starting with Hamiltonian computed in [19], appear technically much more involved and, up to now, are not known.

LEMMA 3.4. *It is possible to find complex domains $\overline{\mathfrak{B}}_i$ with non-empty real part and a canonical, real-analytic change of coordinates*

$$(3.42) \quad \overline{\phi_{\text{int}}^i} : (p_i, q_i, y_i^*, \mathbf{x}_i^*) \in \overline{\mathfrak{B}}_i \rightarrow (\Theta_i, \vartheta_i, y_i, \mathbf{x}_i)$$

where

$$(3.43) \quad \begin{aligned} y_i^* &:= \begin{cases} (\chi_{n-2}^*, \chi_{n-1}^*, \Lambda_{n-1}^*, \Lambda_n^*) & i = n-1 \\ (\chi_{i-1}^*, \chi_i^*, \chi_{i+1}^*, \Lambda_i^*, \Lambda_{i+1}^*) & i = 1, \dots, n-2 \ \& \ n \geq 3 \end{cases} \\ \mathbf{x}_i^* &:= \begin{cases} (\kappa_{n-2}^*, \kappa_{n-1}^*, \ell_{n-1}^*, \ell_n^*) & i = n-1 \\ (\kappa_{i-1}^*, \kappa_i^*, \kappa_{i+1}^*, \ell_i^*, \ell_{i+1}^*) & i = 1, \dots, n-2 \ \& \ n \geq 3 \end{cases} \\ y_i &:= \begin{cases} (\chi_{n-2}, \chi_{n-1}, \Lambda_{n-1}, \Lambda_n) & i = n-1 \\ (\chi_{i-1}, \chi_i, \chi_{i+1}, \Lambda_i, \Lambda_{i+1}) & i = 1, \dots, n-2 \ \& \ n \geq 3 \end{cases} \\ \mathbf{x}_i &:= \begin{cases} (\kappa_{n-2}, \kappa_{n-1}, \ell_{n-1}, \ell_n) & i = n-1 \\ (\kappa_{i-1}, \kappa_i, \kappa_{i+1}, \ell_i, \ell_{i+1}) & i = 1, \dots, n-2 \ \& \ n \geq 3 \end{cases} \end{aligned}$$

such that

$$(3.44) \quad \overline{h_{\text{sec}}^i} := \begin{cases} \overline{f_{\mathcal{P}}^{n-1, n}^{(2)}} \circ \overline{\phi_{\text{int}}^{n-1}} & i = n-1 \\ \overline{f_{\mathcal{P}}^{i, i+1}^{(2)}} \circ \overline{\phi_{\text{int}}^i} & i = 1, \dots, n-2 \ \& \ n \geq 3 \end{cases}$$

depends only on

$$Y_i^* := \begin{cases} \left(\frac{p_{n-1}^2 + q_{n-1}^2}{2}, \Lambda_{n-1}^*, \Lambda_n^*, \chi_{n-2}^*, \chi_{n-1}^* \right) & i = n-1 \\ \left(\frac{p_i^2 + q_i^2}{2}, \Lambda_i^*, \Lambda_{i+1}^*, \chi_{i-1}^*, \chi_i^*, \chi_{i+1}^* \right) & i = 1, \dots, n-2 \ \& \ n \geq 3. \end{cases}$$

The transformation $\overline{\phi_{\text{int}}^i}$ may be chosen so as to verify

$$y_i^* = y_i, \quad (\Theta_i, \vartheta_i, \mathbf{x}_i - \mathbf{x}_i^*) = F_i(p_i, q_i, y_i^*)$$

$$(3.45) \quad \overline{\phi_{\text{int}}^i}(-p_i, -q_i, y_i^*, \mathbf{x}_i^*) = (-\Theta_i, -\vartheta_i, y_i, \mathbf{x}_i)$$

if

$$\overline{\phi_{\text{int}}^i}(p_i, q_i, y_i^*, \mathbf{x}_i^*) = (\Theta_i, \vartheta_i, y_i, \mathbf{x}_i).$$

Lemma 3.4 is proved in the following Chapter 5.2.1.

Global Kolmogorov tori in the planetary problem

In this chapter we show how the \mathcal{P} -map can be used to prove Theorem A. We defer to the next Chapter 5 more technical parts.

4.1. A domain of holomorphy

A typical practice, in order to use perturbation theory techniques, is to extend Hamiltonians governing dynamical systems to the complex field, and then to study their holomorphy properties.

In this section we aim to discuss a domain of holomorphy for the perturbing function $f_{\mathcal{P}}$ in (3.37), regarded as a function of complex coordinates. We shall choose it of the following form

$$\mathfrak{D}_{\mathcal{P}} := \mathbb{T}_{\Theta^+, \vartheta^+} \times (\mathbb{X}_{\theta} \times \overline{\mathbb{T}}_s^n) \times (\mathbb{A}_{\theta} \times \overline{\mathbb{T}}_s^n),$$

where, for given positive numbers

$$\Theta_j^+, \quad \vartheta_j^+, \quad G_i^{\pm}, \quad \Lambda_i^{\pm}, \quad \theta_i, \quad s$$

with $i = 1, \dots, n$, $j = 1, \dots, n-1$,

$$\mathbb{T}_{\Theta^+, \vartheta^+} := \left\{ (\overline{\Theta}, \overline{\vartheta}) = (\Theta_1, \dots, \Theta_{n-1}, \vartheta_1, \dots, \vartheta_{n-1}) \in \mathbb{C}^{n-1} \times \mathbb{T}_{\mathbb{C}}^{n-1} : \right. \\ \left. |\vartheta_j - \pi| \leq \vartheta_j^+, \quad |\Theta_j| \leq \Theta_j^+, \quad \forall j = 1, \dots, n-1 \right\}$$

$$(4.46) \quad \mathbb{X}_{\theta} := \left\{ \chi = (\chi_0, \dots, \chi_{n-1}) \in \mathbb{C}^n : G_j^- \leq |\chi_{j-1} - \chi_j| \leq G_j^+, \right. \\ \left. |\operatorname{Im}(\chi_{j-1} - \chi_j)| \leq \theta_j \forall j = 1, \dots, n \right\}$$

$$\mathbb{A}_{\theta} := \left\{ \Lambda = (\Lambda_1, \dots, \Lambda_n) \in \mathbb{C}^n : \Lambda_j^- \leq |\Lambda_j| \leq \Lambda_j^+, \quad |\operatorname{Im} \Lambda_j| \leq \theta_j \right. \\ \left. \forall j = 1, \dots, n \right\}$$

$$\overline{\mathbb{T}}_s := \mathbb{T} + i[-s, s]$$

with $\chi_n := 0$.

The domain $\mathfrak{D}_{\mathcal{P}}$ will be determined as the intersection of the “collision-less” set, where, as functions of complex variables, the mutual distances of the planets

$$d_{j,\mathcal{P}} := \|x_{\mathcal{P}}^{(j)} - x_{\mathcal{P}}^{(j+1)}\|$$

are far away from zero, with the holomorphy domain of \mathcal{P} , where, again as functions of complex variables, the absolute values $|e_{j,\mathcal{P}}|$ of eccentricities in (2.18)

are bounded away from 0 and 1, those of the inclinations $|\iota_j|, |i_j|$ in (2.14) are away from 0 and, finally, *Kepler equation* (2.2) provides a holomorphic solution.

The latter issue is not a peculiarity of this problem, since it naturally arises in the context of the two-body problem's equations. In the early XX century, T. Levi Civita [24] studied the holomorphy of the solution of Kepler's Equation with respect to the eccentricity. The holomorphy with respect to the mean anomaly has been investigated, using similar arguments as in [24], in [4]. Here, we address the problem of determining the holomorphy with respect to both the arguments.

PROPOSITION 4.1. *Let $\widehat{e} = 0.6627\dots$ be the solution of*

$$(4.47) \quad 0 \leq \rho \leq 1 \quad \& \quad \frac{\rho e^{\sqrt{1+\rho^2}}}{1 + \sqrt{1+\rho^2}} = 1.$$

Then for any $0 < \bar{e} < \widehat{e}$, one can find a positive number $\bar{\ell}$ depending on \bar{e} such that, for any $e = e_1 + ie_2 \in \mathbb{C}$, with $|e| \leq \bar{e}$, the complex Kepler equation

$$\zeta - e \sin \zeta = \ell$$

has a unique solution $\zeta(\ell, e)$ which turns out to be real-analytic for $\ell \in \overline{\mathbb{T}}_{\bar{\ell}}$.

The following result completes the study of the holomorphy of $f_{\mathcal{P}}$.

PROPOSITION 4.2. *Let \widehat{e} be as in Proposition 4.1. For any given $\underline{e}_i, \bar{e}_i$, with*

$$0 < \underline{e}_i < \bar{e}_i < \widehat{e} \quad i = 1, \dots, n$$

it is possible to find positive numbers

$$\mathcal{A}_j, \quad \mathcal{B}_j, \quad \bar{\mathcal{C}}_i > \underline{\mathcal{C}}_i, \quad \bar{d}_j, \quad s \in (0, 1), \quad \sigma \in (0, 1)$$

such that, if the following inequalities are satisfied

$$(4.48) \quad \begin{aligned} & \underline{\mathcal{C}}_i \Lambda_i^+ < G_i^- < G_i^+ < \bar{\mathcal{C}}_i \Lambda_i^-; \\ & \max \left\{ \frac{\theta_i}{\Lambda_i^-}, \frac{\theta_i}{G_i^-}, \sum_{i=1}^{n-1} \left| \sin^{-1} \left(\frac{G_i^+}{G_{i+1}^-} \right) \right|, \frac{\Theta_j^+}{G_n^-}, \sum_{i=1}^{n-1} \frac{G_i^+}{G_n^-}, \vartheta_j^+, |\operatorname{Im} \kappa_j|, |\operatorname{Im} \ell_i| \right\} \leq s \\ & \vartheta_j^+ \leq \min \left\{ \frac{\mathcal{A}}{G_n^+} \sqrt{(G_j^-)^2 - (\underline{\mathcal{C}}_j \Lambda_j^+)^2}, \frac{\mathcal{B}}{G_n^+} \sqrt{(\bar{\mathcal{C}}_j \Lambda_j^-)^2 - (G_j^+)^2} \right\}, \end{aligned}$$

then the eccentricities $e_{i,\mathcal{P}}$, inclinations ι_i, i_i and the mutual distances $d_{i,\mathcal{P}}$ verify

$$(4.49) \quad \underline{e}_i \leq |e_{i,\mathcal{P}}| \leq \bar{e}_i, \quad \max_{i,j} \left\{ |\cos \iota_i|, |\cos i_j| \right\} \leq \sigma, \quad |d_{i,\mathcal{P}}| \geq \bar{d}$$

Proposition 4.1 and Proposition 4.2 are proved in Appendix A.1 and A.2, respectively. We shall use them in the form below. We remark that the super-exponential decay of the semi-major axes ratio will be used only in Chapter 5.2 below.

COROLLARY 4.1 (choice of parameters). *Fix $\underline{e}_i < \bar{e}_i$, $c \in (0, 1)$, and let $\underline{\mathcal{C}}_i <$*

$$\underline{\mathcal{C}}_i^* < \bar{\mathcal{C}}_i^* < \bar{\mathcal{C}}_i, \quad \mathcal{D}_i := \min \left\{ \mathcal{A} \sqrt{(\underline{\mathcal{C}}_i^*)^2 - (\underline{\mathcal{C}}_i)^2}, \mathcal{B} \sqrt{(\bar{\mathcal{C}}_i)^2 - (\bar{\mathcal{C}}_i^*)^2} \right\}, \quad \mathcal{D} :=$$

$\min_{1 \leq j \leq n-1} \frac{D_j}{\bar{c}_n^*} \frac{m_j \sqrt{\mathfrak{M}_j}}{m_n \sqrt{\mathfrak{M}_n}}, \alpha < \frac{s}{\bar{D}}$. Define, for $i = 1, \dots, n$ and $j = 1, \dots, n-1$,

$$(4.50) \quad \begin{aligned} \Lambda_i^\pm &:= m_i \sqrt{\mathfrak{M}_i a_i^\pm}, & G_i^+ &:= \bar{c}_i^* \Lambda_i^-, & G_i^- &:= \underline{c}_i^* \Lambda_i^+, & \Theta_j^+ &:= s G_n^-, \\ \vartheta_j^+ &:= \mathcal{D}_i \frac{\Lambda_i^-}{G_n^+} & \theta_i &:= s \sqrt{\Lambda_i^-} \end{aligned}$$

where a_i^\pm is as in (*). Then $f_{\mathcal{P}}$ is real-analytic in the domain $\mathfrak{D}_{\mathcal{P}}$.

4.2. A normal form for the planetary problem

DEFINITION 4.1 ([2]). Given $m, \nu_1, \dots, \nu_m \in \mathbb{N}$, $\nu := \nu_1 + \dots + \nu_m$, let

$$\mathfrak{L}_0 \supset \mathfrak{L}_1 \supset \mathfrak{L}_2 \supset \dots \supset \mathfrak{L}_m = \{0\}$$

be a decreasing sequence of sub-lattices of \mathbb{Z}^ν defined by

$$(4.51) \quad \mathfrak{L}_0 := \mathbb{Z}^\nu, \quad \mathfrak{L}_i := \{k = (k_1, \dots, k_m) \in \mathbb{Z}^\nu, \quad k_j \in \mathbb{Z}^{\nu_j} : k_1 = \dots = k_i = 0\}$$

with $i = 1, \dots, m$. Next, given $\gamma, \gamma_1, \dots, \gamma_m, \tau \in \mathbb{R}_+$, we define the set $D_{\gamma_1 \dots \gamma_m; \tau}^\nu$ of the $(\gamma_1 \dots \gamma_m; \tau)$ -diophantine numbers via the following formulae:

$$\begin{aligned} D_{\gamma; \tau}^{\nu, K, i} &:= \left\{ \omega \in \mathbb{R}^\nu : |\omega \cdot k| \geq \frac{\gamma}{|k|^\tau} \quad \forall k \in \mathfrak{L}_{i-1} \setminus \mathfrak{L}_i, |k|_1 \leq K \right\} \\ D_{\gamma_1 \dots \gamma_m; \tau}^{\nu, K} &:= \bigcap_{i=1}^m D_{\gamma_i; \tau}^{\nu, K, i} & D_{\gamma_1 \dots \gamma_m; \tau}^\nu &:= \bigcap_{K \in \mathbb{N}} D_{\gamma_1 \dots \gamma_m; \tau}^{\nu, K} \end{aligned}$$

In other words $\omega = (\omega_1, \dots, \omega_m) \in D_{\gamma_1 \dots \gamma_m; \tau}^\nu$ if, for any $k = (k_1, \dots, k_m) \in \mathbb{Z}^\nu \setminus \{0\}$, with $k_j \in \mathbb{Z}^{\nu_j}$,

$$(4.52) \quad |\omega \cdot k| = \left| \sum_{j=1}^m \omega_j \cdot k_j \right| \geq \begin{cases} \frac{\gamma_1}{|k|^\tau} & \text{if } k_1 \neq 0; \\ \frac{\gamma_2}{|k|^\tau} & \text{if } k_1 = 0, \quad k_2 \neq 0; \\ \vdots & \\ \frac{\gamma_m}{|k_m|^\tau} & \text{if } k_1 = \dots = k_{m-1} = 0, \quad \dots, \quad k_m \neq 0. \end{cases}$$

REMARK 4.1. The choice $m = 1, \gamma_1 := \gamma$ gives the usual Diophantine set $\mathcal{D}_{\gamma, \tau}^\nu$. The $m = 2$ -case, $\mathcal{D}_{\gamma_1, \gamma_2, \tau}^\nu$, with $\gamma_1 = O(1)$ and $\gamma_2 = O(\mu)$, where μ is the strength of the planetary masses has been considered in [2] for the proof of the Fundamental Theorem, mentioned in the introduction.

The following result is proven in the next Chapter 5. It is unavoidably detailed.

PROPOSITION 4.3. Let m_j, \mathfrak{M}_j be as in (3.34) and $m_j := \sum_{i=1}^{j-1} m_i$, with $j = 2, \dots, n$. There exists a number \mathfrak{c} , depending only on $n, m_0, \dots, m_n, a_n^\pm, \underline{e}_j, \bar{e}_j$, and a number $0 < \bar{c} < 1$, depending only on n such that, for any fixed positive numbers $\bar{\gamma} < 1 < \bar{K}, \alpha > 0$ verifying

$$(4.53) \quad \bar{K} \leq \frac{\mathfrak{c}}{\alpha^{3/2}}$$

and

$$(4.54) \quad \frac{1}{c} \max \left\{ \mu \left(\frac{a_n^+}{a_1} \right)^5 \frac{\bar{K}^{2\bar{\tau}+2}}{\bar{\gamma}^2}, \frac{\bar{K}^{2(\bar{\tau}+1)} \alpha}{\bar{\gamma}^2} \right\} < 1$$

there exist natural numbers ν_1, \dots, ν_{2n-1} , with $\sum_j \nu_j = 3n - 2$, open sets $B_j^* \subset B_{\varepsilon_j}^2$, $X^* \subset X$, positive real numbers $\gamma_1 > \dots > \gamma_{2n-1} \varepsilon_1, \dots, \varepsilon_{n-1}, \bar{r}_1, \dots, \bar{r}_{n-1}, \tilde{r}_1, \dots, \tilde{r}_n$, a domain

$$\mathfrak{D}_n := B_{\sqrt{2\bar{r}}} \times X_{\bar{r}} \times \mathcal{A}_{\bar{r}} \times \mathbb{T}_{\bar{c}s}^n \times \mathbb{T}_{\bar{c}s}^n$$

a sub-domain of the form

$$\mathfrak{D}_n^* := B_{\sqrt{2\bar{r}}}^* \times X_{\bar{r}}^* \times \mathcal{A}_{\bar{r}} \times \mathbb{T}_{\bar{c}s}^n \times \mathbb{T}_{\bar{c}s}^n$$

verifying

$$(4.55) \quad \text{meas } \mathfrak{D}_n^* \geq \left(1 - \frac{\bar{\gamma}}{c}\right) \text{meas } \mathfrak{D}_n$$

a real-analytic transformation

$$\phi_n : (p, q, \chi, \Lambda, \kappa, \ell) \in \mathfrak{D}_n^* \rightarrow \mathfrak{D}_{\mathcal{P}}$$

which conjugates $H_{\mathcal{P}}$ to

$$H_n(p, q, \chi, \Lambda, \kappa, \ell) := H_{\mathcal{P}} \circ \phi_n = h_{\text{fast,sec}}(p, q, \chi, \Lambda) + \mu f_{\text{exp}}(p, q, \chi, \Lambda, \kappa, \ell)$$

where $f_{\text{exp}}(p, q, \chi, \Lambda, \kappa, \ell)$ is independent of κ_0 , and the following holds.

1. The function $h_{\text{fast,sec}}(p, q, \chi, \Lambda)$ is a sum

$$h_{\text{fast,sec}}(p, q, \chi, \Lambda) = h_{\text{fast}}(\Lambda) + \mu h_{\text{sec}}(p, q, \chi, \Lambda)$$

where, if

$$\hat{y}_i := \left(\frac{p_i^2 + q_i^2}{2}, \dots, \frac{p_{n-1}^2 + q_{n-1}^2}{2}, \chi_{i-1}, \dots, \chi_{n-1}, \Lambda_i, \dots, \Lambda_n \right)$$

then h_{fast} and h_{sec} are given by

$$h_{\text{fast}}(\Lambda) = - \sum_{j=1}^n \frac{m_j^3 \mathfrak{M}_j^2}{2\Lambda_j^2} - \mu \sum_{j=2}^n \frac{\mathfrak{M}_j m_j^2 m_j}{\Lambda_j^2}, \quad h_{\text{sec}}(p, q, \chi, \Lambda) = \sum_{i=1}^{n-1} h_{\text{sec}}^i(\hat{y}_i)$$

where the functions h_{sec}^i have an analytic extension on \mathfrak{D}_n and verify

$$c \frac{(a_{n-j}^+)^2}{(a_{n-j+1}^-)^3} \leq |h_{\text{sec}}^j(\hat{y}_j)| \leq \frac{1}{c} \frac{(a_{n-j}^+)^2}{(a_{n-j+1}^-)^3}.$$

2. The function f_{exp} satisfies

$$|f_{\text{exp}}| \leq \frac{1}{c} \frac{e^{-c\bar{K}}}{a_1}.$$

3. If ζ is \hat{y}_1 deprived of χ_0 , the frequency-map

$$\zeta \rightarrow \omega_{\text{fast,sec}}(\zeta) := \partial_{\zeta} h_{\text{fast,sec}}(\zeta)$$

is a diffeomorphism of $\Pi_{\zeta}(B_{\sqrt{2\bar{\tau}}}^* \times X_{\bar{\tau}}^* \times \mathcal{A}_{\bar{\tau}}^*)$ and, moreover, it satisfies (4.52), with $m = 2n - 1$, $\tau = \bar{\tau} > 2$, and

$$(4.56) \quad \nu_j := \begin{cases} 1 & j = 1, \dots, n \\ 2 & j = 3, n = 2 \\ 3 & j = n + 1, n \geq 3 \\ 2 & n + 2 \leq j \leq 2n - 2, n \geq 4 \\ 1 & j = 2n - 1, n \geq 3 \end{cases}$$

$$\omega_j := \begin{cases} \partial_{\Lambda_j} h_{\text{fast,sec}} & j = 1, \dots, n \\ \partial_{\left(\frac{p_1^2 + q_1^2}{2}, \chi_1\right)} h_{\text{fast,sec}} & j = 3, n = 2 \\ \partial_{\left(\frac{p_{n-1}^2 + q_{n-1}^2}{2}, \chi_{n-2}, \chi_{n-1}\right)} h_{\text{fast,sec}} & j = n + 1, n \geq 3 \\ \partial_{\left(\frac{p_{2n-j}^2 + q_{2n-j}^2}{2}, \chi_{2n-j-1}\right)} h_{\text{fast,sec}} & n + 2 \leq j \leq 2n - 2, n \geq 4 \\ \partial_{\frac{p_1^2 + q_1^2}{2}} h_{\text{fast,sec}} & j = 2n - 1, n \geq 3 \end{cases}$$

$$\gamma_j := \begin{cases} \frac{1}{a_j^-} \bar{\gamma} & 1 \leq j \leq n \\ \frac{\mu(a_{j-n}^+)^2}{(a_{j+1-n}^-)^3} \bar{\gamma} & n + 1 \leq j \leq 2n - 1 \end{cases}$$

4. The mentioned constants are

$$\varepsilon_j := \mathbf{c} \sqrt{\theta_j}, \quad \bar{r}_j := \frac{\theta_j \bar{\gamma}}{K^{\bar{\tau}+1}}, \quad \tilde{r}_i := \mathbf{c} \theta_j$$

with $\bar{\tau} > 2$.

4.3. A “multi-scale” KAM Theorem and proof of Theorem A

In this section we state a “multi-scale” KAM Theorem and next we show how this theorem applies to the Hamiltonian H_n so as to obtain the proof of Theorem A.

THEOREM 4.1 (Multi-scale KAM Theorem). *Let $m, \ell, \nu_1, \dots, \nu_m \in \mathbb{N}$, $\nu := \nu_1 + \dots + \nu_m \geq \ell$, $\tau_* > \nu$, $\gamma_1 \geq \dots \geq \gamma_m > 0$, $0 < 4s \leq \bar{s} < 1$, $\rho_1, \dots, \rho_{\ell}, r_1, \dots, r_{\nu-\ell}$,*

$\varepsilon_1, \dots, \varepsilon_\ell > 0$, $B_1, \dots, B_\ell \subset \mathbb{R}^2$, $D_j := \left\{ \frac{x^2+y^2}{2} \in \mathbb{R} : (x, y) \in B_j \right\} \subset \mathbb{R}$, $B := B_1 \times \dots \times B_\ell \subset \mathbb{R}^{2\ell}$, $D := D_1 \times \dots \times D_\ell \subset \mathbb{R}^\ell$, $C \subset \mathbb{R}^{\nu-\ell}$, $A := D_\rho \times C_r$. Let

$$\mathbf{H}(p, q, I, \psi) = \mathbf{h}(p, q, I) + f(p, q, I, \psi)$$

be real-analytic on $B_{\sqrt{2\rho}} \times C_r \times \mathbb{T}_{\bar{s}+s}^{\nu-\ell}$, where $\mathbf{h}(p, q, I)$ depends on (p, q) only via

$$J(p, q) := \left(\frac{p_1^2 + q_1^2}{2}, \dots, \frac{p_\ell^2 + q_\ell^2}{2} \right).$$

Assume that $\omega_0 := \partial_{(J(p,q), I)} \mathbf{h}$ is a diffeomorphism of A with non singular Hessian matrix $U := \partial_{(J(p,q), I)}^2 \mathbf{h}$ and let U_k denote the $(\nu_k + \dots + \nu_m) \times \nu$ submatrix of U , i.e., the matrix with entries $(U_k)_{ij} = U_{ij}$, for $\nu_1 + \dots + \nu_{k-1} + 1 \leq i \leq \nu$, $1 \leq j \leq \nu$, where $2 \leq k \leq m$. Let

$$M \geq \sup_A \|U\|, \quad M_k \geq \sup_A \|U_k\|, \quad \bar{M} \geq \sup_A \|U^{-1}\|, \quad E \geq \|f\|_{\rho, \bar{s}+s}$$

$$\bar{M}_k \geq \sup_A \|T_k\| \quad \text{if} \quad U^{-1} = \begin{pmatrix} T_1 \\ \vdots \\ T_m \end{pmatrix} \quad 1 \leq k \leq m.$$

Define

$$K := \frac{6}{s} \log_+ \left(\frac{EM_1^2 L}{\gamma_1^2} \right)^{-1} \quad \text{where} \quad \log_+ a := \max\{1, \log a\}$$

$$\hat{\rho}_k := \frac{\gamma_k}{3M_k K^{\tau_*+1}}, \quad \hat{\rho} := \min\{\hat{\rho}_1, \dots, \hat{\rho}_m, \rho_1, \dots, \rho_\ell, r_1, \dots, r_{\nu-\ell}\}$$

$$L := \max\{\bar{M}, M_1^{-1}, \dots, M_m^{-1}\}$$

$$\hat{E} := \frac{EL}{\hat{\rho}^2}.$$

Then one can find two numbers $\hat{c}_\nu > c_\nu$ depending only on ν such that, if the perturbation f is so small that the following ‘‘KAM condition’’ holds

$$\hat{c}_\nu \hat{E} < 1,$$

for any $\omega \in \Omega_* := \omega_0(D) \cap \mathcal{D}_{\gamma_1, \dots, \gamma_m, \tau_*}$, one can find a unique real-analytic embedding

$$\begin{aligned} \phi_\omega : \quad \vartheta = (\hat{\vartheta}, \bar{\vartheta}) \in \mathbb{T}^\nu &\rightarrow (\hat{v}(\vartheta; \omega), \hat{\vartheta} + \hat{u}(\vartheta; \omega), \mathcal{R}_{\bar{\vartheta} + \bar{u}(\vartheta; \omega)} w_1, \dots, \mathcal{R}_{\bar{\vartheta} + \bar{u}(\vartheta; \omega)} w_\ell) \\ &\in \text{Re } C_r \times \mathbb{T}^{\nu-\ell} \times \text{Re } B_{\sqrt{2r}}^{2\ell} \end{aligned}$$

where $r := c_\nu \hat{E} \hat{\rho}$ such that $\mathbb{T}_\omega := \phi_\omega(\mathbb{T}^\nu)$ is a real-analytic ν -dimensional H-invariant torus, on which the H-flow is analytically conjugated to $\vartheta \rightarrow \vartheta + \omega t$. Furthermore, the map $(\vartheta; \omega) \rightarrow \phi_\omega(\vartheta)$ is Lipschitz and one-to-one and the invariant set $\mathbf{K} := \bigcup_{\omega \in \Omega_*} \mathbb{T}_\omega$ satisfies the following measure estimate

$$\text{meas} \left(\text{Re} (D_r) \times \mathbb{T}^n \setminus \mathbf{K} \right) \leq c_\nu \left(\text{meas} (D \setminus D_{\gamma_1, \dots, \gamma_m, \tau_*} \times \mathbb{T}^n) + \text{meas} (\text{Re} (D_r) \setminus D) \times \mathbb{T}^n \right),$$

where $D_{\gamma_1, \dots, \gamma_m, \tau_*}$ denotes the ω_0 -pre-image of $\mathcal{D}_{\gamma_1, \dots, \gamma_m, \tau_*}$ in D . Finally, on $\mathbb{T}^\nu \times \Omega_*$, the following uniform estimates hold

$$\begin{aligned} |v_k(\cdot; \omega) - I_k^0(\omega)| &\leq c_\nu \left(\frac{\bar{M}_k}{\bar{M}} + \frac{M_k}{M_1} \right) \hat{E} \hat{\rho} \\ |u(\cdot; \omega)| &\leq c_\nu \hat{E} s \end{aligned}$$

where v_k denotes the projection of $v = (\hat{v}, \bar{v}) \in \mathbb{R}^{\nu_1} \times \dots \times \mathbb{R}^{\nu_m}$ over \mathbb{R}^{ν_k} , $\bar{v}_k := \frac{|w_k|^2}{2}$ and $I^0(\omega) = (I_1^0(\omega), \dots, I_\nu^0(\omega)) \in D$ is the ω_0 -pre-image of $\omega \in \Omega_*$.

Theorem 4.1 generalizes [6, Proposition 3] in two respects. The first generalization concerns the consideration of $m \geq 2$ scales (in [6] only the case $m = 2$ was treated). The second consists of taking \mathbf{H} depending also on the rectangular variables $(p, q) \in B^{2\ell}$. Such generalizations can be easily obtained, and hence will be not discussed here.

PROOF OF THEOREM A. Let

$$\bar{\gamma} := \bar{\tau} \sqrt{\alpha} (\log \alpha^{-1})^{\bar{\tau}+1}, \quad \bar{K} = \frac{1}{\bar{\tau}} \log \frac{1}{\alpha}$$

where $\bar{\tau}$ is as in (4.55) and $\bar{\tau}$ will be fixed later. We aim to apply Theorem 4.1 to the Hamiltonian H_n of Proposition 4.3, with these choices of $\bar{\gamma}$ and \bar{K} . To this end, we take

$$\begin{aligned} M_j &= \begin{cases} \frac{1}{c_1 a_j^- \theta_j^2} & 1 \leq j \leq n \\ \frac{\mu(a_j^+)^2}{c_1 (a_{j+1}^-)^3 \theta_j^2} & n+1 \leq j \leq 2n-1 \end{cases} & L = \bar{M} = \frac{1}{c_2} \frac{\theta_1^2 (a_2^+)^3}{\mu(a_1^-)^2} \\ E &= \frac{1}{c_3} \frac{\mu}{a_1^-} e^{-c\bar{K}} & K = \frac{1}{c_4} \log_+ \left(\frac{1}{\bar{\gamma}^2} \frac{(a_2)^3}{(a_1^-)^3} e^{-c\bar{K}} \right)^{-1} \\ \hat{\rho}_j &= \begin{cases} c_5 \frac{\bar{\gamma} \theta_j}{K^{\tau_*+1}} & 1 \leq j \leq n \\ c_5 \frac{\bar{\gamma} \theta_{j-n}}{K^{\tau_*+1}} & n+1 \leq j \leq 2n-1 \end{cases} & \hat{\rho} := \frac{\theta_1 \bar{\gamma}}{\hat{K}^{\tau_*+1}} \quad \tau_* > 3n-2 \\ \hat{E} &= \frac{1}{c_6} \frac{1}{\bar{\gamma}^2} \frac{(a_2)^3}{(a_1^-)^3} e^{-c\bar{K}} \hat{K}^{2(\tau_*+1)} \end{aligned}$$

where $\hat{K} := \max\{K, \bar{K}\}$. The number $\frac{1}{\bar{\gamma}^2} \frac{(a_2)^3}{(a_1^-)^3}$ can be bounded by $\frac{1}{\alpha^N}$ for a sufficiently large N depending only on n . Hence, if $\bar{\tau} < \frac{c}{N}$ and $\alpha < c_6$, we have $\hat{E} < 1$ and the theorem is proved. \square

Proofs

In this chapter we provide the proof of Proposition 4.3. This is divided in two steps: normalization of fast angles and of secular coordinates.

5.1. Normalization of fast angles

Let $\overline{f_{\mathcal{P}}^{ij}}, \overline{f_{\mathcal{P}}^{ij}{}^{(k)}}$ as in Lemma 3.1, and let

$$(5.57) \quad \overline{f_{\mathcal{P}}^{ij}{}^{(\geq 2)}} := \overline{f_{\mathcal{P}}^{ij}} - \overline{f_{\mathcal{P}}^{ij}{}^{(0)}}.$$

PROPOSITION 5.1. *There exist two small numbers $\widehat{\mathbf{c}}, \mathbf{c}_1$, where $\widehat{\mathbf{c}}$ depends only on n , while \mathbf{c}_1 depends only on n, m_1, \dots, m_n , such that, if the inequality in (4.53) and*

$$(5.58) \quad \frac{1}{\mathbf{c}} \mu \bar{K} \left(\frac{a_n^+}{a_1^-} \right)^{\frac{3}{2}} < 1$$

hold, one can find a real-analytic and symplectic transformation

$$\phi_{\text{fast}} : (\overline{\Theta}, \overline{\vartheta}, \chi, \Lambda, \kappa, \ell) \in \mathfrak{D}_{\text{fast}} := \mathbb{T}_{\widehat{\mathbf{c}}\Theta^+, \widehat{\mathbf{c}}\vartheta^+} \times \mathbb{X}_{\widehat{\mathbf{c}}\theta} \times \mathcal{A}_{\widehat{\mathbf{c}}\theta} \times \mathbb{T}_{\widehat{\mathbf{c}}s}^n \times \mathbb{T}_{\widehat{\mathbf{c}}s}^n \rightarrow \mathfrak{D}_{\mathcal{P}}$$

which conjugates $\mathbb{H}_{\mathcal{P}}$ to

$$(5.59) \quad \begin{aligned} & \mathbb{H}_{\text{fast,exp}}(\overline{\Theta}, \chi, \Lambda, \overline{\vartheta}, \kappa, \ell) \\ & := \mathbb{H}_{\mathcal{P}} \circ \phi_{\text{fast}} = \mathfrak{h}_{\text{fast}}(\Lambda) + \mu f_{\text{fast}}(\overline{\Theta}, \chi, \Lambda, \overline{\vartheta}, \kappa) + \mu f_{\text{fast,exp}}(\overline{\Theta}, \chi, \Lambda, \overline{\vartheta}, \kappa, \ell) \end{aligned}$$

where $\mathfrak{h}_{\text{fast}}$ is as in Proposition 4.3, and

$$(5.60) \quad f_{\text{fast}} := \sum_{i=1}^{n-1} f_{\text{fast}}^i, \quad f_{\text{fast,exp}} := \sum_{i=1}^{n-1} f_{\text{fast,exp}}^i.$$

Here,

1. The “fast frequency-map”

$$\omega_{\text{fast}} := \partial \mathfrak{h}_{\text{fast}}$$

is a diffeomorphism of \mathcal{A} with non-vanishing Jacobian matrix on $\mathcal{A}_{\widehat{\mathbf{c}}\theta}$ and, moreover,

$$\omega_{\text{fast}} \in \mathcal{D}_{\gamma_{\text{fast}}, \tau}^{\bar{K}, \nu_{\text{fast}}} \quad \forall \Lambda \in \mathcal{A},$$

with

$$\gamma_{\text{fast}} := (\gamma_1, \dots, \gamma_n) \quad \nu_{\text{fast}} := (\nu_1, \dots, \nu_n)$$

and ν_i, γ_i as in (4.56);

2. the functions f_{fast}^i , $f_{\text{fast,exp}}^i$ do not depend on κ_0 ; the f_{fast}^i 's are given by

$$(5.61) \quad f_{\text{fast}}^i = f_{\text{fast}}^i(t_i, y_i, x_i) = \overline{f_{\mathcal{P}}^i}^{(\geq 2)}(t_i, y_i, x_i) + \widetilde{f_{\text{fast}}^i}(t_i, y_i, x_i), \quad i = 1, \dots, n-1,$$

with

$$\begin{aligned} \overline{f_{\mathcal{P}}^i}^{(\geq 2)} &:= \sum_{j=i+1}^n \overline{f_{\mathcal{P}}^{ij}}^{(\geq 2)} \\ t_i &:= (\Theta_i, \dots, \Theta_{n-1}, \vartheta_i, \dots, \vartheta_{n-1}), \\ y_i &:= (\chi_{i-1}, \dots, \chi_{n-1}, \Lambda_i, \dots, \Lambda_n) \\ x_i &:= (\kappa_i, \dots, \kappa_{n-1}). \end{aligned}$$

In particular, $\widetilde{f_{\text{fast}}^i}$ do not depend on ℓ_1, \dots, ℓ_n ;

3. finally, $\widetilde{f_{\text{fast}}^i}$, $f_{\text{exp,fast}}^i$ satisfy the following bounds

$$(5.62) \quad \|\widetilde{f_{\text{fast}}^i}\|_{\mathfrak{D}_{\text{fast}}} \leq \frac{1}{\mathfrak{c}_1} \mu \bar{K} \left(\frac{a_n^+}{a_1^-} \right)^{\frac{3}{2}} \frac{1}{a_{i+1}^-}, \quad \|f_{\text{fast,exp}}^i\|_{\mathfrak{D}_{\text{fast}}} \leq \frac{1}{\mathfrak{c}_1} \frac{e^{-\widehat{\mathfrak{c}} \bar{K} s}}{a_{i+1}^-}.$$

Let L_0, \dots, L_n be defined as \mathfrak{L}_i in (4.51), with $\nu = m = n$ and $\nu_1 = \dots = \nu_n = 1$.

LEMMA 5.1. *If \bar{K} verifies the inequality in (4.53), then one can find a number \mathfrak{c}_3 , depending only on m_0, \dots, m_n , such that*

$$|\omega_{k,\text{fast}}(\Lambda) \cdot k| \geq \frac{\mathfrak{c}_3}{(a_j^+)^{3/2}} \quad \forall k \in L_{j-1} \setminus L_j, \quad |k| \leq \bar{K}, \quad \forall \Lambda \in A_\theta, \quad \forall j = 1, \dots, n.$$

PROOF. For $\Lambda \in A_\theta$, $\omega_{k,\text{fast},j} := \frac{\mathfrak{M}_j^2 m_j^3}{\Lambda_j^3}$ verifies $\frac{\sqrt{\mathfrak{M}_j}}{(a_j^+)^{3/2}} \leq |\omega_{k,\text{fast},j}| \leq \frac{\sqrt{\mathfrak{M}_j}}{(a_j^-)^{3/2}}$.

In the case $j = n$, we find $|\omega_{k,\text{fast}} \cdot k| = |\omega_{k,\text{fast},n} k_n| \geq \frac{\sqrt{\mathfrak{M}_j}}{(a_n^+)^{3/2}}$, since $k_n \neq 0$. Let then $j \neq n$. For $k \in L_{j-1} \setminus L_j$, $k_j \neq 0$, so, inequality (4.53), with $\mathfrak{c}_2 \leq \frac{\min_j \sqrt{\mathfrak{M}_j}}{\max_j \sqrt{\mathfrak{M}_j}}$, and (4.50) imply

$$\bar{K} \leq \frac{\min_j \sqrt{\mathfrak{M}_j}}{\max_j \sqrt{\mathfrak{M}_j}} \min_{1 \leq j \leq n-1} \left(\frac{a_{j+1}^-}{a_j^+} \right)^{3/2}$$

and hence

$$\begin{aligned} |\omega_{k,\text{fast}} \cdot k| &= \left| \sum_{i=j}^n \omega_{k,\text{fast},i} k_i \right| \geq \inf_{A_\theta} |\omega_{k,\text{fast},j}| - \bar{K} \max_{j < i \leq n} \sup_{A_\theta} |\omega_{k,\text{fast},i}| \\ &\geq \frac{\sqrt{\mathfrak{M}_j}}{(a_j^+)^{3/2}} - \bar{K} \frac{\max_{i>j} \sqrt{\mathfrak{M}_i}}{(a_{j+1}^-)^{3/2}} \geq \frac{\sqrt{\mathfrak{M}_j}}{2(a_j^+)^{3/2}} e q \end{aligned}$$

□

PROOF OF PROPOSITION 5.1. The proof proceeds by recursion, in n steps. We describe the h^{th} step of this recursion, with $h = 1, \dots, n$. We start with a Hamiltonian of the form

$$(5.63) \quad H_{h-1} = h_{\text{fast}}^0 + \mu f_{h-1}$$

where h_{fast}^0 is as in (3.38), and a domain

$$\mathfrak{D}_{h-1} = \mathbb{T}_{\Theta^{+(h-1)}, \vartheta^{+(h-1)}} \times \mathbb{X}_{\theta^{(h-1)}} \times \mathcal{A}_{\theta^{(h-1)}} \times \mathbb{T}_{s^{(h-1)}}^n \times \mathbb{T}_{s^{(h-1)}}^n.$$

When $h = 1$, we take $\mathbb{H}_0 := \mathbb{H}_{\mathcal{P}}$, $\Theta_+^{(0)} := \Theta^+$, $\vartheta_+^{(0)} := \vartheta^+$, $\theta^{(0)} := \theta$, $s^{(0)} := s$, $f_0 := f_{\mathcal{P}}$ and we decompose

$$f_0 := \widehat{f}_0 := \sum_{i=1}^{n-1} \widehat{f}_0^i \quad \text{with} \quad f_0^i := \sum_{j=i+1}^n f_{\mathcal{P}}^{ij}.$$

We observe that \widehat{f}_0^i depends on the coordinates

$$\begin{aligned} & \Theta_i, \quad \dots, \quad \Theta_{n-1}, \quad \chi_{i-1}, \quad \dots, \chi_{n-1}, \quad \Lambda_i, \quad \dots, \quad \Lambda_n \\ & \vartheta_i, \quad \dots, \quad \vartheta_{n-1}, \quad \kappa_i, \quad \dots, \quad \kappa_{n-1}, \quad \ell_i, \quad \dots, \quad \ell_n. \end{aligned}$$

For $n \geq 3$ and $2 \leq h \leq n-1$, we assume, inductively, that f_{h-1} is a sum

$$(5.64) \quad f_{h-1} = \widehat{f_{h-1}} + f_{\text{exp}, h-1} = \sum_{1 \leq i \leq n} \widehat{f_{h-1}^i} + \sum_{1 \leq i \leq n} f_{\text{exp}, h-1}^i,$$

where, in turn,

$$\widehat{f_{h-1}^i} = \overline{f_{h-1}^i} + \widetilde{f_{h-1}^i}$$

with $\overline{f_{h-1}^i}$, $\widetilde{f_{h-1}^i}$ depending only on the coordinates

$$\begin{aligned} & \Theta_i, \quad \dots, \quad \Theta_{n-1}, \quad \chi_{i-1}, \quad \dots, \chi_{n-1}, \quad \Lambda_i, \quad \dots, \quad \Lambda_n \\ & \vartheta_i, \quad \dots, \quad \vartheta_{n-1}, \quad \kappa_i, \quad \dots, \quad \kappa_{n-1}, \quad \ell_{i \vee h}, \quad \dots, \quad \ell_n \end{aligned}$$

and $\overline{f_{h-1}^i}$, $\widetilde{f_{h-1}^i}$, $f_{\text{exp}, h-1}^i$ verifying the following bounds and identities

$$(5.65) \quad \begin{aligned} \overline{f_{h-1}^i} &= \Pi_{L_{h-1}} T_{\bar{K}} \widehat{f_{h-2}^i} \\ \|\widetilde{f_{h-1}^i}\|_{\mathfrak{D}_{h-1}} &\leq \mathfrak{C}_{1, h-1} \mu \bar{K} \left(\frac{a_n^+}{a_1}\right)^{\frac{3}{2}} \|\widehat{f_{h-2}^i}\|_{\mathfrak{D}_{h-2}} \\ \|f_{\text{exp}, h-1}^i\|_{\mathfrak{D}_{h-1}} &\leq \mathfrak{C}_{2, h-1} e^{-Ks^{(h)}} \|\widehat{f_{h-2}^i}\|_{\mathfrak{D}_{h-2}}. \end{aligned}$$

Here Π_{L_h} denotes the projection over the module L_h . In any case, $h = 1$, or $2 \leq h \leq n-1$, we focus on the Hamiltonian

$$(5.66) \quad \widehat{\mathbb{H}_{h-1}} = h_{\text{fast}}^0 + \mu \widehat{f_{h-1}} = h_{\text{fast}}^0 + \mu \sum_{i=1}^{n-1} \widehat{f_{h-1}^i}.$$

Our purpose is to apply Proposition D.1 to this Hamiltonian, in the case that the abstract system (D.133) does not depend on the coordinates (p, q) . To this end, we take the coordinates

$$I := \Lambda, \quad \varphi := \ell, \quad \eta := (\bar{\Theta}, \chi), \quad \xi := (\bar{\vartheta}, \kappa),$$

the functions f_i in (D.135) to be the $\widehat{f_{h-1}^{n-i}}$, and

$$\begin{aligned} N &= n-1, & \nu &= n, & m_i &:= 2i \\ (I_1, \dots, I_\nu) &:= (\Lambda_n, \dots, \Lambda_1) \\ (\varphi_1, \dots, \varphi_{\nu_i}) &:= (\ell_n, \dots, \ell_{\max\{n-i, h\}}) \\ (\eta_1, \dots, \eta_{m_i}) &:= (\Theta_{n-1}, \dots, \Theta_{n-i}, \chi_{n-1}, \dots, \chi_{n-i-1}) \\ (\xi_1, \dots, \xi_{m_i}) &:= (\vartheta_{n-1}, \dots, \vartheta_{n-i}, \kappa_{n-1}, \dots, \kappa_{n-i}) \\ u_i &:= (\Lambda_n, \dots, \Lambda_1, \Theta_{n-1}, \dots, \Theta_{n-i}, \chi_{n-1}, \dots, \chi_{n-i-1}, \vartheta_{n-1}, \dots, \vartheta_{n-i}, \\ &\quad \kappa_{n-1}, \dots, \kappa_{n-i}). \end{aligned}$$

The non-resonance assumption (D.134) for $\omega = \omega_{k, \text{fast}} = \partial_\Lambda h_{k, \text{fast}}$, with

$$\mathfrak{J}_i = L_{h-1}, \quad \mathfrak{J} = \cup_i \mathfrak{J}_i = L_{h-1}, \quad \mathfrak{L} = L_h \quad K = \bar{K}$$

is ensured by Lemma 5.1, with

$$\mathfrak{a} = \frac{\mathfrak{c}_3}{(a_h^+)^{3/2}}, \quad A = \mathcal{A}, \quad r = \theta_1^{(h-1)}.$$

Now we have to check condition (D.139). In the case $2 \leq h \leq n-1$ the inductive assumptions (5.65) and assumption (5.58) imply

$$\begin{aligned} \|\widehat{f_{h-1}^i}\|_{\mathfrak{D}_{h-1}} &\leq \|\overline{f_{h-1}^i}\|_{\mathfrak{D}_{h-1}} + \|\widetilde{f_{h-1}^i}\|_{\mathfrak{D}_{h-1}} \leq \left(1 + \mathfrak{C}_1 \mu \bar{K} \left(\frac{a_n^+}{a_1}\right)^{\frac{3}{2}}\right) \|\widehat{f_{h-2}^i}\|_{\mathfrak{D}_{h-2}} \\ (5.67) \quad &\leq \dots \leq (1 + \mathfrak{C}_{1, h-1} \mathfrak{c}_1)^{h-1} \|\widehat{f_0^i}\|_{\mathfrak{D}_0} \leq \frac{\mathfrak{C}_{4, h-1}}{a_i} =: E_i. \end{aligned}$$

An analogous bound holds also for $h=1$. The numbers c_i and d_i in (D.138) may be evaluated as

$$c_i = e(1 + 2ie)/2 \quad d_i = \min\{\theta_1^{(h-1)} s^{(h-1)}, \Theta_i^{+(h-1)} \vartheta_i^{+(h-1)}\} = \mathfrak{c}_2 \theta_1^{(h-1)}.$$

From these bounds it is immediate to see that inequality (D.139) is implied by (5.58), provided $\mathfrak{c}_1 < 2^{-7} \frac{8}{7} (\frac{8}{9})^{n-2} \mathfrak{c}_2 / (\mathfrak{C}_4 c_n)$. Then Proposition D.1 applies. Its thesis implies that $\widehat{H_{h-1}}$ in (5.66) can be conjugated to a suitable $H_h^* = h_{k, \text{fast}} + \mu f_h^*$, where f_h^* verifies equalities and inequalities in (5.64)-(5.65) with h replaced by $h+1$ and $\mathfrak{C}_{1, h-1}, \mathfrak{C}_{2, h-1}$ replaced by suitable $\mathfrak{C}_{1, h}^*, \mathfrak{C}_{2, h}^*$. Then, applying the same transformation to H_{h-1} in (5.63), we shall conjugate H_{h-1} to $H_h = h_{k, \text{fast}} + \mu f_h$, where f_h satisfies the same equalities and inequalities as f_h^* , with suitable $\mathfrak{C}_{1, h} \geq \mathfrak{C}_{1, h}^*, \mathfrak{C}_{2, h} \geq \mathfrak{C}_{2, h}^*$.

After we have performed n steps, we let $\mathfrak{D}_{\text{fast}} := \mathfrak{D}_n$, $H_{\text{fast}, \text{exp}} := H_n$, $f_{\text{fast}}^i := \widehat{f_n^i}$, $\widetilde{f_{\text{fast}}^i} := f_{\text{fast}}^i - \overline{f_{\mathcal{P}}^i}$, $f_{\text{fast}, \text{exp}}^i := f_{\text{exp}, n}^i$, $\widehat{f_{\text{fast}}^i} := \sum_{i=1}^{n-1} \widehat{f^i}$, $\widetilde{f_{\text{fast}}^i} := \sum_{i=1}^{n-1} \widetilde{f^i}$, $f_{\text{fast}, \text{exp}} := \sum_{i=1}^{n-1} f_{\text{fast}, \text{exp}}^i$, with $\overline{f_{\mathcal{P}}^i} := \sum_{j=i+1}^n \overline{f_{\mathcal{P}}^{ij}}$. Therefore,

$$H_{\text{fast}} = h_{\text{fast}}^{(0)} + \mu(\widehat{f_{\text{fast}}} + f_{\text{exp}, \text{fast}}) = h_{\text{fast}}^{(0)} + \mu\left(\sum_{1 \leq i < j \leq n} \overline{f_{\mathcal{P}}^{ij}} + \widetilde{f_{\text{fast}}} + f_{\text{exp}, \text{fast}}\right)$$

reduces to (5.59) and the formulae given below, using (5.57).

It remains to check the bound on the left in (5.62) (the one on the right follows by construction). This follows by telescopic arguments. Indeed,

$$\begin{aligned}
\|\widehat{f_{\text{fast}}^i}\|_{\mathfrak{D}_n} &= \|f_{\text{fast}}^i - \overline{f_{\mathcal{P}}^i}\|_{\mathfrak{D}_n} = \|\widehat{f_n^i} - \overline{f_{\mathcal{P}}^i}\|_{\mathfrak{D}_n} = \|\Pi_{L_n} \widehat{f_n^i} - \Pi_{L_n} f_{\mathcal{P}}^i\|_{\mathfrak{D}_n} \\
&\leq \sum_{h=1}^n \|\Pi_{L_n} \widehat{f_h^i} - \Pi_{L_n} T_{\bar{K}} \widehat{f_{h-1}^i}\|_{\mathfrak{D}_n} \\
&= \sum_{h=1}^n \|\Pi_{L_n} \widehat{f_h^i} - \Pi_{L_n} \Pi_{L_h} T_{\bar{K}} \widehat{f_{h-1}^i}\|_{\mathfrak{D}_n} \\
&\leq \sum_{h=1}^n \|\widehat{f_h^i} - \Pi_{L_h} T_{\bar{K}} \widehat{f_{h-1}^i}\|_{\mathfrak{D}_n} \\
&\leq \sum_{h=1}^n \|\widehat{f_h^i} - \Pi_{L_h} T_{\bar{K}} \widehat{f_{h-1}^i}\|_{\mathfrak{D}_h} \\
&\leq \mu \bar{K} \left(\frac{a_n^+}{a_1^-} \right)^{\frac{3}{2}} \frac{\sum_{h=1}^n \mathfrak{C}_{1,h} \mathfrak{C}_{4,h-1}}{a_{i+1}^-}.
\end{aligned}$$

Here, we have used the second bound in (5.65), (5.67), that $\widehat{f_n^i}$ does not depend on ℓ_1, \dots, ℓ_n , and, finally, $\Pi_{L_n} = \Pi_{L_n} T_{\bar{K}} = \Pi_{L_n} \Pi_{L_h}$, for all $1 \leq h \leq n$. \square

5.2. Secular normalizations

Consider the following truncation

$$H_{\text{fast}}(\bar{\Theta}, \chi, \Lambda, \bar{\vartheta}, \kappa) := h_{\text{fast}}(\Lambda) + \mu f_{\text{fast}}(\bar{\Theta}, \chi, \Lambda, \bar{\vartheta}, \kappa)$$

of the Hamiltonian $H_{\text{fast}, \text{cfp}}$ in (5.59). The purpose of this section is to describe an iterative scheme which, after $(n-1)$ steps, conjugates H_{fast} to a close-to-be integrable system, with an arbitrarily small remainder.

Let us firstly establish the following notation.

- Given a Taylor-Fourier expansion of the form

$$g(p, q, \kappa) = \sum_{\substack{(a,b) \in \mathbb{N}^{2m_1} \\ k \in \mathbb{Z}^{m_2}}} g_{a,b,k} \left(\frac{p - iq}{\sqrt{2}} \right)^a \left(\frac{p + iq}{\sqrt{2i}} \right)^b e^{ik \cdot \kappa} \quad (p, q, \kappa) \in B^{2m_1}(0) \times \mathbb{T}^{m_2}.$$

we denote as

$$\Pi_{p,q,\kappa} g := \sum_{a \in \mathbb{N}^{m_1}} g_{0,a,a} \left(\frac{p^2 + q^2}{2i} \right)^a.$$

PROPOSITION 5.2. *There exists number $\bar{\mathfrak{c}}_h$, depending only on $n, m_0, \dots, m_n, a_n^\pm$ such that, for any $h = 1, \dots, n-1$ and any $\bar{K}, \bar{\gamma} > 0$ such that (4.54) hold with \mathfrak{c} replaced by $\bar{\mathfrak{c}}_h$, one finds open sets*

$$B_j^* \subset B_{\varepsilon_j}^2, \quad G_j^* \subset G_j := \left[G_j^+, G_j^+ \right], \quad j = n-h, \dots, n-1$$

verifying

$$(5.68) \quad \text{meas}(B_j^* \times G_j^*) \geq \left(1 - \frac{\bar{\gamma}}{\bar{\mathfrak{c}}_h}\right) \text{meas}(B_{\varepsilon_j}^2 \times G_j)$$

such that, defining

$$(5.69) \quad \mathbb{T}_{\bar{c}_h \theta}^h := \begin{cases} \left\{ \begin{array}{l} (\Theta_1, \dots, \Theta_{n-h-1}, \vartheta_1, \dots, \vartheta_{n-h-1}) \\ \in \mathbb{C}^{n-1} \times \mathbb{T}_{\mathbb{C}}^{n-1} : \\ |\vartheta_j - \pi| \leq \bar{c}_h \frac{\theta_j}{G_n^+}, \quad |\Theta_j| \leq \bar{c}_h G_n^+ \\ \forall j = 1, \dots, n-h-1 \end{array} \right\} & n \geq 3, 1 \leq h < n-2 \\ \emptyset & \text{otherwise} \end{cases}$$

$$B_{\bar{c}_h \bar{r}}^{*h} := (B_{n-h}^*)_{\bar{c}_h \sqrt{\bar{r}_{n-h}}} \times \dots \times (B_{n-1}^*)_{\bar{c}_h \sqrt{\bar{r}_{n-1}}}$$

$$X_{\bar{c}_h \theta, \bar{c}_h \bar{r}}^{*h} := \left\{ \chi = (\chi_0, \dots, \chi_{n-1}) : \begin{array}{l} \chi_{i-1} - \chi_i \in (G_i^*)_{\bar{c}_h \theta_i}, \quad \chi_{j-1} - \chi_j \in (G_j)_{\bar{c}_h \bar{r}_j} \\ \forall i = 1, \dots, n-h-1, \quad j = n-h, \dots, n, \quad \chi_n := 0 \end{array} \right\}$$

$$\mathfrak{D}_{\bar{c}_h \theta}^h := \mathbb{T}_{\bar{c}_h \theta}^h \times B_{\bar{c}_h \bar{r}}^{*h} \times X_{\bar{c}_h \theta, \bar{c}_h \bar{r}}^{*h} \times A_{\bar{c}_h \bar{r}} \times \mathbb{T}_{\bar{c}_h s}^n \times \mathbb{T}_{\bar{c}_h s}^n$$

a real-analytic transformation

$$\Phi_{\text{sec}, h} : \mathfrak{D}_{\text{sec}}^h \rightarrow \mathfrak{D}_{\text{fast}},$$

may be found, which conjugates f_{fast} to a new function

$$f_{\text{sec}, h} := f_{\text{fast}} \circ \Phi_{\text{sec}, h}$$

enjoying the following properties.

1. Denoting by $(t^{(h)}, z^{(h)}, y^{(h)}, x^{(h)})$, where

$$(5.70) \quad \begin{aligned} t^{(h)} &= (\Theta^{(h)}, \vartheta^{(h)}) = (\Theta_1^{(h)}, \dots, \Theta_{n-h-1}^{(h)}, \vartheta_1^{(h)}, \dots, \vartheta_{n-h-1}^{(h)}) \\ z^{(h)} &= (p^{(h)}, q^{(h)}) = (p_{n-h}^{(h)}, \dots, p_{n-1}^{(h)}, q_{n-h}^{(h)}, \dots, q_{n-1}^{(h)}) \\ y^{(h)} &= (\chi^{(h)}, \Lambda^{(h)}) = (\chi_0^{(h)}, \dots, \chi_{n-1}^{(h)}, \Lambda_1^{(h)}, \dots, \Lambda_n^{(h)}) \\ x^{(h)} &= (\kappa^{(h)}, \ell^{(h)}) = (\kappa_0^{(h)}, \dots, \kappa_{n-1}^{(h)}, \ell_1^{(h)}, \dots, \ell_n^{(h)}), \end{aligned}$$

coordinates on $\mathfrak{D}_{\text{sec}}^h$ then $\Phi_{\text{sec}, h}$ is co-variant with the symmetry:

$$\Phi_{\text{sec}, h}(-t^{(h)}, -z^{(h)}, y^{(h)}, x^{(h)}) = (-t^{(0)}, y^{(0)}, x^{(0)}) \quad \text{if}$$

$$\Phi_{\text{sec}, h}(t^{(h)}, z^{(h)}, y^{(h)}, x^{(h)}) = (t^{(0)}, y^{(0)}, x^{(0)})$$

and hence, $f_{\text{sec}, h}$ is even around

$$t^{(h)} = (0, k\pi), \quad z^{(h)} = 0 \quad k \in \{0, 1\}^{n-h-1}$$

2. Defining

$$\begin{aligned}
 \mathfrak{t}_i^{(h)} &:= \begin{cases} (\Theta_i^{(h)}, \dots, \Theta_{n-h-1}^{(h)}, \vartheta_i^{(h)}, \dots, \vartheta_{n-h-1}^{(h)}) & i \leq n-h-1 \\ \emptyset & \text{otherwise} \end{cases} \\
 \hat{y}_i^{(h)} &= \begin{cases} \left(\frac{(p_i^{(h)})^2 + (q_i^{(h)})^2}{2}, \dots, \frac{(p_{n-1}^{(h)})^2 + (q_{n-1}^{(h)})^2}{2}, \chi_{i-1}^{(h)}, \dots, \chi_{n-1}^{(h)}, \right. \\ \quad \left. \Lambda_i^{(h)}, \dots, \Lambda_n^{(h)} \right) & i \geq n-h \\ \left(\frac{(p_{n-h}^{(h)})^2 + (q_{n-h}^{(h)})^2}{2}, \dots, \frac{(p_{n-1}^{(h)})^2 + (q_{n-1}^{(h)})^2}{2}, \chi_{i-1}^{(h)}, \dots, \chi_{n-1}^{(h)}, \right. \\ \quad \left. \Lambda_i^{(h)}, \dots, \Lambda_n^{(h)} \right) & \text{otherwise} \end{cases} \\
 \hat{x}_i^{(h)} &= \begin{cases} (\kappa_i^{(h)}, \dots, \kappa_{n-h-2}^{(h)}) & n \geq 4 \ \& \ 1 \leq h \leq n-3 \ \& \ 1 \leq i \leq n-h-2 \\ \emptyset & \text{otherwise} \end{cases}
 \end{aligned}
 \tag{5.71}$$

and $\hat{y} := \hat{y}_1$, $\hat{x} := \hat{x}_1$, $f_{\text{sec},h}$ has the form

$$\begin{aligned}
 f_{\text{sec},h}(\mathfrak{t}^{(h)}, \mathfrak{z}^{(h)}, \mathfrak{y}^{(h)}, \mathfrak{x}^{(h)}) &= h_{\text{sec},h}(\hat{y}_{n-h}^{(h)}) + f_{\text{norm},h}(\mathfrak{t}^{(h)}, \hat{y}^{(h)}, \hat{x}^{(h)}) \\
 &+ f_{\text{exp,sec},h}(\mathfrak{t}^{(h)}, \mathfrak{z}^{(h)}, \mathfrak{y}^{(h)}, \mathfrak{x}^{(h)})
 \end{aligned}
 \tag{5.72}$$

with

$$\begin{aligned}
 h_{\text{sec}}(\hat{y}_{n-h}^{(h)}) &= \sum_{i=n-h}^{n-1} h_{\text{sec}}^i(\hat{y}_i^{(h)}) \\
 f_{\text{norm},h}(\mathfrak{t}^{(h)}, \hat{y}^{(h)}, \hat{x}^{(h)}) &= \sum_{i=1}^{n-h-1} f_{\text{norm},h}^i(\mathfrak{t}_i^{(h)}, \hat{y}_i^{(h)}, \hat{x}_i^{(h)})
 \end{aligned}
 \tag{5.73}$$

where

3. the functions h_{sec}^i $f_{\text{norm},h}^i$ may be decomposed as

$$\begin{aligned}
 h_{\text{sec}}^i(\hat{y}_i^{(h)}) &= \overline{h_{\text{sec}}^i}(\hat{y}_i^{(h)}) + \widetilde{h_{\text{sec}}^i}(\hat{y}_i^{(h)}) \\
 f_{\text{norm},h}^i(\mathfrak{t}_i^{(h)}, \hat{y}_i^{(h)}, \hat{x}_i^{(h)}) &= \overline{f_{\text{norm},h}^i}(\mathfrak{t}_i^{(h)}, \hat{y}_i^{(h)}, \hat{x}_i^{(h)}) + \widetilde{f_{\text{norm},h}^i}(\mathfrak{t}_i^{(h)}, \hat{y}_i^{(h)}, \hat{x}_i^{(h)})
 \end{aligned}
 \tag{5.74}$$

where

$$\overline{f_{\text{norm},h}^i} = \sum_{j=i+1}^n \Pi_h(\overline{f_{\mathcal{P}}^{ij}})^{(\geq 2)} \circ \overline{\phi_{\text{int}}^{n-1}} \circ \dots \circ \overline{\phi_{\text{int}}^{n-h}}
 \tag{5.75}$$

and $\overline{h_{\text{sec}}^i}, \overline{\phi_{\text{int}}^i}$ as in Lemma 3.4. The functions $\widetilde{h_{\text{sec},h}}, \widetilde{f_{\text{norm},h}}, f_{\text{exp,sec},h}$ in (5.72) may be bounded as

$$(5.76) \quad \begin{aligned} |\widetilde{h_{\text{sec},h}^i}| &\leq \frac{1}{c_h} \max \left\{ \mu \bar{K} \left(\frac{a_n}{a_1} \right)^{3/2} \frac{1}{a_{i+1}^-}, \frac{\bar{K}^{\bar{\tau}+1} \sqrt{\alpha} (a_i^+)^2}{\bar{\gamma} (a_{i+1}^+)^3}, \frac{\varepsilon_{i+1}^2 (a_i^+)^2}{\theta_{i+1} (a_{i+1}^-)^3} \right\} \\ |\widetilde{f_{\text{norm},h}^i}| &\leq \frac{1}{c_h} \max \left\{ \mu \bar{K} \left(\frac{a_n}{a_1} \right)^{3/2} \frac{1}{a_{i+1}^-}, \frac{\bar{K}^{\bar{\tau}+1} \sqrt{\alpha} (a_i^+)^2}{\bar{\gamma} (a_{i+1}^-)^3} \right\} \\ |f_{\text{exp,sec},h}| &\leq \frac{1}{c_h} \frac{(a_{n-1}^+)^2}{(a_n^-)^3} e^{-c_h \bar{K}} \end{aligned}$$

4. Defining

$$\zeta^{(h)} := \left(\frac{(p_{n-h}^{(h)})^2 + (q_{n-h}^{(h)})^2}{2}, \dots, \frac{(p_{n-1}^{(h)})^2 + (q_{n-1}^{(h)})^2}{2}, \chi_{i-1}^{(h)}, \dots, \chi_{n-1} \right)$$

so that

$$\hat{y}_{n-h}^{(h)} = (\zeta^{(h)}, \Lambda_{n-h}^{(h)}, \dots, \Lambda_n^{(h)})$$

for any $\Lambda_{n-h}^{(h)}, \dots, \Lambda_n^{(h)}$, the map

$$\zeta^{(h)} \rightarrow \omega_{\text{sec},h} := \partial_{\zeta^{(h)}} h_{\text{sec},h}(\zeta^{(h)}, \Lambda^{(h)})$$

is a diffeomorphism of $D_{\bar{\tau}} \times X_{\bar{\tau}}$, with non-vanishing Jacobian matrix. The set $D_{\bar{\tau}}^* \times X_{\bar{\tau}}^*$ consists of the subset of $D_{\bar{\tau}} \times X_{\bar{\tau}}$ such that $\omega_{\text{fast,sec}} \in \mathcal{D}_{\gamma_{\text{sec}}; \bar{\tau}}^{\bar{K}, \nu_{\text{sec}}}$, where, if ν_j, γ_j are as in (4.56),

$$\nu_{\text{sec}} := (\nu_{n+1}, \dots, \nu_{2n-1}) \quad \gamma_{\text{sec}} := (\gamma_{n+1}, \dots, \gamma_{2n-1}).$$

We shall give the complete details of the proof of Proposition 5.2 along the following sections 5.2.1-5.2.4. In this section we just provide main ideas.

SCHEME OF PROOF. The proof is by recursion. The h^{th} step of this recursion starts with

$$f_{\text{sec},h-1} = h_{\text{sec},h-1} + f_{\text{norm},h-1} + f_{\text{exp,sec},h-1},$$

where, for $h = 1$

$$(5.77) \quad h_{\text{sec},0} \equiv 0, \quad f_{\text{exp,sec},0} \equiv 0, \quad f_{\text{sec},0} := f_{\text{norm},0} := f_{\text{fast}},$$

while, for $n \geq 3$ and $h = 2, \dots, n-1$, we assume, inductively, that $h_{\text{sec},h-1}, f_{\text{sec},h-1}$ and $f_{\text{exp,sec},h-1}$ satisfy the theses of Proposition 5.2, with h replaced by $(h-1)$.

The transformation ϕ_{sec}^{n-h} conjugating $f_{\text{sec},h-1}$ to $f_{\text{sec},h}$ will be constructed as a product $\phi_{\text{sec}}^{n-h} = \phi_{\text{int}}^{n-h} \circ \phi_{\text{norm}}^{n-h}$ of an ‘‘integrating’’ and a ‘‘normalizing’’ transformation.

Due to the bound on $f_{\text{exp,sec},h-1}$, it is enough to focus on the truncation

$$(5.78) \quad \widehat{f_{\text{sec},h-1}} := h_{\text{sec},h-1} + f_{\text{norm},h-1} = h_{\text{sec},h-1} + \sum_{i=1}^{n-h} f_{\text{norm},h-1}^i(t_i^{(h-1)}, \hat{y}_i^{(h-1)}, \hat{x}_i^{(h-1)})$$

of $f_{\text{sec},h-1}$. We split

$$f_{\text{norm},h-1} = f_{\text{norm},h-1}^{n-h}(t_{n-h}^{(h-1)}, \hat{y}_{n-h}^{(h-1)}, \hat{x}_{n-h}^{(h-1)}) + \sum_{i=1}^{n-h-1} f_{\text{norm},h-1}^i(t_i^{(h-1)}, \hat{y}_i^{(h-1)}, \hat{x}_i^{(h-1)})$$

and we distinguish two cases.

Case $n \geq 3$, $h = 2, \dots, n-1$. By the inductive assumption (see (5.71) with h replaced by $(h-1)$), the function $f_{\text{norm},h-1}^{n-h}$ depends only on

$$\mathfrak{t}_{n-h}^{(h-1)} = (\Theta_{n-h}^{(h-1)}, \vartheta_{n-h}^{(h-1)}) \quad \text{and} \quad \hat{y}_{n-h}^{(h-1)}$$

therefore is integrable. In Chapter 5.2.2, we shall construct a canonical, real-analytic change of coordinates

$$\begin{aligned} \phi_{\text{int}}^{n-h} : \quad & \mathfrak{D}_{\text{int}}^h \rightarrow \mathfrak{D}_{\text{sec}}^{h-1} \\ & (\mathfrak{t}_*^{(h)}, \mathfrak{z}_*^{(h)}, \mathfrak{y}_*^{(h)}, \mathfrak{x}_*^{(h)}) \rightarrow (\mathfrak{t}^{(h-1)}, \mathfrak{z}^{(h-1)}, \mathfrak{y}^{(h-1)}, \mathfrak{x}^{(h-1)}) \end{aligned}$$

$$(5.79) \quad \mathfrak{D}_{\text{int}}^h := \mathbb{T}_{\hat{\mathfrak{c}}_h \theta}^h \times B_{\hat{\mathfrak{c}}_h \varepsilon_{n-h}}^2 \times B_{\hat{\mathfrak{c}}_h \varepsilon}^{*,h-1} \times \mathbb{X}_{\hat{\mathfrak{c}}_h \theta, \hat{\mathfrak{c}}_h \bar{r}}^{*,h-1} \times \Lambda_{\hat{\mathfrak{c}}_h \bar{r}} \times \mathbb{T}_{\hat{\mathfrak{c}}_h s}^n \times \mathbb{T}_{\hat{\mathfrak{c}}_h s}^n$$

such that

$$(5.80) \quad f_{\text{norm},h-1}^{n-h} \circ \phi_{\text{int}}^{n-h} = \mathfrak{h}_{\text{sec}}^{n-h}(\hat{y}_{*,n-h}^{(h)})$$

depends only on $\hat{y}_{*,n-h}^{(h)}$, where $\hat{y}_{*,i}^{(h)}$ is defined analogously to $\hat{y}_i^{(h)}$ in (5.71). Here,

$$\begin{cases} \mathfrak{t}_*^{(h)} := (\Theta_*^{(h)}, \vartheta_*^{(h)}) \\ \mathfrak{z}_*^{(h)} := (p_*^{(h)}, q_*^{(h)}) \\ \mathfrak{y}_*^{(h)} := (\chi_*^{(h)}, \Lambda_*^{(h)}) \\ \mathfrak{x}_*^{(h)} := (\kappa_*^{(h)}, \ell_*^{(h)}) \end{cases} \quad \begin{cases} \mathfrak{t}^{(h-1)} := (\Theta^{(h-1)}, \vartheta^{(h-1)}) \\ \mathfrak{z}^{(h-1)} := (p^{(h-1)}, q^{(h-1)}) \\ \mathfrak{y}^{(h-1)} := (\chi^{(h-1)}, \Lambda^{(h-1)}) \\ \mathfrak{x}^{(h-1)} := (\kappa^{(h-1)}, \ell^{(h-1)}) \end{cases}$$

are defined analogously to (5.70).

We shall construct ϕ_{int}^{n-h} in such a way that it involves only the coordinates

$$\phi_{\text{int}}^{n-h} : (\mathfrak{z}_{*,n-h}^{(h)}, \mathfrak{y}_{*,n-h}^{(h)}, \mathfrak{x}_{*,n-h}^{(h)}) \rightarrow (\mathfrak{t}_{n-h}^{(h-1)}, \mathfrak{z}_{n-h+1}^{(h-1)}, \mathfrak{y}_{n-h}^{(h-1)}, \mathfrak{x}_{n-h}^{(h-1)})$$

with

$$\begin{aligned} \mathfrak{z}_{*,n-h}^{(h)} &:= (p_{*,n-h}^{(h)}, \dots, p_{n-1}^{(h)}, q_{*,n-h}^{(h)}, \dots, q_{n-1}^{(h)}) \\ \mathfrak{y}_{*,n-h}^{(h)} &:= (\chi_{*,n-h-1}^{(h)}, \dots, \chi_{*,n-1}^{(h)}, \Lambda_{*,n-h}^{(h)}, \dots, \Lambda_n^{(h)}) \\ \mathfrak{x}_{*,n-h}^{(h)} &:= (\kappa_{*,n-h-1}^{(h)}, \dots, \kappa_{*,n-1}^{(h)}, \ell_{*,n-h}^{(h)}, \dots, \ell_n^{(h)}) \\ \mathfrak{t}_{n-h}^{(h-1)} &:= (\Theta_{n-h}^{(h-1)}, \vartheta_{n-h}^{(h-1)}) \\ \mathfrak{z}_{n-h+1}^{(h-1)} &:= (p_{n-h+1}^{(h-1)}, \dots, p_{n-1}^{(h-1)}, q_{n-h+1}^{(h-1)}, \dots, q_{n-1}^{(h-1)}) \\ \mathfrak{y}_{n-h}^{(h-1)} &:= (\chi_{n-h-1}^{(h-1)}, \dots, \chi_{n-1}^{(h-1)}, \Lambda_{n-h}^{(h-1)}, \dots, \Lambda_n^{(h-1)}) \\ \mathfrak{x}_{n-h}^{(h-1)} &:= (\kappa_{n-h-1}^{(h-1)}, \dots, \kappa_{n-1}^{(h-1)}, \ell_{n-h}^{(h-1)}, \dots, \ell_n^{(h-1)}) \end{aligned} \quad (5.81)$$

and has the form

$$(5.82) \quad \phi_{\text{int}}^{n-h} : \begin{cases} \Theta_{n-h}^{(h-1)} = F_{\text{int}}^{(h)}(p_{*,n-h}^{(h)}, q_{*,n-h}^{(h)}, \tilde{y}_*^{(h)}) \\ \vartheta_{n-h}^{(h-1)} - \pi = G_{\text{int}}^{(h)}(p_{*,n-h}^{(h)}, q_{*,n-h}^{(h)}, \tilde{y}_*^{(h)}) \\ \hat{z}_j^{(h-1)} = \hat{z}_{*,j}^{(h)} e^{i\psi_{\text{int},j}^{(h)}}(p_{*,n-h}^{(h)}, q_{*,n-h}^{(h)}, \tilde{y}_*^{(h)}) \\ y_{n-h}^{(h-1)} = y_{*,n-h}^{(h)} \\ x_{n-h}^{(h-1)} = x_{*,n-h}^{(h)} + \varphi_{\text{int}}^{(h)}(p_{*,n-h}^{(h)}, q_{*,n-h}^{(h)}, \tilde{y}_*^{(h)}) \end{cases}$$

with $F_{\text{int}}^{(h)}$, $G_{\text{int}}^{(h)}$ odd, $\psi_{\text{int},j}^{(h)}$, $\varphi_{\text{int}}^{(h)}$ even in $(p_{*,n-h}^{(h)}, q_{*,n-h}^{(h)})$,

$$(5.83) \quad \begin{aligned} \tilde{y}_*^{(h)} &:= \left(\frac{(p_{*,n-h+1}^{(h)})^2 + (q_{*,n-h+1}^{(h)})^2}{2}, \dots, \frac{(p_{*,1}^{(h)})^2 + (q_{*,1}^{(h)})^2}{2}, y_{*,n-h}^{(h)} \right) \\ \hat{z}_j^{(h-1)} &:= (p_j^{(h-1)}, q_j^{(h-1)}) := p_j^{(h-1)} + iq_j^{(h-1)} \\ \hat{z}_{*,j}^{(h)} &:= (p_{*,j}^{(h)}, q_{*,j}^{(h)}) := p_{*,j}^{(h)} + iq_{*,j}^{(h)} \end{aligned}$$

with $j = n - h + 1, \dots, n - 1$, for $n \geq 3$, $h \geq 2$ and $y_{*,n-h}^{(h)}$ as in (5.81).

In particular, observe that ϕ_{int}^{n-h} enjoys the following properties:

- it is co-variant with the symmetry: if

$$\phi_{\text{int}}^{n-h}(t_*^{(h)}, z_*^{(h)}, y_*^{(h)}, x_*^{(h)}) = (t^{(h-1)}, z^{(h-1)}, y^{(h-1)}, x^{(h-1)}),$$

then

$$\phi_{\text{int}}^{n-h}(-t_*^{(h)}, -z_*^{(h)}, y_*^{(h)}, x_*^{(h)}) = (-t^{(h-1)}, -z^{(h-1)}, y^{(h-1)}, x^{(h-1)});$$

- leaves the “actions”

$$\tilde{y}_*^{(h)} = \tilde{y}^{(h-1)}$$

invariant, where $\tilde{y}^{(h)}$ is as in (5.83), and

$$\tilde{y}^{(h-1)} := \left(\frac{(p_{n-h+1}^{(h-1)})^2 + (q_{n-h+1}^{(h-1)})^2}{2}, \dots, \frac{(p_1^{(h-1)})^2 + (q_1^{(h-1)})^2}{2}, y_{n-h}^{(h-1)} \right)$$

is defined analogously;

- leaves the averages with respect to the x-coordinates invariant. Namely, for any real-analytic function g on $\mathfrak{D}_{\text{sec}}^{h-1}$,

$$\Pi_{x_*^{(h)}}(g \circ \phi_{\text{int}}^{n-h}) = (\Pi_{x^{(h-1)}}g) \circ \phi_{\text{int}}^{n-h}.$$

Applying ϕ_{int}^{n-h} to $\widehat{f_{\text{sec},h-1}}$ in (5.78), we obtain

$$\begin{aligned} f_{\text{sec,int},h-1} &:= \widehat{f_{\text{sec},h-1}} \circ \phi_{\text{int}}^{n-h} \\ &= h_{\text{sec},h-1} + h_{\text{sec}}^{n-h} + \sum_{i=1}^{n-h-1} f_{\text{norm,int},h-1}^i(t_{*,i}^{(h)}, \tilde{y}_{*,i}^{(h)}, \tilde{x}_{*,i}^{(h)}) \\ &= \sum_{i=n-h}^{n-1} h_{\text{sec},h}^i(\hat{y}_{*,i}^{(h)}) + \sum_{i=1}^{n-h-1} f_{\text{norm,int},h-1}^i(t_{*,i}^{(h)}, \tilde{y}_{*,i}^{(h)}, \tilde{x}_{*,i}^{(h)}) \end{aligned}$$

with

$$(5.84) \quad \mathfrak{h}_{\text{sec},h} := \mathfrak{h}_{\text{sec},h-1} + \mathfrak{h}_{\text{sec}}^{n-h}, \quad f_{\text{norm,int},h-1}^i := f_{\text{norm},h-1}^i \circ \phi_{\text{int}}^{n-h}$$

and (as it follows from (5.71) with $h-1$ replacing h and (5.82)) $f_{\text{norm,int},h-1}^i$ depends only on the arguments

$$(5.85) \quad \begin{aligned} \mathfrak{t}_{*,i}^{(h)} &:= (\Theta_{*,i}^{(h)}, \dots, \Theta_{*,n-h-1}^{(h)}, \vartheta_{*,i}^{(h)}, \dots, \vartheta_{*,n-h-1}^{(h)}) \\ \tilde{\mathfrak{y}}_{*,i}^{(h)} &:= (p_{*,n-h}^{(h)}, q_{*,n-h}^{(h)}, \frac{(p_{*,n-h+1}^{(h)})^2 + (q_{*,n-h+1}^{(h)})^2}{2}, \dots, \frac{(p_{*,n-1}^{(h)})^2 + (q_{*,n-1}^{(h)})^2}{2}, \\ &\quad \chi_{*,i-1}^{(h)}, \dots, \chi_{*,n-1}^{(h)}, \Lambda_{*,i}^{(h)}, \dots, \Lambda_{*,n}^{(h)}) \\ \tilde{\mathfrak{x}}_{*,i}^{(h)} &:= \begin{cases} (\kappa_{*,i}^{(h)}, \dots, \kappa_{*,n-h-1}^{(h)}) & n \geq 4 \ \& \ 1 \leq h-1 \leq n-3 \\ \emptyset & \text{otherwise} \end{cases} \end{aligned}$$

The next step will be to retain the dependence on $(p_{n-h}^{(h)}, q_{n-h}^{(h)})$ only via $\frac{(p_{n-h}^{(h)})^2 + (q_{n-h}^{(h)})^2}{2}$ and, for $h < n-1$, to eliminate from $f_{\text{sec,int},h-1}$ the dependence upon the angle $k_{*,n-h-1}^{(h)}$, up to an exponential remainder. Namely, we look for another canonical, real-analytic change of coordinates

$$(5.86) \quad \begin{aligned} \phi_{\text{norm}}^{n-h} : \quad \mathfrak{D}_{\text{sec}}^h &\rightarrow \mathfrak{D}_{\text{int}}^h \\ (\mathfrak{t}^{(h)}, \mathfrak{z}^{(h)}, \mathfrak{y}^{(h)}, \mathfrak{x}^{(h)}) &\rightarrow (\mathfrak{t}_*^{(h)}, \mathfrak{z}_*^{(h)}, \mathfrak{y}_*^{(h)}, \mathfrak{x}_*^{(h)}) \end{aligned}$$

so as to conjugate $f_{\text{sec,int},h-1}$ to a new Hamiltonian

$$(5.87) \quad \widehat{f_{\text{sec},h}} := f_{\text{sec,int},h-1} \circ \phi_{\text{norm}}^{n-h} = \mathfrak{h}_{\text{sec},h} + \sum_{i=1}^{n-h-1} f_{\text{norm},h}^i(\mathfrak{t}_i^{(h)}, \hat{\mathfrak{y}}_i^{(h)}, \hat{\mathfrak{x}}_i^{(h)}) + \widehat{f_{\text{exp,sec},h}}$$

where $f_{\text{norm},h}^i$ and $\widehat{f_{\text{exp,sec},h}}$ satisfy (5.74)-(5.76). We choose $\mathfrak{D}_{\text{sec}}^h$ as the subset of $\mathfrak{D}_{\text{int}}^h$ where the map

$$\omega_{\text{sec},h} := \begin{cases} \frac{\partial}{\partial \frac{(p_{n-h}^{(h)})^2 + (q_{n-h}^{(h)})^2}{2}, \chi_{n-h-1}^{(h)}} \mathfrak{h}_{\text{sec},h} & h = 2, \dots, n-2 \ \& \ n \geq 4 \\ \frac{\partial}{\partial \frac{(p_1^{(n-1)})^2 + (q_1^{(n-1)})^2}{2}} \mathfrak{h}_{\text{sec},n-1} & h = n-1 \end{cases}$$

does not verify resonances up to order \bar{K} , and next we apply a suitable normal form theory (Proposition D.1). We shall choose ϕ_{norm}^{n-h} in such a way that

- it is co-variant with the symmetry: if

$$\phi_{\text{norm}}^{n-h}(\mathfrak{t}^{(h)}, \mathfrak{z}^{(h)}, \mathfrak{y}^{(h)}, \mathfrak{x}^{(h)}) = (\mathfrak{t}_*^{(h)}, \mathfrak{z}_*^{(h)}, \mathfrak{y}_*^{(h)}, \mathfrak{x}_*^{(h)}),$$

then

$$(5.88) \quad \phi_{\text{norm}}^{n-h}(-\mathfrak{t}^{(h)}, -\mathfrak{z}^{(h)}, \mathfrak{y}^{(h)}, \mathfrak{x}^{(h)}) = (-\mathfrak{t}_*^{(h)}, -\mathfrak{z}_*^{(h)}, \mathfrak{y}_*^{(h)}, \mathfrak{x}_*^{(h)}),$$

- leaves the “actions”

$$\mathfrak{y}_{*,n-h}^{(h)} = \mathfrak{y}_{n-h}^{(h)}$$

unvaried, where

$$(5.89) \quad \begin{aligned} y_{n-h}^{(h)} &:= \left(\frac{(p_{n-h+1}^{(h)})^2 + (q_{n-h+1}^{(h)})^2}{2}, \dots, \frac{(p_1^{(h)})^2 + (q_1^{(h)})^2}{2}, \chi_{n-h}^{(h)}, \dots, \chi_{n-1}^{(h)}, \right. \\ &\quad \left. \Lambda_1^{(h)}, \dots, \Lambda_n^{(h)} \right); \\ y_{*,n-h}^{(h)} &:= \left(\frac{(p_{*,n-h+1}^{(h)})^2 + (q_{*,n-h+1}^{(h)})^2}{2}, \dots, \frac{(p_{*,1}^{(h)})^2 + (q_{*,1}^{(h)})^2}{2}, \chi_{*,n-h}^{(h)}, \dots, \chi_{*,n-1}^{(h)}, \right. \\ &\quad \left. \Lambda_{*,1}^{(h)}, \dots, \Lambda_{*,n}^{(h)} \right); \end{aligned}$$

• verifies

$$(5.90) \quad \Pi_{z_{*,n-h+1}^{(h)}, x_{*,n-h+1}^{(h)}} (g \circ \phi_{\text{norm}}^{n-h}) = \left(\Pi_{z_{n-h+1}^{(h)}, x_{n-h+1}^{(h)}} g \right) \circ \phi_{\text{norm}}^{n-h}.$$

The thesis of Proposition 5.2 at rank h follows, with

$$f_{\text{sec},h} := \widehat{f_{\text{sec},h}} + f_{\text{exp,sec},h-1} \circ \phi_{\text{sec}}^{n-h}, \quad f_{\text{exp,sec},h} := \widehat{f_{\text{exp,sec},h}} + f_{\text{exp,sec},h-1} \circ \phi_{\text{sec}}^{n-h}.$$

Case $h = 1$. The proof of this case uses similar ideas as the proof of the case $2 \leq h \leq n-1$ for $n \geq 3$. However, due to subtle differences between the two cases (compare, e.g., the inductive assumption on $f_{\text{norm},h-1}^{n-h}$ in (5.71) for $h \geq 2$ with Eq. (5.91); the definition of h_{sec}^{n-h} , ϕ_{int}^{n-h} for $h \geq 2$ in (5.80), with the definition of h_{sec}^{n-1} , ϕ_{int}^{n-1} in (5.93) and (5.96)), for sake of precision, we briefly discuss also this case.

Let $f_{\text{sec},0}$ be as in (5.77). In view of (5.60) and (5.61), we can split

$$(5.91) \quad f_{\text{sec},0} = \overline{f_{\mathcal{P}}^{n-1,n}^{(2)}} + \overline{f_{\mathcal{P}}^{n-1}^{(\geq 3)}} + \widetilde{f_{\text{fast}}^{n-1}} + \sum_{i=1}^{n-2} f_{\text{fast}}^i$$

where

$$\overline{f_{\mathcal{P}}^{n-1}^{(\geq 3)}} := \overline{f_{\mathcal{P}}^{n-1}^{(\geq 2)}} - \overline{f_{\mathcal{P}}^{n-1}^{(2)}}$$

and the summand appears only when $n \geq 3$. As for $\overline{f_{\mathcal{P}}^{n-1,n}^{(2)}}$, by Lemmata 3.4 (see also Lemma 5.2), we find a domain \mathfrak{B}_{n-1} (defined in Eq. (5.97) below), a real-analytic and canonical transformation

$$(5.92) \quad \begin{aligned} \phi_{\text{int}}^{n-1} : (z_{*,n-1}^{(1)}, y_{*,n-1}^{(1)}, x_{*,n-1}^{(1)}) &\in \overline{\mathfrak{B}_{n-1}} \rightarrow (z_{n-1}^{(0)}, y_{n-1}^{(0)}, x_{n-1}^{(0)}) \\ &\in \overline{\mathfrak{D}_{n-1}} := \phi_{\text{int}}^{n-1}(\overline{\mathfrak{B}_{n-1}}) \end{aligned}$$

of the form (5.82), with $h = 1$ (but neglecting the coordinates $\hat{z}_j^{(0)}$, $\hat{z}_{j,*}^{(1)}$) such that

$$(5.93) \quad \overline{f_{\mathcal{P}}^{n-1}^{(2)}} \circ \phi_{\text{int}}^{n-1} = \overline{h_{\text{sec}}^{n-1}}(\hat{y}_{*,n-1}^{(1)})$$

depends only on

$$(5.94) \quad \hat{y}_{*,n-1}^{(1)} = \left(\frac{(p_{*,n-1}^{(1)})^2 + (q_{*,n-1}^{(1)})^2}{2}, \chi_{*,n-2}^{(1)}, \chi_{*,n-1}^{(1)}, \Lambda_{*,n-1}^{(1)}, \Lambda_{*,n}^{(1)} \right).$$

In (5.92), we have let

$$\left\{ \begin{array}{l} z_{*,n-1}^{(1)} := (p_{*,n-1}^{(1)}, q_{*,n-1}^{(1)}) \\ y_{*,n-1}^{(1)} := (\chi_{*,n-2}^{(1)}, \chi_{*,n-1}^{(1)}, \Lambda_{*,n-1}^{(1)}, \Lambda_{*,n}^{(1)}) \\ x_{*,n-1}^{(1)} := (\kappa_{*,n-2}^{(1)}, \kappa_{*,n-1}^{(1)}) \end{array} \right. \quad \left\{ \begin{array}{l} t_{n-1}^{(0)} := (\Theta_{n-1}^{(0)}, \vartheta_{n-1}^{(0)}) \\ y_{n-1}^{(0)} := (\chi_{n-2}^{(0)}, \chi_{n-1}^{(0)}, \Lambda_{n-1}^{(0)}, \Lambda_n^{(0)}) \\ x_{n-1}^{(0)} := (\kappa_{n-2}^{(0)}, \kappa_{n-1}^{(0)}). \end{array} \right.$$

We let

$$\begin{cases} \mathfrak{t}^{(0)} := (\Theta^{(0)}, \vartheta^{(0)}) \\ \mathfrak{y}^{(0)} := (\chi^{(0)}, \Lambda^{(0)}) \\ \mathfrak{x}^{(0)} := (\kappa^{(0)}, \ell^{(0)}) \end{cases} \quad \begin{cases} \mathfrak{t}_*^{(1)} := (\Theta_*^{(1)}, \vartheta_*^{(1)}) \\ \mathfrak{z}_*^{(1)} := (p_*^{(1)}, q_*^{(1)}) \\ \mathfrak{y}_*^{(1)} := (\chi_*^{(1)}, \Lambda_*^{(1)}) \\ \mathfrak{x}_*^{(1)} := (\kappa_*^{(1)}, \ell_*^{(1)}) \end{cases}$$

analogously to (5.70), with $h = 0, 1$, and then we regard the map in (5.92) as a map

$$\phi_{\text{int}}^{n-1} : (\mathfrak{t}_*^{(1)}, \mathfrak{z}_*^{(1)}, \mathfrak{y}_*^{(1)}, \mathfrak{x}_*^{(1)}) \in \mathfrak{D}_{\text{int}}^1 \rightarrow (\mathfrak{t}^{(0)}, \mathfrak{y}^{(0)}, \mathfrak{x}^{(0)})$$

on the set

$$\mathfrak{D}_{\text{int}}^1 := \left\{ (\mathfrak{t}_*^{(1)}, \mathfrak{z}_*^{(1)}, \mathfrak{y}_*^{(1)}, \mathfrak{x}_*^{(1)}) : (\mathfrak{z}_{*,n-1}^{(1)}, \mathfrak{y}_{*,n-1}^{(1)}, \mathfrak{x}_{*,n-1}^{(1)}) \in \overline{\mathfrak{B}_{n-1}} \right\}$$

where ϕ_{int}^{n-1} is defined on the extra-coordinates via the identity. $\mathfrak{D}_{\text{int}}^1$ has the form in (5.79), with $h = 1$. Applying this extension to $f_{\text{sec},0}$ in (5.91) we obtain

$$f_{\text{sec,int},0} := f_{\text{sec},0} \circ \overline{\phi_{\text{int}}^{n-1}} = \overline{h_{\text{sec}}^{n-1}}(\hat{y}_{*,n-1}^{(1)}) + \sum_{i=1}^{n-1} f_{\text{norm,int},0}^i(\mathfrak{t}_{*,i}^{(1)}, \tilde{y}_{*,i}^{(1)}, \tilde{x}_{*,i}^{(1)})$$

where

$$f_{\text{norm,int},0}^{n-1} := (\overline{f_{\mathcal{P}}^{n-1}}(\geq 3) + \widetilde{f_{\text{fast}}^{n-1}}) \circ \overline{\phi_{\text{int}}^{n-1}}, \quad f_{\text{norm,int},0}^i := \widehat{f_{\text{fast}}^i} \circ \overline{\phi_{\text{int}}^{n-1}}$$

and, as a consequence of (5.61) and of (5.82), with $h = 1$, $f_{\text{norm,int},0}^i$ depends only on the arguments

$$\begin{aligned} \mathfrak{t}_{*,i}^{(1)} &:= (\Theta_{*,i}^{(1)}, \dots, \Theta_{*,n-2}^{(1)}, \vartheta_{*,i}^{(1)}, \dots, \vartheta_{*,n-2}^{(1)}) \\ \tilde{y}_{*,i}^{(1)} &:= (p_{*,n-1}^{(1)}, q_{*,n-1}^{(1)}, \chi_{*,i-1}^{(1)}, \dots, \chi_{*,n-1}^{(1)}, \Lambda_{*,i}^{(1)}, \dots, \Lambda_{*,n}^{(1)}) \\ \tilde{x}_{*,i}^{(1)} &:= (\kappa_{*,i}^{(1)}, \dots, \kappa_{*,n-1}^{(1)}). \end{aligned}$$

Note, in particular, that $f_{\text{norm,int},0}^{n-1}$ is a function of (5.95)

$$(\mathfrak{t}_{*,n-1}, \mathfrak{y}_{*,n-1}, \mathfrak{x}_{*,n-1}) = (p_{*,n-1}^{(1)}, q_{*,n-1}^{(1)}, \chi_{*,n-2}^{(1)}, \chi_{*,n-1}^{(1)}, \Lambda_{*,n-1}^{(1)}, \Lambda_{*,n}^{(1)}, \kappa_{*,n-1}^{(1)}).$$

In view of the fact that $\overline{h_{\text{sec}}^{n-1}}$ depends on the actions in (5.94), we aim to eliminate from $f_{\text{sec,int},0}$ the dependence on the following angles

$$\begin{cases} \kappa_{*,1} & \text{if } n = 2 \\ \kappa_{*,n-2}, \kappa_{*,n-1} & \text{if } n \geq 3 \end{cases}$$

and to retain the dependence on $(p_{*,n-1}^{(1)}, q_{*,n-1}^{(1)})$ only via $\frac{(p_{*,n-1}^{(1)})^2 + (q_{*,n-1}^{(1)})^2}{2}$. Then we choose a domain $\mathfrak{D}_{\text{sec}}^1 \subset \mathfrak{D}_{\text{int}}^1$ as in (5.69) where the frequency

$$\omega_{\text{sec},1} := \begin{cases} \frac{\partial}{\partial \frac{(p_{*,n-1}^{(1)})^2 + (q_{*,n-1}^{(1)})^2}{2}} \overline{h_{\text{sec}}^{n-1}} \Big|_{\mathcal{X}_{*,n-1}^{(1)}} & n = 2 \\ \frac{\partial}{\partial \frac{(p_{*,n-1}^{(1)})^2 + (q_{*,n-1}^{(1)})^2}{2}} \overline{h_{\text{sec}}^{n-1}} \Big|_{\mathcal{X}_{*,n-2}^{(1)}, \mathcal{X}_{*,n-1}^{(1)}} & n \geq 3 \end{cases}$$

is non-resonant up to the order \bar{K} and on this domain we construct a real-analytic transformation ϕ_{norm}^{n-1} as in (5.86) which conjugates $\overline{f_{\text{sec},1}}$ to a Hamiltonian

$$f_{\text{sec},1} := f_{\text{sec,int},0} \circ \phi_{\text{norm}}^{n-1} = \overline{h_{\text{sec}}^{n-1}}(\hat{y}_{n-1}^{(1)}) + \sum_{i=1}^{n-1} f_{\text{norm},1}^i(\hat{v}_i^{(1)}, \hat{y}_i^{(1)}, \hat{x}_i^{(1)}) + f_{\text{exp,sec},1}$$

Now, since (as it follows from (5.95)), $f_{\text{norm},1}^{n-1}$ is actually a function of $\hat{y}_{n-1}^{(1)}$ only, this step is proved, with

$$(5.96) \quad h_{\text{sec}}^{n-1}(\hat{y}_{n-1}^{(1)}) := \overline{h_{\text{sec}}^{n-1}}(\hat{y}_{n-1}^{(1)}) + f_{\text{norm},1}^{n-1}(\hat{y}_{n-1}^{(1)}).$$

□

5.2.1. Construction of ϕ_{int}^{n-1} . The following lemma completes Lemma 3.4. In particular, it provides the transformation $\phi_{\text{int}}^{n-1} = \overline{\phi_{\text{int}}^{n-1}}$ in (5.93).

LEMMA 5.2. *Let $i = 1, \dots, n-1$. Let \mathcal{A}, X, θ in (4.46) be chosen in such a way that*

$$(5.97) \quad \inf_{\mathfrak{D}_p} |g| > 0, \quad \sup_{\mathfrak{D}_p} |\arg g| < \frac{\pi}{4}$$

$$\forall g \in \left\{ \chi_{i-1}, \quad \chi_i, \quad \chi_{i-1} + \chi_i, \quad 5\chi_{i-1}\Lambda_i^2 - (\chi_{i-1} - \chi_i)^2(4\chi_{i-1} - \chi_i) \right\}.$$

Then, the domains $\overline{\mathfrak{B}_i}$ in (3.42), the functions $\overline{h_{\text{sec}}^i}$ and the transformations $\overline{\phi_{\text{int}}^i}$ can be taken as follows

$$(5.98) \quad \begin{aligned} \overline{\mathfrak{B}_i} &= \begin{cases} B_{\varepsilon_i}^2 \times \mathcal{A}_{\theta_i}^i \times \chi_{\theta_i}^i \times \mathbb{T}_{s_i}^4 & i = n-1 \\ B_{\varepsilon_i}^2 \times \mathcal{A}_{\theta_i}^i \times \chi_{\theta_i}^i \times \mathbb{T}_{s_i}^5 & i = 1, \dots, n-2 \ \& \ n \geq 3 \end{cases} \\ \overline{\phi_{\text{int}}^i} &: \begin{cases} \Theta_i = \frac{p_i}{\beta_i} + f_i(p_i, q_i, y_i^*) \\ \vartheta_i - \pi = \beta_i q_i + g_i(p_i, q_i, y_i^*) \\ y_i = y_i^* \\ x_i = x_i^* + \varphi_i(p_i, q_i, y_i^*) \end{cases} \\ \overline{h_{\text{sec}}^i} &= \mathcal{A}_i \left[E_i + \Omega_i \frac{p_i^2 + q_i^2}{2} + \tau_i \left(\frac{p_i^2 + q_i^2}{2} \right)^2 + O(p_i, q_i)^6 \right] \end{aligned}$$

where $X_{\bar{\theta}_i}^i \times A_{\bar{\theta}_i}^i$ denote the projection of the set $X_{\bar{\theta}} \times A_{\bar{\theta}}$ over the coordinates y_i in (3.43), $\bar{\theta} := \theta/2$, $\bar{s} := s/2$, f_i, g_i are $O(p_i, q_i)^3$, odd in (p_i, q_i) , φ_i is $O(p_i, q_i)^2$, and

$$\begin{aligned}
\varepsilon_i &= c_i \sqrt{\bar{\theta}_i} \\
\beta_i &:= \sqrt[4]{\frac{5\chi_{i-1}\Lambda_i^2 - (\chi_{i-1} - \chi_i)^2(4\chi_{i-1} - \chi_i)}{\chi_{i-1}^2\chi_i^2(\chi_{i-1} + \chi_i)}} \\
\mathcal{A}_i &:= m_i m_{i+1} \frac{a_i^2}{4a_{i+1}^3} \\
E_i &:= -\frac{\Lambda_{i+1}^3}{2(\chi_i - \chi_{i+1})^3} \left(5 - 3\frac{(\chi_{i-1} - \chi_i)^2}{\Lambda_i^2} \right) \\
\Omega_i &:= \frac{3\Lambda_{i+1}^3}{\chi_i\Lambda_i^2(\chi_i - \chi_{i+1})^3} \sqrt{(5\chi_{i-1}\Lambda_i^2 - (\chi_{i-1} - \chi_i)^2(4\chi_{i-1} - \chi_i))(\chi_{i-1} + \chi_i)} \\
\tau_i &:= \frac{\Lambda_{i+1}^3}{\chi_i^2(\chi_i - \chi_{i+1})^3} \left[-\frac{9}{16} \frac{(\chi_{i-1} - \chi_i)^2(3\chi_{i-1} - \chi_i)(5\chi_{i-1} + \chi_i)}{\chi_{i-1}^3\chi_i\Lambda_i^2\beta_i^4} \right. \\
(5.99) \quad &\left. - \frac{3}{8} \frac{2\chi_{i-1}^3 + 9\chi_{i-1}^2\chi_i + 2\chi_{i-1}\chi_i^2 + \chi_i^3}{\chi_{i-1}\Lambda_i^2} - \frac{3}{16} \frac{\chi_{i-1}\chi_i^2}{\Lambda_i^2} (4\chi_{i-1} + \chi_i)\beta_i^4 \right]
\end{aligned}$$

with $\chi_n \equiv 0$, \bar{c}_i depending at most on the ratios a_i^+/a_i^- , the masses m_1, \dots, m_n and, as usual, $\sqrt[n]{z}$ denoting the principal determination of the n^{th} root of a complex number z .

PROOF. Since the formula for $\overline{f_{\mathcal{P}}^{n-1, n}}$ coincides with the one for $\overline{f_{\mathcal{P}}^{n-1, n}}$ taking $\chi_n \equiv 0$, we shall only work on the terms $\overline{f_{\mathcal{P}}^{i, i+1}}$'s.

Let y_i be as in (3.43), and let

$$(5.100) \quad \mathfrak{D}_i : (\Theta_i, \vartheta_i) \in \mathbb{T}_{\Theta_i^+, \vartheta_i^+}^i \quad y_i \in \mathcal{A}_{\theta_i}^i \times X_{\theta_i}^i \quad x_i \in \mathbb{T}_s^{m_i}$$

where $\mathbb{T}_{\Theta_i^+, \vartheta_i^+}^i$ is the projection of $\mathbb{T}_{\Theta^+, \vartheta^+}$ over the coordinates (Θ_i, ϑ_i) , while m_i is 4 or 5, accordingly to (5.98). We shall obtain the transformation $\overline{\phi_{\text{int}}^i}$ in (3.42) as a product $\overline{\phi_{\text{int}}^i} = \overline{\phi_{\text{diag}}^i} \circ \overline{\phi_{\text{bir}}^i}$, where $\overline{\phi_{\text{diag}}^i}$ and $\overline{\phi_{\text{bir}}^i}$ are described below.

A Taylor expansion of $\overline{f_{\mathcal{P}}^{i, i+1}}$ around $(\Theta_i, \vartheta_i) = (0, \pi)$ gives

$$(5.101) \quad \overline{f_{\mathcal{P}}^{i, i+1}} = \mathcal{A}_i \left[E_i + \Omega_i \frac{\beta_i^2 \Theta_i^2 + \frac{(\vartheta_i - \pi)^2}{\beta_i^2}}{2} + \mathcal{R}_i \right]$$

where $\mathcal{A}_i, E_i, \beta_i, \Omega_i$ are as in (5.99). Note that β_i, Ω_i are well defined under the assumption (5.97). The expansion in (5.101) shows that $(\Theta_i, \vartheta_i) = (0, \pi)$ is an elliptic equilibrium point for $\overline{f_{\mathcal{P}}^{i, i+1}}$. The remainder \mathcal{R}_i is given by

$$\begin{aligned}
\mathcal{R}_i &= F \left[-\frac{3}{2} \frac{4\Theta_i^2 - \chi_i^2}{\Lambda_i^2} \left(\frac{(\chi_i^2 - \chi_{i-1}^2)^2}{(\sqrt{\chi_i^2 - \Theta_i^2} + \sqrt{\chi_{i-1}^2 - \Theta_i^2})^2} \right. \right. \\
&\quad \left. \left. + 2\sqrt{(\chi_i^2 - \Theta_i^2)(\chi_{i-1}^2 - \Theta_i^2)}(1 + \cos \vartheta_i) \right) + \frac{1}{2} \frac{(\chi_i^2 - \Theta_i^2)(\chi_{i-1}^2 - \Theta_i^2)}{\Lambda_i^2} \sin^2 \vartheta_i \right]
\end{aligned}$$

where the symbol F on the left means that only terms of the fourth order in $(\Theta_i, \vartheta_i - \pi)$ have to be included. The lower order expansion of \mathcal{R}_i is

$$\mathcal{R}_i = \tau_{1,i}\Theta_i^4 + \tau_{2,i}(\vartheta_i - \pi)^2\Theta_i^2 + \tau_{3,i}(\vartheta_i - \pi)^4 + \mathcal{O}(\Theta_i, \vartheta_i - \pi)^6$$

with

$$\begin{aligned}\tau_{1,i} &:= \tau_1(y_i) := -\frac{3(\chi_{i-1} - \chi_i)^2(3\chi_{i-1} - \chi_i)(5\chi_{i-1} + \chi_i)}{8\chi_{i-1}^3\chi_i\Lambda_i^2} \\ \tau_{2,i} &:= \tau_2(y_i) := -\frac{3(2\chi_{i-1}^3 + 9\chi_{i-1}^2\chi_i + 2\chi_{i-1}\chi_i^2 + \chi_i^3)}{4\chi_{i-1}\Lambda_i^2} \\ \tau_{3,i} &:= \tau_3(y_i) := -\frac{\chi_{i-1}\chi_i^2}{8\Lambda_i^2}(4\chi_{i-1} + \chi_i).\end{aligned}$$

We introduce the generating function

$$S_{\text{diag},i}(\tilde{p}_i, \tilde{y}_i, \vartheta_i, \mathbf{x}_i) = \frac{\tilde{p}_i(\vartheta_i - \pi)}{\tilde{\beta}_i} + \tilde{y}_i \mathbf{x}_i.$$

It generates the canonical transformation

$$\overline{\phi_{\text{diag}}^i} : \quad \Theta_i = \frac{\tilde{p}_i}{\tilde{\beta}_i} \quad \vartheta_i - \pi = \tilde{\beta}_i \tilde{q}_i, \quad y_i = \tilde{y}_i, \quad \mathbf{x}_i = \tilde{\mathbf{x}}_i + \frac{\partial_{y_i} \beta_i(\tilde{y}_i)}{\beta_i(\tilde{y}_i)} \tilde{p}_i \tilde{q}_i$$

which transforms $\overline{f_{\mathcal{P}}^{i,i+1}}$ into

$$(5.102) \quad \overline{f_{\text{diag},i}} = \overline{f_{\mathcal{P}}^{i,i+1}} \circ \overline{\phi_{\text{diag}}^i} = \tilde{\mathcal{A}}_i \left[\tilde{\mathbf{E}}_i + \tilde{\Omega}_i \frac{\tilde{p}_i^2 + \tilde{q}_i}{2} + \tilde{\mathcal{R}}_i \right]$$

with

$$\tilde{\beta}_i := \beta(\tilde{y}_i), \quad \tilde{\mathcal{A}}_i := \mathcal{A}_i(\tilde{y}_i), \quad \tilde{\mathbf{E}}_i := C(\tilde{y}_i), \quad \tilde{\Omega}_i := \Omega(\tilde{y}_i),$$

$$\tilde{\mathcal{R}}_i := \mathcal{R}_i \circ \phi_{\text{diag}}^i = \tilde{\tau}_{1,i} \tilde{p}_i^4 + \tilde{\tau}_{2,i} \tilde{p}_i^2 \tilde{q}_i^2 + \tilde{\tau}_{3,i} \tilde{q}_i^4 + \mathcal{O}(\tilde{p}_i, \tilde{q}_i)^6$$

$$\tilde{\tau}_{1,i} := \frac{\tau_1(\tilde{y}_i)}{\tilde{\beta}_i^4}, \quad \tilde{\tau}_{2,i} := \tau_2(\tilde{y}_i), \quad \tilde{\tau}_{3,i} := \tau_3(\tilde{y}_i) \tilde{\beta}_i^4$$

To compute the domain of $\overline{\phi_{\text{diag}}^i}$, we use the following inequalities, which readily follow from the definitions:

$$\hat{\mathbf{c}}_i \frac{\sqrt{\theta_i}}{\mathbf{G}_n^+} \leq |\beta_i| \leq \frac{1}{\hat{\mathbf{c}}_i} \frac{\sqrt{\theta_i}}{\mathbf{G}_n^-}$$

and

$$\left| \frac{\partial_{y_i} \beta_i(\tilde{y}_i)}{\beta_i(\tilde{y}_i)} \right| \leq \frac{1}{\hat{\mathbf{c}}_i \theta_i}.$$

We then see that, choosing a suitable $\tilde{\mathbf{c}}_i \leq \hat{\mathbf{c}}_i$, and the domain

$$\widetilde{\mathfrak{B}}_i : \quad |(\tilde{p}_i, \tilde{q}_i)| \leq \tilde{\varepsilon}_i = \tilde{\mathbf{c}}_i \sqrt{\theta_i} \quad \tilde{y}_i \in \mathcal{A}_{\theta_i}^i \times \mathbf{X}_{\theta_i}^i \quad \tilde{\mathbf{x}}_i \in \mathbb{T}_{\frac{3}{4}s}^{m_i}$$

inequalities¹ (5.100) are verified, as desired. Now we look for another canonical transformation

$$\overline{\phi_{\text{bit}}^i} : \quad (p_i^*, q_i^*, y_i^*, \mathbf{x}_i^*) \rightarrow (\tilde{p}_i, \tilde{q}_i, \tilde{y}_i, \tilde{\mathbf{x}}_i) \quad (y_i^* = \tilde{y}_i)$$

¹Compare (4.50).

defined in an analogous domain

$$\overline{\mathfrak{B}_i^*} := \overline{\mathfrak{B}_i} : \quad |(p_i^*, q_i^*)| \leq \varepsilon_i = \mathbf{c}_i^* \sqrt{\theta_i} \quad y_i^* \in \mathcal{A}_{\theta_i}^i \times X_{\theta_i}^i \quad x_i^* \in \mathbb{T}_{\frac{\varepsilon}{2}}^{m_i}$$

with $\mathbf{c}_i^* := \mathbf{c}_i \leq \tilde{\mathbf{c}}_i/2$, such that

$$\overline{f_{\text{diag},i}} \circ \overline{\phi_{\text{bit}}^i} = \overline{h_{\text{sec}}^i}$$

satisfies the thesis of the lemma. We aim to apply Theorem D.1, with

$$h = \tilde{E}_i + \tilde{\Omega}_i \frac{\tilde{p}_i^2 + \tilde{q}_i}{2}, \quad f = \mathcal{R}_i, \quad \varepsilon = 2\mathbf{c}_i^* \sqrt{\theta_i}, \quad \bar{\varepsilon} = \mathbf{c}_i^* \sqrt{\theta_i}.$$

We have to check that inequalities (D.152) are satisfied. We can take \mathbf{a} and \mathbf{e} as it follows from the following inequalities, which, in turn, are easily implied by the definitions

$$\begin{aligned} \inf_{\mathfrak{B}_i^*} |\partial h| &= \inf_{\mathfrak{B}_i^*} |\tilde{\Omega}_i| \geq \frac{\tilde{\mathbf{c}}_i |\mathbf{G}_n^-|^2}{\theta_i} =: \mathbf{a} \\ \sup_{\mathfrak{B}_i^*} |\tilde{\mathcal{R}}_i| &\leq \sup_{\mathfrak{D}_i^*} |\mathcal{R}_i| \leq \frac{1}{\tilde{\mathbf{c}}_i} \max \sup_{\mathfrak{D}_i^*} \left\{ \frac{(\Theta_i^*)^4}{(\mathbf{G}_n^-)^2}, (\vartheta_i^* - \pi)^2 \Theta_i^2, (\mathbf{G}_n^+)^2 (\vartheta_i^* - \pi)^4 \right\} \\ &\leq \frac{(\mathbf{c}^*)^4 (\mathbf{G}_n^+)^2}{\tilde{\mathbf{c}}_i} =: \mathbf{e} \end{aligned}$$

Here, we have used that, for $|(p_i^*, q_i^*)| \leq 2\mathbf{c}^* \sqrt{\theta_i}$, $(\Theta_i^*, \vartheta_i^*) := (\overline{\phi_{\text{diag}}^i})^{-1}(p_i^*, q_i^*)$ verifies

$$\begin{aligned} |\Theta_i^*| &= \frac{|p_i^*|}{|\beta_i|} \leq 2\mathbf{c}^* \sqrt{\theta_i} \frac{\mathbf{G}_n^+}{\tilde{\mathbf{c}}_i \sqrt{\theta_i}} = 2 \frac{\mathbf{c}^* \mathbf{G}_n^+}{\tilde{\mathbf{c}}_i} \\ |\vartheta_i^* - \pi| &= |q_i^*| |\beta_i| \leq \frac{2\mathbf{c}^* \sqrt{\theta_i}}{\mathbf{c}_1} \frac{\sqrt{\theta_i}}{\mathbf{G}_n^-} = 2 \frac{\mathbf{c}^*}{\tilde{\mathbf{c}}_i} \frac{\theta_i}{\mathbf{G}_n^-}. \end{aligned}$$

We then have that condition (D.152) holds, provided one takes

$$\mathbf{c}^* := \min \left\{ \frac{\mathbf{G}_n^-}{\mathbf{G}_n^+} \sqrt{\tilde{\mathbf{c}}_i \mathbf{c}_i}, \frac{\tilde{\mathbf{c}}_i}{2} \right\}.$$

From (5.102), one easily computes that the fourth order term of $\overline{h_{\text{sec}}^i}$ corresponds to be as in (5.98), with

$$\tau_i = \frac{3}{2} \tau_{1,i}^* + \frac{1}{2} \tau_{2,i}^* + \frac{3}{2} \tau_{3,i}^* \quad \tau_{j,i}^* := \tilde{\tau}_{j,i}(y_i^*).$$

Finally, properties (3.45) easily follow from the construction. \square

5.2.2. Construction of $\phi_{\text{int}}^1, \dots, \phi_{\text{int}}^{n-2}$ ($n \geq 3$). We have to solve (5.80), assuming that Proposition 5.2 holds, up to rank $h-1$. Accordingly to (5.74), (5.75) and letting

$$\overline{\Phi_{\text{int}}^{n-h+1}} := \overline{\phi_{\text{int}}^{n-h+1}} \circ \dots \circ \overline{\phi_{\text{int}}^{n-1}}$$

we may split

$$\begin{aligned}
f_{\text{norm},h-1}^{n-h} &= \sum_{j=n-h+1}^n \Pi_{h-1}(\overline{f_{\mathcal{P}}^{n-h,j}}^{(\geq 2)} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) + \widetilde{f_{\text{sec},h-1}^{n-h}} \\
&= \Pi_{h-1}(\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) + \Pi_{h-1}(\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(\geq 3)} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) \\
&+ \sum_{j=n-h+2}^n \Pi_{h-1}(\overline{f_{\mathcal{P}}^{n-h,j}}^{(\geq 2)} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) + \widetilde{f_{\text{sec},h-1}^{n-h}} \\
&= \overline{\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)}} + \Pi_{h-1}(\widetilde{\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)}} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) \\
&+ \Pi_{h-1}(\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(\geq 3)} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) \\
&+ \sum_{j=n-h+2}^n \Pi_{h-1}(\overline{f_{\mathcal{P}}^{n-h,j}}^{(\geq 2)} \circ \overline{\Phi_{\text{int}}^{n-h+1}}) + \widetilde{f_{\text{sec},h-1}^{n-h}}
\end{aligned}$$

where

$$\begin{aligned}
\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(\geq 3)} &:= \overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(\geq 2)} - \overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)} \\
\widetilde{\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)}} &:= \overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)} - \overline{\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)}}
\end{aligned}$$

and $\overline{\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)}}$ as in Lemma 3.4. Note that we have used that $\overline{f_{\mathcal{P}}^{n-h,n-h+1}}^{(2)}$ is left unvaried by $\overline{\Phi_{\text{int}}^{n-h+1}}$. Let $\overline{\mathfrak{B}_{n-h}}$, $\overline{\phi_{\text{int}}^{n-h}}$ be as in Lemmata 3.4, with the symbols $(\Theta_{n-h}, \vartheta_{n-h})$, \mathcal{Y}_{n-h} , \mathcal{X}_{n-h} of that lemma corresponding to

$$\begin{aligned}
\overline{\mathfrak{t}_{n-h}}^{(h-1)} &:= (\Theta_{n-h}^{(h-1)}, \vartheta_{n-h}^{(h-1)}) \\
\overline{\mathcal{Y}_{n-h}}^{(h-1)} &:= (\chi_{n-h-1}^{(h-1)}, \chi_{n-h}^{(h-1)}, \chi_{n-h+1}^{(h-1)}, \Lambda_{n-h}^{(h-1)}, \Lambda_{n-h+1}^{(h-1)}) \\
\overline{\mathcal{X}_{n-h}}^{(h-1)} &:= (\kappa_{n-h-1}^{(h-1)}, \kappa_{n-h}^{(h-1)}, \kappa_{n-h+1}^{(h-1)}, \ell_{n-h}^{(h-1)}, \ell_{n-h+1}^{(h-1)})
\end{aligned}$$

and the symbols (p_{n-h}, q_{n-h}) , \mathcal{Y}_{n-h}^* , \mathcal{X}_{n-h}^* to

$$\begin{aligned}
\overline{\mathfrak{z}_{n-h}}^{*(h)} &:= (p_{n-h}^{*(h)}, q_{n-h}^{*(h)}) \\
\overline{\mathcal{Y}_{n-h}}^{*(h)} &:= (\chi_{n-h-1}^{*(h)}, \chi_{n-h}^{*(h)}, \chi_{n-h+1}^{*(h)}, \Lambda_{n-h}^{*(h)}, \Lambda_{n-h+1}^{*(h)}) \\
\overline{\mathcal{X}_{n-h}}^{*(h)} &:= (\kappa_{n-h-1}^{*(h)}, \kappa_{n-h}^{*(h)}, \kappa_{n-h+1}^{*(h)}, \ell_{n-h}^{*(h)}, \ell_{n-h+1}^{*(h)}).
\end{aligned}$$

Defining

$$\begin{aligned}
\mathfrak{t}^{*(h)} &:= (\Theta^{*(h)}, \vartheta^{*(h)}) \\
\mathfrak{z}^{*(h)} &:= (p^{*(h)}, q^{*(h)}) \\
\mathfrak{y}^{*(h)} &:= (\chi^{*(h)}, \Lambda^{*(h)}) \\
\mathfrak{x}^{*(h)} &:= (\kappa^{*(h)}, \ell^{*(h)})
\end{aligned}$$

in an analogous way as in (5.70), we regard $\overline{\phi_{\text{int}}^{n-h}}$ as a map on the set

$$\overline{\mathfrak{D}_{\text{int}}^h} := \left\{ (\mathfrak{t}^{*(h)}, \mathfrak{z}^{*(h)}, \mathfrak{y}^{*(h)}, \mathfrak{x}^{*(h)}) : (\overline{\mathfrak{z}_{n-h}}^{*(h)}, \overline{\mathcal{Y}_{n-h}}^{*(h)}, \overline{\mathcal{X}_{n-h}}^{*(h)}) \in \overline{\mathfrak{B}_{n-h}} \right\}$$

extended via the identity on the extra-coordinates. We then have that $\overline{\phi_{\text{int}}^{n-h}}$ transforms $f_{\text{sec},h-1}^{n-h}$ into

$$\overline{f_{\text{sec},h-1}^{n-h}} := f_{\text{sec},h-1}^{n-h} \circ \overline{\phi_{\text{int}}^{n-h}} = \overline{h_{\text{sec}}^{n-h}} + \overline{f_{\text{sec}}^{n-h}}$$

where

$$\begin{aligned} \overline{f_{\text{sec}}^{n-h}} &:= \Pi_{h-1} \left(\overline{f_{\mathcal{P}}^{n-h,n-h+1}} \circ \overline{\Phi_{\text{int}}^{n-h}} \right) \\ &+ \Pi_{h-1} \left(\overline{f_{\mathcal{P}}^{n-h,n-h+1}} \circ \overline{\Phi_{\text{int}}^{n-h}} \right)^{(\geq 3)} + \sum_{j=n-h+2}^n \Pi_{h-1} \left(\overline{f_{\mathcal{P}}^{n-h,j}} \circ \overline{\Phi_{\text{int}}^{n-h}} \right)^{(\geq 2)} \\ (5.103) \quad &+ \overline{f_{\text{sec},h-1}^{n-h}} \circ \overline{\phi_{\text{int}}^{n-h}}. \end{aligned}$$

Here, we have used $\overline{\Phi_{\text{int}}^{n-h}} = \overline{\Phi_{\text{int}}^{n-h+1}} \circ \overline{\phi_{\text{int}}^{n-h}}$; that Π_{h-1} and $\overline{\phi_{\text{int}}^{n-h}}$ commute and observe that $\overline{\mathfrak{D}_{\text{int}}^h}$ has the form of $\mathfrak{D}_{\text{int}}^h$ in (5.79), with \hat{c}_h replaced by a suitable \hat{c}'_h of the same form. The function $\overline{f_{\text{sec}}^{n-h}}$ satisfies the following two properties:

- It depends only on

$$(p_{n-h}^{*(h)}, q_{n-h}^{*(h)}, \tilde{y}^{*(h)})$$

where $\tilde{y}^{*(h)}$ is defined analogously to (5.83);

- is uniformly bounded by the right hand side of the first inequality in (5.76) (this follows from the definition in (5.103));
- is even for

$$(p_{n-h}^{*(h)}, q_{n-h}^{*(h)}) \rightarrow -(p_{n-h}^{*(h)}, q_{n-h}^{*(h)}).$$

Proceeding in a similar way as we did for the construction of $\overline{\phi_{\text{bir}}^i}$ in the proof of Lemma 5.2, we may apply Theorem D.1, with

$$\begin{aligned} h &= \overline{h_{\text{sec}}^{n-h}}, \quad f = \overline{f_{\text{sec}}^{n-h}}, \quad (P, Q) = (p_{n-h}^{*(h)}, q_{n-h}^{*(h)}) \\ (P', Q') &= (\hat{z}_{n-h}^{*(h)}, y = y_{n-h}^{*(h)}, \quad x = x_{n-h}^{*(h)}). \end{aligned}$$

with $y_{n-h}^{*(h)}, x_{n-h}^{*(h)}$ defined analogously to $y_{*,n-h}^{(h)}, x_{*,n-h}^{(h)}$ in (5.81) and $\hat{z}_{n-h}^{*(h)}$ defined analogously to $\hat{z}_{*,n-h}^{(h)}$ in (5.83). We then find another domain $\mathfrak{D}_{\text{int}}^h$ as in (5.79) and another real-analytic transformation

$$\phi_{*,\text{int}}^{n-h} : (t_*^{(h)}, z_*^{(h)}, y_*^{(h)}, x_*^{(h)}) \in \mathfrak{D}_{\text{int}}^h \rightarrow (t^{*(h)}, z^{*(h)}, y^{*(h)}, x^{*(h)}) \in \overline{\mathfrak{D}_{\text{int},h}}$$

such that

$$\overline{f_{\text{sec},h-1}^{n-h}} := \overline{f_{\text{sec},h-1}^{n-h}} \circ \overline{\phi_{*,\text{int}}^{n-h}} = f_{\text{sec},h-1}^{n-h} \circ \overline{\phi_{\text{int}}^{n-h}} \circ \overline{\phi_{*,\text{int}}^{n-h}} = \overline{h_{\text{sec}}^{n-h}}$$

as desired, depends only on $\hat{y}_{*,n-h}^{(h)}$ in (5.71), and hence (5.80) is satisfied. That $\overline{\phi_{*,\text{int}}^{n-h}}$ may be also chosen of a form analogous to (5.82), with $\Theta_{n-h}^{(h-1)}, \vartheta_{n-h}^{(h-1)}, \hat{z}^{(h-1)}, y_{n-h}^{(h-1)}, x_{n-h}^{(h-1)}$ replaced by $p^{*(h)}, q^{*(h)}, \hat{z}^{*(h)}, y^{*(h)}, x^{*(h)}$ also easily follows from the properties above. Therefore the composition

$$\phi_{\text{int}}^{n-h} := \overline{\phi_{\text{int}}^{n-h}} \circ \overline{\phi_{*,\text{int}}^{n-h}}$$

has again the form in (5.82) and satisfies (5.80), as wanted. \square

5.2.3. Construction of $\phi_{\text{norm}}^1, \dots, \phi_{\text{norm}}^{n-2}$ ($n \geq 3$). In this section we aim to determine, for $n \geq 3$ and $1 \leq h \leq n-2$, a transformation ϕ_{norm}^{n-h} solving (5.86)-(5.87), assuming the Proposition 5.2 holds up to rank $(h-1)$ and that ϕ_{int}^{n-h} has been constructed.

We switch from the coordinates $(\chi_*^{(h)}, \kappa_*^{(h)})$ defined implicitly via the right hand side of (5.86) to the auxiliary coordinates

$$\mathbf{G}_{\text{aux}}^{(h)} = (\mathbf{G}_{\text{aux},1}^{(h)}, \dots, \mathbf{G}_{\text{aux},n}^{(h)}), \quad \mathbf{g}_{\text{aux}}^{(h)} = (\mathbf{g}_{\text{aux},1}^{(h)}, \dots, \mathbf{g}_{\text{aux},n}^{(h)})$$

defined via the linear transformation

$$(5.104) \quad \phi_{\text{aux}}^{n-h} : \begin{cases} \chi_{*,i-1}^{(h)} = \mathbf{G}_{\text{aux},i}^{(h)} + \dots + \mathbf{G}_{\text{aux},n}^{(h)} \\ \kappa_{*,i-1}^{(h)} = \mathbf{g}_{\text{aux},i}^{(h)} - \mathbf{g}_{\text{aux},i-1}^{(h)} \end{cases}$$

with $1 \leq i \leq n$ and $\mathbf{g}_{\text{aux},0} := 0$. We regard ϕ_{aux}^{n-h} as a transformation on all the coordinates, extending it as the identity on the remaining ones. We denote the new coordinates as

$$\begin{aligned} \mathbf{t}_{\text{aux}}^{(h)} &:= \begin{cases} (\Theta_{\text{aux},1}^{(h)}, \dots, \Theta_{\text{aux},n-h-1}^{(h)}, \vartheta_{\text{aux},1}^{(h)}, \dots, \vartheta_{\text{aux},n-h-1}^{(h)}) \\ n \geq 4, 2 \leq h \leq n-2 \\ \emptyset \quad \text{otherwise} \end{cases} \\ \mathbf{z}_{\text{aux}}^{(h)} &:= (p_{\text{aux},n-h}^{(h)}, \dots, p_{\text{aux},n-1}^{(h)}, q_{\text{aux},n-h}^{(h)}, \dots, q_{\text{aux},n-1}^{(h)}) \\ \mathbf{y}_{\text{aux}}^{(h)} &:= (\mathbf{G}_{\text{aux},1}^{(h)}, \dots, \mathbf{G}_{\text{aux},n}^{(h)}, \Lambda_{\text{aux},1}^{(h)}, \dots, \Lambda_{\text{aux},n}^{(h)}) \\ \mathbf{x}_{\text{aux}}^{(h)} &:= (\mathbf{g}_{\text{aux},1}^{(h)}, \dots, \mathbf{g}_{\text{aux},n}^{(h)}, \ell_{\text{aux},1}^{(h)}, \dots, \ell_{\text{aux},n}^{(h)}) \end{aligned}$$

the new Hamiltonian as

$$(5.105) \quad f_{\text{sec,int,aux},h-1}(\mathbf{t}_{\text{aux}}^{(h)}, \mathbf{z}_{\text{aux}}^{(h)}, \mathbf{y}_{\text{aux}}^{(h)}, \mathbf{x}_{\text{aux}}^{(h)}) := f_{\text{sec,int},h-1} \circ \phi_{\text{aux}}^{n-h}(\mathbf{t}_{\text{aux}}^{(h)}, \mathbf{z}_{\text{aux}}^{(h)}, \mathbf{y}_{\text{aux}}^{(h)}, \mathbf{x}_{\text{aux}}^{(h)}).$$

Now we define the domain where we want to consider $f_{\text{sec,int,aux},h-1}$. Firstly, we let

$$\mathfrak{D}_{\text{int,aux}}^h := \left\{ (\mathbf{t}_{\text{aux}}^{(h)}, \mathbf{z}_{\text{aux}}^{(h)}, \mathbf{y}_{\text{aux}}^{(h)}, \mathbf{x}_{\text{aux}}^{(h)}) : (\mathbf{t}_*^{(h)}, \mathbf{z}_*^{(h)}, \mathbf{y}_*^{(h)}, \mathbf{x}_*^{(h)}) \in \mathfrak{D}_{\text{int}}^h \right\}$$

where $\mathfrak{D}_{\text{int}}^h$ is defined in (5.79). Then $\mathfrak{D}_{\text{int,aux}}^h$ is given by

$$\mathfrak{D}_{\text{int,aux}}^h = \mathbb{T}_{\hat{\mathbf{c}}_h \theta}^h \times B_{\hat{\mathbf{c}}_h \varepsilon_n}^2 \times B_{\hat{\mathbf{c}}_h \varepsilon}^{*,h-1} \times (\mathbf{G}_*)_{\bar{\mathbf{c}}_h \theta, \bar{\mathbf{c}}_h \bar{\tau}} \times \Lambda_{\bar{\mathbf{c}}_h \bar{\tau}^{(h)}} \times \mathbb{T}_{\bar{\mathbf{c}}_h s}^n \times \mathbb{T}_{\bar{\mathbf{c}}_h s}^n,$$

with

$$\begin{aligned} (\mathbf{G}_*)_{\bar{\mathbf{c}}_h \theta, \bar{\mathbf{c}}_h \bar{\tau}} &:= (\mathbf{G}_1)_{\hat{\mathbf{c}}_1 \theta_1} \times \dots \times (\mathbf{G}_{n-h})_{\hat{\mathbf{c}}_{n-h} \theta_{n-h}} \\ &\quad \times (\mathbf{G}_{n-h+1}^*)_{\hat{\mathbf{c}}_{n-h+1} \bar{\tau}_{n-h+1}} \times \dots \times (\mathbf{G}_{n-1}^*)_{\hat{\mathbf{c}}_{n-1} \bar{\tau}_{n-1}}. \end{aligned}$$

Next, for $1 \leq h' \leq h$ and any fixed $\bar{\gamma}, \bar{K} > 0$ and $\bar{\tau} > 2$, we define

$$(5.106) \quad \omega_{\text{sec}}^{n-h'}(\hat{y}_{\text{aur},n-h'}^{(h)}) := \begin{cases} \partial_{\frac{(p_{\text{aur},n-1}^{(1)})^2 + (q_{\text{aur},n-1}^{(1)})^2}{2}, G_{\text{aur},n-1}^{(1)}, G_{\text{aur},n}^{(1)}}} h_{\text{sec}}^{n-1}(\hat{y}_{\text{aur},n-1}^{(h)}) & n \geq 3, h' = 1, \\ & 2 \leq h \leq n-1 \\ \partial_{\frac{(p_{\text{aur},n-h'}^{(h')})^2 + (q_{\text{aur},n-h'}^{(h')})^2}{2}, G_{\text{aur},n-h'}^{(h')}}} h_{\text{sec}}^{n-h'}(\hat{y}_{\text{aur},n-h'}^{(h)}) & n \geq 3, \\ & 2 \leq h' \leq h \leq n-1, \\ & (h', h) \neq (n-1, n-1) \\ \partial_{\frac{(p_{\text{aur},1}^{(n-1)})^2 + (q_{\text{aur},1}^{(n-1)})^2}{2}} h_{\text{sec}}^{n-h}(\hat{y}_{\text{aur},1}^{(n-1)}) & h' = h = n-1. \end{cases}$$

We then choose the following sub-domain of $\mathfrak{D}_{\text{int,aur}}^h$

$$(5.107) \quad \mathfrak{D}_{\text{sec,aur}}^h := \left\{ (t_{\text{norm,aur}}^{(h)}, z_{\text{norm,aur}}^{(h)}, y_{\text{norm,aur}}^{(h)}, x_{\text{norm,aur}}^{(h)}) \in \mathfrak{D}_{\text{int,aur}}^h : \right. \\ \left. |\omega_{\text{sec}}^{n-h'} \cdot \mathbf{k}| \geq \frac{(a_{n-h'}^+)^2}{(a_{n-h'+1}^-)^3 \theta_{n-h}} \frac{\bar{\gamma}}{\bar{K}^{\bar{\tau}}}, \right. \\ \left. \forall \mathbf{k} \in \mathbb{Z}^j \setminus \{0\}, |\mathbf{k}|_1 \leq \bar{K}, \quad \forall 2 \leq h' \leq h \right\}.$$

Here j is chosen to be 3, 2 or 1 accordingly to the three cases above. The set $\mathfrak{D}_{\text{int,aur}}^h$ is non-empty, if $\bar{\gamma}$ is chosen suitably small. Indeed, if we put

$$\hat{y}_{\text{aur},n-h}^{(h)} := \left(\frac{(p_{\text{aur},n-h}^{(h)})^2 + (q_{\text{aur},n-h}^{(h)})^2}{2}, G_{\text{aur},n-h}^{(h)}, \Lambda_{\text{aur},n-h}^{(h)}, \hat{y}_{\text{aur},n-h+1}^{(h)} \right)$$

then standard quantitative arguments show that, for any fixed value

$$(\bar{\Lambda}_{\text{aur},n-h}^{(h)}, \hat{y}_{\text{aur},n-h+1}^{(h)}) \in \Pi_{\Lambda_{\text{aur},n-h}^{(h)}, \hat{y}_{\text{aur},n-h+1}^{(h)}} \mathfrak{D}_{\text{int,aur}}^h,$$

the measure of the set $\mathcal{N}_{n-h} \subset B_{\mathfrak{c}_h \varepsilon_{n-h}}^2 \times G_{n-h}$ of $(p_{\text{aur},n-h}^{(h)}, q_{\text{aur},n-h}^{(h)}, G_{\text{aur},n-h}^{(h)})$ where the inequality in (5.107) does not hold may be bounded as

$$\text{meas } \mathcal{N}_{n-h} \leq \frac{\bar{\gamma}}{\mathfrak{c}} \text{meas } (B_{\mathfrak{c}_h \varepsilon_{n-h}}^2 \times G_{n-h}),$$

(where \mathfrak{c} depends only on the semi-axes ratio and the masses), hence (5.68) follows. This is because $\omega_{\text{sec}}^{n-h'}(\hat{y}_{\text{aur},n-h'}^{(h)})$ is a diffeomorphism (Compare Appendix C).

Now we inspect the form of $f_{\text{sec,int,aur},h-1}$ in (5.105). Introducing the following symbols

$$\begin{aligned}
t_{\text{aur},i}^{(h)} &:= \begin{cases} (\Theta_{\text{aur},i}^{(h)}, \dots, \Theta_{\text{aur},n-h-1}^{(h)}, \vartheta_{\text{aur},i}^{(h)}, \dots, \vartheta_{\text{aur},n-h-1}^{(h)}) \\ n \geq 4, 2 \leq h \leq n-2, 1 \leq i \leq n-h-1 \\ \emptyset \quad \text{otherwise} \end{cases} \\
y_{\text{aur},i}^{(h)} &:= (G_{\text{aur},i}^{(h)}, \dots, G_{\text{aur},n}^{(h)}, \Lambda_{\text{aur},1}^{(h)}, \dots, \Lambda_{\text{aur},n}^{(h)}) \\
x_{\text{aur},i}^{(h)} &:= (g_{\text{aur},i}^{(h)}, \dots, g_{\text{aur},n}^{(h)}, \ell_{\text{aur},1}^{(h)}, \dots, \ell_{\text{aur},n}^{(h)}) \\
\hat{y}_{\text{aur},i}^{(h)} &:= \left(\frac{(p_{\text{aur},i}^{(h)})^2 + (q_{\text{aur},i}^{(h)})^2}{2}, \dots, \frac{(p_{\text{aur},n-1}^{(h)})^2 + (q_{\text{aur},n-1}^{(h)})^2}{2}, G_{\text{aur},i}^{(h)}, \dots, \right. \\
&\quad \left. G_{\text{aur},n}^{(h)}, \Lambda_{\text{aur},i}^{(h)}, \dots, \Lambda_{\text{aur},n}^{(h)} \right) \\
\hat{x}_{\text{aur},i}^{(h)} &:= \begin{cases} (g_{\text{aur},i+1}^{(h)} - g_{\text{aur},i}^{(h)}, \dots, g_{\text{aur},n-h}^{(h)} - g_{\text{aur},n-h-1}^{(h)}) \\ n \geq 4 \ \& \ 1 \leq h-1 \leq n-3 \\ \emptyset \quad \text{otherwise} \end{cases} \\
\hat{X}_{\text{aur},i}^{(h)} &:= \begin{cases} (G_{\text{aur},i}^{(h)}, \dots, G_{\text{aur},n-h}^{(h)}) & n \geq 4 \ \& \ 1 \leq h-1 \leq n-3 \\ \emptyset \quad \text{otherwise} \end{cases} \\
z_{\text{aur},n-h}^{(h)} &:= (p_{\text{aur},n-h}^{(h)}, q_{\text{aur},n-h}^{(h)}), \quad \hat{z}_{\text{norm},j}^{(h)} := p_{\text{norm},j}^{(h)} + iq_{\text{norm},j}^{(h)} \\
\tilde{y}_{\text{aur},i}^{(h)} &:= \left(\frac{(p_{\text{aur},n-h+1}^{(h)})^2 + (q_{\text{aur},n-h+1}^{(h)})^2}{2}, \dots, \frac{(p_{\text{aur},n-1}^{(h)})^2 + (q_{\text{aur},n-1}^{(h)})^2}{2}, \right. \\
&\quad \left. G_{\text{aur},i}^{(h)}, \dots, G_{\text{aur},n}^{(h)}, \Lambda_{\text{aur},i}^{(h)}, \dots, \Lambda_{\text{aur},n}^{(h)} \right) \\
\tilde{y}_{\text{aur}}^{(h)} &:= \tilde{y}_{\text{aur},1}^{(h)}, \quad \hat{x}_{\text{aur}}^{(h)} := \hat{x}_{\text{aur},1}^{(h)}, \quad \hat{X}_{\text{aur}}^{(h)} := \hat{X}_{\text{aur},1}^{(h)},
\end{aligned}$$

by means of (5.85), we have

$$\begin{aligned}
&f_{\text{sec,int,aur},h-1}(t_{\text{aur}}^{(h)}, z_{\text{aur}}^{(h)}, y_{\text{aur}}^{(h)}, x_{\text{aur}}^{(h)}) \\
&= h_{\text{sec},h}(\hat{y}^{(h)}) + f_{\text{norm,int,aur},h-1}(t_{\text{aur}}^{(h)}, z_{\text{aur},n-h}^{(h)}, \tilde{y}_{\text{aur}}^{(h)}, \hat{x}_{\text{aur}}^{(h)}) \\
(5.108) \quad &= \sum_{i=n-h}^{n-1} h_{\text{sec}}^i(\hat{y}_i^{(h)}) \\
&\quad + \sum_{i=1}^{n-h-1} f_{\text{norm,int,aur},h-1}^i(t_{\text{aur},i}^{(h)}, z_{\text{aur},n-h}^{(h)}, \tilde{y}_{\text{aur},i}^{(h)}, \hat{x}_{\text{aur},i}^{(h)})
\end{aligned}$$

where we have let

$$(5.109) \quad f_{\text{norm,int,aur},h-1} := f_{\text{norm,int},h-1} \circ \phi_{\text{aur}}^{n-h}, \quad f_{\text{norm,int,aur},h-1}^i := f_{\text{norm,int},h-1}^i \circ \phi_{\text{aur}}^{n-h}.$$

On the domain $\mathfrak{D}_{\text{sec,aur}}^h$ specified in (5.107), we aim to construct and real-analytic and canonical transformation

$$\begin{aligned}
(5.110) \quad \phi_{\text{norm,aur}}^{n-h} &: (t_{\text{norm,aur}}^{(h)}, z_{\text{norm,aur}}^{(h)}, y_{\text{norm,aur}}^{(h)}, x_{\text{norm,aur}}^{(h)}) \\
&\quad \in \mathfrak{D}_{\text{sec,aur}}^h \rightarrow (t_{\text{aur}}^{(h)}, z_{\text{aur}}^{(h)}, y_{\text{aur}}^{(h)}, x_{\text{aur}}^{(h)}) \in \mathfrak{D}_{\text{int,aur}}^h
\end{aligned}$$

such that the transformed Hamiltonian

$$f_{\text{sec,aur},h} := f_{\text{sec,int,aur},h-1} \circ \phi_{\text{norm,aur}}^{n-h}$$

has the form

$$\begin{aligned} f_{\text{sec,aur},h} &= h_{\text{sec},h}(\hat{y}_{\text{norm,aur}}^{(h)}) + f_{\text{norm,aur},h}(t_{\text{norm,aur}}^{(h)}, \hat{y}_{\text{norm,aur}}^{(h)}, \hat{x}_{\text{norm,aur}}^{(h)}) \\ &= \sum_{i=n-h}^{n-1} h_{\text{sec}}^i(\hat{y}_{\text{norm,aur},i}^{(h)}) \\ &+ \sum_{i=1}^{n-h-1} f_{\text{norm,aur},h}^i(t_{\text{norm,aur},i}^{(h)}, \hat{y}_{\text{norm,aur},i}^{(h)}, \hat{x}_{\text{norm,aur},i}^{(h)}) \\ &+ f_{\text{exp,sec,aur},h}(t_{\text{norm,aur}}^{(h)}, z_{\text{norm,aur}}^{(h)}, y_{\text{norm,aur}}^{(h)}, x_{\text{norm,aur}}^{(h)}) \end{aligned}$$

where

$$\hat{x}_{\text{norm,aur},i}^{(h)} := \begin{cases} (g_{\text{norm,aur},i+1}^{(h)} - g_{\text{norm,aur},i}^{(h)}, \dots, g_{\text{norm,aur},n-h-1}^{(h)} \\ -g_{\text{norm,aur},n-h-2}^{(h)}) & \text{if } n \geq 4 \text{ \& } 1 \leq h-1 \leq n-3 \\ \emptyset & \text{otherwise} \end{cases}$$

$$\hat{y}_{\text{norm,aur},i}^{(h)} := \left(\frac{(p_{\text{norm,aur},n-h}^{(h)})^2 + (q_{\text{norm,aur},n-h}^{(h)})^2}{2}, \dots, \right.$$

$$\left. \frac{(p_{\text{norm,aur},n-1}^{(h)})^2 + (q_{\text{norm,aur},n-1}^{(h)})^2}{2}, G_{\text{norm,aur},i}^{(h)}, \dots, \right.$$

$$\left. G_{\text{norm,aur},n}^{(h)}, \Lambda_{\text{norm,aur},i}^{(h)}, \dots, \Lambda_{\text{norm,aur},n}^{(h)} \right)$$

and $f_{\text{exp,sec,aur},h}$ satisfies the bound for $f_{\text{exp,sec},h}$ in (5.76). This will conclude the proof, up to apply the inverse transformation of (5.104), with $G_{\text{aur},i}^{(h)}$, $g_{\text{aur},i}^{(h)}$, $\chi_{*,i}^{(h)}$, $\kappa_{*,i}^{(h)}$ replaced by $G_{\text{norm,aur},i}^{(h)}$, $g_{\text{norm,aur},i}^{(h)}$, $\chi_i^{(h)}$, $\kappa_i^{(h)}$, and to take

$$\mathfrak{D}_{\text{sec}}^h := \phi_{\text{aur}}^{n-h}(\mathfrak{D}_{\text{sec,aur}}^h).$$

We shall obtain the transformation $\phi_{\text{norm,aur}}^{n-h}$ in (5.110) via an application of Proposition D.1. Before doing it, we just remark that, since, in our particular case, $f_{\text{norm,int,aur},h-1}$ depends on $z_{\text{aur}}^{(h)}$, $y_{\text{aur}}^{(h)}$, $x_{\text{aur}}^{(h)}$ only via $z_{\text{aur},n-h}^{(h)}$, $\tilde{y}_{\text{aur}}^{(h)}$, $\hat{x}_{\text{aur}}^{(h)}$ and is even in $(t_{\text{aur}}^{(h)}, z_{\text{aur},n-h}^{(h)})$, the proof of Proposition D.1 can be easily handled to show that

$\phi_{\text{norm},\text{au}\mathbb{F}}^{n-h}$ can be chosen of the form

$$\phi_{\text{norm},\text{au}\mathbb{F}}^{n-h} : \left\{ \begin{array}{l} \Theta_{\text{au}\mathbb{F},j}^{(h)} = \mathbb{F}_{\text{norm},\text{au}\mathbb{F},j}^{(h)}(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}, \tilde{\mathbb{Y}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}) \\ \vartheta_{\text{au}\mathbb{F},j}^{(h)} - \pi = \mathbb{G}_{\text{norm},\text{au}\mathbb{F},j}^{(h)}(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}, \tilde{\mathbb{Y}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}) \\ \quad j = 1, \dots, n-h-1 \\ \mathbb{Z}_{\text{au}\mathbb{F},n-h}^{(h)} = \mathbb{Z}_{\text{norm},\text{au}\mathbb{F}}^{(h)}(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}, \tilde{\mathbb{Y}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}) \\ (\hat{\mathbb{X}}_{\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{au}\mathbb{F}}^{(h)}) = \mathbb{X}_{\text{norm},\text{au}\mathbb{F}}^{(h)}(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}, \tilde{\mathbb{Y}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}) \\ \hat{\mathbb{Z}}_{\text{norm},j}^{(h)} = \hat{\mathbb{Z}}_{\text{norm},\text{au}\mathbb{F},j}^{(h)} e^{i\psi_{\text{norm},\text{au}\mathbb{F},j}^{(h)}}(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}, \tilde{\mathbb{Y}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}) \\ \quad j = n-h+1, \dots, n-1 \\ \mathbb{Y}_{\text{au}\mathbb{F},n-h+1}^{(h)} = \mathbb{Y}_{\text{norm},\text{au}\mathbb{F},n-h+1}^{(h)} \\ \mathbb{X}_{\text{au}\mathbb{F},n-h+1}^{(h)} = \mathbb{X}_{\text{norm},\text{au}\mathbb{F},n-h+1}^{(h)} \\ \quad + \varphi_{\text{norm},\text{au}\mathbb{F}}^{(h)}(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}, \tilde{\mathbb{Y}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \hat{\mathbb{X}}_{\text{norm},\text{au}\mathbb{F}}^{(h)}) \end{array} \right.$$

where $\mathbb{F}_{\text{norm},\text{au}\mathbb{F}}^{(h)}$, $\mathbb{G}_{\text{norm},\text{au}\mathbb{F}}^{(h)}$ and $\mathbb{Z}_{\text{norm},\text{au}\mathbb{F}}^{(h)}$ are odd; $\mathbb{X}_{\text{norm},\text{au}\mathbb{F}}^{(h)}$, $\psi_{\text{norm},\text{au}\mathbb{F},j}^{(h)}$ and $\varphi_{\text{norm},\text{au}\mathbb{F}}^{(h)}$ are even under the change

$$(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}) \rightarrow -(\mathfrak{t}_{\text{norm},\text{au}\mathbb{F}}^{(h)}, \mathbb{Z}_{\text{norm},\text{au}\mathbb{F},n-h}^{(h)}).$$

Then (5.88)–(5.90) follow.

Now we proceed with proving the existence of $\phi_{\text{norm},\text{au}\mathbb{F}}^{n-h}$. We can choose, in (D.133), (D.135) and (D.136),

$$(5.111) \quad \begin{aligned} \nu_i &= 2(h+1), \quad \ell_i = h, \quad m_i = 3i, \quad i = 1, \dots, n-h-1 = N \\ \mathfrak{h}(p, q, I) &= \sum_{i=n-h}^{n-1} \mathfrak{h}_{\text{src}}^i(\hat{\mathbb{Y}}_i^{(h)}), \quad \mathfrak{f}(p, q, I, \varphi, \eta, \xi) = \sum_{i=1}^{n-h-1} \mathfrak{f}^i(u_i, p, q, \varphi) \\ \mathfrak{f}^i(u_i, p, q, \varphi) &:= \mathfrak{f}_{\text{norm},\text{int},\text{au}\mathbb{F},h-1}^{n-h-i}(\mathfrak{t}_{\text{au}\mathbb{F},n-h-i}^{(h)}, \tilde{\mathbb{Y}}_{\text{au}\mathbb{F},n-h-i}^{(h)}, \hat{\mathbb{X}}_{\text{au}\mathbb{F},n-h-i}^{(h)}) \\ \mathfrak{Z} &:= \mathfrak{Z}_i := \left\{ (k', k'', k''') \in \mathbb{Z}^h \times \mathbb{Z}^{h+1} \times \mathbb{Z}^{h+1} : k'_{n-h+1} = \dots = k'_{n-1} = 0 \right. \\ &\quad \left. k''_{n-h+1} = \dots = k''_n = 0, k'''_{n-h} = \dots = k'''_n = 0, k'_1 + \dots + k''_{n-h} = 0 \right\} \\ \mathfrak{L} &:= \left\{ (k', k'', k''') \in \mathfrak{Z} : k'_{n-h} = k''_{n-h} = 0 \right\} \end{aligned}$$

where we have re-named

$$\begin{aligned}
(p, q) &:= (p_{\text{aur}, n-h}^{(h)}, q_{\text{aur}}^{(h)}) = (p_{\text{aur}, n-h}^{(h)}, \dots, p_{\text{aur}, n-1}^{(h)}, q_{\text{aur}, n-h}^{(h)}, \dots, q_{\text{aur}, n-1}^{(h)}) \\
I &:= (G_{\text{aur}, n-h}^{(h)}, \dots, G_{\text{aur}, n}^{(h)}, \Lambda_{\text{aur}, n-h}^{(h)}, \dots, \Lambda_{\text{aur}, n}^{(h)}) \\
\varphi &:= (g_{\text{aur}, n-h}^{(h)}, \dots, g_{\text{aur}, n}^{(h)}, \ell_{\text{aur}, n-h}^{(h)}, \dots, \ell_{\text{aur}, n}^{(h)}) \\
u_i &:= (I, \eta^i, \xi^i), \quad \eta := \eta^1, \quad \xi := \xi^1
\end{aligned}$$

with

$$\begin{aligned}
\eta^i &:= (\Theta_{\text{aur}, n-h-i}, \dots, \Theta_{\text{aur}, n-1}, G_{\text{aur}, n-h-i}, \dots, G_{\text{aur}, n-h-1}, \\
&\quad \Lambda_{\text{aur}, n-h-i}, \dots, \Lambda_{\text{aur}, n-h-1}) \\
\xi^i &:= (\vartheta_{\text{aur}, n-h-i}, \dots, \vartheta_{\text{aur}, n-1}, g_{\text{aur}, n-h-i}, \dots, g_{\text{aur}, n-h-1}, \\
&\quad \ell_{\text{aur}, n-h-i}, \dots, \ell_{\text{aur}, n-h-1}).
\end{aligned}$$

In order to verify that Proposition D.1 can be applied, we have to check conditions (D.134) and (D.139). Due to the choices of \mathfrak{Z} , \mathfrak{L} and to the fact that only the function h_{sec}^{n-h} in the summand for h_{sec} in (5.108) depends on

$$(p_{\text{aur}, n-h}^{(h)}, q_{\text{aur}, n-h}^{(h)}, G_{\text{aur}, n-h}^{(h)}),$$

it is sufficient to check that condition (D.134) holds with

$$\omega = \omega_{\text{sec}}^{n-h}, \quad (k', k) \in \mathbb{Z}^2 \setminus \{0\}, \quad K = \bar{K}.$$

But due to the choice of $\mathfrak{D}_{\text{int}, \text{aur}}^h$ in (5.107), we have that (D.134) is verified, with

$$\mathfrak{a} = \frac{(a_{n-h}^+)^2}{(a_{n-h+1}^-)^3} \frac{\bar{\gamma}}{\bar{K}^{\bar{\tau}}}, \quad r = \mathfrak{c}_h \frac{\theta_{n-h} \bar{\gamma}}{\bar{K}^{\bar{\tau}+1}}, \quad \varepsilon = \mathfrak{c}_h \sqrt{\theta_{n-h}}.$$

It remains to check the inequalities in (D.139). In view of the definition of f^i following from the formulae (5.84), (5.109) and (5.111), of the definition of $f_{\text{norm}, h-1}^i$ in (5.74), the definition of $\widetilde{f_{\text{norm}, h-1}^i}$, the bound for $\widetilde{f_{\text{norm}, h-1}^i}$ in (5.76), and first inequality in (4.54), we see that the former of the inequalities in (D.139) is satisfied with

(5.112)

$$E_i = \frac{1}{\mathfrak{c}_h} \max \left\{ \frac{(a_{n-h-i}^+)^2}{(a_{n-h-i+1}^+)^3}, \mu \bar{K} \left(\frac{a_n^+}{a_1^+} \right)^{\frac{3}{2}} \frac{1}{a_{n-h-i+1}^-} \right\} \quad i = 1, \dots, n-h-1.$$

In order to check that also the second inequality in (D.139) is satisfied, we previously note that the number d_i in (D.138) can be taken to be

$$d_i = \mathfrak{c}_h \min \left\{ \frac{\theta_{n-h} \bar{\gamma}}{\bar{K}^{\bar{\tau}+1}}, \theta_{n-h-i} \right\}, \quad i = 1, \dots, n-h-1.$$

Inserting then the above values for K , \mathfrak{a} , E_i and d_i into the left hand side of the second inequality in (D.139), we find that this can be bounded by

$$\frac{1}{\hat{c}_h} \max \left\{ \frac{\bar{K}^{2\bar{\tau}+2} (a_{n-h-i}^+)^2 (a_{n-h+1}^-)^3}{\bar{\gamma}^2 (a_{n-h}^+)^2 (a_{n-h-i+1}^-)^3}, \frac{\bar{K}^{\bar{\tau}+1} (a_{n-h-i}^+)^2 (a_{n-h+1}^-)^3}{\bar{\gamma} (a_{n-h}^+)^2 (a_{n-h-i+1}^-)^3} \frac{\theta_{n-h}}{\theta_{n-h-i}} \right. \\ \left. \frac{\bar{K}^{2\bar{\tau}+2} \mu \bar{K} \left(\frac{a_n^+}{a_1}\right)^{\frac{3}{2}} (a_{n-h+1}^-)^3}{\bar{\gamma}^2 (a_{n-h}^+)^2 a_{n-h-i+1}^-}, \frac{\bar{K}^{\bar{\tau}+1} \mu \bar{K} \left(\frac{a_n^+}{a_1}\right)^{\frac{3}{2}} (a_{n-h+1}^-)^3}{\bar{\gamma} (a_{n-h}^+)^2 a_{n-h-i+1}^-} \frac{\theta_{n-h}}{\theta_{n-h-i}} \right\}$$

Using (4.50), one easily finds that this quantity does not exceed

$$(5.113) \quad \frac{1}{\hat{c}_h} \max \left\{ \mu \left(\frac{a_n}{a_1}\right)^5 \frac{\bar{K}^{2\bar{\tau}+2}}{\bar{\gamma}^2}, \frac{\bar{K}^{\bar{\tau}+1} \sqrt{\alpha}}{\bar{\gamma}} \right\} < 1.$$

where \hat{c}_h depends only on the ratio a_n^-/a_n^+ and the masses and the inequality follows from (4.54). This concludes the proof of this case. \square

5.2.4. Construction of ϕ_{norm}^{n-1} . The arguments we have used in the previous section to construct $\phi_{\text{norm}}^1, \dots, \phi_{\text{norm}}^{n-2}$ also fit for the case of ϕ_{norm}^{n-1} , therefore we shall not repeat them. We only limit to remark that, for this case, Equations (5.106), (5.111), (5.112) and (5.113) have to be replaced with

$$\omega_{\text{sec}}^{n-1}(\hat{y}_{\text{aur},n-1}^{(1)}) := \begin{cases} \frac{\partial_{(p_{\text{aur},n-1}^{(1)})^2 + (q_{\text{aur},n-1}^{(1)})^2}, G_{\text{aur},n-1}^{(1)}, G_{\text{aur},n}^{(1)}}{2} h_{\text{sec}}^{n-1}(\hat{y}_{\text{aur},n-1}^{(1)}) & n \geq 3 \\ \partial_{G_{\text{aur},2}^{(1)}} h_{\text{sec}}^2(\hat{y}_{\text{aur},1}^{(1)}) & n = 2, \end{cases}$$

$$f^i = f_{\text{norm,int,aur},0}^{n-i}(t_{\text{aur},i}^{(1)}, \hat{y}_{\text{aur},i}^{(1)}, \hat{x}_{\text{aur},i}^{(1)}), \quad d_i = c_1 \min \left\{ \frac{\theta_{n-1} \bar{\gamma}}{\bar{K}^{\bar{\tau}+1}}, \theta_{n-i-1} \right\}$$

$$i = 1, \dots, n-1, \quad \theta_0 := \theta_1$$

$$E_i = \frac{1}{\hat{c}_1} \begin{cases} \max \left\{ \mu \left(\frac{a_n^+}{a_1}\right)^{\frac{3}{2}} \frac{1}{a_n^-}, \frac{(a_{n-1}^+)^3}{(a_n^-)^4} \right\} & i = 1 \\ \max \left\{ \mu \bar{K} \left(\frac{a_n^+}{a_1}\right)^{\frac{3}{2}} \frac{1}{a_{n-i+1}^-}, \frac{(a_{n-i}^+)^2}{(a_{n-i+1}^-)^3} \right\} & n \geq 3, i = 2, \dots, n-1 \end{cases}$$

$$\frac{1}{\hat{c}_1} \max \left\{ \mu \left(\frac{a_n}{a_1}\right)^5 \frac{\bar{K}^{2\bar{\tau}+2}}{\bar{\gamma}^2}, \frac{\bar{K}^{2(\bar{\tau}+1)} \alpha}{\bar{\gamma}^2} \right\}.$$

\square

APPENDIX A

Computing the domain of holomorphy

A.1. On the analyticity of the solution of Kepler equation

Here is a refinement of Proposition 4.1.

PROPOSITION A.1. *Let \hat{e} be as in (4.47). For any $0 < \bar{e} < \hat{e}$ there exists $\bar{\eta} = \bar{\eta}(\bar{e})$ such that, for any $\bar{\eta} < \eta < 1$ and any $e \in \mathbb{C}$ with $|e| \leq \bar{e}$, there exist two positive numbers $\bar{\zeta} = \bar{\zeta}(\eta, e)$, $\bar{\ell} = \bar{\ell}(\eta, \bar{e})$ such that the map*

$$(A.114) \quad \zeta \in \bar{\mathbb{T}}_{\bar{\zeta}} \rightarrow K(\zeta, e) := \zeta - e \sin \zeta$$

is injective, its image verifies

$$K(\bar{\mathbb{T}}_{\bar{\zeta}}, e) \supset \bar{\mathbb{T}}_{\bar{\ell}} \quad \forall e \in \mathbb{C} : |e| \leq \bar{e}.$$

The inverse function

$$\ell \in \bar{\mathbb{T}}_{\bar{\ell}} \rightarrow \zeta(\ell, e) := K^{-1}(\ell, e) \in \bar{\mathbb{T}}_{\bar{\zeta}_\eta(e)}$$

verifies

$$(A.115) \quad |1 - e \cos \zeta(\ell, e)| \geq 1 - \eta$$

Therefore, $\zeta(\ell, e)$ is real-analytic for $\ell \in \bar{\mathbb{T}}_{\bar{\ell}}$.

The proof of Proposition A.1 is elementary and goes along the same lines of [24]. Therefore, we shall present it skipping some detail.

LEMMA A.1. *Let \hat{e} be as in Proposition 4.1. For any $0 < \bar{e} < \hat{e}$ there exists a unique $\bar{\eta} = \bar{\eta}(\bar{e}) \in (\bar{e}, 1)$ such that*

$$\begin{aligned} \forall \eta \in [\bar{\eta}, 1) : \quad \bar{\ell}_\eta(\bar{e}) &:= \log \left[\frac{\eta}{\bar{e}} + \sqrt{1 + \frac{\eta^2}{\bar{e}^2}} \right] - \sqrt{\eta^2 + \bar{e}^2} \geq 0, \\ \bar{\ell}_\eta(\bar{e}) = 0 &\iff \eta = \bar{\eta}. \end{aligned}$$

PROOF. By definition of \hat{e} , and since the function $\rho \in [0, 1] \rightarrow \frac{\rho e^{\sqrt{1+\rho^2}}}{1+\sqrt{1+\rho^2}}$ increases with ρ , we have

$$\frac{\bar{e} e^{\sqrt{1+\bar{e}^2}}}{1 + \sqrt{1 + \bar{e}^2}} < 1.$$

Consider now the function

$$\eta \in (0, 1] \rightarrow g_\rho(\eta) := \frac{\rho e^{\sqrt{\eta^2 + \rho^2}}}{\eta + \sqrt{\eta^2 + \rho^2}}.$$

This function decreases with η for any $\rho \in (0, 1]$. Since

$$g_{\bar{e}}(0) = e^{\bar{e}} > 1, \quad g_{\bar{e}}(1) = \frac{\bar{e} e^{\sqrt{1+\bar{e}^2}}}{1 + \sqrt{1 + \bar{e}^2}} < 1$$

we find a unique $\bar{\eta} = \bar{\eta}(\bar{e}) \in [0, 1]$ such that

$$g_{\bar{e}}(\eta) < 1 \quad \forall \bar{\eta} < \eta < 1, \quad g_{\bar{e}}(\eta(\bar{e})) = 1.$$

Since also

$$g_{\bar{e}}(\bar{e}) = \frac{e^{\bar{e}\sqrt{2}}}{1 + \sqrt{2}} \geq \frac{e^{\sqrt{2}}}{1 + \sqrt{2}} > 1$$

we actually have

$$\bar{e} < \bar{\eta} < 1.$$

□

PROOF OF PROPOSITION A.1. We shall prove Proposition A.1 with

$$(A.116) \quad \begin{aligned} \bar{\zeta}(\eta, \mathbf{e}) &:= \log \frac{\sqrt{\eta^2 + \mathbf{e}_2^2} + \sqrt{\eta^2 - \mathbf{e}_1^2}}{\sqrt{\mathbf{e}_1^2 + \mathbf{e}_2^2}} \\ \bar{\ell}(\eta, \bar{e}) &:= \log \left[\frac{\eta}{\bar{e}} + \sqrt{1 + \frac{\eta^2}{\bar{e}^2}} \right] - \sqrt{\eta^2 + \bar{e}^2} \end{aligned}$$

where $\mathbf{e} = \mathbf{e}_1 + i\mathbf{e}_2$. Observe that $\bar{\ell}(\eta, \bar{e}) > 0$ by Lemma A.1. Moreover, since

$$\mathbf{e}_1 \leq |\mathbf{e}| \leq \bar{e} < \bar{\eta} < \eta$$

we have that $\bar{\zeta}(\eta, \mathbf{e})$ is well defined and positive¹:

$$\bar{\zeta}(\eta, \mathbf{e}) \geq \log \frac{\eta}{\bar{e}} > 0.$$

We split Equation (A.114) into its real and imaginary part

$$\begin{cases} K_1(\zeta_1, \zeta_2, \mathbf{e}_1, \mathbf{e}_2) := \zeta_1 - (\mathbf{e}_1 \sin \zeta_1 \cosh \zeta_2 - \mathbf{e}_2 \cos \zeta_1 \sinh \zeta_2) = \ell_1 \\ K_2(\zeta_1, \zeta_2, \mathbf{e}_1, \mathbf{e}_2) := \zeta_2 - (\mathbf{e}_1 \cos \zeta_1 \sinh \zeta_2 + \mathbf{e}_2 \sin \zeta_1 \cosh \zeta_2) = \ell_2 \end{cases}$$

(with $\zeta = \zeta_1 + i\zeta_2$, $\ell = \ell_1 + i\ell_2$). The equation for the real part gives a unique solution

$$\zeta_1 = Z_1(\mathbf{e}_1, \mathbf{e}_2, \zeta_2, \ell_1)$$

provided

$$(A.117) \quad |\mathbf{e}_1| \leq \eta, \quad |\zeta_2| \leq \bar{\zeta}(\eta, \mathbf{e})$$

since it reduces to an ordinary real Kepler equation

$$\begin{aligned} \zeta_1 - E_1(\mathbf{e}_1, \mathbf{e}_2, \zeta_2) \sin(\zeta_1 - \phi_1(\mathbf{e}_1, \mathbf{e}_2, \zeta_2)) &= \ell_1 & \text{if } E_1(\mathbf{e}_1, \mathbf{e}_2, \zeta_2) \neq 0 \\ \zeta_1 &= \ell_1 & \text{otherwise} \end{aligned}$$

with

$$\begin{aligned} E_1(\mathbf{e}_1, \mathbf{e}_2, \zeta_2) &:= \sqrt{\mathbf{e}_1^2 \cosh^2 \zeta_2 + \mathbf{e}_2^2 \sinh^2 \zeta_2} \\ \phi_1(\mathbf{e}_1, \mathbf{e}_2, \zeta_2) &: \quad E_1 \cos \phi_1 = \mathbf{e}_1 \cosh \zeta_2, \quad E_1 \sin \phi_1 = \mathbf{e}_2 \sinh \zeta_2. \end{aligned}$$

and, under condition (A.117), one has

$$(A.118) \quad E_1 \leq \eta < 1.$$

¹Actually, $\bar{\zeta}(\eta, \mathbf{e})$, as a function of $(\mathbf{e}_1, \mathbf{e}_2)$, reaches its positive minimum

$$\bar{\zeta}_{\min} = \log \left[\frac{\eta}{\bar{e}} + \sqrt{1 + \frac{\eta^2}{\bar{e}^2}} \right] > \log(1 + \sqrt{2})$$

for $(\mathbf{e}_1, \mathbf{e}_2) = (0, \bar{e})$.

Observe that this solution $Z_1(e_1, e_2, \zeta_2, \ell_1)$ verifies

$$(A.119) \quad Z_1(e_1, e_2, -\zeta_2, \ell_1) = -Z_1(e_1, e_2, \zeta_2, \ell_1) \pmod{2\pi}.$$

On the other hand, the function

$$\zeta_2 \rightarrow K_2(e_1, e_2, \zeta_2, \ell_1) := K_2(Z_1(e_1, e_2, \zeta_2, \ell_1), \zeta_2, e_1, e_2)$$

is strictly increasing, therefore, it maps the interval $[-\bar{\zeta}(\eta, e), \bar{\zeta}(\eta, e)]$, onto the interval $[-L_2(\eta, e, \ell_1), L_2(\eta, e, \ell_1)]$, where $L_2(\eta, e, \ell_1) := K_2(e_1, e_2, \bar{\zeta}(\eta, e), \ell_1)$ (note that $K_2(e_1, e_2, -\bar{\zeta}(\eta, e), \ell_1) = -K_2(e_1, e_2, \bar{\zeta}(\eta, e), \ell_1)$ because of (A.119)). We have thus proved that the map (A.114) maps bijectively the strip $\overline{\mathbb{T}}_{\bar{\zeta}(\eta, e)}$ onto the set

$$\ell = \ell_1 + i\ell_2 \in \mathbb{C} : \quad \ell_1 \in \mathbb{T}, \quad \ell_2 \in [-L_2(\eta, e, \ell_1), L_2(\eta, e, \ell_1)].$$

But the curve

$$\ell_2 = L_2(\eta, e, \ell_1) \quad \ell_1 \in [0, 2\pi)$$

is concave, its minimum points are cusps, where L_2 attains the value

$$L_{2,\min}(\eta, e) = \bar{\zeta}(\eta, e) - \sqrt{\eta^2 - e_1^2 + e_2^2}.$$

The minimum of this quantity while $|e| \leq \bar{e}$ is just $\bar{\ell}(\eta, \bar{e})$ in (A.116). Inequality in (A.115) follows from

$$|1 - e \cos \zeta| \geq |\operatorname{Re}(1 - e \cos \zeta)| \geq 1 - |\operatorname{Re}(e \cos \zeta)|$$

and (by (A.118))

$$|\operatorname{Re}(e \cos \zeta)| = |E_1(e_1, e_2, \zeta_2) \cos(\zeta_1 - \phi_1(e_1, e_2, \zeta_2))| \leq E_1 \leq \eta.$$

□

A.2. Proof of Proposition 4.2

Define

$$\underline{\delta}_j := \sqrt{1 - \bar{e}_j^2}, \quad \bar{\delta}_j := \sqrt{1 - e_j^2}.$$

Assume (4.48), with

$$A := (1 - \sigma^2) \sqrt{\frac{1}{(1 + \sigma)^3(1 + \sigma^2)^4}}, \quad B := \sqrt{\frac{1}{(1 - \sigma^2)(1 + \sigma)^3(1 + \sigma^2)}}$$

$$\bar{C}_i := \begin{cases} C_1(\sigma) \bar{\delta}_i & i = 1, \dots, n-1 \\ \bar{\delta}_n & i = n \end{cases},$$

$$\underline{C}_i := \begin{cases} C_2(\sigma) \sqrt{\underline{\delta}_i^2 + 2g(\sigma)^2 \bar{\delta}_i^2} & i = 1, \dots, n-1 \\ \sqrt{\underline{\delta}_i^2 + 2g(\sigma)^2 \bar{\delta}_n^2} & i = n \end{cases}$$

$$s = \sigma(1 - \sigma)$$

where

$$(A.120) \quad C_1(\sigma) := \sqrt{1 - \sigma^2}, \quad C_2(\sigma) := \sqrt{\frac{(1 + \sigma^2)^3}{(1 - \sigma^2)^2}}$$

and σ, g are chosen as follows: $g(\sigma')$ is a suitable positive function, depending at most on the ratios $\frac{\Lambda_j^+}{\Lambda_j^-}, \frac{G_i^+}{G_i^-}$, such² that

(A.121)

$$g(\sigma') \rightarrow 0 \quad \text{as } \sigma' \rightarrow 0, \quad \text{and} \quad \left| \sin \arg \frac{\|C_{\mathcal{P}}^{(j)}\|^2}{\Lambda_j^2} \right| \leq g(\sigma'), \quad j = 1, \dots, n,$$

provided

$$\max \left\{ |\arg(\Lambda_i)|, |\arg(\chi_j)|, |\arg(\Theta_j)|, |\arg(\vartheta_j)| \right\} \leq \sigma'$$

while σ is so small that, if $\bar{\ell}_1, \dots, \bar{\ell}_n$ are as in Proposition 4.1, with $\bar{\epsilon}$ replaced by $\bar{\epsilon}_1, \dots, \bar{\epsilon}_n$, then

$$\sigma \leq \min \left\{ \frac{3}{4}, \bar{\ell}_1, \dots, \bar{\ell}_n \right\}$$

and the following inequality is satisfied

$$\frac{C_1(\sigma)}{C_2(\sigma)} \frac{\bar{\delta}_j}{\sqrt{\underline{\delta}_j^2 + \sqrt{2}g(\sigma)\bar{\delta}_j}} > 1 \quad \forall i = 1, \dots, n.$$

Note that this inequality is satisfied for σ suitably small, since, by definition,

$$\bar{\delta}_j > \underline{\delta}_j, \quad C_1(\sigma') \uparrow 1, \quad C_2(\sigma') \downarrow 1, \quad g(\sigma') \downarrow 0 \quad \text{as } \sigma' \rightarrow 0.$$

Definitions and assumptions in (4.48) imply, since $\sigma(1 - \sigma) < \sigma$,

$$(1 - \sigma)G_n^- < |\chi_i| < G_n^+(1 + \sigma)$$

$$|\tan \arg(\chi_{i-1} - \chi_i)| \leq \frac{\max |\operatorname{Im}(\chi_{i-1} - \chi_i)|}{\min |\operatorname{Re}(\chi_{i-1} - \chi_i)|} \leq \frac{\theta_i}{G_i^-} \leq \sigma \leq 1$$

$$(A.122) \quad |\arg \chi_i| \leq |\arg \chi_{n-1}| + \sum_{j=i+1}^{n-1} \left| \sin^{-1} \frac{|\chi_{j-1} - \chi_j|}{|\chi_j - \chi_{j+1}|} \right| \leq \sigma \leq \frac{\pi}{3}$$

The previous inequalities imply that, firstly

$$\left| \frac{\Theta_j}{\chi_{j-1}} \right| \leq \frac{\sigma(1 - \sigma)G_n^-}{(1 - \sigma)G_n^-} \leq \sigma$$

and, similarly,

$$\left| \frac{\Theta_j}{\chi_j} \right| \leq \sigma$$

therefore, the inequality for i_j, ι_i is (4.49) follows. Secondly, the definitions of $\Theta_i^+, \vartheta_i^+$ imply that conditions (A.127) are met and hence Lemma A.2 applies. By the thesis (A.128), we have³, for $j = 1, \dots, n - 1$,

²Since, for $j = 1, \dots, n$, $\|C_{\mathcal{P}}^{(j)}\|^2$ depends only on $\chi_{j-1}, \chi_j, \Theta_j$ and ϑ_j as in (2.17) and all such coordinates, together also with Λ_j , have their anomalies bounded by σ' , we can always find such a function $g(\sigma')$.

³Beware that, if $z = (z_1, z_2, z_3) \in \mathbb{C}^3$, we denote

$$\|z\|^2 := z_1^2 + z_2^2 + z_3^2.$$

For a given $z \in \mathbb{C}$, the symbol $|z|$ denotes the usual modulus of $z \in \mathbb{C}$:

$$|z| := \sqrt{(\operatorname{Re} z)^2 + (\operatorname{Im} z)^2}.$$

$$\begin{aligned}
\|C_{\mathcal{P}}^{(j)}\|^2 &\leq \frac{|\chi_{j-1} - \chi_j|^2}{1 - \sigma^2} + (1 + \sigma)(1 + \sigma^2)|\chi_{j-1}||\chi_j||\vartheta_j - \pi|^2 \\
&\leq \frac{(G_j^+)^2}{\underline{C}_j^2} + \frac{(G_n^+)^2}{\underline{C}_j^2 B^2} |\vartheta_j - \pi|^2 \\
\text{(A.123)} \quad &\leq \bar{\delta}_j^2 (\Lambda_j^-)^2.
\end{aligned}$$

For $j = n$,

$$\|C_{\mathcal{P}}^{(j)}\|^2 = |\chi_{n-1}|^2 \leq (G_n^+)^2 < \bar{\delta}_n^2 (\Lambda_n^-)^2.$$

We suddenly have the left bound in (4.49):

$$1 - |e_{i,\mathcal{P}}^2| \leq |1 - e_{i,\mathcal{P}}^2| = \left| \frac{\|C_{\mathcal{P}}^{(i)}\|^2}{\Lambda_i^2} \right| \leq \bar{\delta}_i^2 = 1 - \underline{\epsilon}_i^2,$$

for $i = 1, \dots, n$. Now we check the right bound. To this end, previously check the following inequality

$$\text{(A.124)} \quad \left| |\chi_{j-1}| - |\chi_j| \right| \geq \frac{1 - \sigma^2}{1 + \sigma^2} G_j^-.$$

Because of the second inequality in (A.122),

$$|\arg [(\chi_{j-1} - \chi_j)(\bar{\chi}_{m-1} - \bar{\chi}_m)]| \leq 2 \tan^{-1} \sigma.$$

Then we have

$$\text{Re} [(\chi_{j-1} - \chi_j)(\bar{\chi}_{m-1} - \bar{\chi}_m)] \geq \frac{1 - \sigma^2}{1 + \sigma^2} |\chi_{j-1} - \chi_j| |\bar{\chi}_{m-1} - \bar{\chi}_m|.$$

Taking the sum for $m = j + 1, \dots, n$, gives

$$\begin{aligned}
\text{Re} (\chi_{j-1} - \chi_j) \bar{\chi}_j &\geq \frac{1 - \sigma^2}{1 + \sigma^2} |\chi_{j-1} - \chi_j| \sum_{m=j+1}^n |\bar{\chi}_{m-1} - \bar{\chi}_m| \\
&\geq \frac{1 - \sigma^2}{1 + \sigma^2} |\chi_{j-1} - \chi_j| |\bar{\chi}_j| \\
&\geq \frac{1 - \sigma^2}{1 + \sigma^2} G_j^- |\bar{\chi}_j|
\end{aligned}$$

So, Lemma A.3 with

$$A = \chi_{j-1}, \quad B = \chi_j, \quad \Delta = G_j^-, \quad a = \frac{1 - \sigma^2}{1 + \sigma^2}$$

gives (A.124). Then the thesis (A.129) of Lemma A.2 and the definition of ϑ_j provide, for $j = 1, \dots, n - 1$,

$$\text{(A.125)} \quad \|C_{\mathcal{P}}^{(j)}\|^2 \geq \frac{1}{\mathcal{A}^2 \underline{C}_j^2} \left[\mathcal{A}^2 (G_j^-)^2 - (G_n^+)^2 |\vartheta_j - \pi|^2 \geq \right] \geq (\underline{\delta}_j^2 + \sqrt{2} g(\sigma) \bar{\delta}_i) (\Lambda_j^+)^2$$

where $g(\sigma)$ is as in (A.121). Again, this inequality is implied by the definition of ϑ_j^+ in (4.48) and the ones of \mathcal{A} and C_2 in (A.120). By (A.121), (A.123) and (A.125),

for $j = 1, \dots, n$, we have

$$\begin{aligned}
 |e_{j,\mathcal{P}}|^2 &= \sqrt{\left(1 - \operatorname{Re} \frac{\|C_{\mathcal{P}}^{(j)}\|^2}{\Lambda_j^2}\right)^2 + \left(\operatorname{Im} \frac{\|C_{\mathcal{P}}^{(j)}\|^2}{\Lambda_j^2}\right)^2} \\
 &\leq \sqrt{\left(1 - \left|\frac{\|C_{\mathcal{P}}^{(j)}\|^2}{\Lambda_j^2}\right|\right)^2 + 2\left|\operatorname{Im} \frac{\|C_{\mathcal{P}}^{(j)}\|^2}{\Lambda_j^2}\right|} \\
 \text{(A.126)} \quad &\leq \sqrt{\left(1 - \underline{\delta}_j^2 - \sqrt{2}g(\sigma)\bar{\delta}_j\right)^2 + 2\bar{\delta}_j^2 g(\sigma)^2} \leq 1 - \underline{\delta}_j^2 = \bar{e}_j^2.
 \end{aligned}$$

For $j = n$,

$$\|C_{\mathcal{P}}^{(n)}\|^2 = |\chi_{n-1}|^2 \geq (\underline{\delta}_n^2 + \sqrt{2}g(\sigma)\bar{\delta}_n)(\Lambda_n^+)^2$$

again implies (A.126) with $j = n$.

The proof of the inequality on the right in (4.49) proceeds in a similar way. Indeed, starting with

$$\begin{aligned}
 |d_{i,\mathcal{P}}|^2 &= \left| \|x_{\mathcal{P}}^{(i+1)}\|^2 - 2x_{\mathcal{P}}^{(i)} \cdot x_{\mathcal{P}}^{(i+1)} + \|x_{\mathcal{P}}^{(i)}\|^2 \right| \\
 &\geq \left| \|x_{\mathcal{P}}^{(i+1)}\|^2 \right| - 2\left| x_{\mathcal{P}}^{(i)} \cdot x_{\mathcal{P}}^{(i+1)} \right| - \left| \|x_{\mathcal{P}}^{(i)}\|^2 \right|
 \end{aligned}$$

and using (as it follows from Proposition A.1)

$$\left| \|x_{\mathcal{P}}^{(i+1)}\|^2 \right| = |a_{i+1}^2(1 - e_{i+1,\mathcal{P}} \cos \zeta_{i+1})^2| \geq (1 - \eta_{i+1})^2 (a_{i+1}^-)^2$$

and analogous arguments as above to evaluate $\left| x_{\mathcal{P}}^{(i)} \cdot x_{\mathcal{P}}^{(i+1)} \right|$ and $\left| \|x_{\mathcal{P}}^{(i)}\|^2 \right|$, one easily finds the ansatz. \square

Estimates.

LEMMA A.2. *Fix a number $\sigma > 0$. Assume that, for $1 \leq j \leq n-1$,*

$$\begin{aligned}
 \text{(A.127)} \quad &\operatorname{Re} \bar{\chi}_j(\chi_{j-1} - \chi_j) > 0, \quad |\Theta_j| \leq \sigma \min\{|\chi_{j-1}|, |\chi_j|\}, \quad |\operatorname{Im}(\vartheta_j - \pi)| \leq \log(1 + \sigma).
 \end{aligned}$$

Then

$$\text{(A.128)} \quad \left\| \|C_{\mathcal{P}}^{(j)}\|^2 \right\| \leq \frac{|\chi_{j-1} - \chi_j|^2}{1 - \sigma^2} + (1 + \sigma)(1 + \sigma^2)|\chi_{j-1}||\chi_j||\vartheta_j - \pi|^2$$

$$\text{(A.129)} \quad \left\| \|C_{\mathcal{P}}^{(j)}\|^2 \right\| \geq \frac{|\chi_{j-1}| - |\chi_j|}{1 + \sigma^2} - (1 + \sigma)(1 + \sigma^2)|\chi_{j-1}||\chi_j||\vartheta_j - \pi|^2$$

PROOF. We use the formula (2.19). By Taylor's, given $a, b, z \in \mathbb{C}$, with $|z| \leq \sigma \min_{t \in [0,1]} |a + t(b-a)|$

$$\begin{aligned} \left| \sqrt{b^2 - z^2} - \sqrt{a^2 - z^2} \right| &= \left| \int_0^1 \frac{d}{dt} \sqrt{(a + t(b-a))^2 - z^2} dt \right| \\ &= \left| (b-a) \int_0^1 \frac{a + t(b-a)}{\sqrt{(a + t(b-a))^2 - z^2}} dt \right| \\ &\leq |b-a| \int_0^1 \frac{|a + t(b-a)|}{\sqrt{|a + t(b-a)|^2 - |z|^2}} dt \\ &\leq \frac{|b-a|}{\sqrt{1-\sigma^2}} \end{aligned}$$

We use this formula with $b := \chi_{j-1}$, $a := \chi_j$, $z := \Theta_j$, with the observation that, for $\operatorname{Re} \bar{\chi}_j (\chi_{j-1} - \chi_j) > 0$, the function

$t \in [0, 1] \rightarrow |\chi_j + t(\chi_{j-1} - \chi_j)|^2 = |\chi_j|^2 + 2t \operatorname{Re} \bar{\chi}_j (\chi_{j-1} - \chi_j) + t^2 |\chi_{j-1} - \chi_j|^2$ reaches its minimum, given by $\min\{|\chi_{j-1}|^2, |\chi_j|^2\}$, for $t = 0$ or $t = 1$. Developing also the function $w \in \mathbb{C} \rightarrow \cos w$ around $w = \pi$, with $\varrho := w - \pi = \varrho_1 + i\varrho_2$ and $|\varrho_2| \leq \log(1 + \sigma)$

$$\begin{aligned} |\cos w + 1| &= \left| \int_0^1 (1-t) \frac{d^2}{dt^2} \cos(\pi + t(w - \pi)) dt \right| = \frac{1}{2} |\varrho|^2 \sup_{|\varrho'| \leq \varrho} |\cos(\pi + \varrho')| \\ &\leq \frac{1}{2} |\varrho|^2 e^{|\varrho_2|} \leq \frac{1}{2} |\varrho|^2 (1 + \sigma) \end{aligned}$$

and using again the second inequality in (A.127), then inequality in (A.128) follows. The inequality in (A.129) is obtained via the second inequality in (A.127) and

$$\begin{aligned} \left| \sqrt{\chi_j^2 - \Theta_j^2} - \sqrt{\chi_{j-1}^2 - \Theta_j^2} \right| &= \frac{|\chi_{j-1}^2 - \chi_j^2|}{\left| \sqrt{\chi_j^2 - \Theta_j^2} + \sqrt{\chi_{j-1}^2 - \Theta_j^2} \right|} \\ &\geq \frac{||\chi_{j-1}|^2 - |\chi_j|^2|}{\left| \sqrt{\chi_j^2 - \Theta_j^2} + \sqrt{\chi_{j-1}^2 - \Theta_j^2} \right|} \\ &\geq \frac{||\chi_{j-1}| - |\chi_j||}{\sqrt{1 + \sigma^2}}. \end{aligned}$$

□

LEMMA A.3. *If $A, B \in \mathbb{C}$ and $a, \Delta \in \mathbb{R}_+$ verify $|A-B| \geq \Delta$ and $\operatorname{Re} \bar{B}(A-B) \geq a|B|\Delta$, where $0 < a < 1$, then $||A| - |B|| > a\Delta$.*

PROOF. Let $D := A - B$. Then $||A| - |B|| = ||B + D| - |B|| \leq a\Delta$ implies

$$|B|^2 + |D|^2 + 2 \operatorname{Re} \bar{B}D = |B + D|^2 \leq (|B| + a\Delta)^2 = |B|^2 + a^2(\Delta)^2 + 2a|B|\Delta.$$

This contradicts assumptions $|D| \geq \Delta > a\Delta$ and $\operatorname{Re} \bar{B}D \geq a|B|\Delta$. □

APPENDIX B

Proof of Lemma 3.2

In this chapter, we prove the formulae (3.40) and (3.41) given in Lemma 3.2. We recall the following result

PROPOSITION B.1 ([28]). *Let $\mathfrak{X} = \mathfrak{X}_1 \times \cdots \times \mathfrak{X}_n \subset \mathbb{R}^5 \times \cdots \times \mathbb{R}^5$ and let*

$$(\ell_k, \mathbf{X}_k) \in \mathbb{T}^1 \times \mathfrak{X}_k \rightarrow (y_\phi^{(k)}(\ell_k, \mathbf{X}_k), x_\phi^{(k)}(\ell_k, \mathbf{X}_k)) \in \mathbb{R}^3 \times \mathbb{R}^3 \quad k = 1, \dots, n$$

be mappings such that, for $1 \leq i < j \leq n$

(A) *the map*

$$\phi_{ij} : (\ell_i, \ell_j, \mathbf{X}_i, \mathbf{X}_j) \rightarrow (y_\phi^{(i)}, y_\phi^{(2)}, x_\phi^{(j)}, x_\phi^{(2)})$$

is symplectomorphism of $\mathbb{T}^2 \times \mathfrak{X}_i \times \mathfrak{X}_j$ into \mathbb{R}^{12} .

(B) *The map $(\ell_j, \mathbf{X}_j) \rightarrow (y_\phi^{(2)}(\ell_j, \mathbf{X}_j), x_\phi^{(2)}(\ell_j, \mathbf{X}_j))$ verifies*

$$\frac{\|y_\phi^{(2)}(\ell_j, \mathbf{X}_j)\|^2}{2m_j} - \frac{m_j \mathfrak{M}_j}{\|x_\phi^{(2)}(\ell_j, \mathbf{X}_j)\|} = -\frac{m_j^3 \mathfrak{M}_j^2}{2\Lambda_j^2};$$

where Λ_j is the variable conjugated to ℓ_j in this symplectomorphism.

Then the function

$$\begin{aligned} \mathbf{P}^{(i)}(\ell_i, \mathbf{X}) &:= -\frac{1}{2\pi} \int_{\mathbb{T}} d\ell_j \\ &\frac{3(x_\phi^{(i)}(\ell_i, \mathbf{X}_i) \cdot x_\phi^{(j)}(\ell_j, \mathbf{X}_j))^2 - \|x_\phi^{(i)}(\ell_i, \mathbf{X}_i)\|^2 \|x_\phi^{(j)}(\ell_j, \mathbf{X}_j)\|^2}{2\|x_\phi^{(j)}(\ell_j, \mathbf{X}_j)\|^5} \end{aligned}$$

is given by

$$(B.130) \quad \mathbf{P}^{(i)} = \frac{\mathfrak{M}_j m_j^2}{4} \frac{3(x_\phi^{(i)} \cdot C_\phi^{(j)})^2 - \|x_\phi^{(i)}\|^2 \|C_\phi^{(j)}\|^2}{\|C_\phi^{(j)}\|^4} \frac{1}{2\pi} \int_{\mathbb{T}} \frac{d\ell_j}{\|x_\phi^{(j)}\|^2}.$$

with $C_\phi^{(j)}(\mathbf{X}) := x_\phi^{(j)}(\ell_j, \mathbf{X}) \times y_\phi^{(j)}(\ell_j, \mathbf{X})$.

Even though the (i, j) projections of the \mathcal{P} -map do not verify assumption (A), one has

COROLLARY B.1. *The formula (B.130) applies also to the \mathcal{P} -map, or, more in general, to any Kepler map \mathcal{K} related to the map $\mathcal{D}el$ in Definition 2.5 via*

$$\mathbf{X}_{\mathcal{D}el} = \mathfrak{F}(\mathbf{X}).$$

PROOF. *Del* verifies (A) and (B). □

In particular, we have an expression for the second-order term of the doubly averaged Newtonian potential

$$\begin{aligned} \overline{f_{\mathcal{K}}^{ij}}^{(2)} &:= -\frac{m_i m_j}{(2\pi)^2} \int_{\mathbb{T}^2} d\ell_i d\ell_j \\ &\quad \frac{3(x_{\mathcal{K}}^{(i)}(\ell_i, \mathbf{X}_{\mathcal{K}}) \cdot x_{\mathcal{K}}^{(j)}(\ell_j, \mathbf{X}_{\mathcal{K}}))^2 - \|x_{\mathcal{K}}^{(i)}(\ell_i, \mathbf{X}_{\mathcal{K}})\|^2 \|x_{\mathcal{K}}^{(j)}(\ell_j, \mathbf{X}_{\mathcal{K}})\|^2}{2\|x_{\mathcal{K}}^{(j)}(\ell_j, \mathbf{X}_{\mathcal{K}})\|^5}. \end{aligned}$$

COROLLARY B.2. *For any \mathcal{K} as in Corollary B.1,*

$$\begin{aligned} \overline{f_{\mathcal{K}}^{ij}}^{(2)} &= m_i m_j \frac{a_i^2}{4a_j^3} \frac{\Lambda_j^3}{\|C_{\mathcal{K}}^{(j)}\|^5} \left[-\left(\frac{5}{2} - \frac{3}{2} \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2}\right) \|C_{\mathcal{K}}^{(j)}\|^2 \right. \\ \text{(B.131)} \quad &+ \left. \frac{3}{2} \left(5 - 4 \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2}\right) (P_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)})^2 + \frac{3}{2} \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2} (Q_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)})^2 \right] \end{aligned}$$

PROOF. Lemma B.1 implies that

$$\begin{aligned} \overline{f_{\mathcal{K}}^{ij}}^{(2)} &= m_i m_j \frac{M_j m_j^2}{4} \frac{\frac{1}{2\pi} \int_{\mathbb{T}} \left(3(x_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)})^2 - \|x_{\mathcal{K}}^{(i)}\|^2 \|C_{\mathcal{K}}^{(j)}\|^2\right) d\ell_i}{\|C_{\mathcal{K}}^{(j)}\|^4} \\ &\quad \times \frac{1}{2\pi} \int_{\mathbb{T}} \frac{d\ell_j}{\|x_{\mathcal{K}}^{(j)}\|^2}. \end{aligned}$$

By (2.1)

$$\begin{aligned} x_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)} &= (a_{i,\mathcal{K}} P_{\mathcal{K}}^{(i)} + b_{i,\mathcal{K}} Q_{\mathcal{K}}^{(i)}) \cdot C_{\mathcal{K}}^{(j)} \\ &= a_{i,\mathcal{K}} P_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)} + b_{i,\mathcal{K}} Q_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)} \end{aligned}$$

Therefore, squaring, ℓ_i -averaging and using

$$\begin{aligned} \frac{1}{2\pi} \int_{\mathbb{T}} (a_{i,\mathcal{K}})^2 d\ell_i &= \frac{a_i^2}{2} \left(5 - 4 \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2}\right) \\ \frac{1}{2\pi} \int_{\mathbb{T}} (b_{i,\mathcal{K}})^2 d\ell_i &= \frac{a_i^2}{2} \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2} \\ \frac{1}{2\pi} \int_{\mathbb{T}} a_{i,\mathcal{K}} b_{i,\mathcal{K}} d\ell_i &= 0 \end{aligned}$$

we obtain

$$\begin{aligned} \frac{1}{2\pi} \int_{\mathbb{T}} (x_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)})^2 d\ell_i &= \frac{a_i^2}{2} \left(5 - 4 \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2}\right) (P_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)})^2 \\ &+ \frac{a_i^2}{2} \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2} (Q_{\mathcal{K}}^{(i)} \cdot C_{\mathcal{K}}^{(j)})^2. \end{aligned}$$

Using finally

$$\frac{1}{2\pi} \int_{\mathbb{T}} \|x_{\mathcal{K}}^{(i)}\|^2 d\ell_i = a_i^2 \left(\frac{5}{2} - \frac{3}{2} \frac{\|C_{\mathcal{K}}^{(i)}\|^2}{\Lambda_i^2}\right), \quad \frac{1}{2\pi} \int_{\mathbb{T}} \frac{d\ell_j}{\|x_{\mathcal{K}}^{(j)}\|^2} = \frac{1}{a_j^2} \frac{\Lambda_j}{\|C_{\mathcal{K}}^{(j)}\|}$$

we obtain (B.131). □

Now we may proceed with proving the formulae in (3.40) and (3.41).

PROOF OF (3.40). We apply Corollary B.2 with $\mathcal{K} = \mathcal{P}$, $i = n - 1$, $j = n$. Using $\|C_{\mathcal{P}}^{(n)}\| = \chi_{n-1}$ (see (2.17)), $C_{\mathcal{P}}^{(n)} = S_{\mathcal{P}}^{(n)}$ and Eq. (2.3), Proposition 2.1, and Remark 2.2, we have

$$\begin{aligned} P_{\mathcal{P}}^{(n-1)} \cdot S_{\mathcal{P}}^{(n)} &= \Theta_{n-1} \\ Q_{\mathcal{P}}^{(n-1)} \cdot S_{\mathcal{P}}^{(n)} &= \frac{1}{\|C_{\mathcal{P}}^{(n-1)}\|} ((S_{\mathcal{P}}^{(n-1)} - S_{\mathcal{P}}^{(n)}) \times P_{\mathcal{P}}^{(n-1)}) \cdot S_{\mathcal{P}}^{(n)} \\ &= \frac{1}{\|C_{\mathcal{P}}^{(n-1)}\|} S_{\mathcal{P}}^{(n-1)} \times P_{\mathcal{P}}^{(n-1)} \cdot S_{\mathcal{P}}^{(n)} \\ &= \frac{1}{\|C_{\mathcal{P}}^{(n-1)}\|} \sqrt{(\chi_{n-1}^2 - \Theta_{n-1}^2)(\chi_{n-2}^2 - \Theta_{n-1}^2)} \sin \vartheta_{n-1}. \end{aligned}$$

PROOF OF (3.41). By Corollary B.2 with $\mathcal{K} = \mathcal{P}$, $j = i + 1$, we find, for $\frac{f_{\mathcal{P}}^{i,i+1}(2)}{(2)}$ an expression as in (B.131), replacing $(n - 1, n)$ with $(i, i + 1)$. □

(B.132)

$$\begin{aligned} P_{\mathcal{P}}^{(i)} \cdot C_{\mathcal{P}}^{(i+1)} &= P_{\mathcal{P}}^{(i)} \cdot (S_{\mathcal{P}}^{(i+1)} - S_{\mathcal{P}}^{(i+2)}) = \Theta_i - P_{\mathcal{P}}^{(i)} \cdot S_{\mathcal{P}}^{(i+2)} \\ Q_{\mathcal{P}}^{(i)} \cdot C_{\mathcal{P}}^{(i+1)} &= Q_{\mathcal{P}}^{(i)} \cdot (S_{\mathcal{P}}^{(i+1)} - S_{\mathcal{P}}^{(i+2)}) = \frac{1}{\|C_{\mathcal{P}}^{(i)}\|} (\sqrt{(\chi_i^2 - \Theta_i^2)(\chi_{i-1}^2 - \Theta_i^2)} \sin \vartheta_i \\ &\quad - S_{\mathcal{P}}^{(i)} \times P_{\mathcal{P}}^{(i)} \cdot S_{\mathcal{P}}^{(i+2)} - P_{\mathcal{P}}^{(i)} \times S_{\mathcal{P}}^{(i+1)} \cdot S_{\mathcal{P}}^{(i+2)}). \end{aligned}$$

Now, when $(\Theta_{i+1}, \vartheta_{i+1}) = (0, \pi)$, $\|C_{\mathcal{P}}^{(i+1)}\|$ reduces to

$$\|C_{\mathcal{P}}^{(i+1)}\| = \chi_i - \chi_{i+1},$$

(provided $\arg(\chi_i - \chi_{i+1}) \in (-\frac{\pi}{2}, \frac{\pi}{2}] \bmod 2\pi$) and $S_{\mathcal{P}}^{(i+2)} \parallel S_{\mathcal{P}}^{(i+1)}$, so

$$S_{\mathcal{P}}^{(i+2)} = \frac{\chi_{i+1}}{\chi_i} S_{\mathcal{P}}^{(i+1)}$$

and hence, the extra-terms in (B.132) reduce to

$$\begin{aligned} P_{\mathcal{P}}^{(i)} \cdot S_{\mathcal{P}}^{(i+2)} &= \Theta_i \frac{\chi_{i+1}}{\chi_i} \\ S_{\mathcal{P}}^{(i)} \times P_{\mathcal{P}}^{(i)} \cdot S_{\mathcal{P}}^{(i+2)} &= \frac{\chi_{i+1}}{\chi_i} \sqrt{\chi_{i-1}^2 - \Theta_i^2} \sqrt{\chi_i^2 - \Theta_i^2} \sin \vartheta_i \\ P_{\mathcal{P}}^{(i)} \times S_{\mathcal{P}}^{(i+1)} \cdot S_{\mathcal{P}}^{(i+2)} &= 0. \end{aligned}$$

Then (3.41) readily follows. □

Checking the non-degeneracy condition

In this chapter we prove statement 4 of Proposition 5.2.

Due to the form of $\widehat{h_{\text{sec}}}$ in (5.73)–(5.74) and to the bound for $\widehat{h_{\text{sec},h}^i}$ in (5.76), it is sufficient to prove that the maps

$$\zeta_i^{(h)} \rightarrow \overline{\omega_{\text{sec}}^i} := \partial_{\zeta_i^{(h)}} \overline{h_{\text{sec}}^i}(\zeta_i^{(h)}, \Lambda_{n-h}^{(h)}, \Lambda_{n-h+1}^{(h)})$$

in (5.74), where

$$\zeta_i^{(h)} = \begin{cases} \left(\frac{(p_1^{(h)})^2 + (q_1^{(h)})^2}{2}, \chi_1^{(h)} \right) & i = 1 & \& n = 2 \\ \left(\frac{(p_{n-1}^{(h)})^2 + (q_{n-1}^{(h)})^2}{2}, \chi_{n-2}^{(h)}, \chi_{n-1}^{(h)} \right) & i = n - 1 & \& n \geq 3 \\ \left(\frac{(p_i^{(h)})^2 + (q_i^{(h)})^2}{2}, \chi_{i-1}^{(h)} \right) & i = 2, \dots, n - 2 & \& n \geq 4 \\ \left(\frac{(p_1^{(h)})^2 + (q_1^{(h)})^2}{2} \right) & i = 1 & \& n \geq 3 \end{cases}$$

are diffeomorphisms, with non-vanishing Hessian matrices. We shall do this verifications for just one of the cases above, and we choose the second case in the list, $i = n - 1$, for $n \geq 3$. The explicit expression of $\overline{h_{\text{sec}}^{n-1}}$ is given in (5.98)–(5.99). We neglect the coefficient \mathcal{A}_{n-1} (which does not depend on $\zeta_{n-1}^{(h)}$) and we denote

$$\widehat{h_{\text{sec}}^{n-1}} = E_{n-1} + \Omega_{n-1} \frac{p_{n-1}^2 + q_{n-1}^2}{2} + \tau_{n-1} \left(\frac{p_{n-1}^2 + q_{n-1}^2}{2} \right)^2 + O(p_{n-1}, q_{n-1})^6]$$

the function $\overline{h_{\text{sec}}^{n-1}}$ thus rescaled, and $\widehat{\omega_{\text{int}}^{n-1}}$ its gradient with respect to

$$\left(\frac{(p_{n-1}^{(h)})^2 + (q_{n-1}^{(h)})^2}{2}, \chi_{n-2}, \chi_{n-1} \right).$$

A perturbative argument shows that, under the choices of Corollary 4.1, the frequency-map with respect to (χ_{n-2}, χ_{n-1}) associated to

$$E_{n-1} = -\frac{\Lambda_n^3}{2\chi_{n-1}^3} \left(5 - 3 \frac{(\chi_{n-2} - \chi_{n-1})^2}{\Lambda_{n-1}^2} \right)$$

is an injection of its domain and hence, by another perturbative argument, so is the gradient of $\widehat{h_{\text{sec}}^{n-1}}$ with respect to the same coordinates, for any fixed value of $\frac{p_{n-1}^2 + q_{n-1}^2}{2}$. On the other hand, since τ_{n-1} does not vanish under the same

assumptions of Corollary 4.1, $\widehat{\omega_{\text{int}}^{n-1}}$ is an injection. The computation shows that the Jacobian of $\widehat{\omega_{\text{int}}^{n-1}}$ does not vanish. \square

Some results from perturbation theory

D.1. A multi-scale normal form theorem

The purpose of this chapter is to present a normal form result which takes into account different scale lengths. It is a particularization of [31, Normal Form Lemma, p. 192] and uses the same techniques of that paper.

Following [31], the notations are as follows.

- If $A \subset \mathbb{R}^\nu$ is open and connected, $\mathbb{T} := \mathbb{R}/(2\pi\mathbb{Z})$ is the usual flat torus, r, s are positive numbers, we denote as $A_r := \bigcup_{x \in A} \left\{ z \in \mathbb{C}^\nu : z \in B_r^\nu(x) \right\}$ the complex r -neighborhood of A . \mathbb{T}_s^ν will denote the complex set $\mathbb{T} + i[-s, s]$. As usual, $B_r^\nu(x)$ denotes the ball in \mathbb{C}^ν with radius r centered at x , accordingly to a prefixed norm $|\cdot|$ of \mathbb{C}^ν .
- If $f = f(u, p, q, \varphi)$ is real-analytic for $(u, p, q, \varphi) \in W_{v,s,\varepsilon} = U_v \times B_\varepsilon^{2\ell} \times \mathbb{T}_s^\nu$, and affords the Taylor-Fourier expansion

$$f = \sum_{k \in \mathbb{Z}^m} f_{k,\alpha,\beta}(u) e^{ik \cdot \varphi} \prod_{j=1}^{\ell} \left(\frac{p_j - iq_j}{\sqrt{2}} \right)^{\alpha_j} \left(\frac{p_j + iq_j}{i\sqrt{2}} \right)^{\beta_j},$$

we denote as $\|f\|_{v,s,\varepsilon}$ its “sup-(Taylor, Fourier) norm”:

$$\|f\|_{v,s,\varepsilon} := \sum_{\substack{(a,b) \in \mathbb{N}^{2\ell} \\ k \in \mathbb{Z}^\nu}} \sup_{u \in U_v} |f_{\alpha,\beta,k}(u)| e^{|k|s} \varepsilon^{-|(\alpha,\beta)|}$$

with $|k| := |k|_1$, $|(\alpha, \beta)| := |\alpha|_1 + |\beta|_1$.

- If f is as in the previous item, $K > 0$ and $\mathfrak{L} = \mathfrak{L}_1 \times \mathfrak{L}_2$ is a sub-lattice of $\mathbb{Z}^\nu \times \mathbb{Z}^\ell$, $T_K f$ and $\Pi_{\mathfrak{L}} f$ denote, respectively, the K -truncation and the \mathfrak{L} -projection of f :

$$T_K f := \sum_{|(\alpha,\beta)| \leq K, |k| \leq K} f_{\alpha,\beta,k}(u) e^{ik \cdot \varphi} \prod_{j=1}^{\ell} \left(\frac{p_j - iq_j}{\sqrt{2}} \right)^{\alpha_j} \left(\frac{p_j + iq_j}{i\sqrt{2}} \right)^{\beta_j}$$

$$\Pi_{\mathfrak{L}} f := \sum_{\substack{k \in \mathfrak{L}_1 \\ \alpha - \beta \in \mathfrak{L}_2}} f_{\alpha,\beta,k}(u) e^{ik \cdot \varphi} \prod_{j=1}^{\ell} \left(\frac{p_j - iq_j}{\sqrt{2}} \right)^{\alpha_j} \left(\frac{p_j + iq_j}{i\sqrt{2}} \right)^{\beta_j}.$$

PROPOSITION D.1 (Multi-scale normal form). *Let*

$$\nu, \quad \ell, \quad 1 \leq m_1 < \dots < m_N = m$$

be natural numbers;

$$A \subset \mathbb{R}^\nu, \quad B \subset \mathbb{R}^{2\ell}, \quad C_1, C_1' \subset \mathbb{R}^{m_1}, \quad C_2, C_2' \subset \mathbb{R}^{m_2 - m_1}, \quad \dots, \quad C_N, C_N' \subset \mathbb{R}^{m_N - m_{N-1}},$$

be open and connected sets;

$$r, s, \varepsilon, \rho_1 \geq \rho_2 \cdots \geq \rho_N, \rho'_1 \geq \rho'_2 \cdots \geq \rho'_N$$

positive numbers. Put

$$v_i := (r, \rho_1, \dots, \rho_i, \rho'_1, \dots, \rho'_i), \quad v := v_N$$

$$U_{v_i}^{(i)} := A_r \times C_{1\rho_1} \times \cdots \times C_{i\rho_i} \times C'_{1\rho'_1} \times \cdots \times C'_{i\rho'_i}, \quad U_v := U_{v_N}^{(N)}$$

$$W_{v_i, s, \varepsilon}^{(i)} := U_{v_i}^{(i)} \times \mathbb{T}_s^\nu \times B_\varepsilon, \quad W_{v, s, \varepsilon} := W_{v_N, s, \varepsilon}^{(N)}$$

with $i = 1, \dots, N$.

Let \mathbf{a} , $K > 0$ with $0 < s < 6 \log 6/5$ and $Ks \geq 12$; let also \mathfrak{L} and $\mathfrak{Z}_1, \dots, \mathfrak{Z}_N$ be sub-lattices of $\mathbb{Z}^\ell \times \mathbb{Z}^\nu$ and let $\mathfrak{Z} := \mathfrak{Z}_1 \cup \cdots \cup \mathfrak{Z}_N$.

Let

$$(D.133) \quad H(u, \varphi, p, q) = h(p, q, I) + f(u, \varphi, p, q)$$

be real-analytic for $(u, \varphi, p, q) \in W_{v, s, \varepsilon}$, where $u := (I, \eta, \xi) = (I_1, \dots, I_\nu, \eta_1, \dots, \eta_m, \xi_1, \dots, \xi_m)$. Suppose that

- (i) h depends on (p, q) only via $\frac{p_i^2 + q_i^2}{2}$, with the frequency map $\omega = (\omega_1, \dots, \omega_\ell, \omega_{\ell+1}, \dots, \omega_{\ell+\nu})$ defined via

$$\omega_i := \begin{cases} \frac{\partial_{p_i^2 + q_i^2} h}{2} & 1 \leq i \leq \ell \\ \partial_{I_{i-\ell}} h & \ell + 1 \leq i \leq \ell + \nu \end{cases}$$

verifying

$$(D.134) \quad |\omega(p, q, I) \cdot (k', k)| \geq \mathbf{a} \quad \forall (k', k) \in \mathfrak{Z} \setminus \mathfrak{L}, |(k', k)| \leq K$$

and all $(p, q, I) \in B_\varepsilon^{2\ell} \times A_r$;

- (ii) f is a sum

$$(D.135) \quad f = \sum_{i=1}^N f_i(u_i, \varphi, p, q)$$

where f_i is real-analytic on $W_{v_i, s, \varepsilon}^{(i)}$ and has the form

$$(D.136)$$

$$f_i(u_i, \varphi, p, q) = \sum_{(\alpha^- \alpha^+, k) \in \mathfrak{Z}_i} f_{k, \alpha^-, \alpha^+}^i(u_i) \prod_{j=1}^{\nu} e^{ik_j \varphi_j} \prod_{k=1}^{\ell} \left(\frac{p_k - iq_k}{\sqrt{2}} \right)^{\alpha_k^-} \left(\frac{p_k + iq_k}{\sqrt{2}i} \right)^{\alpha_k^+}$$

with

$$(D.137) \quad u_i := (I, \eta^i, \xi^i) := (I_1, \dots, I_\nu, \eta_1, \dots, \eta_{m_i}, \xi_1, \dots, \xi_{m_i});$$

- (iii) the following “smallness” conditions hold. If

$$(D.138) \quad c_i := e(1 + \ell_i e + m_i e)/2, \quad d_i := \min\{rs, \varepsilon^2, \rho_i \rho'_i\}$$

with e denoting Neper number, then

$$(D.139) \quad \|f_i\|_{W_{v_i, s, \varepsilon}^{(i)}} \leq E_i, \quad \sum_{i=1}^N \frac{7}{6} \left(\frac{9}{8}\right)^{i-1} \frac{2^7 c_i K s}{ad_i} E_i < 1.$$

Then, one can find a real-analytic and symplectic transformation

$$\Phi : W_{v/6^N, s/6^N, \varepsilon/6^N} \rightarrow W_{v, \sigma, \varepsilon}$$

which conjugates H to

$$H_*(u, \varphi, p, q) := H \circ \Phi = \mathfrak{h}(I, p, q) + \sum_{i=1}^N g_i(u_i, \varphi, p, q) + \sum_{i=1}^N f_i^*(u, \varphi, p, q),$$

where g_i, f_i verify

$$\begin{aligned} g_i &= \Pi_{\mathfrak{I}_i \cap \Sigma} T_K g_i \\ \|g_i - \Pi_{\mathfrak{I}_i \cap \Sigma} T_K f_i\|_{v_i/6^N, \sigma/6^N, \varepsilon/6^N} &\leq \left(\frac{9}{8}\right)^{2(i-1)} \frac{2^7 c_i \|f_i\|_{v_i, s, \varepsilon}^2}{\mathfrak{ad}_i} \\ &\quad + \frac{7}{6} \left(\frac{9}{8}\right)^{2(i-1)} \sum_{j=1}^{i-1} \frac{2^7 c_j \|f_j\|_{v_j, s, \varepsilon}}{\mathfrak{ad}_j} \|f_i\|_{v_i, s, \varepsilon} \\ &\quad + \sum_{k=1}^{i-1} \left(\frac{9}{8}\right)^{i-1-k} \frac{2^4 c_k \|f_k\|_{v_k, s, \varepsilon} K s}{\mathfrak{ad}_k} \|f_i\|_{v_i, s, \varepsilon} \\ \|f_i^*\|_{v_i/6^N, s/6^N, \varepsilon/6^N} &\leq \left(\frac{9}{8}\right)^{N-1} e^{-Ks/6^i} \|f_i\|_{v_i, s, \varepsilon} \end{aligned}$$

Finally, Φ is close to the identity in the following sense. Given F , real-analytic on $W_{v_i/6^N, s/6^N, \varepsilon/6^N}^{(i)}$,

$$\|F \circ \Phi - F\|_{v/6^N, s/6^N, \varepsilon/6^N} \leq \sum_{k=1}^N \left(\frac{9}{8}\right)^{N-k} \frac{2^4 c_k \|f_k\|_{v_k, s, \varepsilon} K s}{\mathfrak{ad}_{k,i}} \|F\|_{v_i/6^N, s/6^N, \varepsilon/6^N}$$

with $d_{k,i} := \max\{d_k, d_i\}$.

The proof of Proposition D.1 is based on the following

LEMMA D.1. Let $\bar{N} \in \mathbb{N}$, $\nu, \ell, m_i, A, B, C_i, C'_i, r, s, \rho_i, \rho'_i, U_{v_i}^{(i)}, W_{v_i, s, \varepsilon}^{(i)}, d_i$, with $i = 1, \dots, \bar{N} + 1$, be as in Proposition D.1;

$$v := (r, \rho_1, \dots, \rho_{\bar{N}+1}, \rho'_1, \dots, \rho'_{\bar{N}+1}), U_v := U_{v_{\bar{N}+1}}^{(\bar{N}+1)}, W_{v, s, \varepsilon} := W_{v_{\bar{N}+1}, s, \varepsilon}^{(\bar{N}+1)}.$$

Let

$$(D.140) \quad H(p, q, I, \varphi, \eta, \xi) = \mathfrak{h}(p, q, I) + g(p, q, I, \varphi, \eta, \xi) + f(p, q, I, \varphi, \eta, \xi)$$

be real-analytic for $(u, \varphi, p, q) \in W_{v, s, \varepsilon}$. Suppose assumption (i) of Proposition D.1 and, moreover, the following ones

(ii) g is a sum

$$(D.141) \quad g = \sum_{i=1}^{\bar{N}} g_i(u_i, \varphi, p, q)$$

where g_i is real-analytic on $W_{v_i, s, \varepsilon}^{(i)}$ and u_i is as in (D.137);

(iii) $g_1, \dots, g_{\bar{N}}$ and f satisfy

$$g_i = \Pi_{\Sigma} g_i, \quad f = \Pi_{\mathfrak{I}} f$$

and

$$(D.142) \quad \sum_{i=1}^{\bar{N}} \frac{2^7 c_i K s}{\mathbf{ad}_i} \|g_i\|_{v_i, s_i, \varepsilon_i} < 1, \quad \|f\|_{v, s, \varepsilon} < \frac{\mathbf{ad}_{\bar{N}+1}}{2^7 c_{\bar{N}+1} K s}.$$

Then, one can find a real-analytic and symplectic transformation

$$\Phi : (u', \varphi', p', q') \in W_{v/6, s/6, \varepsilon/6} \rightarrow (u, \varphi, p, q) \in W_{v, s, \varepsilon}$$

such that

$$H_* := H \circ \Phi = h + g + g_* + f_*,$$

where $g_* = \Pi_{\mathfrak{Z} \cap \mathfrak{L}} T_K g_*$ is $\mathfrak{Z} \cap \mathfrak{L}$ -resonant and the following bounds hold

$$\begin{aligned} \|g_* - T_K \Pi_{\mathfrak{Z} \cap \mathfrak{L}} f\|_{v/6, s/6, \varepsilon/6} &\leq \left(\frac{2^7 c_{\bar{N}+1} \|f\|_{v, s, \varepsilon}}{\mathbf{ad}_{\bar{N}+1}} + \sum_{i=1}^n \frac{2^7 c_i \|g_i\|_{v_i, s_i, \varepsilon_i}}{\mathbf{ad}_i} \right) \|f\|_{v, s, \varepsilon} \\ &\leq \frac{\|f\|_{v, s, \varepsilon}}{6} \\ \|f_*\|_{v/6, s/6, \varepsilon/6} &\leq e^{-Ks/6} \|f\|_{v, s, \varepsilon}. \end{aligned}$$

Finally, Φ is close to the identity in the following sense: for any F which is real-analytic on $W_{v, s, \varepsilon}^{(i)}$,

$$(D.143) \quad \|F \circ \Phi - \Phi\|_{v/6, s/6, \varepsilon/6} \leq \frac{2^4 c_{\bar{N}+1} \|f\|_{v, s, \varepsilon} K s}{\mathbf{ad}_i} \|F\|_{v_i, s_i, \varepsilon_i} < \frac{1}{8} \|F\|_{v, s, \varepsilon}.$$

The following Lemma is a trivial extension¹ of [31, Iterative Lemma]. Its proof is omitted.

LEMMA D.2. Let $s = (s_1, \dots, s_\nu)$, $r = (r_1, \dots, r_\nu)$, $\varepsilon = (\varepsilon_1, \dots, \varepsilon_\ell)$, $\rho = (\rho_1, \dots, \rho_m)$, $\rho' = (\rho'_1, \dots, \rho'_m)$, $v := (r, \rho, \rho')$, $\hat{v} := (\hat{r}, \hat{\rho}, \hat{\rho}') < v/2$, $\hat{s} < s/2$, $\hat{\varepsilon} < \varepsilon/2$,

$$\delta := \min_{\substack{i=1, \dots, \nu \\ j=1, \dots, \ell \\ k=1, \dots, m}} \{\hat{r}_i \hat{s}_i, \hat{\varepsilon}_j^2, \hat{\rho}_k \hat{\rho}'_k\}.$$

Let

$$H(u, \varphi, p, q) = h(I, p, q) + g(u, \varphi, p, q) + f(u, \varphi, p, q) \quad g(u, \varphi, p, q) = \sum_{i=1}^m g_i(u, \varphi, p, q)$$

be real-analytic on $W_{v, s, \varepsilon}$. Assume that inequality (D.134) and

$$(D.144) \quad \|f\|_{v, s, \varepsilon} < \frac{\mathbf{ad}^\delta}{c}$$

are satisfied. Then one can find a real-analytic and symplectic transformation

$$\Phi : W_{v-2\hat{v}, s-2\hat{s}, \varepsilon-2\hat{\varepsilon}} \rightarrow W_{v, s, \varepsilon}$$

¹In order to obtain the extension it is sufficient to replace ϕ of [31, Appendix A] with

$$\phi = \sum_{\substack{(\alpha-\beta, k) \in \mathfrak{R} \setminus \mathfrak{L} \\ |(\alpha, \beta)| \leq K, |k| \leq K}} \frac{f_{k, \alpha, \beta}(u)}{i(\alpha - \beta, k) \cdot \omega} e^{ik \cdot \varphi} \prod_{j=1}^{\ell} \left(\frac{p_j - iq_j}{\sqrt{2}} \right)^{\alpha_j} \left(\frac{p_j + iq_j}{i\sqrt{2}} \right)^{\beta_j}$$

defined by the time-one flow² $X_\phi^1 f := f \circ \Phi$ of a suitable ϕ verifying

$$\|\phi\|_{v,s,\varepsilon} \leq \frac{\|f\|_{v,s,\varepsilon}}{\mathbf{a}}$$

such that

$$H_+ := H \circ \Phi = h + g + \Pi_{\mathcal{L} \cap \mathfrak{J}} f + f_+$$

and, moreover, the following bounds hold

$$\begin{aligned} \|f_+\|_{v-2\hat{v},s-2\hat{s},\varepsilon-2\hat{\varepsilon}} &\leq \left(1 - \frac{c}{\mathbf{a}\bar{d}}\|f\|_{v,s,\varepsilon}\right)^{-1} \left[\frac{c}{\mathbf{a}\bar{d}}\|f\|_{v,s,\varepsilon}^2 \right. \\ &\quad \left. + e^{-K\hat{s}}\|f\|_{v,s,\varepsilon} + \left(\frac{\varepsilon - \hat{\varepsilon}}{\varepsilon}\right)^K \|f\|_{v,s,\varepsilon} + \|\{\phi, g\}\|_{v-\hat{v},s-\hat{s},\varepsilon-\hat{\varepsilon}} \right] \end{aligned}$$

Finally, for any real-analytic function F on $W_{v,s,\varepsilon}$,

$$\|F \circ \Phi - F\|_{v-2\hat{v},s-2\hat{s},\varepsilon-2\hat{\varepsilon}} \leq \frac{\|\{\phi, F\}\|_{v-\hat{v},s-\hat{s},\varepsilon-\hat{\varepsilon}}}{1 - \frac{c\|f\|_{v,s,\varepsilon}}{\mathbf{a}\bar{d}}}.$$

PROOF OF LEMMA D.1. Following [31], the proof is obtained via iterate applications of Lemma D.2.

To avoid too many indices, we shall prove this lemma taking, in (D.141), $\bar{N} = 1$; the extension to $\bar{N} \geq 1$ being straightforward. Namely, we take

$$(D.145) \quad \begin{aligned} \rho_1 = \cdots = \rho_{m_1} = \bar{\rho}, \quad \rho'_1 = \cdots = \rho'_{m_1} = \bar{\rho}' \\ \rho_{m_1+1} = \cdots = \rho_m = \rho, \quad \rho'_{m_1+1} = \cdots = \rho'_m = \rho \end{aligned}$$

where $1 \leq m_1 < m$. Letting

$$\begin{aligned} v &:= (r, \rho, \rho'), \quad \bar{v} := (r, \bar{\rho}, \bar{\rho}'), \quad E := \|f\|_{v,s,\varepsilon}, \quad G := \|g\|_{\bar{v},s,\varepsilon}, \\ \bar{c} &= c_1, \quad c = c_2, \quad \bar{d} := \min\{rs, \varepsilon^2, \bar{\rho}\bar{\rho}'\}, \quad d := \{rs, \varepsilon^2, \rho\rho'\}, \end{aligned}$$

we rewrite the assumptions in (D.142) as

$$(D.146) \quad \frac{2^7 \bar{c} G K s}{\mathbf{a}\bar{d}} < 1, \quad \frac{2^7 c E K s}{\mathbf{a}d} < 1.$$

The inequality on the right clearly implies (D.144). So, we apply Lemma D.2 to the Hamiltonian (D.140), taking $r_1 = \cdots = r_\nu = r$, $s_1 = \cdots = s_\nu = s$, $\varepsilon_1 = \cdots = \varepsilon_\ell = \varepsilon$, ρ_k, ρ'_k as in (D.145) and

$$\begin{aligned} \hat{v} = \hat{v}_0 &:= v/6, \quad \hat{s} = \hat{s}_0 := s/6, \quad \hat{\varepsilon} = \hat{\varepsilon}_0 := \varepsilon/6 \\ \bar{\hat{v}} = \bar{\hat{v}}_0 &:= \bar{v}/6, \quad \bar{\hat{s}} := \bar{\hat{s}}_0 := \bar{s}/6, \quad \bar{\hat{\varepsilon}} := \bar{\hat{\varepsilon}}_0 := \bar{\varepsilon}/6 \\ \delta &:= \{\hat{r}\hat{s}, \hat{\varepsilon}^2, \hat{\rho}\hat{\rho}'\} = \frac{d}{36}. \end{aligned}$$

Letting

$$v_1 := v - 2\hat{v}_0 = 3/4v, \quad s_1 := s - 2\hat{s}_0 = 2/3s, \quad \varepsilon_1 := \varepsilon - 2\hat{\varepsilon}_0 = 2/3\varepsilon$$

by Lemma D.2, we find a canonical transformation $\Phi_0 = X_{\phi_0}$ which is real-analytic on $W_{v_1, s_1, \varepsilon_1}$ and conjugates H to $H_1 = h + g + g_1 + f_1$, where $g_1 = \Pi_{\mathcal{L} \cap \mathfrak{J}} T_K f$ and

²The time-one flow generated by ϕ is defined as the differential operator

$$X_\phi^1 := \sum_{k=0}^{\infty} \frac{L_\phi^k}{k!}$$

where $L_\phi^0 f := f$ and $L_\phi^k f := \{\phi, L_\phi^{k-1} f\}$, with $k = 1, 2, \dots$.

$$\begin{aligned} \|f_1\|_{v_1, s_1, \varepsilon_1} &\leq \left(1 - \frac{36cE}{ad}\right)^{-1} \left[\frac{36cE}{ad} + e^{-Ks/6} + \left(\frac{5}{6}\right)^K \right] E \\ &+ \left(1 - \frac{36cE}{ad}\right)^{-1} \frac{36\bar{c}G}{a\bar{d}} E \end{aligned}$$

where

$$\bar{\delta} := \min\{\hat{r}\hat{s}, \hat{\varepsilon}^2, \hat{\rho}\hat{\rho}'\} = \frac{\bar{d}}{36}.$$

Here, we have used

$$\begin{aligned} \|\{\phi, g\}_{I, \varphi, \eta, \xi}\|_{v-\hat{v}, s-\hat{s}, \varepsilon-\hat{\varepsilon}} &= \|\{\phi, g\}_{I, \varphi, \eta^1, \xi^1}\|_{\bar{v}-\hat{v}, s-\hat{s}, \varepsilon-\hat{\varepsilon}} \\ \text{(D.147)} \qquad \qquad \qquad &\leq \frac{\bar{c}G}{a\bar{\delta}} = 36 \frac{\bar{c}G}{ad} \end{aligned}$$

since g depends on η, ξ only via $\eta^1 = (\eta_1, \dots, \eta_{m_1})$, $\xi^1 = (\xi_1, \dots, \xi_{m_1})$. It is sufficient to consider the case

$$e^{-Ks/6} + \left(\frac{5}{6}\right)^K \leq \frac{18cE}{ad}$$

since otherwise the Lemma is proved. In such case, using (D.146) we can write

$$\begin{aligned} E_1 = \|f_1\|_{v_1, s_1, \varepsilon_1} &\leq \frac{32}{23} \left(\frac{9}{32} \frac{2^7 cEKs}{ad} + \frac{9}{64} \frac{2^7 cEKs}{ad} + \frac{9}{32} \frac{2^7 \bar{c}GKs}{a\bar{d}} \right) \frac{E}{Ks} \\ \text{(D.148)} \qquad \qquad \qquad &< \frac{E}{Ks} \max\left\{ \frac{2^7 cEKs}{ad}, \frac{2^7 \bar{c}GKs}{a\bar{d}} \right\} < \frac{E}{4} \end{aligned}$$

Let

$$L := \left\lceil \frac{Ks}{12 \log 2} \right\rceil.$$

Note that

$$\text{(D.149)} \qquad \qquad \qquad L \geq 1, \quad Ks > 8L,$$

since we have assumed $Ks \geq 12$. We want to prove that Lemma D.2 can be applied L times with parameters

$$\text{(D.150)} \qquad \hat{v}_i = \frac{v}{4L}, \quad \hat{\varepsilon}_i = \frac{\varepsilon}{4L}, \quad \hat{s}_i = \frac{s}{4L}, \quad \delta_i = \frac{d}{16L^2}, \quad i = 1, \dots, L.$$

For $L = 1$, this follows from (D.148):

$$E_1 := \|f_1\|_{v_1, s_1, \varepsilon_1} \leq \frac{E}{Ks} \leq 2^{-7} \frac{ad}{c(Ks)^2} < 2^{-13} \frac{a\delta_1}{c}$$

which is implied by the inequality in (D.148) and assumption (D.142). We then assume $L \geq 2$. Suppose, by induction, that, for a certain $1 \leq i \leq L-1$, and any $1 \leq j \leq i$, we have conjugated H to

$$H_j = h + g + \bar{g}_j + f_j$$

where $\bar{g}_j = \sum_{k=0}^{j-1} \Pi_{\mathcal{L} \cap \mathfrak{Z}} T_K f_k$

$$\text{(D.151)} \qquad E_j := \|f_j\|_{v_j, s_j, \varepsilon_j} \leq \min\left\{ \frac{E}{4^j}, 2^{-6} \frac{a\delta_j}{c} \right\}$$

where $\hat{v}_0, \hat{s}_0, \hat{\varepsilon}_0$ are as above, $v_0 := v, s_0 := s, \varepsilon_0 := \varepsilon$ and $v_j = v_{j-1} - 2\hat{v}_{j-1}$. Then by Lemma D.2, on the domain $W_{v_{j+1}, s_{j+1}, \varepsilon_{j+1}}$, we fined a real-analytic transformation $\Phi_i = X_{\phi_i}$, which conjugates H_i to

$$H_{i+1} = h + g + \bar{g}_{i+1} + f_{i+1}$$

where $\bar{g}_{i+1} = \bar{g}_i + \Pi_{\mathfrak{L} \cap \mathfrak{K}} f_i = \sum_{k=0}^i \Pi_{\mathfrak{L} \cap \mathfrak{Z}} T_K f_k$. We prove that (D.151) is satisfied for $j = i+1$. Using³ the assumption on the right in (D.146), (D.148), the inequality for Ks in (D.149) and the definition of δ_i in (D.150), we have

$$\|\{\bar{g}_i, \phi_i\}\|_{v_i - \hat{v}_i, s_i - \hat{s}_i, \varepsilon_i - \hat{\varepsilon}_i} \leq \left[\frac{c}{\mathfrak{a}\delta_i} \left(E_1 + \frac{E}{L} \right) \right] E_i \leq \left[\frac{c}{\mathfrak{a}\delta_i} \frac{E}{Ks} + \frac{c}{\mathfrak{a}\delta_i} \frac{E}{L} \right] E_i < \frac{E_i}{32}.$$

Moreover, by a similar argument as in (D.147) and since g is actually real-analytic in the larger domain

$$W_{\bar{v}, s, \varepsilon} \supset W_{\bar{v}_i - \hat{v}_i + \bar{v}, s_i - \hat{s}_i + \bar{s}, \varepsilon_i - \hat{\varepsilon}_i + \bar{\varepsilon}},$$

we have

$$\|\{g, \phi_i\}\|_{v_i - \hat{v}_i, s_i - \hat{s}_i, \varepsilon_i - \hat{\varepsilon}_i} = \|\{g, \phi_i\}\|_{\bar{v}_i - \hat{v}_i, s_i - \hat{s}_i, \varepsilon_i - \hat{\varepsilon}_i} \leq \frac{\bar{c}E_i G}{\mathfrak{a}\delta_i L} < \frac{E_i}{64},$$

where

$$\bar{\delta}_i := \min\{\hat{r}_i \hat{s}_i, \bar{\rho}_i \hat{\rho}'_i\} = \frac{\bar{d}}{16L^2}, \quad i = 1, \dots, L.$$

Then we find⁴

$$\begin{aligned} E_{i+1} = \|f_{i+1}\|_{v_{i+1}, s_{i+1}, \varepsilon_{i+1}} &\leq \left(1 - \frac{cE_i}{\mathfrak{a}\delta_1}\right)^{-1} \left[\frac{cE_i}{\mathfrak{a}\delta_1} + e^{-K\hat{s}_i} + \left(\frac{\varepsilon_i - \hat{\varepsilon}_i}{\varepsilon_i}\right)^K \right] E_i \\ &+ \left(1 - \frac{cE_i}{\mathfrak{a}\delta_1}\right)^{-1} \|\{\bar{g}_i, \phi_i\}\|_{v_i - \hat{v}_i, s_i - \hat{s}_i, \varepsilon_i - \hat{\varepsilon}_i} \\ &+ \left(1 - \frac{cE_i}{\mathfrak{a}\delta_1}\right)^{-1} \|\{g, \phi_i\}\|_{\bar{v}_i - \hat{v}_i, s_i - \hat{s}_i, \varepsilon_i - \hat{\varepsilon}_i} \\ &\leq \frac{64}{63} \left[\frac{1}{64} + \frac{1}{8} + \left(\frac{4}{7}\right)^{16} + \frac{1}{32} + \frac{1}{64} \right] E_i \\ &< \frac{E_i}{4} < E_1 < 2^{-6} \frac{\mathfrak{a}\delta_1}{c}. \end{aligned}$$

since $i \geq 1$. Then we let $\Phi := \Phi_0 \circ \dots \circ \Phi_L, H_* := H \circ \Phi = h + g + \bar{g}_{L+1} + f_{L+1}, g_* := g_{L+1}, f_* := f_{L+1}$ and we have, by telescopic inequalities and (D.148),

$$\begin{aligned} \|g_* - \Pi_{\mathfrak{L} \cap \mathfrak{K}} T_K f\|_{v/6, s/6, \varepsilon/6} &= \sum_{i=1}^L \|\Pi_{\mathfrak{L} \cap \mathfrak{K}} T_K f_i\| \leq \sum_{i=1}^L E_i \leq E_1 \sum_{i=1}^L \frac{1}{4^{i-1}} \\ &= \frac{4}{3} E_1 \leq \left(\frac{2^7 cE}{\mathfrak{a}d} + \frac{2^7 \bar{c}G}{\mathfrak{a}d} \right) E \end{aligned}$$

³For the proof of inequality $\|\{g_i, \phi_i\}\|_{v_i - \hat{v}_i, s_i - \hat{s}_i, \varepsilon_i - \hat{\varepsilon}_i} \leq \frac{cE_i}{\mathfrak{a}\delta_1} \left(E_1 + \frac{E}{L} \right)$, compare [31, Proof of the Normal Form Lemma].

⁴Since $K > 8L$ and $L \geq 2$, one has $(1 - \frac{3}{2L})^K \leq \frac{1}{(1 + \frac{3}{2L})^{8L}}$ with the r.h.s bounded above by $(4/7)^{16}$ (it decreases to e^{-12} as $L \rightarrow +\infty$).

Now we prove (D.143). Let $F \in W_{\bar{v},s,\varepsilon}$, $F_{-1} := F$, $F_i := F \circ \Phi_0 \circ \dots \circ \Phi_i$, $i = 0, \dots, L$. Then

$$\begin{aligned} \|F \circ \Phi - F\|_{\bar{v}/6,s/6,\varepsilon/6} &= \|F_L - F\|_{\bar{v}_{L+1},s_{L+1},\varepsilon_{L+1}} \\ &\leq \sum_{i=0}^L \|F_{i-1} \circ \Phi_i - F_{i-1}\|_{\bar{v}_{i+1},s_{i+1},\varepsilon_{i+1}} \\ &\leq \sum_{i=0}^L \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i} \|F\|_{\bar{v}_i,s_i,\varepsilon_i} \leq \frac{\sum_{i=0}^L \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i}}{\prod_{i=0}^L (1 - \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i})} \|F\|_{\bar{v},s,\varepsilon} \\ &\leq \sum_{i=0}^L \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i} e^{\frac{5}{4} \sum_{i=0}^L \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i}} \|F\|_{\bar{v},s,\varepsilon} \leq \frac{2^5 \bar{c}E_0 Ks}{\mathfrak{a}d} \|F\|_{\bar{v},s,\varepsilon} \end{aligned}$$

where we have used $\frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i} < 1/24$ that, for $0 \leq x \leq 1/24$, $\log(1-x)^{-1} < \frac{5}{4}x$ and

$$\begin{aligned} \sum_{i=0}^L \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i} &= \frac{\bar{c}E_0}{\mathfrak{a}\bar{\delta}_0} + \sum_{i=1}^L \frac{\bar{c}E_i}{\mathfrak{a}\bar{\delta}_i} \leq \frac{2^6 \bar{c}E_0}{\mathfrak{a}d} + \frac{\bar{c}E_1}{\mathfrak{a}\bar{\delta}_1} \sum_{i=1}^L \frac{1}{4^{i-1}} \\ &\leq \frac{2^6 \bar{c}E_0}{\mathfrak{a}d} + \frac{4}{3} \frac{\bar{c}E_1}{\mathfrak{a}\bar{\delta}_1} < \frac{2^4 \bar{c}E_0 Ks}{\mathfrak{a}d}. \end{aligned}$$

The proof for $F \in W_{v,s,\varepsilon}$ is similar. \square

PROOF OF PROPOSITION D.1. For simplicity of notations, we prove Proposition D.1 in the case $\nu = \ell = 1$; the generalization to any ν, ℓ being straightforward. Consider the Hamiltonian

$$H_0(u_1, \varphi, p, q) := h(I, p, q) + f_1(u_1, \varphi, p, q), \quad (u_1, \varphi, p, q) \in W_{v_1,s,\varepsilon}^{(1)}.$$

To this Hamiltonian let us apply Lemma D.1, with $g \equiv 0$, so as to conjugate it to

$$H_1 := H_0 \circ \Phi_1 = h + g_1 + f_{*1}^{(1)}, \quad (u_1, \varphi, p, q) \in W_{v_1/6,s/6,\varepsilon/6}^{(1)}$$

where $g_1, f_{*1}^{(1)}$ correspond to g_*, f_* , hence satisfy

$$\begin{aligned} \|f_{*1}^{(1)}\|_{v_1/6,s/6,\varepsilon/6} &\leq e^{-Ks/6} \|f_1^{(i)}\|_{v_1,s,\varepsilon} \\ \|g_1\|_{v_1/6,s/6,\varepsilon/6} &\leq \frac{7}{6} \|f_1\|_{v_1,s,\varepsilon} \\ \|g_1 - \Pi_{\mathcal{L} \cap \mathcal{I}^3} T_K f_1\|_{v_1/6,s/6,\varepsilon/6} &\leq \frac{2^7 c_1 \|f_1\|_{v_1,s,\varepsilon}^2}{\mathfrak{a}d_1} \end{aligned}$$

Then we have

$$H^{(1)}(u, \varphi, p, q) := H \circ \Phi_1 = H_0 \circ \Phi_1 + \sum_{j=2}^N f_j \circ \Phi_1 = h + g_1 + f_{*1}^{(1)} + \sum_{j=2}^N f_j^{(1)}$$

where $f_j^{(1)} := f_j \circ \Phi_1$. Assume, inductively, that, for some $1 \leq i \leq N-1$ and any $1 \leq j \leq i$ we have conjugated H to

$$H^{(j)}(u, \varphi, p, q) = H \circ \Phi_1 \circ \dots \circ \Phi_j = h + \sum_{k=1}^j g_k + \sum_{k=1}^j f_{k*}^{(j)} + \sum_{k=j+1}^N f_k^{(j)}$$

where

$$\Phi_j : W_{v/6^j,s/6^j,\varepsilon/6^j}^{(j)} \rightarrow W_{v/6^{j-1},s/6^{j-1},\varepsilon/6^{j-1}}^{(j-1)}$$

transforms

$$H_{j-1} := h + \sum_{k=1}^{j-1} g_k + f_j^{(j-1)}$$

into

$$H_{j-1} \circ \Phi_j = h + \sum_{k=1}^j g_k + f_{*j}^{(j)}.$$

The Hamiltonian

$$H_i(u_{i+1}, \varphi, p, q) := h + \sum_{k=1}^i g_k(u_k, \varphi, p, q) + f_{i+1}^{(i)}(u_{i+1}, \varphi, p, q)$$

is real-analytic for $(u_{i+1}, \varphi, p, q) \in W_{v_{i+1}/6^i, s/6^i, \varepsilon/6^i}^{(i+1)}$ and satisfies the assumptions of Lemma D.1, with $\bar{N} = i$. Then one can find $\Phi_{i+1} : W_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}}^{(i+1)} \rightarrow W_{v_{i+1}/6^i, s/6^i, \varepsilon/6^i}^{(i+1)}$ such that $H_i \circ \Phi_{i+1} = h + \sum_{k=1}^{i+1} g_k + f_{*i+1}^{(i+1)}$, where

$$\begin{aligned} & \|f_{*i+1}^{(i+1)}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \leq e^{-Ks/6^{i+1}} \|f_{i+1}^{(i)}\|_{v_{i+1}/6^i, s/6^i, \varepsilon/6^i} \\ & \leq \left(\frac{9}{8}\right)^i e^{-Ks/6^{i+1}} \|f_{i+1}\|_{v_{i+1}, s, \varepsilon} \\ & \|g_{i+1}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \leq \frac{7}{6} \|f_{i+1}^{(i)}\|_{v_{i+1}/6^i, s/6^i, \varepsilon/6^i} \leq \frac{7}{6} \left(\frac{9}{8}\right)^i \|f_{i+1}\|_{v_{i+1}, s, \varepsilon} \\ & \|g_{i+1} - \Pi_{\mathfrak{L}\cap\mathfrak{N}} T_K f_{i+1}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \leq \|g_{i+1} - \Pi_{\mathfrak{L}\cap\mathfrak{N}} T_K f_{i+1}^{(i)}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \quad + \|\Pi_{\mathfrak{L}\cap\mathfrak{N}} T_K f_{i+1}^{(i)} - \Pi_{\mathfrak{L}\cap\mathfrak{N}} T_K f_{i+1}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \leq \|g_{i+1} - \Pi_{\mathfrak{L}\cap\mathfrak{N}} T_K f_{i+1}^{(i)}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \quad + \|f_{i+1}^{(i)} - f_{i+1}\|_{v_{i+1}/6^{i+1}, s/6^{i+1}, \varepsilon/6^{i+1}} \\ & \leq \left(\frac{9}{8}\right)^{2i} \frac{2^7 c_{i+1} \|f_{i+1}\|_{v_{i+1}, s, \varepsilon}^2}{\mathfrak{ad}_{i+1}} \\ & \quad + \frac{7}{6} \left(\frac{9}{8}\right)^{2i} \sum_{j=1}^i \frac{2^7 c_j \|f_j\|_{v_j, s, \varepsilon}}{\mathfrak{ad}_j} \|f_{i+1}\|_{v_{i+1}, s, \varepsilon} \\ & \quad + \sum_{k=1}^i \left(\frac{9}{8}\right)^{i-k} \frac{2^4 c_k \|f_k\|_{v_k, s, \varepsilon} Ks}{\mathfrak{ad}_k} \|f_{i+1}\|_{v_{i+1}, s, \varepsilon} \end{aligned}$$

with $f_{k*}^{(i+1)} := f_{k*}^{(i)} \circ \Phi_{i+1}$ for $1 \leq k \leq i+1$ and $f_k^{(i+1)} := f_k^{(i)} \circ \Phi_{i+1}$ for $i+2 \leq k \leq N$. Then we find

$$\begin{aligned} H^{(i+1)} &:= H^{(i)} \circ \Phi_{i+1} = \left(h + \sum_{k=1}^i g_k + \sum_{k=1}^i f_{k*}^{(i)} + \sum_{k=i+1}^N f_k^{(i)} \right) \circ \Phi_{i+1} \\ &= H_i \circ \Phi_{i+1} + \left(\sum_{k=1}^i f_{k*}^{(i)} + \sum_{k=i+2}^N f_k^{(i)} \right) \circ \Phi_{i+1} \\ &= h + \sum_{k=1}^{i+1} g_k + \sum_{k=1}^{i+1} f_{k*}^{(i+1)} + \sum_{k=i+2}^N f_k^{(i+1)} \end{aligned}$$

and hence, after N steps,

$$H^{(N)} := H \circ \Phi_1 \cdots \circ \Phi_N = h + \sum_{k=1}^N g_k + \sum_{k=1}^{i+1} f_{k*}^{(N)}$$

satisfies the thesis of Proposition D.1. \square

D.2. A slightly-perturbed integrable system

The following result is well known in the literature of close-to be integrable systems, hence its proof is omitted. Note that it deals with an integrable system, close to another integrable one.

THEOREM D.1. *One can find a number \mathbf{c}_0 such that, for any real-analytic, one-dimensional, system*

$$H(P, Q) = h\left(\frac{P^2 + Q^2}{2}\right) + f(P, Q) \quad (P, Q) \in \mathfrak{B} = B_\varepsilon^2(0) \subset \mathbb{C}^2$$

and any $0 < \bar{\varepsilon} < \varepsilon$, such that

$$(D.152) \quad \inf_{B_\varepsilon^2} |\partial h| \geq \mathbf{a}, \quad \sup_{B_\varepsilon^2} |f| \leq \mathbf{e}, \quad \frac{1}{\mathbf{c}_0} \frac{\mathbf{e}}{\mathbf{a}\bar{\varepsilon}^2} < 1,$$

one can find a real-analytic transformation

$$\phi_* : (P_*, Q_*) \in B_{\varepsilon - \bar{\varepsilon}}^2 \rightarrow (P, Q) \in B_\varepsilon^2$$

which conjugates H to a function $H_* = H \circ \phi_*$ depending only on $\frac{P_*^2 + Q_*^2}{2}$. The assertion can be extended to the case that h, f are functions of other canonical coordinates (P', Q', y, x) , depending on them only via $Y = (y, \frac{P_1'^2 + Q_1'^2}{2}, \dots, \frac{P_m'^2 + Q_m'^2}{2})$, with $y \in Y_\rho$, $(P'_j, Q'_j) \in B_{\varepsilon'_j}^2$. In this case, letting $(P_*, Q_*) \rightarrow \phi_*(P_*, Q_*; Y)$ the transformation obtained for any fixed value of Y , there exists a canonical, real-analytic, transformation Φ_* of the form

$$\Phi_* : (P, Q) = \phi_*(P_*, Q_*; Y_*) \quad y = y_*, \quad x = x_* + \varphi(Y_*), \quad P'_j + iQ'_j = e^{i\psi_j(Y_*)} (P'_{*j} + iQ'_{*j})$$

which conjugates H to a function $H_* = H \circ \Phi_*$ depending only on $\frac{P_*^2 + Q_*^2}{2}$ and Y_* . In this case, the functions φ_j, ψ_j verify

$$|\varphi_j| \leq \frac{1}{\mathbf{c}_0} \frac{\mathbf{e}}{\mathbf{a}\rho_j}, \quad |\psi_j| \leq \frac{1}{\mathbf{c}_0} \frac{\mathbf{e}}{\mathbf{a}\varepsilon_j'^2}.$$

More on the geometrical structure of the \mathcal{P} -coordinates, compared to Deprit's coordinates

In this chapter we aim to point out differences and similarities between the \mathcal{P} -coordinates and the coordinates denoted as $(\Psi, \Gamma, \Lambda, \psi, \gamma, \ell)$ in [7, 9, 27].

We recall that the “planetary” coordinates $(\Psi, \Gamma, \Lambda, \psi, \gamma, \ell)$ may be derived (after a canonical transformation) from a more general set of canonical coordinates studied by A. Deprit. In their planetary form, the coordinates $(\Psi, \Gamma, \Lambda, \psi, \gamma, \ell)$ have been rediscovered¹ by the author during her PhD, under the strong motivation of their application to the planetary problem [9, 27].

Let us recall their definition², in the spirit of *Kepler maps* (Definition 2.2).

Let $C_{\mathcal{E}}^{(i)}, S_{\mathcal{E}}^{(i)}$ be as in (2.8) of Chapter 2 and define the $\mathcal{D}ep$ -nodes

$$(E.153) \quad n_i := \begin{cases} k^{(3)} \times S_{\mathcal{E}}^{(1)} & i = 0 \\ S_{\mathcal{E}}^{(i)} \times S_{\mathcal{E}}^{(i+1)} = -S_{\mathcal{E}}^{(i)} \times C_{\mathcal{E}}^{(i)} & i = 1, \dots, n-1. \\ -n_{n-1} & i = n \end{cases}$$

Then let

$$\mathcal{E}_{\mathcal{D}ep} := \{((\mathfrak{E}_1, \dots, \mathfrak{E}_n) \subset E^3 \times \dots \times E^3) : 0 < e_i < 1, \quad n_{i-1} \neq 0 \quad \forall i = 1, \dots, n\}.$$

On $\mathcal{E}_{\mathcal{D}ep}$, define the map

$$\tau_{\mathcal{D}ep}^{-1} : (\mathfrak{E}_1, \dots, \mathfrak{E}_n) \in \mathcal{E}_{\mathcal{D}ep} \rightarrow X_{\mathcal{D}ep} \in \mathfrak{X}_{\mathcal{D}ep} = \tau_{\mathcal{D}ep}^{-1}(\mathcal{E}_{\mathcal{D}ep})$$

where

$$X_{\mathcal{D}ep} = (\Psi, \Gamma, \Lambda, \psi, \gamma) \in \mathbb{R}^n \times \mathbb{R}_+^n \times \mathbb{R}_+^n \times \mathbb{T}^n \times \mathbb{T}^n$$

where

$$\Psi = (\Psi_{-1}, \Psi_0, \bar{\Psi}) \in \mathbb{R}_+ \times \mathbb{R}_+ \times \mathbb{R}_+^{n-2} \quad \psi = (\psi_{-1}, \psi_0, \bar{\psi}) \in \mathbb{T} \times \mathbb{T} \times \mathbb{T}^{n-2}$$

$$\Gamma = (\Gamma_1, \dots, \Gamma_n) \in \mathbb{R}_+^n \quad \gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{T}^n$$

$$\Lambda = (\Lambda_1, \dots, \Lambda_n) \in \mathbb{R}_+^n$$

with

$$\bar{\Psi} = (\Psi_1, \dots, \Psi_{n-2}) \quad \bar{\psi} = (\psi_1, \dots, \psi_{n-2})$$

¹The proof of their symplectic character found in [27] has been published in [7]. Another proof has been given in [36].

²For sake of uniformity, we use slightly different notations with respect to the ones in [7], actually closer to the ones of the paper [12]).

are defined as follows. The coordinates Λ_j are as in (2.11), while $(\Psi, \Gamma, \psi, \gamma)$ are defined as

$$(E.154) \quad \begin{aligned} \Psi_{i-2} &= \begin{cases} Z := S_{\mathcal{E}}^{(1)} \cdot k^{(3)} \\ |S_{\mathcal{E}}^{(i)}| \end{cases} & \psi_{i-2} &= \begin{cases} \zeta := \alpha_{k^{(3)}}(k^{(1)}, n_0) & i = 1 \\ \alpha_{S_{\mathcal{E}}^{(i-1)}}(n_{i-2}, n_{i-1}) & 2 \leq i \leq n \end{cases} \\ \Gamma_i &:= |C_{\mathcal{E}}^{(i)}| & \gamma_i &:= \alpha_{C_{\mathcal{E}}^{(i)}}(n_i, P^{(i)}) & 1 \leq i \leq n \end{aligned}$$

Then $\tau_{\mathcal{D}ep}^{-1}$ is a bijection [7, 12, 27, 36].

DEFINITION E.1. We call *Deprit's map*, or *Dep map*, the Kepler map

$$\mathcal{D}ep : \quad \text{Dep} = (X_{\mathcal{D}ep}, \ell) \in \mathcal{D}_{\mathcal{D}ep} = \mathfrak{X}_{\mathcal{D}ep} \times \mathbb{T}^n \rightarrow (y, x) \in \mathbb{R}^{3n} \times \mathbb{R}^{3n}$$

associated to $\tau_{\mathcal{D}ep}$.

Comparing \mathcal{P} and $\mathcal{D}ep$

a) Both the \mathcal{P} and $\mathcal{D}ep$ -coordinates reduce the system to $(3n-2)$ degrees of freedom. They share the following three coordinates (two actions and an angle)

$$\Psi_{-1} = Z = \Theta_0, \quad \psi_{-1} = \zeta = \vartheta_0, \quad \Psi_0 = G = \chi_0$$

which are integrals of the system. As a consequence, the coordinates (Z, ζ) and, respectively,

$$\mathfrak{g} := \psi_0, \quad \mathfrak{g} := \kappa_0$$

do not appear into the Hamiltonian. Note that $\mathcal{D}ep$ and \mathcal{P} share also the fixed node $n_0 = \nu_1$.

b) The angle \mathfrak{g} for the set $\mathcal{D}ep$ describes the motion of the node n_1 in (E.153) and, by the cyclic character of \mathfrak{g} , this motion is negligible. Its counterpart in the set \mathcal{P} is the node n_1 in (2.10), the negligible motion of which is governed by \mathfrak{g} .

c) Compare the diagrams in (2.20) and (2.21) with the two ones associated to the $\mathcal{D}ep$ -map, respectively:

$$\begin{array}{cccccc}
& n_0 & n_1 & \vdots & n_{n-2} & n_{n-1} \\
& \uparrow & \uparrow & \vdots & \uparrow & \uparrow \\
k^{(3)} & \rightarrow S_{\mathcal{E}}^{(1)} & \rightarrow S_{\mathcal{E}}^{(2)} & \rightarrow \cdots & \rightarrow S_{\mathcal{E}}^{(n-1)} & \rightarrow S_{\mathcal{E}}^{(n)} = C_{\mathcal{E}}^{(n)} \\
& \downarrow & \downarrow & \vdots & \downarrow & \\
& C_{\mathcal{E}}^{(1)} & C_{\mathcal{E}}^{(2)} & \vdots & C_{\mathcal{E}}^{(n-1)} & \\
& \Downarrow & \Downarrow & \vdots & \Downarrow & \\
& -n_1 & -n_2 & \vdots & -n_{n-1} &
\end{array}$$

and

$$\begin{array}{cccccc}
F_0 & \rightarrow & F_1^* & \rightarrow & \cdots & \rightarrow & F_i^* & \rightarrow & \cdots & \rightarrow & F_n^* = G_n^* \\
& & \downarrow & & \vdots & & \downarrow & & \vdots & & \downarrow \\
& & G_1^* & & & & G_i^* & & & & G_n^*
\end{array}$$

where

$$F_i^* = (n_{i-1}, \cdot, S_{\mathcal{E}}^{(i)}) \quad G_i^* = (-n_i, \cdot, C_{\mathcal{E}}^{(i)}) \quad i = 1, \dots, n.$$

Note that, analogously to (2.20), n_i in (E.153) is the skew-product of its two previous vectors in the tree (2.20).

d) While $\mathcal{D}ep$ is not defined for the planar problem, \mathcal{P} is, and, in that case, the coordinates $(\Theta, \chi, \vartheta, \kappa)$ in (2.11) reduce to³

$$\Theta_i = \begin{cases} \chi_0 \\ 0 \end{cases} \quad \vartheta_i = \begin{cases} 0 \\ \pi \end{cases} \quad \kappa_i = \begin{cases} \arg P^{(1)} - \frac{\pi}{2} \\ \widehat{P^{(i)}P^{(i+1)}} + \pi \end{cases} \quad \begin{matrix} i = 0 \\ i = 1, \dots, n-1 \end{matrix}$$

$$\chi_i = \sum_{j=i+1}^n \|C_{\mathcal{E}}^{(j)}\|$$

while the (Λ, ℓ) remain unchanged.

e) The \mathcal{P} -map is singular when some eccentricity e_i vanishes or some of the following relations hold

$$S_{\mathcal{E}}^{(1)} \parallel k^{(3)} \quad P^{(i)} \parallel S_{\mathcal{E}}^{(i)} \quad S_{\mathcal{E}}^{(i+1)} \parallel P^{(i)}.$$

³Here by “planar case” we mean $C_{\mathcal{E}}^{(1)} \parallel \cdots \parallel C_{\mathcal{E}}^{(n)} \parallel k^{(3)}$. Note that, to be more precise, ϑ_0 and κ_0 would not exist in that case (since $\nu_1 = 0$). However, since they are both cyclic angles, we can fix them to an arbitrary value. The choice above corresponds to replace ν_1 with $k^{(1)}$.

The former of such relations is negligible, while the other ones have no physical meaning. Therefore, the only physically relevant singularities of \mathcal{P} are for zero-eccentric motions.

The $\mathcal{D}ep$ -map is singular when some eccentricity e_i vanishes or some of the following relations hold

$$S_{\mathcal{E}}^{(1)} \parallel k^{(3)} \quad S_{\mathcal{E}}^{(i+1)} \parallel S_{\mathcal{E}}^{(i)} \quad i = 1, \dots, n-1.$$

The configurations $S_{\mathcal{E}}^{(i)} \parallel S_{\mathcal{E}}^{(i+1)}$ have a relevant physical meaning, since the planar case corresponds to the intersection of all such configurations. A complete regularization of *all* the singularities of the $\mathcal{D}ep$ -map has been obtained in [9, 27], which allowed to overcome the problem of the *rotational degeneracy* (see [8] for information) of the planetary problem and to construct the Brkhoff normal form of it. It works at expenses of one extra-degree of freedom.

f) The Euclidean lengths $\|C_{\mathcal{E}}^{(i)}\|$ of the planets' angular momenta are the actions Γ_i among $\mathcal{D}ep$ -coordinates: see (E.154). In terms of the \mathcal{P} -coordinates they have more involved expressions in (2.17). As mentioned in the previous item, this makes more difficult regularizing singular configurations with zero eccentricity. The formula simplifies in the planar case:

$$\|C_{\mathcal{E}}^{(i)}\| = \begin{cases} |\chi_{i-1} - \chi_i| & i = 1, \dots, n-1 \\ \chi_{n-1} & i = n \end{cases}$$

where $|w| := \sqrt{w^2}$, for a given $w \in \mathbb{C}$.

g) Reflections are not well described in the framework of the $\mathcal{D}ep$ -reduction: Compare, e.g., [29, Section 4.4]. Instead, in the framework of the \mathcal{P} -reduction, the transformation

$$(\bar{\Theta}, \bar{\vartheta}) \rightarrow (-\bar{\Theta}, 2k\pi - \bar{\vartheta}) \quad k \in \mathbb{Z}^{n-1}$$

corresponds to changing the sign of the second component of any $y^{(i)}$ and any $x^{(i)}$. Therefore, any of the points

$$(\bar{\Theta}, \bar{\vartheta}) = (0, k\pi) \quad k \in \mathbb{Z}^{n-1}$$

is an equilibrium point for the Hamiltonian, corresponding to a co-planar configuration. Compare Proposition 2.2.

Bibliography

- [1] V. I. Arnol'd, *Proof of a theorem of A. N. Kolmogorov on the preservation of conditionally periodic motions under a small perturbation of the Hamiltonian* (Russian), *Uspehi Mat. Nauk* **18** (1963), no. 5 (113), 13–40. MR0163025
- [2] V. I. Arnol'd, *Small denominators and problems of stability of motion in classical and celestial mechanics* (Russian), *Uspehi Mat. Nauk* **18** (1963), no. 6 (114), 91–192. MR0170705
- [3] F. Boigey, *Élimination des nœuds dans le problème newtonien des quatre corps* (French, with English summary), *Celestial Mech.* **27** (1982), no. 4, 399–414, DOI 10.1007/BF01228562. MR670321
- [4] A. Celletti and G. Pinzari, *Four classical methods for determining planetary elliptic elements: a comparison*, *Celestial Mech. Dynam. Astronom.* **93** (2005), no. 1-4, 1–52, DOI 10.1007/s10569-005-8663-8. MR2186799
- [5] L. Chierchia. The Planetary N-Body Problem. *UNESCO Encyclopedia of Life Support Systems*, 6.119.55, 2012.
- [6] L. Chierchia and G. Pinzari, *Properly-degenerate KAM theory (following V. I. Arnold)*, *Discrete Contin. Dyn. Syst. Ser. S* **3** (2010), no. 4, 545–578, DOI 10.3934/dcdss.2010.3.545. MR2684064
- [7] L. Chierchia and G. Pinzari, *Deprit's reduction of the nodes revisited*, *Celestial Mech. Dynam. Astronom.* **109** (2011), no. 3, 285–301, DOI 10.1007/s10569-010-9329-8. MR2777135
- [8] L. Chierchia and G. Pinzari, *Planetary Birkhoff normal forms*, *J. Mod. Dyn.* **5** (2011), no. 4, 623–664. MR2903753
- [9] L. Chierchia and G. Pinzari, *The planetary N-body problem: symplectic foliation, reductions and invariant tori*, *Invent. Math.* **186** (2011), no. 1, 1–77, DOI 10.1007/s00222-011-0313-z. MR2836051
- [10] L. Chierchia and G. Pinzari. Metric stability of the planetary n-body problem. *Proceedings of the International Congress of Mathematicians*, 2014.
- [11] A. Delshams, V. Kaloshin, A. de la Rosa, and T. M. Seara. Global instability in the elliptic restricted three body problem. *arXiv: 1501.01214*, 2015.
- [12] A. Deprit, *Elimination of the nodes in problems of n bodies*, *Celestial Mech.* **30** (1983), no. 2, 181–195, DOI 10.1007/BF01234305. MR711657
- [13] J. Féjóz. Work in progress.
- [14] J. Féjóz, *Démonstration du 'théorème d'Arnold' sur la stabilité du système planétaire (d'après Herman)* (French, with English and French summaries), *Ergodic Theory Dynam. Systems* **24** (2004), no. 5, 1521–1582, DOI 10.1017/S0143385704000410. MR2104595
- [15] J. Féjóz, *On action-angle coordinates and the Poincaré coordinates*, *Regul. Chaotic Dyn.* **18** (2013), no. 6, 703–718, DOI 10.1134/S1560354713060105. MR3146588
- [16] J. Féjóz. On "Arnold's theorem" in celestial mechanics -a summary with an appendix on the poincaré coordinates. *Discrete and Continuous Dynamical Systems*, 33:3555-3565, 2013.
- [17] J. Féjóz, M. Guardia, V. Kaloshin, and P. Roldan. Kirkwood gaps and diffusion along mean motion resonances in the restricted planar three body problem. *J. Eur. Math. Soc.*, 2014.
- [18] S. Ferrer and C. Osácar, *Harrington's Hamiltonian in the stellar problem of three bodies: reductions, relative equilibria and bifurcations*, *Celestial Mech. Dynam. Astronom.* **58** (1994), no. 3, 245–275, DOI 10.1007/BF00691977. MR1271458
- [19] R. S. Harrington. The stellar three-body problem. *Celestial Mech. and Dyn. Astronomy*, 1(2):200-209, 1969.
- [20] M. R. Herman. Torsion du problème planétaire, edited by J. Féjóz in 2009. Available in the electronic 'Archives Michel Herman' at http://www.college-de-france.fr/default/EN/all/equ_dif/archives_michel_herman.htm.

- [21] C. G. J. Jacobi. Sur l'élimination des noeuds dans le problème des trois corps. *Astronomische Nachrichten*, Bd XX:81-102, 1842.
- [22] A. N. Kolmogorov, *On conservation of conditionally periodic motions for a small change in Hamilton's function* (Russian), *Dokl. Akad. Nauk SSSR (N.S.)* **98** (1954), 527–530. MR0068687
- [23] J. Laskar and P. Robutel, *Stability of the planetary three-body problem. I. Expansion of the planetary Hamiltonian*, *Celestial Mech. Dynam. Astronom.* **62** (1995), no. 3, 193–217, DOI 10.1007/BF00692088. MR1364477
- [24] T. Levi-Civita. Sopra la equazione di Kepler. *Astronomische Nachrichten*, 165(20):313-314, 1904.
- [25] F. Malige, P. Robutel, and J. Laskar, *Partial reduction in the n -body planetary problem using the angular momentum integral*, *Celestial Mech. Dynam. Astronom.* **84** (2002), no. 3, 283–316, DOI 10.1023/A:1020392219443. MR1935017
- [26] J. Moser, *On invariant curves of area-preserving mappings of an annulus*, *Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. II* **1962** (1962), 1–20. MR0147741
- [27] G. Pinzari. *On the Kolmogorov set for many-body problems*. PhD thesis, Università Roma Tre, April 2009.
- [28] G. Pinzari, *Aspects of the planetary Birkhoff normal form*, *Regul. Chaotic Dyn.* **18** (2013), no. 6, 860–906, DOI 10.1134/S1560354713060178. MR3146595
- [29] G. Pinzari, *Canonical coordinates for the planetary problem*, *Acta Appl. Math.* **137** (2015), 205–232, DOI 10.1007/s10440-014-9996-7. MR3343379
- [30] H. Poincaré. *Les méthodes nouvelles de la mécanique céleste*. Gauthier-Villars, Paris, 1892.
- [31] J. Pöschel, *Nekhoroshev estimates for quasi-convex Hamiltonian systems*, *Math. Z.* **213** (1993), no. 2, 187–216, DOI 10.1007/BF03025718. MR1221713
- [32] R. Radau, *Sur une transformation des équations différentielles de la dynamique* (French), *Ann. Sci. École Norm. Sup.* **5** (1868), 311–375. MR1508550
- [33] P. Robutel, *Stability of the planetary three-body problem. II. KAM theory and existence of quasiperiodic motions*, *Celestial Mech. Dynam. Astronom.* **62** (1995), no. 3, 219–261, DOI 10.1007/BF00692089. MR1364478
- [34] H. Rüssmann, *Invariant tori in non-degenerate nearly integrable Hamiltonian systems*, *Regul. Chaotic Dyn.* **6** (2001), no. 2, 119–204, DOI 10.1070/RD2001v006n02ABEH000169. MR1843664
- [35] F. Tisserand. *Traité de mécanique céleste*. Gauthier-Villars, I, 1889-1896
- [36] L. Zhao, *Partial reduction and Delaunay/Deprit variables*, *Celestial Mech. Dynam. Astronom.* **120** (2014), no. 4, 423–432, DOI 10.1007/s10569-014-9584-1. MR3277249

Editorial Information

To be published in the *Memoirs*, a paper must be correct, new, nontrivial, and significant. Further, it must be well written and of interest to a substantial number of mathematicians. Piecemeal results, such as an inconclusive step toward an unproved major theorem or a minor variation on a known result, are in general not acceptable for publication.

Papers appearing in *Memoirs* are generally at least 80 and not more than 200 published pages in length. Papers less than 80 or more than 200 published pages require the approval of the Managing Editor of the Transactions/Memoirs Editorial Board. Published pages are the same size as those generated in the style files provided for $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$ or $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$.

Information on the backlog for this journal can be found on the AMS website starting from <http://www.ams.org/memo>.

A Consent to Publish is required before we can begin processing your paper. After a paper is accepted for publication, the Providence office will send a Consent to Publish and Copyright Agreement to all authors of the paper. By submitting a paper to the *Memoirs*, authors certify that the results have not been submitted to nor are they under consideration for publication by another journal, conference proceedings, or similar publication.

Information for Authors

Memoirs is an author-prepared publication. Once formatted for print and on-line publication, articles will be published as is with the addition of AMS-prepared frontmatter and backmatter. Articles are not copyedited; however, confirmation copy will be sent to the authors.

Initial submission. The AMS uses Centralized Manuscript Processing for initial submissions. Authors should submit a PDF file using the Initial Manuscript Submission form found at www.ams.org/submission/memo, or send one copy of the manuscript to the following address: Centralized Manuscript Processing, MEMOIRS OF THE AMS, 201 Charles Street, Providence, RI 02904-2294 USA. If a paper copy is being forwarded to the AMS, indicate that it is for *Memoirs* and include the name of the corresponding author, contact information such as email address or mailing address, and the name of an appropriate Editor to review the paper (see the list of Editors below).

The paper must contain a *descriptive title* and an *abstract* that summarizes the article in language suitable for workers in the general field (algebra, analysis, etc.). The *descriptive title* should be short, but informative; useless or vague phrases such as “some remarks about” or “concerning” should be avoided. The *abstract* should be at least one complete sentence, and at most 300 words. Included with the footnotes to the paper should be the 2010 *Mathematics Subject Classification* representing the primary and secondary subjects of the article. The classifications are accessible from www.ams.org/msc/. The Mathematics Subject Classification footnote may be followed by a list of *key words and phrases* describing the subject matter of the article and taken from it. Journal abbreviations used in bibliographies are listed in the latest *Mathematical Reviews* annual index. The series abbreviations are also accessible from www.ams.org/msnhtml/serials.pdf. To help in preparing and verifying references, the AMS offers MR Lookup, a Reference Tool for Linking, at www.ams.org/mrlookup/.

Electronically prepared manuscripts. The AMS encourages electronically prepared manuscripts, with a strong preference for $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$. To this end, the Society has prepared $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$ author packages for each AMS publication. Author packages include instructions for preparing electronic manuscripts, samples, and a style file that generates the particular design specifications of that publication series. Though $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{T}\mathcal{E}\mathcal{X}$ is the highly preferred format of $\mathcal{T}\mathcal{E}\mathcal{X}$, author packages are also available in $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$.

Authors may retrieve an author package for *Memoirs of the AMS* from www.ams.org/journals/memo/memoauthorpac.html. The *AMS Author Handbook* is available in PDF format from the author package link. The author package can also be obtained free

of charge by sending email to tech-support@ams.org or from the Publication Division, American Mathematical Society, 201 Charles St., Providence, RI 02904-2294, USA. When requesting an author package, please specify $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{L}^{\mathcal{A}}\mathcal{T}\mathcal{E}\mathcal{X}$ or $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\mathcal{T}\mathcal{E}\mathcal{X}$ and the publication in which your paper will appear. Please be sure to include your complete mailing address.

After acceptance. The source files for the final version of the electronic manuscript should be sent to the Providence office immediately after the paper has been accepted for publication. The author should also submit a PDF of the final version of the paper to the editor, who will forward a copy to the Providence office.

Accepted electronically prepared files can be submitted via the web at www.ams.org/submit-book-journal/, sent via FTP, or sent on CD to the Electronic Prepress Department, American Mathematical Society, 201 Charles Street, Providence, RI 02904-2294 USA. $\mathcal{T}\mathcal{E}\mathcal{X}$ source files and graphic files can be transferred over the Internet by FTP to the Internet node [ftp.ams.org](ftp://ftp.ams.org) (130.44.1.100). When sending a manuscript electronically via CD, please be sure to include a message indicating that the paper is for the *Memoirs*.

Electronic graphics. Comprehensive instructions on preparing graphics are available at www.ams.org/authors/journals.html. A few of the major requirements are given here.

Submit files for graphics as EPS (Encapsulated PostScript) files. This includes graphics originated via a graphics application as well as scanned photographs or other computer-generated images. If this is not possible, TIFF files are acceptable as long as they can be opened in Adobe Photoshop or Illustrator.

Authors using graphics packages for the creation of electronic art should also avoid the use of any lines thinner than 0.5 points in width. Many graphics packages allow the user to specify a “hairline” for a very thin line. Hairlines often look acceptable when proofed on a typical laser printer. However, when produced on a high-resolution laser imagesetter, hairlines become nearly invisible and will be lost entirely in the final printing process.

Screens should be set to values between 15% and 85%. Screens which fall outside of this range are too light or too dark to print correctly. Variations of screens within a graphic should be no less than 10%.

Any graphics created in color will be rendered in grayscale for the printed version unless color printing is authorized by the Managing Editor and the Publisher. In general, color graphics will appear in color in the online version.

Inquiries. Any inquiries concerning a paper that has been accepted for publication should be sent to memo-query@ams.org or directly to the Electronic Prepress Department, American Mathematical Society, 201 Charles St., Providence, RI 02904-2294 USA.

Editors

This journal is designed particularly for long research papers, normally at least 80 pages in length, and groups of cognate papers in pure and applied mathematics. Papers intended for publication in the *Memoirs* should be addressed to one of the following editors. The AMS uses Centralized Manuscript Processing for initial submissions to AMS journals. Authors should follow instructions listed on the Initial Submission page found at www.ams.org/memo/memosubmit.html.

1. GEOMETRY, TOPOLOGY & LOGIC

Coordinating Editor: Richard Canary, Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-1043 USA; e-mail: canary@umich.edu

Algebraic topology, Michael Hill, Department of Mathematics, University of California Los Angeles, Los Angeles, CA 90095 USA; e-mail: mikehill@math.ucla.edu

Differential geometry, Chiu-Chu Melissa Liu, Department of Mathematics, Columbia University, New York, NY 10027 USA; e-mail: ccliu@math.columbia.edu

Logic, Noam Greenberg, School of Mathematics and Statistics, Victoria University of Wellington, Wellington 6140, New Zealand; e-mail: greenberg@msor.vuw.ac.nz

Low-dimensional topology and geometric structures, Richard Canary

2. ALGEBRA AND NUMBER THEORY

Coordinating Editor: Henri Darmon, Department of Mathematics, McGill University, Montreal, Quebec H3A 0G4, Canada; e-mail: darmon@math.mcgill.ca

Algebra, Michael Larsen, Department of Mathematics, Rawles Hall, Indiana University, 831 E 3rd St., Bloomington, IN 47405 USA; e-mail: mjlarsen@indiana.edu

Algebraic geometry, Lucia Caporaso, Department of Mathematics and Physics, Roma Tre University, Largo San Leonardo Murialdo, I-00146 Rome, Italy; e-mail: LCedit@mat.uniroma3.it

Arithmetic geometry, Ted C. Chinburg, Department of Mathematics, University of Pennsylvania, Philadelphia, PA 19104-6395 USA; e-mail: ted@math.upenn.edu

Commutative algebra, Irena Peeva, Department of Mathematics, Cornell University, Ithaca, NY 14853 USA; e-mail: irena@math.cornell.edu

Number theory, Henri Darmon

3. GEOMETRIC ANALYSIS & PDE

Coordinating Editor: Tatiana Toro, Department of Mathematics, University of Washington, Box 354350, Seattle, WA 98195-4350 USA; e-mail: toro@uw.edu

Geometric analysis, Tatiana Toro

Harmonic analysis and partial differential equations, Monica Visan, Department of Mathematics, University of California Los Angeles, 520 Portola Plaza, Los Angeles, CA 90095 USA; e-mail: visan@math.ucla.edu

Partial differential equations and functional analysis, Alexander A. Kiselev, Department of Mathematics, MS-136, Rice University, 6100 Main Street, Houston, TX 77005 USA; e-mail: kiselev@rice.edu

Real analysis and partial differential equations, Wilhelm Schlag, Department of Mathematics, The University of Chicago, 5734 South University Avenue, Chicago, IL 60637 USA; e-mail: schlag@math.uchicago.edu

4. ERGODIC THEORY, DYNAMICAL SYSTEMS & COMBINATORICS

Coordinating Editor: Vitaly Bergelson, Department of Mathematics, Ohio State University, 231 W. 18th Avenue, Columbus, OH 43210 USA; e-mail: vitaly@math.ohio-state.edu

Algebraic and enumerative combinatorics, Jim Haglund, Department of Mathematics, University of Pennsylvania, Philadelphia, PA 19104 USA; e-mail: jhaglund@math.upenn.edu

Probability theory, Robin Pemantle, Department of Mathematics, University of Pennsylvania, 209 S. 33rd Street, Philadelphia, PA 19104 USA; e-mail: pemantle@math.upenn.edu

Dynamical systems and ergodic theory, Ian Melbourne, Mathematics Institute, University of Warwick, Coventry CV4 7AL, United Kingdom; e-mail: I.Melbourne@warwick.ac.uk

Ergodic theory and combinatorics, Vitaly Bergelson

5. ANALYSIS, LIE THEORY & PROBABILITY

Coordinating Editor: Stefaan Vaes, Department of Mathematics, Katholieke Universiteit Leuven, Celestijnenlaan 200B, B-3001 Leuven, Belgium; e-mail: stefaan.vaes@wis.kuleuven.be

Functional analysis and operator algebras, Stefaan Vaes

Harmonic analysis and complex analysis, Malabika Pramanik, Department of Mathematics, University of British Columbia, 1984 Mathematics Road, Vancouver, British Columbia V6T 1Z2, Canada; e-mail: malabika@math.ubc.ca

Langlands conjectures, Marie-France Vigneras, 8 Rue des Ecoles, 75005 Paris, France; e-mail: marie-france.vigneras@imj-prg.fr

Probability and statistics, Patrick J. Fitzsimmons, Department of Mathematics, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0112 USA; e-mail: pfitzsim@ucsd.edu

All other communications to the editors, should be addressed to the Managing Editor, ALE-JANDRO ADEM, Department of Mathematics, The University of British Columbia, Room 121, 1984 Mathematics Road, Vancouver, B.C., Canada V6T 1Z2; e-mail: adem@math.ubc.ca

SELECTED PUBLISHED TITLES IN THIS SERIES

- 1212 **Roelof Bruggeman, Youngju Choie, and Nikolaos Diamantis**, Holomorphic Automorphic Forms and Cohomology, 2018
- 1211 **Shouhei Honda**, Elliptic PDEs on Compact Ricci Limit Spaces and Applications, 2018
- 1210 **Zhou Gang, Dan Knopf, and Israel Michael Sigal**, Neckpinch Dynamics for Asymmetric Surfaces Evolving by Mean Curvature Flow, 2018
- 1209 **Cristian Anghel, Iustin Coandă, and Nicolae Manolache**, Globally Generated Vector Bundles with Small c_1 on Projective Spaces, 2018
- 1208 **Matthew Harrison-Trainor**, Degree Spectra of Relations on a Cone, 2018
- 1207 **Alastair J. Litterick**, On Non-Generic Finite Subgroups of Exceptional Algebraic Groups, 2018
- 1206 **Anne-Laure Dalibard and Laure Saint-Raymond**, Mathematical Study of Degenerate Boundary Layers: A Large Scale Ocean Circulation Problem, 2018
- 1205 **Charles Collot**, Type II Blow Up Manifolds for the Energy Supercritical Semilinear Wave Equation, 2018
- 1204 **Rui Palma**, Crossed Products by Hecke Pairs, 2018
- 1203 **Xiao Xiong, Quanhua Xu, and Zhi Yin**, Sobolev, Besov and Triebel-Lizorkin Spaces on Quantum Tori, 2018
- 1202 **Naiara V. de Paulo and Pedro A. S. Salomão**, Systems of Transversal Sections Near Critical Energy Levels of Hamiltonian Systems in \mathbb{R}^4 , 2018
- 1201 **Bernhelm Booß-Bavnbek and Chaofeng Zhu**, The Maslov Index in Symplectic Banach Spaces, 2018
- 1200 **Francis Nier**, Boundary Conditions and Subelliptic Estimates for Geometric Kramers-Fokker-Planck Operators on Manifolds with Boundaries, 2018
- 1199 **Ramon Antoine, Francesc Perera, and Hannes Thiel**, Tensor Products and Regularity Properties of Cuntz Semigroups, 2018
- 1198 **Colette Mœglin and J.-L. Waldspurger**, La Formule des Traces Locale Tordue, 2018
- 1197 **Stefano Bianchini and Sara Daneri**, On Sudakov's Type Decomposition of Transference Plans with Norm Costs, 2018
- 1196 **Nicola Gigli**, Nonsmooth Differential Geometry—An Approach Tailored for Spaces with Ricci Curvature Bounded from Below, 2018
- 1195 **Pablo Shmerkin and Ville Suomala**, Spatially Independent Martingales, Intersections, and Applications, 2018
- 1194 **A. M. Mason and N. C. Snaith**, Orthogonal and Symplectic n -level Densities, 2018
- 1193 **James Damon and Ellen Gasparovic**, Medial/Skeletal Linking Structures for Multi-Region Configurations, 2017
- 1192 **R. Lawther**, Maximal Abelian Sets of Roots, 2017
- 1191 **Ben Webster**, Knot Invariants and Higher Representation Theory, 2017
- 1190 **Agelos Georgakopoulos**, The Planar Cubic Cayley Graphs, 2017
- 1189 **John McCuan**, The Stability of Cylindrical Pendant Drops, 2017
- 1188 **Aaron Hoffman, Hermen Hupkes, and E. S. Van Vleck**, Entire Solutions for Bistable Lattice Differential Equations with Obstacles, 2017
- 1187 **Denis R. Hirschfeldt, Karen Lange, and Richard A. Shore**, Induction, Bounding, Weak Combinatorial Principles, and the Homogeneous Model Theorem, 2017
- 1186 **Mikhail Ershov, Andrei Jaikin-Zapirain, and Martin Kassabov**, Property (T) for Groups Graded by Root Systems, 2017
- 1185 **Jörg-Uwe Löbus**, Absolute Continuity Under Time Shift of Trajectories and Related Stochastic Calculus, 2017

For a complete list of titles in this series, visit the
AMS Bookstore at www.ams.org/bookstore/memoseries/.

ISBN 978-1-4704-4102-9



MEMO/255/1218

