Estimation and model-based combination of causality networks among large US Banks and Insurance companies*

Giovani Bonaccolto[†] Massimiliano Caporin [‡] Roberto Panzica [§]

This version: July 9, 2018

Abstract

Causality is a widely-used concept in theoretical and empirical economics. The recent financial economics literature has used Granger causality to detect the presence of contemporaneous links between financial institutions and, in turn, to obtain a network structure. Subsequent studies combined the estimated networks with traditional pricing or risk measurement models to improve their fit to empirical data. In this paper, we provide two contributions: we show how to use a linear factor model as a device for estimating a combination of several networks that monitor the links across variables from different viewpoints; and we demonstrate that Granger causality should be combined with quantile-based causality when the focus is on risk propagation. The empirical evidence supports the latter claim.

Keywords: Granger causality; quantile causality; multi-layer network; network combination **J.E.L.** codes: C58, C31, C32, G01

^{*}We gratefully acknowledge research support from the Research Center SAFE, funded by the State of Hessen initiative for research LOEWE. We thank for the comments provided by the participants of the Computational and Financial Econometrics conference (CFE) held in Sevilla in December 2016, the Italian Conference of Econometrics and Empirical Economics held in Messina in January 2017. The second author acknowledge financial support from the Global Risk Institute in Financial Services and the Louis Bachelier Institute under the project Systemic Risk, the Studies in Nonlinear Dynamics and Econometrics Conference held in Paris in March 2017, the seminar at the Complutense University of Madrid held in February 2017

[†]University of Enna, Italy

[‡]University of Padova, Italy; Corresponding author: Department of Statistical Sciences, Via C. Battisti 241, Padova, Italy, massimiliano.caporin@unipd.it

[§]Goethe University Frankfurt, Germany

Estimation and model-based combination of causality networks among large US Banks and Insurance companies

July 9, 2018

Abstract

Causality is a widely-used concept in theoretical and empirical economics. The recent financial economics literature has used Granger causality to detect the presence of contemporaneous links between financial institutions and, in turn, to obtain a network structure. Subsequent studies combined the estimated networks with traditional pricing or risk measurement models to improve their fit to empirical data. In this paper, we provide two contributions: we show how to use a linear factor model as a device for estimating a combination of several networks that monitor the links across variables from different viewpoints; and we demonstrate that Granger causality should be combined with quantile-based causality when the focus is on risk propagation. The empirical evidence supports the latter claim.

Keywords: Granger causality; quantile causality; multi-layer network; network combination **J.E.L.** codes: C58, C31, C32, G01

1 Introduction

A growing strand of the financial economics literature has investigated the role of links between financial institutions, which serve as channels along which shocks spread through the financial system, so this strand of the financial economics literature is related to studies on financial contagion. The relationships between financial institutions may vary in nature and, as a consequence, the way in which we interpret these links may differ, depending on which type of relations are of interest to us, or which type of link we are monitoring or measuring. As examples, we cite the input-output relationship (where firms use the output of other companies as input, in their own production function), the ownership relationship (companies that hold assets of other companies), or the links measured on the grounds of stock market prices, such as causality-based connections.

The early literature on financial networks focused first on ascertaining the channel through which financial contagion was spreading within the financial system. The main objective was therefore to identify the networks by using appropriate criteria to detect the links between institutions. Following such an approach meant that the networks were, by definition, unique. Acemoglu et al. (2012) used a network structure based on input-output relationships to show that aggregate fluctuations may originate from microeconomic shocks to firms. Kelly et al. (2013), on the other hand, showed how stock firm volatility relates to customer-supplier connectedness. Billio et al. (2014) used contingent claim analysis and network measures to underscore the links between sovereign, banks and insurances. There is a constantly growing number of works proposing competing or alternative approaches for estimating the networks existing between groups of financial institutions, markets, countries and (not necessarily financial) assets. Among the many contributions in this area, we refer to Billio et al. (2012), Diebold and Yilmaz (2014, 2015), Hautsch et al. (2012, 2013, 2014, 2015), Barigozzi and Brownlees (2014), Ozdagli and Weber (2015), and Corsi et al. (2015), which have in common that they all refer to a financial or economic playground.

Despite the growing interest in the financial network topic, and the increasing understanding that there might be different channels over which financial contagion spreads, the literature on combinations of financial or economic networks is still very limited. In fact, the different approaches taken to estimating networks enable different potential channels for the spread of contagion or risk to come to light that might co-exist within the same financial system. It is therefore fundamentally important to allow for the possibility of combining them to obtain a more complete picture of risk spreading within a financial system.

This paper provides a possible solution based on a multiplex network, or a collection of networks (called layers) existing between a set of subjects, that we formally define in Section 2. Ideally, the constituents of the multiplex network represent the outcomes of different approaches to the estimation or identification of links between the subjects analyzed. Postulating the existence of a multiplex network, we take a step forward and consider the combination of the information contained in the various layers of the network for the purpose of analyzing their impact within a financial economic model. In this case, we focus on the linear factor model augmented with network dependence introduced by Billio et al. (2016). We show how the model can be generalized to accommodate the presence of a multiplex network, and we provide a model

parametrization that offers two useful interpretations of the models estimation output. First, the model parameters will enable the statistical significance of the information contained in each multiplex network layer to emerge. This will highlight the relevance of the various approaches to network measurement/identification within the model, with consequences on the financial interpretation associated with the model. These analyses might be relevant for policy purposes and market monitoring, as they may identify the main risk-spreading channel.

Second, the estimated model parameters will reveal a composite network obtained by combining the multiplex network layers. The model-based combination will thus provide an overall picture of the links existing between the variables analyzed, as measured using the different approaches (i.e. those behind the various layers of the network), as well as accounting for their respective relevance.

The model-based combination of networks is the first main contribution provided by the present paper. The second contribution that we make to the financial network literature concerns the estimation of networks. As previously mentioned, several authors have already discussed possible ways to obtain a network of financial assets. Among the various methods, the Granger causality approach of Billio et al. (2012) is taken as our starting point: Granger causality is used within a collection of bivariate Vector Auto Regressive (VAR) models to identify the statistically significant links across the modelled variables; then the network is used to identify the shock propagation channels and establish the systemic relevance of financial companies.

When focusing on the risk dimension, Granger causality within a VAR model detects mean causality whether it is on the financial contagion, or on the diffusion of systemic shocks, whereas we normally focus on variances or, more generally, on systemic and/or systematic risk measures. We consequently do not believe that using Granger causality will produce the most appropriate picture for interpreting the risk of the estimated financial networks. This prompts us to discuss a set of alternative approaches to estimating financial networks based on causality relationships that go beyond the classical Granger causality. We survey a collection of methods that have a common denominator: they all resort to the quantile regression approach of Koenker and Bassett (1978), in either a parametric or a non-parametric representation. Using these methods, we will show how we can identify causality among quantiles of the modelled variables. These quantile-based causality detection methods will lead to the estimation of causality networks mimicking the approach of Billio et al. (2012) but oriented towards a risk dimension. This is immediate when focusing on left quantiles of a random variable. As a further approach, we will also show that the non-parametric quantile causality test of Jeong et al. (2012) represents a further alternative to traditional Granger causality for building causality networks. It is important to highlight that, although the implemented methods focus on the same object, that is, a given τ -th quantile of the response variable, they differ in capturing the causal impact of the explanatory variable and, therefore, provide a different source of information. In particular, in traditional quantile regression (Koenker and Bassett (1978), see Section A.2) we detect the linear impact of the explanatory variable without weighting its support, i.e. using its entire distribution. In the non-parametric causality test (Jeong et al. (2012), see Section A.4) we allow for a non-linear impact. Finally, in the quantile-on-quantile case (see Section A.3) we allow for linear impact on the quantile of the response variable but focusing attention in a neighborhood

of the θ -th quantile of the explanatory variable.

We stress that, by focusing on the quantiles of the modelled variables, we attach a clear risk interpretation to the estimated networks: on the one hand, the left quantiles are closely related to traditional risk measures such as the Value-at-Risk and the Expected Shortfall; on the other, the networks will monitor the spreading of shocks or the links between the modelled variables when they are in a high-risk state, and not when they are in a mean (or median) state, as captured by Granger causality. Our purpose is consistent with two other recent contributions, from Hong et al. (2009) and Corsi et al. (2015), that focus on causality among tail events. The Granger causality in risk in Hong et al. (2009), and also used by Corsi et al. (2015), is based on the occurrence of tail events and detects the possible causality among such events. We might associate the tail with a conditional quantile and estimate it with quantile regressions. We differ from the previous papers in that we focus on the causality within the assessment of the conditional quantiles, not on the occurrence of tail events. We thus consider the causality on the risk measures rather than the causality on the occurrence of extreme events.

We suggest different approaches to constructing causality networks, which involve monitoring the links across assets with views that might complement the (mean-based) Granger causality. These competing networks will represent the input for our model-based network combination. Using our model, we will be able to identify the relevance and the role of the various causality networks.

We empirically validate our two main proposals concerning the use of quantile causality to infer the network structure across a set of (financial) variables, and the model-based combination of causality networks. By using different datasets (US industrial portfolio returns, and a set of large banks and insurance companies), we first provide evidence of the different network structures that we can estimate from Granger causality and quantile causality. We show how the networks differ across methods and over two different samples relating to the global financial crisis (2006-2008) and to the years 2011-2015. Our results suggest that quantile causality networks are denser than Granger causality networks, a finding of relevance to systemic risk interpretation because a denser network is indicative of a much larger set of links, and thus a possibly greater systemic effect of shocks. The networks based on the non-parametric quantile causality test of Jeong et al. (2012) also have a clear core-periphery structure, which might help us to identify the more systemically relevant companies or sectors. Moreover, by resorting to the model-based combination approach, we show how different networks have a different impact and relevance on the various datasets, and we estimate a composite network. The model parameters indicate that quantile causality networks are much more relevant than Granger causality. In addition, the linear factor model augmented with a plurality of networks provides residuals that are much less correlated (on average) than traditional linear factor models, or the network augmented linear factor model of Billio et al. (2015), which is based on the Granger causality network alone. This latter finding might be implicated in explaining why the idiosyncratic risk is priced in the cross section. Since the quantile-based networks are the more relevant, it comes as no surprise that the composite network has a structure much closer to that of the quantile causality networks.

The paper proceeds as follows. Section 2 discusses the approach for obtaining a model-based

composite network, extending the model of Billio et al. (2015). Section 3 presents the various approaches for estimating a causality network starting from the Granger causality method and then moving to quantile-based causality detection. Section 4 provides the empirical evidence with respect to both the estimation of quantile causality networks and the model-based combination of networks. This section also includes some robustness checks. Section 5 concludes. A Web Appendix accompanies the paper, which contains further tables and graphs, and some methodological notes.

2 Model-based network combination

In this section, we introduce how we obtain a model-based combination of a collection of networks. At this stage, we assume that all the networks are available. We therefore do not estimate the structure of the various networks, but concentrate only on their optimal combination. We start with a formal definition of a simple network and of a multiplex or multilayer network.

A network or graph G = (V, E) is a collection of vertexes V and edges E, where the edges represent the links between the vertexes,¹ with $E \subseteq (V \times V)$. Networks are generally represented by using the adjacency matrix W, a binary matrix where each element $w_{i,j}$ can take only two values, 1 and 0. When $w_{i,j}$ is 1, the node j is linked to node i, with an information flow from i to j. A value of zero identifies the absence of a link. We do not consider self-loops, so the diagonal of the W matrix is identically equal to zero. If W is symmetric, we associate with each node a measure called degree, that counts the number of edges for the node concerned. Note that, when focusing on K assets, W has a dimension $K \times K$.

The W matrix might not be symmetric because we can have a link from i to j, but the opposite might not hold. If the matrix W is symmetric, the associated network is undirected as the information contained in the direction of the links becomes redundant; in this case, edges represent mutual relationships between nodes. If W is not symmetric, the network is directed and there may be two edges between a pair of nodes; in this case, unlike the undirected case, we distinguish between indegree and outdegree. In particular, outdegree is the number of outgoing links starting from a given node, while indegree is the sum of the ingoing links on that node. If we relax the assumption of only two possible values for $w_{i,j}$ and allow for values within the (positive) real axis, the network becomes weighted (as opposed to unweighted) because the connection also has a size/relevance. In unweighted networks, each edge simply represents the presence of a connection, while in weighted networks edges also contain information on the strength of the relation between nodes.

As an example, the matrix in (1) represents the directed and unweighted network existing between a set of 5 nodes. Figure 1 provides a graphical representation of the network.

¹The terms *vertex* and *nodes* are equivalent, and both are used interchangeably in this work. In the same way, *edges* and *links* take on the same meaning.




Figure 1: Network associated to the Matrix W

$$W = \begin{bmatrix} 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}. \tag{1}$$

It is worth noting the parallelism between the adjacency matrix and the spatial (proximity) matrix used in spatial econometrics. The latter describes how the subjects of an analysis are spatially related. In such a different framework, the distance between subjects is (generally) measured on a physical (natural) playground, and connections correspond to neighborhood relationships. Therefore, if the element $w_{i,j}$ equals 1, the element j is a neighbor of the element i, following the spatial econometric nomenclature. The spatial matrices in the standard application are symmetric because neighboring relationships are symmetric. It is also common practice to focus on proximity matrices that are row-normalized. These matrices can therefore be interpreted as directed weighted networks, given that the symmetrical relations might become asymmetrical after normalization.

Within a financial framework, networks are generally directed and weighted when they are measured by means of balance sheet quantities. Billio et al. (2016) used BIS cross-holdings, Diebold and Yilmaz (2014) developed an approach based on variance decompositions of target series, Acemoglu et al. (2012) used the Input-Output matrix. The network might also be directed and unweighted like those obtained using Granger causality in Billio et al. (2012), or Granger Causality in the tails Corsi et al. (2015), or it might even be unweighted and undirected as in the economic sector case of Caporin and Paruolo (2015).

In the following, we assume that the strength of the edges is normalized at each node, as in a row normalization of the adjacency matrix W. We also assume that we are dealing with directed networks (the most general case that we could consider).

Assuming that we have d different networks, the multiplex (or multilayer) network is a collection of the d networks organized into d different planes, or layers. In principle, the number of nodes needs not be the same across networks, so the total number of nodes in a multilayer network equals $\sum_{i=1}^{d} V_i$, where V_i is the number of nodes or vertexes in the i-th layer.

Assuming that the graphs (associated with the networks) are simple (there are no loops), the maximum number of edges (connections between nodes) equals $\prod_{i=1}^{d} \frac{V_i(V_i-1)}{2}$, since connections across layers are also allowed.

In a financial framework, the multilayer network has an even simpler structure: the nodes are the same across layers because we have measures of connections over, say, the same financial institutions, but we use different approaches to estimate the networks. We also assume that there are no connections between layers, leaving us a collection of networks that are unrelated, but all referring to the same nodes.

The literature on multiplex networks is relatively recent and, to the best of our knowledge, only a few papers combine networks in finance, focusing particularly on contagion and topological properties. Since institutions are linked in very different ways, it is useful to find a way to process all this information. Bargigli et al. (2015) analyzed the Italian interbank market using a multiplex network, distinguishing between five layers: unsecured overnight transactions, unsecured short-term transactions (up to 12 months), unsecured long-term transactions, secured short-term transactions and secured long-term transactions; they also considered the aggregated network, computed as the accumulation of all the layers. According to the authors, each layer had a different topological property and persistence over time and, among all the layers, only the overnight market was representative of the aggregate network.

Léon et al. (2014) studied the Colombian sovereign security market and built a multiplex network with three different layers corresponding to the three Colombian environments trading and registering sovereign securities. They found a strong nonlinear effect for the aggregate network value, attributable to the fact that the principal layer did not transfer its properties to the aggregate network or multiplex.

Montagna and Kok (2013) developed an agent-based multi-layered interbank network model using a sample of large EU banks, highlighting that there were non-negligible non-linearity effects influencing shock propagation to the individual banks. In other words, when different layers were considered simultaneously, the contagion effect was larger than the sum of the contagion-induced losses when the network layers were analyzed separately.

Molina-Balboa et al. (2014) analyze the multiplex structure of the bank exposures within the Mexican banking system. By associating layers with different financial instruments and the authors provides insight on the interdependence among banks with relevant implications from a systemic risk perspective.

The present contribution aims to combine financial networks by building on a variant of a financial economic model, the linear factor model augmented with a spatial link following the approach of Billio et al. (2016).

The advantage of our model lies in that the combination of the various layers of the multiplex network is weighted with a set of coefficients appearing in the models parametrization. As well as representing the elements for a convex linear combination of the layers, the coefficients would consequently enable the significance and relevance of each layer to be assessed.

In the next subsection we briefly review the model of Billio et al. (2016), and introduce the models generalization in the presence of a multilayer network. Then we discuss the approach used to derive a composite network from the models estimates.

2.1 A linear factor model with multilayer spatial dependence

To obtain a composite network, we generalize the model of Billio et al. (2016), a revisited multifactor model capable of taking into account the contemporaneous links between the assets (as measured by a network) together with the presence of common factors. In this model, the returns equation equals

$$A(R_t - \mathbb{E}[R_t]) = \bar{\beta}F_t + \eta_t \tag{2}$$

where A is the matrix of the contemporaneous relationships between endogenous variables, i.e. the returns, coexisting with the exposure to a set of common factors F_t ; R_t , $\mathbb{E}[R_t]$ are, respectively, the asset returns and the expected asset returns, $\bar{\beta}$ is the structural asset exposure to the common factor and η_t are the structural idiosyncratic residuals with a diagonal covariance matrix Σ_{η} .

The standard multifactor model can be seen as the reduced form of equation (2). Under the invertibility of matrix A, the model has the following reduced-form representation

$$R_t = \mathbb{E}[R_t] + A^{-1}\bar{\beta}F_t + A^{-1}\eta_t$$

= $\mathbb{E}[R_t] + \beta^*F_t + \epsilon_t^*$ (3)

Starting from the reduced form, and in the presence of contemporaneous links across assets, Billio et al. (2016) make the point that: i) the reduced-form residuals $\epsilon_t^{\star} = A^{-1}\eta_t$ are cross-correlated; and ii) the reduced-form betas, $\beta^{\star} = A^{-1}\bar{\beta}$, are a nonlinear function of the structural betas and of the contemporaneous relation matrix A.

The intuition of Billio et al. (2016) is to provide a structure for A driven by the existence of a network representing the (contemporaneous) links across the assets. They therefore propose to make the matrix A an affine function of a network, $A = (I - \rho W)$, where ρ is a scalar parameter indicating the spatial dependence of the returns on the network, while W is an exogenous spatial matrix, or adjacency matrix of a network. Such a choice also enables us to cope with the identification issues of the simultaneous equation system in (2).

Following Anselin (1988) and LeSage and Pace (2009), the model thus includes a spatial autoregressive component. The estimation of the spatial parameter ρ follows from concentrated likelihood methods.

Billio et al. (2016) also allow for a more flexible parametrization of the A matrix. In particular, they consider two relevant extensions:

• the use of asset-specific impacts from the network, which corresponds to a contemporaneous parameter matrix A with equation-specific network impacts, namely $A = I - \mathcal{R}W$,

where $\mathcal{R} = diag(\rho_1, \rho_2, \dots, \rho_K)$ is a diagonal matrix, and the coefficient ρ_i represents the impact of the network on each asset; and

• the possibility of the spatial matrix, or network, varying in time, thus leading to W_t ; this is particularly relevant in finance because W is not related to a *physical* distance between subjects, but measures proximity or connections starting from variables liable to temporal differences.

We refer the reader to the paper by Billio et al. (2016) for further details on the models interpretation.

In the present paper, for the sake of simplicity, we assume the networks are time-invariant, though the approach that we introduce can easily be extended to the case of dynamic networks. In the presence of a multilayer network, we generalize the model of Billio et al. (2016) to accommodate the presence of several adjacency matrices.

In the spatial econometrics literature, we have some examples associated with the existence of richer spatial dynamics. In particular, Brandsma and Ketellapper (1979) propose a higher-order spatial autoregressive model, introducing two spatial matrices and consequently two spatial autocorrelation parameters: the first matrix based on the first-order neighbors, the second derived from the higher-order neighboring relations.

In the framework of Billio et al. (2016), the matrix of simultaneous relations A, in the presence of many networks, is easily generalized to

$$A = I - \sum_{j=1}^{d} \rho_j W_j. \tag{4}$$

In equation (4), we have d different scalar coefficients, ρ_j , capturing the impact of the network W_j on the contemporaneous links across the assets.

Such a parametrization would give rise to a common impact of each network on the whole collection of modelled asset returns, so we propose to combine the existence of many networks with a restricted parametrization, accommodating the presence of the various networks, while also allowing for heterogeneous impacts of the networks on the assets. In detail, we suggest the following model specification:

$$\left[I - \mathcal{R}\left(\sum_{j=1}^{d} \delta_{j} W_{j}\right)\right] R_{t} = AR_{t} = \mathbb{E}\left[R_{t}\right] + \beta F_{t} + \eta_{t}$$
(5)

where d is the number of layers or networks, \mathcal{R} is a diagonal matrix, and the δ_j coefficients satisfy the following constraints:

$$\sum_{j=1}^{m} \delta_j = 1, \quad \delta_j \ge 0, j = 1, 2, \dots m.$$
 (6)

The parameter δ_j controls the impact and role of each layer of the multiplex network, while the parameters in \mathcal{R} determine the heterogeneous impact of the networks on the asset returns.

We suggest estimating the model using concentrated maximum likelihood, according to standard practice in spatial econometrics. If matrix A is known, the factor loadings β and the innovation covariance matrix Σ_{η} can be obtained with traditional least square estimators. We can thus concentrate out the latter parameters and use concentrated maximum likelihood to estimate the parameters entering A. The maximization must account for the constraints that we impose on the δ_j parameters.

The model contains a sort of spatial autoregression, the structure of which is driven by several spatial proximity matrices. A standard requirement for estimations of spatial econometric models is the invertibility of matrix A. When the number of networks is restricted to one, or d=1, the non-singularity of A^{-1} is ensured when $\lambda_{min}^{-1} < \rho < \lambda_{max}^{-1}$, where λ_{max} and λ_{min} are, respectively, the maximum and minimum eigenvalues of the spatial/adjacency matrix W.

In the presence of numerous proximity matrices, the non-singularity of A might be addressed with analytical tools, but only in specific cases. For instance, Lee and Liu (2010) and Elhorst et al. (2012) discuss the case of two matrices W, referring to the estimation of a second-order spatial autoregressive model. These results are not easy to extend to our specification, however, we control for the invertibility of A within the estimation step.

The coexistence of many networks in the augmented linear factor model of Billio et al. (2016) has consequences on the identification of the models parameters too, particularly for those in \mathcal{R} .

The identification of the parameters entering the matrix A is guaranteed by the constraints on the δ_j parameters, the row normalization in the W_j matrices, and a further set of necessary (but simple and intuitive) conditions on the adjacency matrices:

- i. $W_i \neq 0, \quad j = 1, 2, \dots, d;$
- ii. $W_i \neq W_i$, i, j = 1, 2, ..., d, $i \neq j$;
- iii. $\rho(W_{i,.}) > 0$ where $\rho(\mathcal{A})$ is the rank of matrix \mathcal{A} , $W_{i,.} = [W'_1|_{.,i} \quad W'_2|_{.,i} \quad \dots \quad W'_d|_{.,i}]$ and $W'_i|_{.,i}$ denotes the i-th row of W_i .

Condition [i] simply rules out the case of absence of edges between a collection of nodes. Condition [ii] excludes the case where two networks, measured by two different approaches are identical. If the latter case realize, we loose identification as the two networks are indistinguishable, and one of the two must be excluded from the analysis. The first two conditions are relevant for the identification of the δ_j parameters. Condition [iii] refers to the parameters in \mathcal{R} , and ensures that in each row of the combined network

$$W^* = \sum_{j=1}^d \delta_j W_j. \tag{7}$$

we do have at least a non-null element. The parameter $\rho_{i,i}$ of the diagonal matrix \mathcal{R} is identified if the i-th row of W^* has at least one non-null element. In fact, if a given row of W^* contains only null elements, the corresponding parameter in \mathcal{R} is not identified. Such a case corresponds to asset i having no spatial dependence on the other assets in the model, so we set the corresponding parameters in \mathcal{R} to zero in the case of null rows in W^* .

Finally, we note that the model is overidentified, given that the covariance matrix of the innovation, Σ_{η} , is diagonal, thus leaving space for any richer parametrizations in (5), with heterogeneous impacts on returns from groups of layers, for instance.

2.2 Relevance and combination of networks

The model offers two important insights. The first, and most relevant, concerns the feasibility of interpreting the summation of the adjacency matrices, matrix W^* , as a new adjacency matrix representing a composite network. This is a direct consequence of the constraints on the coefficients leading to a convex linear combination of adjacency matrices, which are all rownormalized. The composite matrix will therefore also be a row-normalized matrix representing a combination of a collection of networks. The combination will also be directed and weighted because the initial networks are directed and weighted. The coefficients δ_j measure the relevance of each network layer in obtaining the composite network. The model thus generates both a composite network and an insight on the impact of each approach used to construct the various layers.

Secondly, the parameters included in \mathcal{R} represent the heterogeneous reaction of the assets to the composite network, thus making the multiple-network model comparable, in terms of its economic interpretation, to the network of Billio et al. (2016), but with the important difference that its underlying network is composite.

2.3 Causality testing and causality networks

The concept of causality and the tools used to test for its existence date back to the seminal contributions of Granger (1969, 1980, 1988). The purpose of his original contribution was to identify causal relationships in mean across economic variables, and his work attracted considerable interest in the econometrics literature, see Geweke (1984) and Hoover (2001). The concept was later extended to the analysis of causality among variances: Granger et al. (1986) provided a first definition of variance causality, while Comte and Lieberman (2000) extended and generalized the concept. On the testing side, Cheung and Ng (1996) proposed a test based on cross-correlations, Hafner (2003) introduced a procedure based on the likelihood of competing models, and Hafner and Herwartz (2008) presented a Wald-type approach.

More recently, there have been contributions on causality between two random variables that go well beyond the first- and second-order moments. We refer in particular to causality among quantiles such as the non-parametric test of Jeong et al. (2012) and the work of Lee and Yang (2014), and to the work of Candelon and Topkavi (2016) on causality in distributions.

Within a financial framework, the interest in causality and the use of Granger causality tests to estimate a network stems from the work done by Billio et al. (2012). The authors associated the statistical evidence of Granger causality in the mean with the existence of an edge connecting two assets (two nodes) of a financial network. In times of market turmoil, however, or when our interest lies in estimating systemic risk spreading after local or specific shocks, the causality in the mean might not be enough, and should be combined with a causality testing that covers the risk dimension.

Appendix A provides a brief review of the Granger causality approach adopted by Billio et al. (2012), then suggesst a number of competing methods we might consider for testing the existence of causality among quantiles of return distributions. In particular, the Appendix discusses the methods we will then use in the empirical analyses: the already mentioned approach of Billio et al. (2012) based on Granger causality; a Granger-type causality approach building on the baseline Quantile Regression (see Koenker (2005)); a generalization of the latter accounting also for the location of the conditioning variable over its support, that Sim and Zhou (2015) call Quantile-on-Quantile; finally, the non-parametric quantile causality test introduced by Jeong et al. (2012).

As discussed in the introduction, the lower quantiles of returns acquire a more relevant role within a financial framework, because they correspond to Value-at-Risk levels. The availability of a number of competing methods for estimating causality networks enables the construction of a multiplex network, in which each layer is associated with a specific network. The multiplex network based on causality links can then be used for the model-based construction of composite networks, as discussed in the previous section.

The same criteria apply to the estimation of a single causality network, irrespective of the approach used to identify causality links. It is also important to note that we estimate a causality network starting from historical information on K assets, generally focusing on asset returns.

The K assets thus represent the nodes in the networks, and our purpose is to identify edges connecting each pair of nodes. If all assets are connected, and we exclude self-loops, we will have a total number of $K^2 - K$ edges in which case the associated adjacency matrix is symmetric if the network is unweighted.

In all the cases we discuss below, for any pair of assets i and j, the existence of an edge connecting the nodes i and j with a specific direction (say, from i to j) is established from a statistical test. The null hypothesis of the various tests corresponds to the absence of a causality link, while the null hypothesis is rejected in the presence of some form of causality. The type of causality detected clearly depends on the hypothesis tested.

For a given test statistic \mathcal{M} , and for any pair of assets i and j, we define the edge from i to j as

$$w_{i,j} = \begin{cases} 0 & \mathcal{M} \text{ null hypothesis rejected} \\ 1 & \mathcal{M} \text{ null hypothesis accepted} \end{cases}$$
 (8)

where $w_{i,j}$ is the element in position i, j within an adjacency matrix. Given the use of the test statistic \mathcal{M} , we denote the corresponding adjacency matrix as $W_{\mathcal{M}}$. The outcome of the test statistic, either rejecting or accepting the null, thus identifies the edges of a directed and unweighted network. We do not discuss here the possible generalization to a more flexible form of network construction that accounts for the strength of the causal relation too: such an extension is clearly possible but is left to future research. It should be noted that the networks will be row-normalized before they are used in the model presented in the previous section.

3 Empirical analyses

3.1 Data description

The empirical analyses are conducted on three different datasets: the first concerns the 48 industry portfolios on the Kenneth R. French website; the second includes the 25 US banks with the greatest market value; the third the 25 US insurance firms with the highest market value.² Using the three datasets, we can test for the presence of causality links from a general economic standpoint, or by looking at the companies that were mostly affected by the recent financial crisis.

For the three datasets, we use the daily returns over two different samples. The first spans the open-market business days between 3 January 2006 and 31 December 2008. The second covers the period between 3 January 2011 and 31 December 2015. The former is characterized by 755 daily returns for each time series, while the latter records 1258 observations for each industry portfolio or firm. The two samples behave very differently, since the first includes the years of the global financial crisis, while the second covers a period of upward trend in the financial market, excluding the initial recovery period after the crisis. Our purpose is to test whether there is a different dependence structure among the variables during periods of market distress vis-à-vis periods with a positive market trend. We proceed with our estimation of causality networks by focusing on the daily returns.

Just as we selected the 25 US banks and the 25 US insurance firms with the highest market value as at the beginning of 2006, for the first time interval considered, the choice of banks and insurance companies to consider in the second period was based on their market value as at the beginning of 2011. This means that most, but not all of these financial institutions are included in both of the samples. Appendix B contains a list of the selected banks and insurance companies. We stress that our objective is to obtain an overview of the changes occurring in the links across nodes (banks, insurance companies, or economic sectors) when we change the approach used to ascertain the existence of nodes, switching from Granger causality to quantile-based causality, to arrive at the optimal combination of competing available networks.

Tables C.9-C.10 in Appendix C show the main descriptive statistics computed for the banks in the first and second time intervals, respectively; Tables C.11-C.12 concern the insurance companies and the statistics for the 48 industry portfolios are given in Tables C.13-C.14. In all cases, the average returns are lower in 2006-2008 than in 2011-2015. This was expected and is due to the effects of the global financial crisis. The first time interval is also characterized by a greater uncertainty, averaging higher standard deviations and interquartile ranges; this was also expected. The risk affecting the years 2006-2008 is also apparent from the corresponding minimum and maximum returns. The distributions of these returns are fat-tailed (especially in the first period), consistently with the stylized facts highlighted by Cont (2001). The distributions are also highly skewed. Finally, we used the Ljung-Box test for mean serial dependence on both the returns and the squared returns, using 1, 5 and 10 lags. As expected, the outcomes show no serial dependence on returns, while squared returns (a proxy of variances) are highly correlated, suggesting the presence of heteroskedasticity. In short, the descriptive analysis shows significant

²The series of industry portfolios are available at http://mba.tuck.dartmouth.edu/pages/faculty/ken.french/, the data on the banks and insurance firms were obtained from Thomson Reuters Datastream.

differences between the years 2006-2008 and 2011-2015, that are clearly a consequence of the turmoil affecting the US economy during the former period. We now take a closer look at the dependence across companies or economic sectors by focusing on causality analyses.

3.2 Granger and quantile causality networks

In building the causality networks, we extend the existing literature by combining the classic Granger testing approach with methods based on analyses of dependence across quantiles, as described in Section A.1.

The methods based on quantile regression and quantile causality depend on the quantile at which the analysis is performed, so we must first specify which quantiles we use to conduct the dependence or causality analysis. Our analysis illustrates the changes occurring in the causality networks when we switch from Granger causality to quantile causality, without identifying the optimal quantile over which to measure quantile causality. We therefore decided to fix the quantiles of interest a priori: we chose three quantiles, 0.1, 0.5 and 0.9, corresponding to the tails and the center of the distributions of the returns. Note that, when considering the approach of Sim and Zhou (2015), where we have quantile causality depending on two different quantiles, we set both quantiles at the same level (see Appendix A for details on the quantile causality methods we consider).

In the evaluation of quantile regression parameters, we estimate the standard errors with the xy-pair bootstrap method, Efron and Tibshirani (1993), with 5000 replications. This approach provides accurate results without assuming any particular distribution for the error term. When a kernel function is involved (see Appedinx A for details), we set it always to the Gaussian one. Following Balcilar et al. (2017) and Bonaccolto et al. (2018), we estimate the optimal bandwidths involved in the Jeong et al. (2012) test using the least squares cross-validation method of Rudemo (1982) and Bowman (1984). We also need to specify the lag orders for both the dependent and the explanatory variables. To limit the computational burden, we fix a-priori both lag orders to be 1. Finally, as for the quantile-on-quantile method, the objective functions depend also on a bandwidth (see Appedinx A for details). Larger values of the bandwidth make smaller the variance of the estimates, but, on the other hand, would increase their bias. The opposite holds when h takes lower values. Optimizing the value of h by using ad-hoc statistical methods is still an open issue in a quantile-on-quantile framework; we do not analyze this aspect in the present study and leave it for future research. In contrast, we follow Sim and Zhou (2015) and use a plug-in bandwidth equal to 0.1. We remind that the bandwidth applies to a quantile level. Our choice allows avoiding estimating the parameters with a small number of data points (as it is the case with smaller bandwidths) and, at the same time, to maintain the focus around the quantiles of interest. We set the confidence level at 5% for judging the significance of the coefficients and testing the null hypothesis behind the tests for Granger or quantile causality.

To control for the possible impact of the common dependence on the market, we test for Granger causality as well as for the various forms of quantile causality, also including the supercomposite US market index within the corresponding models.³

 $^{^3}$ We obtain the market return from the Fama/French 3 Factors dataset provided by Kenneth R. French and available at http://mba.tuck.dartmouth.edu/pages/faculty/ken.french/.

In the remainder of this paper, we identify the various networks that we estimate by means of acronyms. For networks estimated using quantile regression approaches (the baseline case, the quantile-on-quantile, and the non-parametric test), the acronym consists of four or five characters. The first and second identify the method: QR for baseline quantile regression; Qo for quantile-on-quantile; QN for non-parametric quantile. The third and fourth characters identify the quantile: for instance, QR10 refers to the network estimated by baseline quantile regression at the 10% quantile. For the quantile-on-quantile cases, the reported quantile is used for both the dependent and the explanatory variables. For standard Granger causality networks we use GR. Then we add a final letter F to the acronym if we allow for the presence of a common factor in our estimation of the causality networks.

Figures 2-4 graphically represent selected networks. Appendix D includes graphs for all the estimated causality networks.

Figure 2 includes four different networks among the largest US banks: the Granger causality network and three different quantile causality networks estimated by focusing on the 10% quantile. We indicate the baseline quantile network, the quantile-on-quantile network and the non-parametric quantile causality network. All the networks refer to the financial crisis period (2006 to 2008), and are estimated without any common factors. Figures 3 and 4 show similar networks estimated for the 25 largest US insurance companies and the US industry portfolios, respectively. We stress that we report results for the 10% quantile because we wish to shed some light on the possible differences between Granger causality (which focuses on the mean) and quantile causality (which places more emphasis on the tails). Networks that are based on quantile causality in the mean and also pay attention to the upper tail are available in Appendix D.

Figure 2 shows that the quantile causality network extracted using the baseline quantile regression method has a majority of isolated nodes, whereas the network extracted with a non-parametric quantile causality test suggests the presence of a much greater density. The comparison between the networks of the banks and those of the insurance companies and industry portfolios shows that the former are less connected whatever methods is used. The non-parametric quantile causality test also generates networks characterized by a similar topology for all institutions and portfolios; in particular, it is easy to distinguish between a kernel and a periphery.⁴

Figure 5 shows the dataset of 25 banks over time, comparing the networks estimated for 2006-2008 with those obtained for 2011-2015, using two specific causality network estimation methods: Granger causality and non-parametric quantile causality (10% quantile). We can see some changes in the Granger network structure when moving from the earlier to the later period. These changes are less clear when we consider non-parametric quantile causality, in which case the network still presents a kernel-periphery structure.

These preliminary graphical analyses suggest that there are potentially relevant differences between the networks estimated using Granger causality as opposed to quantile-based causality approaches. The latter have a more clear focus on risk than the former. Before moving on to the combination of the estimated networks, we run a comparison of the estimated networks using

 $^{^4}$ We use the Fruchterman & Reingold algorithm for network visualization, Fruchterman and Reingold (1991).




Figure 2: The figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 banks ordered for market capitalization and listed in Appendix B. Panel a) reports the network extracted by the standard Granger causality approach of Billio et al. (2012). Panel b) reports the Network extracted from a baseline quantile regression methodology, at the 10% quantile. Panel c) plots the network extracted from a quantile-on-quantile methodology at the 10% quantile. Panel d) graphs the network estimated by a non-parametric quantile causality methodology at the 10% quantile. All the estimates exclude the presence of common factors. We use the Fruchterman & Reingold algorithm for network visualization.




Figure 3: The figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 insurances ordered for market capitalization and listed in Appendix B. Panel a) reports the network extracted by the standard Granger causality approach of Billio et al. (2012). Panel b) reports the Network extracted from a baseline quantile regression methodology, at the 10% quantile. Panel c) plots the network extracted from a quantile-on-quantile methodology at the 10% quantile. Panel d) graphs the network estimated by a non-parametric quantile causality methodology at the 10% quantile. All the estimates exclude the presence of common factors. We use the Fruchterman & Reingold algorithm for network visualization.




Figure 4: The figure visualizes 4 different networks for the period 2006-2008 relative to the 48 Industry portfolios obtained from the Kenneth French website. Panel a) reports the network extracted by the standard Granger causality approach of Billio et al. (2012). Panel b) reports the Network extracted from a baseline quantile regression methodology, at the 10% quantile. Panel c) plots the network extracted from a quantile-on-quantile methodology at the 10% quantile. Panel d) graphs the network estimated by a non-parametric quantile causality methodology at the 10% quantile. All the estimates exclude the presence of common factors. We use the Fruchterman & Reingold algorithm for network visualization.




Figure 5: The figure displays the network of the 25 banks ordered for market capitalization (see Appendix B by contrasting methods and samples. Panels a) and b) report the networks extracted by using the standard Granger causality methodology for the 2006-2008 and 2011-2015 samples, respectively. Panels c) and d) report the networks extracted by using the the non-parametric quantile causality test at the 10% quantile for samples 2006-2008 and 2011-2015, respectively. All the estimates exclude the presence of common factors. We use the Fruchterman & Reingold algorithm for network visualization.

summary measures.

We consider four different indicators: Density, Assortativity, and two versions of Eigenvector centrality. Density monitors the number of connections between nodes. A higher density is the sign of a large number of connections between nodes, which are consequently more closely related to one another. Assortativity captures the nodes' tendency to connect with other nodes having similar properties; it can take values in the range of (-1,1). For values close to 1, the network has an assortative behavior, with nodes being connected to other nodes that have similar degrees; for values nearing zero, the network becomes similar to a random graph. A disassortative behavior, when high-degree nodes point to low-degree nodes, corresponds to assortativity values close to -1.5 Eigenvector centrality monitors the relevance of each node as a function of the relevance of neighboring nodes. We consider two versions: one based on the non-normalized adjacency matrix, and one based on the row-normalized adjacency matrix. We also normalize the eigenvector centrality of each node to the maximum score obtained by the most connected node. The eigenvector centrality value thus ranges from zero to one, and can easily be compared across networks. In the summary tables, we focus on the average eigenvector centrality across nodes. Changes in the eigenvector centrality are a sign of changes in the network structure. Appendix E provides additional details on the network measures considered. Tables from 1 to 3 contain the summary measures.

Table 1 contains the network measures for the banks dataset. We can see that, on average, the density is slightly smaller during the crisis, while the disassortative behavior (seen in both periods) is higher. Granger's causality networks (with/without the common factor) have small densities that remain almost constant in the two samples, and a disassortative behavior, and the average eigenvector centrality increases in the most recent period. The values for the quantile-based networks are quite different, as the previous graphical analysis suggests. In particular, the networks derived by means of the non-parametric quantile causality test have the highest density, combined with a disassortative pattern. These two elements could explain the kernel-periphery structure exhibited in the Figure 2.

A high density combined with a disassortative pattern is also associated with very high eigenvector centrality averages, which indicate that the nodes' relevance is quite evenly distributed, a somewhat expected result given that we focus on the largest banks. Notably, the picture is quite different in the Granger causality networks, which are characterized by a lower average eigenvector centrality. From a systemic risk perspective, this finding suggests that, while the Granger causality analysis provides evidence of few relevant nodes that are more central to the network structure, a non-parametric quantile causality test leads to the construction of a denser network where the risk is more evenly distributed across nodes, and many nodes (more than those emerging from Granger causality) are systemically relevant. As the graphs suggest, the non-parametric quantile causality networks also differ from those based on other quantile-based approaches. In particular, the baseline quantile case provides summary measures more closely resembling those of Granger causality, while the quantile-on-quantile cases have density measures higher than with Granger causality but lower than with non-parametric quantile causality,

⁵The few blank spaces appearing in the following tables of network summary measures correspond to indeterminate forms. The indeterminate forms of the assortativity and eigenvector centrality measures are discussed in the Appendix E.

Table 1: This table reports summary measures for the networks estimated from competing causality methodologies over the 25 Banks listed in Appendix B. QB stands for baseline quantile, Qo for quantile-on-quantile, QN for non-parametric quantile causality, GR for Granger causality. Numbers identify the reference quantile for quantile-based causality networks, 10 for 10%, 50 for the median and 90 for 90%. An F at the end of the acronyms specifies that the underlying models included the market index as a common factor.

	Density		Assort	ativity	Eigenvector Centrality		
	06-08	11-15	06-08	11-15	06-08	11-15	
GR	0.12	0.07	-0.37	-0.22	0.12	0.26	
QB10	0.03	0.04	0.05	-0.13			
QB50	0.05	0.07	0.03	-0.06	0.16		
QB90	0.12	0.13	-0.44	-0.22	0.25	0.11	
Qo10	0.16	0.36	-0.37	-0.28	0.23	0.45	
Qo50	0.06	0.03	-0.43	0.00	0.34		
Qo90	0.17	0.35	-0.53	-0.07	0.25	0.44	
QN10	0.51	0.63	-0.11	-0.20	0.93	0.83	
QN50	0.29	0.33	-0.30	-0.28	0.75	0.72	
QN90	0.57	0.52	-0.15	-0.13	0.90	0.83	
GRF	0.07	0.07	-0.01	-0.24	0.22	0.31	
QB10F	0.04	0.04	-0.26	0.13	0.08		
QB50F	0.04	0.09	-0.23	-0.12	0.14	0.29	
QB90F	0.10	0.09	-0.37	-0.34	0.21	0.26	
Qo10F	0.22	0.50	-0.29	-0.24	0.24	0.50	
Qo50F	0.06	0.03	-0.07	-0.25	0.18		
Qo90F	0.10	0.51	-0.50	-0.20	0.18	0.57	
QN10F	0.52	0.70	-0.14	-0.23	0.88	0.84	
QN50F	0.34	0.38	-0.22	-0.21	0.80	0.81	
QN90F	0.64	0.64	-0.09	-0.14	0.96	0.89	
Average Standard Deviation	$\begin{array}{c} 0.21 \\ 0.20 \end{array}$	$0.28 \\ 0.24$	$-0.24 \\ 0.17$	-0.17 0.11	$\begin{array}{c} 0.41 \\ 0.33 \end{array}$	$0.54 \\ 0.26$	

Table 2: This table reports summary measures for the networks estimated from competing causality methodologies over the 25 Insurance companies listed in Appendix B. QB stands for baseline quantile, Qo for quantile-on-quantile, QN for non-parametric quantile causality, GR for Granger causality. Numbers identify the reference quantile for quantile-based causality networks, 10 for 10%, 50 for the median and 90 for 90%. An F at the end of the acronyms specifies that the underlying models included the market index as a common factor.

	Density		Assort	ativity	Eigenvector Centrality		
	06-08	11-15	06-08	11-15	06-08	11-15	
GR	0.20	0.03	-0.29	-0.37	0.34		
QB10	0.07	0.06	-0.36	0.15	0.15	0.22	
QB50	0.07	0.14	-0.17	-0.15	0.32	0.36	
QB90	0.04	0.07	-0.44	-0.03			
Qo10	0.17	0.25	-0.31	-0.29	0.29	0.33	
Qo50	0.03	0.05	0.00	-0.10			
Qo90	0.22	0.21	-0.42	-0.08	0.38	0.30	
QN10	0.76	0.56		-0.23	0.96	0.90	
QN50	0.41	0.50	-0.35	-0.18	0.70	0.85	
QN90	0.76	0.59	-0.02	-0.20	0.96	0.82	
GRF	0.15	0.04	-0.31	-0.40	0.26		
QB10F	0.04	0.08	-0.49	-0.12		0.22	
QB50F	0.05	0.12	-0.06	-0.05		0.27	
QB90F	0.04	0.10	-0.44	-0.08		0.37	
Qo10F	0.08	0.43	-0.06	-0.28	0.18	0.44	
Qo50F	0.04	0.05	0.13	0.21		0.21	
Qo90F	0.10	0.45	-0.43	-0.31	0.24	0.46	
QN10F	0.76	0.66		-0.11	0.96	0.93	
QN50F	0.54	0.55	-0.13	-0.11	0.92	0.93	
QN90F	0.76	0.62		-0.18	0.96	0.86	
Average Standard Deviation	$0.26 \\ 0.29$	$0.28 \\ 0.23$	-0.24 0.19	-0.14 0.15	$\begin{array}{c} 0.54 \\ 0.34 \end{array}$	$0.53 \\ 0.29$	

and they show the most disassortative behaviors. We note some interesting differences across quantiles, with higher densities on the extreme quantiles (10% and 90%) than on the median case, which comes the closest to the Granger causality. Notably, this would mean that the network structure changes if we move away from the mean (median), an important aspect to bear in mind if our purpose is to analyze the spread of risk in times of market turmoil. Finally, the introduction of a common market factor seems to have a limited impact.

Table 2 indicates that the insurance companies dataset behaves in much the same way as the banks dataset (thus confirming our previous comments), with a marked change in the network structure when moving from Granger causality to quantile causality. The most relevant difference concerns network density, which is much higher in the tails than in the mean (which coincides with Granger causality). We also find the highest average eigenvector centrality coinciding with non-parametric quantile causality.

Table 3 shows the summary measures for the causality network estimated from the 48 indus-

Table 3: This table reports summary measures for the networks estimated from competing causality methodologies over the 48 Industry portfolios recovered from the Kenneth French website. QB stands for baseline quantile, Qo for quantile-on-quantile, QN for non-parametric quantile causality, GR for Granger causality. Numbers identify the reference quantile for quantile-based causality networks, 10 for 10%, 50 for the median and 90 for 90%. An F at the end of the acronyms specifies that the underlying models included the market index as a common factor.

	Density		Assort	ativity	Eigenvector Centrality		
	06-08	11-15	06-08	11-15	06-08	11-15	
GR	0.14	0.02	-0.49	-0.74	0.15	0.36	
QB10	0.09	0.06	-0.39	-0.20	0.24	0.24	
QB50	0.16	0.06	-0.27	-0.13	0.21	0.27	
QB90	0.14	0.07	-0.24	-0.37	0.29	0.29	
Qo10	0.20	0.52	-0.06	-0.10	0.29	0.60	
Qo50	0.08	0.06	0.09	-0.13	0.25	0.13	
Qo90	0.14	0.48	-0.29	-0.15	0.26	0.56	
QN10	0.52	0.42	-0.25	-0.14	0.88	0.90	
QN50	0.32	0.28	-0.22	-0.23	0.82	0.83	
QN90	0.57	0.48	-0.14	-0.18	0.93	0.90	
GRF	0.15	0.02	-0.33	-0.71	0.18	0.16	
QB10F	0.14	0.04	-0.11	-0.11	0.17	0.18	
QB50F	0.15	0.06	-0.17	-0.21	0.20	0.16	
QB90F	0.09	0.08	-0.12	0.02	0.17	0.18	
Qo10F	0.39	0.79	-0.28	-0.12	0.43	0.79	
Qo50F	0.09	0.07	-0.19	-0.10	0.13	0.10	
Qo90F	0.35	0.82	-0.27	-0.09	0.39	0.82	
QN10F	0.59	0.47	-0.12	-0.18	0.95	0.88	
QN50F	0.33	0.33	-0.18	-0.22	0.85	0.81	
QN90F	0.59	0.51	-0.15	-0.11	0.93	0.93	
Average Standard Deviation	0.26 0.18	0.28 0.26	$-0.21 \\ 0.12$	-0.21 0.19	$0.44 \\ 0.32$	$0.51 \\ 0.32$	

try portfolios. Here again, we find relevant differences between the summary measures for the Granger and quantile causality networks. This holds particularly for network density and average eigenvector centrality. We note, however, that the structure of the non-parametric quantile causality networks differs considerably from the other quantile causality networks in all three datasets.

The graphical analysis and summary measures confirm that, changing our approach to estimating a causality network, coincides with important changes in the network structure. To examine the possible relevance of either the Granger causality network or the quantile causality networks, we therefore proceed, in the following section, with the estimation of a composite network.

3.3 The composite causality network

Starting from the availability of causality networks estimated using four different methods (Granger, Quantile Regression, Quantile-on-Quantile and Non-parametric Quantile causality), and over different quantiles (10%, the median and 90%) for three of them, we now proceed to estimate composite networks.

We estimate the model in equation (5), accounting for the presence of common factors, for which we follow the usual practice and introduce the market factor, the size factor, the book-to-market factor, and the momentum factor, following Fama and French (1993), Fama and French (1995), and Carhart (1997). We have no information a priori to suggest a possible preference for particular network, so we estimate the composite network starting from four different layers, where the quantile-based networks are associated with the same reference quantile. We provide a sensitivity analysis on the effect of excluding a single network in the robustness checks section.

Then we estimate the linear factor model, focusing on weekly returns. While it is more common to consider the monthly frequency when estimating factor models, this would leave us with only 36 observations in the first sample, while we have 60 monthly returns in the second. Hence our decision to focus on the weekly frequency, which enables us to increase the number of returns for the factor model (augmented with network dependence) to 108 in the first sample and 316 in the second. This choice has advantages in terms of model estimation and parameter inference. We do not consider daily data as they would be bound to require the introduction of heteroskedastic dynamics in the variance of the residuals, adding to the complexity of the estimation due to the well-known curse of dimensionality. Estimates of the network combination on monthly returns are nonetheless provided in the robustness checks sections.

The estimates of the linear factor model with network dependence generate different outputs. First, there are the model parameters: the weight of each network, as measured by the δ_i parameters, and the impact of the composite network on the asset returns, the parameters included in \mathcal{R} . Second, comes the composite network, obtained by combining the primitive networks weighted with the δ_i coefficients. Finally, there are the model residuals, which contain information useful for assessing the advantages of moving to a composite network as opposed to the benchmark cases of no network dependence (where the contemporaneous link matrix A is an identity), or a network dependence captured by Granger causality. The latter is a valuable benchmark as we would like to underscore the potential improvement associated with measuring asset links going beyond the mean (i.e. on the quantiles). To compare models, we use the residuals average correlation: if we see improvements, we expect to have residuals characterized by a smaller correlation level, i.e. more whitened residuals.

Table 4 shows the parameters estimated for the combined network in the three datasets and over sample periods. Bearing in mind that the sum of these parameters is 1 and they are all positive, a higher value of a parameter indicates a greater relevance of the associated network. We also assess the significance of the estimated parameters. Within the banks dataset, we note that the non-parametric quantile causality network is the most relevant during the financial crisis period and across all quantiles. Notably, the estimated δ parameter for this network ranges between 0.736 and 0.883, and it is always statistically significant when the estimation of the networks takes the presence of a common market factor into account.

Table 4: The table reports the δ of model (5) that represent the weights for networks combination. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based network, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks. Columns 3 to 6 refer to the crisis sample while columns 7 to 10 to the most recent sample. The second row identifies the four different networks which are optimally combined: baseline quantile causality - QB; quantile-on-quantile causality Qo; non-parametric quantile causality - QN; Granger causality. Parameters are, by construction, positive and sum up to one (within each row and within each period). A star identifies parameters significant at the 5% confidence level.

			2006	-2008		2010-2015					
Quantile	Factor	QB	Qo	$\mathbf{Q}\mathbf{N}$	GR	QB	Qo	$\mathbf{Q}\mathbf{N}$	$\mathbf{G}\mathbf{R}$		
					anks						
10%	N	0.061	0.055	0.875	0.010	0.000	0.630*	0.354*	0.017*		
50%	N	0.146	0.108	0.736*	0.010	0.091*	0.065*	0.773*	0.071		
90%	N	0.005	0.008	0.883	0.104	0.000	0.706*	0.243	0.051*		
10%	Y	0.022	0.073	0.827*	0.077	0.000	0.538*	0.428*	0.034*		
50%	Y	0.052	0.057	0.842*	0.050	0.107	0.072*	0.649*	0.173*		
90%	Y	0.134	0.000*	0.762*	0.104*	0.000	0.876*	0.121	0.003*		
				25 I	nsurance	e Compa	anies				
10%	N	0.000	0.022*	0.951*	0.027	0.058	0.831	0.111	0.000		
50%	N	0.865*	0.000	0.032*	0.103	0.059	0.045	0.886*	0.010		
90%	N	0.000	0.028	0.972*	0.000	0.000	0.364*	0.436*	0.201		
10%	Y	0.001	0.006	0.948*	0.045	0.000	0.488	0.512	0.000		
50%	Y	0.731*	0.054	0.188*	0.027*	0.051	0.040	0.837*	0.072		
90%	Y	0.000	0.046	0.954*	0.000	0.000	0.554	0.446	0.000		
				48	Industr	y Portfo	lios				
10%	N	0.000	1.000*	0.000	0.000	0.032	0.968*	0.000	0.000		
50%	N	0.212	0.000	0.478	0.310	0.021	0.031*	0.848*	0.100		
90%	N	0.108	0.892*	0.000	0.000	0.000	1.000*	0.000	0.000		
10%	Y	0.178	0.471	0.000	0.351	0.036	0.695*	0.228	0.041		
50%	Y	0.123	0.000	0.000	0.877*	0.092	0.007*	0.836*	0.064		
90%	Y	0.000	0.810	0.000	0.190	0.107	0.808	0.030	0.055		

We stress that, from a systemic risk perspective, this result suggests that the risk is more widespread across banks, as the non-parametric quantile network is much denser than the other causality networks. This greater relevance of the non-parametric quantile causality network also emerges for the median and for the lower and upper tails. In all cases, the Granger causality network has a very small weight (which is non-significant in five cases out of six). Switching to the more recent period considered, both the non-parametric quantile causality network and the quantile-on-quantile causality network are relevant, with and without the introduction of common market factors. Here again, although it is more statistically significant, the Granger causality network has smaller weights than the quantile-based causality networks. Overall, in both samples, the baseline quantile regression network has the least relevant role. Finally, in

the second sample, we see a clear change in the combined network parameters when moving from median networks to the use of networks estimated on either the lower or upper tails. In particular, the impact of the quantile-on-quantile networks increases, thus supporting the importance of looking at quantile causality too when accounting for interdependence across assets in a linear factor model, and showing that a double conditioning (on both the dependent and the explanatory variables) is important when estimating causality at quantile level.

Table 5: The table reports summary measures of the composite networks estimated from model (5) over the different datasets, sample periods, and reference quantiles. In all cases the composite networks is formed by the combination of four networks: the baseline quantile causality network, the quantile-on-quantile causality network, the non-parametric quantile causality network, the Granger causality network. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based networks, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks.

Measi	ıres	Den	sity	Assort	ativity	Eigenvector Centrality		Eigenvector Centrality Adj Weighted	
Quantile	Factor	06-08	11-15	06-08	11-15	06-08	11-15	06-08	11-15
					25 E	Banks			
10%	N	0.63	0.79	-0.30	-0.23	0.70	0.83	0.35	0.47
50%	N	0.45	0.43	-0.28	-0.28	0.62	0.61	0.24	0.43
90%	N	0.69	0.76	-0.32	-0.25	0.70	0.77	0.32	0.57
10%	Y	0.65	0.87	-0.36	-0.25	0.67	0.87	0.39	0.64
50%	Y	0.45	0.48	-0.31	-0.36	0.65	0.69	0.31	0.41
90%	Y	0.72	0.85	-0.28	-0.25	0.75	0.85	0.39	0.64
				25 I	nsuranc	e Comp	anies		
10%	N	0.88	0.72	-0.17	-0.29	0.88	0.74	0.66	0.45
50%	N	0.59	0.61	-0.22	-0.21	0.72	0.85	0.37	0.60
90%	N	0.87	0.72	-0.15	-0.24	0.91	0.77	0.66	0.50
10%	Y	0.82	0.82	-0.22	-0.30	0.85	0.82	0.55	0.59
50%	Y	0.66	0.64	-0.25	-0.23	0.79	0.84	0.55	0.51
90%	Y	0.83	0.83	-0.22	-0.26	0.83	0.83	0.60	0.59
				48	Industr	y Portfo	olios		
10%	N	0.71	0.74	-0.22	-0.26	0.77	0.78	0.46	0.62
50%	N	0.50	0.37	-0.30	-0.29	0.58	0.51	0.29	0.19
90%	N	0.71	0.74	-0.21	-0.24	0.76	0.76	0.38	0.62
10%	Y	0.83	0.89	-0.24	-0.24	0.83	0.90	0.51	0.69
50%	Y	0.53	0.42	-0.35	-0.37	0.56	0.55	0.24	0.23
90%	Y	0.80	0.92	-0.29	-0.15	0.80	0.92	0.47	0.70

The insurance companies dataset produces somewhat similar results. In both periods, Granger causality networks are the least relevant, being associated with small coefficients (and only one in twelve is statistically significant). In the first period considered, non-parametric quantile causality is the most relevant, although baseline quantile causality network receives a

much larger weight in the median case (unlike the picture emerging from the banks dataset). In the second sample, both non-parametric quantile causality and quantile-on-quantile causality are relevant, with a change in the estimated parameters when moving from networks estimated on the median to networks estimated on the 10% or 90% quantiles.

Finally, for the industry portfolio dataset, the results are more heterogeneous in the first sample, while in the second they are consistent with the two previous cases. In the first period, the significance is limited in many cases, particularly for the quantile-based causality networks estimated at the median. In the second period, there is a marked difference between the quantile-based causality estimated at the median and those estimated on the tails. Be that as it may, Granger causality networks receive the smallest weight (and are never statistically significant).

Table 5 shows the summary measures for the combined networks. The various composite networks are similar in all the quantities we report (for a given sample period, and a given dataset). The heterogeneity identified is much smaller than was seen for the primitive networks. Figures 6 to 8 show the combined networks for the period 2006 - 2008.

We link this finding to the use of a linear factor model augmented with network dependence. The contemporaneous relation across the modelled variables captures the correlation across these variables and goes beyond what we associate with common factors. It might be that the various composite networks capture the dependence across the returns (beyond common factors in a similar way. The differences we find depend partly on the heterogeneity across networks and partly on the different weights assigned to the primitive networks.




Figure 6: The figure visualizes the network for the Banks companies dataset. The network is extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method, during the period 2006-2008. In this case we do not allow the presence of the common factor for network estimations.

To shed further light on the differences between the composite networks and generate some evidence of the improvement gained by the linear factor model augmented with network combination, we provide descriptive analyses -in Tables 6 to 8- of the correlations between the model residuals (5). The tables include two benchmarks: a standard linear factor model, the Carhart




Figure 7: The figure visualizes the network for the Insurers companies dataset. The network is extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method, during the period 2006-2008. In this case we do not allow the presence of the common factor for network estimations.




Figure 8: The figure visualizes the network for the Industry portfolios dataset. The network is extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method, during the period 2006-2008. In this case we do not allow the presence of the common factor for network estimations.

(1997) 4-factors CAPM; and a linear factor augmented with network dependence where the latter is the Granger causality network (with or without the inclusion of a common factor in estimating causal relationships). For the datasets concerning the banks and insurance companies, the results are much the same. The residuals of the 4-factors CAPM have the largest median correlations, and a distribution of these correlations shifted to the right, with a clear prevalence

of positive values. The introduction of network dependence, when measured by Granger causality, improves the model fit, but the predominance of positive values in the correlation remains (see the small fraction of correlations below -0.1), and the median correlation remains above 0.1. With the introduction of a plurality of networks, the model fit improves considerably: the residual correlations are centered at zero, with a higher fraction of correlations below -0.1, and a marked reduction in the presence of large positive residual correlations. We thus conclude that our approach based on combining several networks within a network augmented linear factor model constitutes an improvement over the use of simple Granger causality. The improvement achieved by the latter, over the multifactor model, was already documented in Billio et al. (2015).

Table 6: The table reports residual correlation descriptive analyses for the Banks dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

Model	N Networks	Factor	Min	Max	q10	q50	q90	% elements ≤ -0.1			
2006-2008											
Q(10%)	4	N	-0.279	0.584	-0.106	0.071	0.282	11.7%			
Q(50%)	4	N	-0.345	0.596	-0.109	0.062	0.311	10.7%			
Q(90%)	4	N	-0.269	0.600	-0.146	0.050	0.276	16.0%			
Q(10%)	4	Y	-0.297	0.566	-0.135	0.035	0.238	17.7%			
Q(50%)	4	Y	-0.349	0.503	-0.119	0.047	0.242	14.7%			
Q(90%)	4	Y	-0.297	0.455	-0.161	0.020	0.218	18.3%			
G	1	N	-0.381	0.670	-0.128	0.111	0.372	13.3%			
G	1	Y	-0.394	0.641	-0.128	0.089	0.378	11.7%			
4-F-CAPM	_		-0.358	0.678	-0.108	0.257	0.502	10.7%			
			201	1-2015							
Q(10%)	4	N	-0.263	0.422	-0.122	-0.013	0.135	15.3%			
Q(50%)	4	N	-0.397	0.537	-0.129	0.055	0.224	13.7%			
Q(90%)	4	N	-0.253	0.431	-0.127	0.010	0.138	15.7%			
Q(10%)	4	Y	-0.283	0.503	-0.132	-0.008	0.140	19.0%			
Q(50%)	4	Y	-0.330	0.505	-0.126	0.041	0.197	14.7%			
Q(90%)	4	Y	-0.304	0.508	-0.140	0.001	0.132	17.7%			
Granger	1	N	-0.211	0.598	-0.027	0.151	0.362	3.0%			
Granger	1	Y	-0.211	0.598	-0.027	0.146	0.361	3.3%			
Multifactor	_		-0.122	0.669	0.054	0.266	0.487	0.3%			

For the industry portfolio dataset, the results are less clear, since the various approaches produced very similar findings. This is in line with the heterogeneous results seen for the weights of the various networks, where we were unable to identify a clear preference. We interpret this as evidence to suggest that network augmented linear factor models might usefully improve on

Table 7: The table reports residual correlation descriptive analyses for the Insurance Companies dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

Model	N Networks	Factor	Min	Max	q10	q50	q90	% elements ≤ -0.1		
2006-2008										
${Q(10\%)}$	4	N	-0.437	0.686	-0.205	0.003	0.250	27.7%		
Q(50%)	4	N	-0.429	0.669	-0.167	0.039	0.258	15.3%		
Q(90%)	4	N	-0.448	0.676	-0.207	0.009	0.255	25.7%		
Q(10%)	4	Y	-0.436	0.684	-0.203	0.004	0.246	28.0%		
Q(50%)	4	Y	-0.417	0.473	-0.176	0.023	0.226	19.3%		
Q(90%)	4	Y	-0.464	0.671	-0.214	0.001	0.253	27.0%		
G	1	N	-0.466	0.831	-0.166	0.044	0.316	17.7%		
G	1	Y	-0.460	0.714	-0.166	0.048	0.323	15.0%		
4-F-CAPM	_	_	-0.370	0.847	-0.149	0.089	0.422	14.0%		
			201	1-2015						
Q(10%)	4	N	-0.290	0.504	-0.121	0.009	0.157	14.7%		
$\mathrm{Q}(50\%)$	4	N	-0.258	0.548	-0.149	0.019	0.202	16.7%		
Q(90%)	4	N	-0.289	0.348	-0.137	-0.006	0.167	19.3%		
Q(10%)	4	Y	-0.333	0.503	-0.153	-0.012	0.189	20.7%		
Q(50%)	4	Y	-0.267	0.592	-0.156	0.016	0.192	18.3%		
Q(90%)	4	Y	-0.299	0.398	-0.152	-0.001	0.199	21.3%		
G	1	N	-0.136	0.658	-0.005	0.121	0.317	1.7%		
G	1	Y	-0.136	0.658	-0.014	0.116	0.315	1.7%		
4-F-CAPM	_		-0.103	0.658	0.017	0.138	0.332	0.3%		

traditional linear factor models when we fit the model over single assets and not over portfolios. The aggregation of assets into portfolios probably distorts the dependence structure across assets and limits its impact. There might be a different reason linking the datasets of the banks and insurance companies to the potential presence of a sector-specific factor in the analysis - but if that were the case, we should have seen no such clear improvement in the residual correlations of the composite network cases. We consequently believe that the idea of a missing factor is inconsistent with our findings. Further analyses are needed on this topic, but we leave them to future research.

We close this section with a few comments on the coefficients for monitoring the impact of the composite networks on the various assets, as included in the diagonal of matrix \mathcal{R} in equation (5). Appendix J shows the plots of the coefficients for each bank, insurance company and industrial portfolio, across the various combined networks. Overall, we note that the coefficients are usually positive and significant for banks and insurance companies, while the networks impact is more

Table 8: The table reports residual correlation descriptive analyses for the Industry portfolios dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

Model	N Networks	Factor	Min	Max	q10	q50	q90	% elements ≤ -0.1			
	2006-2008										
${}$ Q(10%)	4	N	-0.465	0.611	-0.191	-0.001	0.201	25.6%			
Q(50%)	4	N	-0.369	0.663	-0.161	0.018	0.249	20.6%			
Q(90%)	4	N	-0.443	0.608	-0.195	-0.006	0.206	25.9%			
Q(10%)	4	Y	-0.417	0.625	-0.187	0.006	0.220	24.1%			
Q(50%)	4	Y	-0.454	0.627	-0.159	0.011	0.233	20.4%			
Q(90%)	4	Y	-0.508	0.592	-0.187	-0.007	0.215	23.8%			
G	1	N	-0.384	0.687	-0.146	0.025	0.257	18.6%			
G	1	Y	-0.401	0.687	-0.157	0.011	0.239	19.8%			
4-F-CAPM	_		-0.486	0.733	-0.219	0.003	0.267	27.1%			
			201	1-2015							
${Q(10\%)}$	4	N	-0.463	0.567	-0.140	-0.008	0.135	19.7%			
Q(50%)	4	N	-0.454	0.518	-0.136	0.005	0.169	16.7%			
Q(90%)	4	N	-0.471	0.572	-0.144	-0.013	0.139	19.9%			
Q(10%)	4	Y	-0.432	0.497	-0.147	-0.004	0.148	18.9%			
Q(50%)	4	Y	-0.470	0.511	-0.135	0.001	0.155	16.6%			
Q(90%)	4	Y	-0.438	0.565	-0.143	-0.006	0.154	17.9%			
G	1	N	-0.470	0.577	-0.140	0.007	0.199	18.0%			
G	1	Y	-0.503	0.577	-0.139	0.006	0.177	17.7%			
4-F-CAPM			-0.456	0.592	-0.147	0.007	0.208	18.4%			

heterogeneous and of limited significance for industrial portfolios. These results deserve a more thorough analysis, but this goes beyond the scope of the present paper.

3.4 Robustness checks

The previously-reported results focus on the combination of all four different networks, where the model providing the optimal combination makes use of weekly data and allows for a heterogeneous impact of the composite network on the assets. Here we provide some additional comments on variations to the model-based combination design. In particular, we estimate the model on monthly data, we consider combining just three networks, and we control for the optimal combination when the network's impact is homogeneous across assets.

Section G contains tables assessing the model-based network combination when we exclude a network. We consider two specific cases: the first excludes the network estimated from Granger causality in order to highlight the relevance of competing quantile causality networks; the second

disregards the less relevant quantile causality network (the so-called baseline quantile causality). All the analyses are based on weekly data. The results confirm the relevance of the non-parametric approach for the purpose of obtaining a quantile causality that, in several cases, generates the largest and statistically most significant coefficients. The baseline quantile causality is the least relevant, even when the Granger causality network is excluded from the layers. The results for the industry portfolios are the most heterogeneous in terms of network relevance, and they show little improvement with respect to the contraction of the residual correlations. For the banks and insurance companies, there is a marked gain in moving from linear factor models, or Granger causality augmented factor models, to a model that accounts for the presence of quantile causality; the average residual correlation decreases considerably.

Section H contains estimates of the network combination modelled on monthly data (as opposed to the weekly data considered in the previous section).⁶ We confirmed the importance of non-parametric quantile causality and the limited impact of Granger causality. Once again, the outcome is clearer for the datasets concerning banks and insurance companies than the industry portfolio dataset. In addition, contrary to the evidence emerging from the weekly data, the contraction on the residual correlations for the insurance company dataset is less evident during the financial crisis, while it is striking in the second period.

Given the heterogeneity of the network's impact on returns, as mentioned at the end of the previous section, we check the model's performance when we impose a common reaction of the returns to exposure to the network. Section I illustrates the outcomes for the composite network and the impact of the network on returns in the three datasets and two sample periods, when the model is estimated on weekly data. The only notable changes concern the composite network parameters: we now find a larger number of statistically significant coefficients. In addition, Granger causality seems to become more relevant, in the insurance companies dataset at least. The non-parametric quantile causality network nonetheless retains its role and receives the highest coefficients. Moving to the model-based combination confirms the previous results and enables a reduction in the average residual correlations in the banks and insurance companies datasets.

4 Concluding remarks

Causality networks have attracted some attention in the financial economics literature in recent years for the purposes of systemic risk interpretation. We show that the structure of causality networks changes considerably if we move from the mean, commonly associated with traditional Granger causality, to the quantiles. In the latter case, the causality across assets can be measured by means of quantile regression-based approaches. Here we illustrate three different possibilities: the simple quantile regression, its generalization to the quantile-on-quantile approach of Sim and Zhou (2015) and the non-parametric test due to Jeong et al. (2012). We show that changing the approach to network estimation gives rise to networks in which a systemic risk interpretation points to the existence of dense networks with a broader systemic relevance. By focusing on a linear factor model augmented with a multi-layer network dependence, we also demonstrate

⁶In this case too, we ran a sensitivity analysis on the effects of excluding a network. The results are available on request.

that causality networks are useful for capturing this dependence across financial returns, going beyond what can be explained by a market factor. This paves the way to further analyses, and has an associated impact on diversification analyses and portfolio construction, as well as on asset pricing.

References

- Acemoglu, D., Carvalho, D. M., Ozdaglar, A., 2012. The network origins of aggregate fluctuations. Econometrica 80 (5), 1977–2016.
- Anselin, L., 1988. Spatial Econometrics: Methods and Models. Springer Netherlands.
- Balcilar, M., Bekiros, S., Gupta, R., 2017. The role of news-based uncertainty indices in predicting oil markets: a hybrid nonparametric quantile causality method. Empirical Economics 53, 879–889.
- Bargigli, L., di Iasio, G., Infante, L., Lillo, F., Pierobon, F., 2015. The multiplex structure of interbank networks. Quantitative Finance 15 (4), 673–691.
- Barigozzi, M., Brownlees, C., 2014. Nets: Network estimation for time series, working Papers 723, Barcelona Graduate School of Economics.
- Billio, M., Caporin, M., Frattarolo, L., Pelizzon, L., 2016. Networks in risk spillovers: A multivariate garch perspective, university Ca' Foscari of Venice, Dept. of Economics Research Paper Series No. No. 03/WP/2016.
- Billio, M., Caporin, M., Panzica, R., Pelizzon, L., 2015. The impact of network connectivity on factor exposures, asset pricing and portfolio diversification, mimeo.
- Billio, M., Getmansky, M., Lo, A. W., Pelizzon, L., 2012. Econometric measures of connectedness and systemic risk in the finance and insurance sectors. Journal of Financial Economics 104 (3), 535–559.
- Billio, M., Gray, D., Getmansky, M., Lo, A., Merton, R., Pelizzon, L., 2014. Sovereign, bank and insurance credit spreads: Connectedness and system networks, mIT Working Paper.
- Bonaccolto, G., Caporin, M., Gupta, R., 2018. The dynamic impact of uncertainty in causing and forecasting the distribution of oil returns and risk. Physica A: Statistical Mechanics and its Applications 507, 446–469.
- Bonacich, P., 1987. Power and centrality: A family of measures. American journal of sociology, 1170–1182.
- Bowman, A., 1984. An alternative method of cross-validation for the smoothing of kernel density estimates. Biometrika 71, 353–360.
- Brandsma, A. S., Ketellapper, R. H., 1979. A biparametric approach to spatial autocorrelation. Environment and Planning A 11, 51–58.

- Candelon, B., Topkavi, S., 2016. A nonparametric test for granger causality in distribution with application to financial contagion. Journal of Business and Economic Statistics 34 (2), 240–253.
- Caporin, M., Paruolo, P., 2015. Proximity-structured multivariate volatility models. Econometric Reviews 34 (5), 559–593.
- Carhart, M., 1997. On the persistence of mutual fund performance. Journal of Finance 52, 57–82.
- Cheung, Y. W., Ng, L. K., 1996. A causality-in-variance test and its application to financial market prices. Journal of Econometrics 72, 33–48.
- Comte, F., Lieberman, O., 2000. Second-order noncausality in multivariate garch processes. Journal of Time Series Analysis 21 (5), 535–557.
- Cont, R., 2001. Empirical properties of asset returns: stylized facts and statistical issues. Quantitative Finance 1, 223–236.
- Corsi, F., Lillo, F., Pirino, D., 2015. Measuring flight-to-quality with granger-causality tail risk networks. Available at SSRN 2576078.
- Diebold, F. X., Yilmaz, K., 2014. On the network topology of variance decompositions: measuring the connectedness of financial firms. Journal of Econometrics 182 (1), 119–134.
- Diebold, F. X., Yilmaz, K., 2015. Financial and Macroeconomic Connectedness. Oxford University Press.
- Efron, B., Tibshirani, R., 1993. An introduction to the bootstrap. Chapman & Hall, New York.
- Elhorst, J. P., Lacombe, D. J., Piras, G., 2012. On model specification and parameter space definitions in higher order spatial econometric models. Regional Science and Urban Economics 42 (1), 211–220.
- Engle, R. F., Manganelli, S., 2004. Caviar: Conditional autoregressive value at risk by regression quantiles. Journal of Business & Economic Statistics, American Statistical Association 22, 367–381.
- Fama, E., French, K., 1993. Common risk factors in the returns of stocks and bonds. Journal of Financial Economics 33 (3-53).
- Fama, E., French, K., 1995. Size and book-to-market factors in earnings and returns. Journal of Finance 50 (1), 131–155.
- Fruchterman, T. M., Reingold, E. M., 1991. Graph drawing by force-directed placement. Software: Practice and experience 21 (11), 1129–1164.
- Geweke, J., 1984. Inference and causality in economic time series. In Handbook of Econometrics 2, 1101–1144.
- Granger, C. W. J., 1969. Investigating causal relations by econometric models and cross spectral methods. Econometrica 37, 424–438.

- Granger, C. W. J., 1980. Testing for causality: a personal viewpoint. Journal of Economics Dynamic and Control 2 (4), 329–352.
- Granger, C. W. J., 1988. Some recent developments in a concept of causality. Journal of Econometrics 39, 199–211.
- Granger, C. W. J., Robins, R. P., Engle, R. F., 1986. Wholesale retail prices: bivariate time series modeling with forecastable error variances. In Model reliability. Boston MA: MIT Press, 1–17.
- Hafner, C. M., 2003. Fourth moments of multivariate garch models. Journal of Financial Econometrics 1 (1), 26–54.
- Hafner, C. M., Herwartz, H., 2008. Testing for causality in variance using multivariate garch models. Annales d'Économie et de Statistique 89, 215–241.
- Hautsch, N., Kyj, L. M., Malec, P., 2013. Do high-frequency data improve highdimensional portfolio allocations?, technical report, Sonderforschungsbereich 649, Humboldt University, Berlin, Germany.
- Hautsch, N., Kyj, L. M., Oomen, R. C., 2012. A blocking and regularization approach to high-dimensional realized covariance estimation. Journal of Applied Econometrics 27 (4), 625–645.
- Hautsch, N., Schaumburg, J., Schienle, M., 2014. Forecasting systemic impact in financial networks. International Journal of Forecasting 30 (3), 781–794.
- Hautsch, N., Schaumburg, J., Schienle, M., 2015. Financial network systemic risk contributions. Review of Finance 19 (2), 685–738.
- Hong, Y., Liu, Y., S., W., 2009. Granger causality in risk and detection of extreme risk spillover between financial markets. Journal of Econometrics 150, 271–287.
- Hoover, K. D., 2001. Causality in Macroeconomics. Cambridge University Press.
- Jeong, K., Hardle, W. K., Song, S., 2012. A consistent nonparametric test for causality in quantile. Econometric Theory 28, 861–887.
- Kelly, B., Lustig, H., Nieuwerburgh, S. V., 2013. Firm volatility in granular networks, mimeo. Booth School of Business, UCLA-Anderson and Stern School of Business.
- Koenker, R., 2005. Quantile Regression. New York: Cambridge University Press.
- Koenker, R., Bassett, G., 1978. Regression quantiles. Econometrica 46 (1), 33–50.
- Koenker, R., Bassett, G., 1982. Robust tests for heteroscedasticity based on regression quantiles. Econometrica 50, 43–61.
- Lee, L. F., Liu, X., 2010. Efficient gmm estimation of higher order spatial autoregressive models with autoregressive distrurbances. Econometric Theory 26, 187–230.

- Lee, T., Yang, W., 2014. Granger-causality in quantiles between financial markets: Using copula approach. International Review of Financial Analysis 33, 70–78.
- Léon, C., Berndsen, R., Renneboog, L., 2014. Financial stability and interacting networks of financial institutions and market infrastructures, centER discussionpaper Series, 69.
- LeSage, J., Pace, R. K., 2009. Introduction to Spatial Econometrics. Boca Raton, FL, USA, Chapman and Hall/CRC, Taylor and Francs Group.
- Lütkepohl, H., 2005. New introduction to multiple time series analysis. Springer Science & Business Media.
- Molina-Balboa, J., Martinez-Jaramillo, S., van del Leij, M., Lopez-Gallo, F., 2014. A multiplex network analysis of the mexican banking system: link persistence, overlap and waiting times. Journa of Network Theory in Finance 1 (1), 99–138.
- Montagna, M., Kok, C., 2013. Multi-layered interbank model for assessing systemic risk, kiel Institute for the World Economy Working Paper No. 1873 and ECB Working Paper.
- Newman, M., 2010. Networks: an introduction. Oxford university press.
- Newman, M. E., 2002. Assortative mixing in networks. Physical review letters 89 (20), 208701.
- Newman, M. E., 2003. Mixing patterns in networks. Physical Review E 67 (2), 026126.
- Ozdagli, A., Weber, M., 2015. Monetary policy through production networks: Evidence from the stock market, working Paper, University of Chicago, Chicago, IL.
- Rudemo, M., 1982. Empirical choice of histograms and kernel density estimators. Scandinavian Journal of Statistics 9, 65–78.
- Sim, N., Zhou, H., 2015. Oil prices, us stock return, and the dependence between their quantiles. Journal of Banking & Finance 55, 1–8.
- Van Mieghem, P., 2010. Graph spectra for complex networks. Cambridge University Press.
- Wasserman, S., Faust, K., 1994. Social network analysis: Methods and applications. Vol. 8. Cambridge university press.
- White, H., Kim, T., Manganelli, S., 2015. VAR for VaR: Measuring tail dependence using multivariate regression quantiles. Journal of Econometrics 187, 169–188.

Appendix

A Estimating causality networks

A.1 Granger causality networks

We start from the discussion of Granger causality testing, and the estimation of a Granger causality network in which case the framework for testing the presence of causality in mean between two variables x_t and y_t is the Vector Auto Regressive (VAR). We focus on the simplest VAR model with a single lag, that is

$$\begin{bmatrix} y_t \\ x_t \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \begin{bmatrix} y_{t-1} \\ x_{t-1} \end{bmatrix} + \begin{bmatrix} \varepsilon_{y,t} \\ \varepsilon_{x,t} \end{bmatrix}$$
(A.1)

where we assume, for the sake of simplicity, that the innovation term is identically and independently distributed.

Testing for causality from x_t to y_t is tantamount to testing for the significance of the coefficient b_{12} , while the significance of b_{21} provides information on the causality from y_t to x_t . The framework can easily be generalized to the presence of more than one lag, in which case causality is associated with the null hypothesis of zero restrictions on a subset of the model parameters. Furthermore, we might extend the approach for testing causality among a number of variables larger than two. In the latter case, the characterization of causality tests is properly defined on the moving average representation of the model; see Lütkepohl (2005). Test statistics for hypotheses on VAR coefficients might be recovered by resorting to a likelihood ratio framework. Consequently, the test statistics have the usual Chi-square density.

The testing equation can also be extended by introducing further lagged or contemporaneous explanatory variables that affect both the dependent variables. This would bring to light a form of common behavior that might hide or distort the identification of causal relations. Notably, by replacing the series levels with their squares, or by focusing on realized volatility sequences, the Granger causality framework also enables us to test for causality between *risk* measures.

Billio et al. (2012) were the first to adopt Granger causality to estimate a relationship-based network among financial institutions. The connections between nodes in their network denote the presence of a causality relation in the Granger sense. The intuition of Billio et al. (2012) is that Granger causality relationships might be interpreted as a channel through which microeconomic shocks would spread between financial institutions.

The approach used by Billio et al. (2012) starts from a collection of series of K asset returns over a daily sample of size T, which we denote by $R_{i,t}$, with i = 1, 2, ..., N and t = 1, 2, ..., T. Returns are first filtered by means of a GARCH(1,1) model to eliminate the well-known heteroskedastic behavior. The authors thus estimate

$$R_{i,t} = \mu_i + \eta_{i,t} \quad i = 1, 2, \dots N$$
 (A.2)

where μ_i is the unconditional mean and $\eta_{i,t}$ is the innovation for asset *i*. Following the standard literature, $\eta_{it} = \sigma_{it}\epsilon_{it}$, where σ_t is the conditional standard deviation. The conditional

variance follows a simple GARCH(1,1) process

$$\sigma_{i,t}^2 = \omega_i + \alpha_i \eta_{i,t-1}^2 + \beta_i \sigma_{i,t-1}^2 \quad i = 1, 2, \dots N$$
(A.3)

with $\omega_i \geq 0$, $\alpha_{i,1} \geq 0$, $\beta_i \geq 0$, and $\alpha_i + \beta_i < 1$. As usual in the GARCH literature, $\varepsilon_{i,t}$ is assumed to be a sequence of i.i.d random variables with zero mean and unit variance. After the model estimation, we can therefore obtain the so-called standardized residuals $\varepsilon_{i,t} = \frac{\eta_{i,t}}{\sigma_{i,t}}$.

The next step develops Granger's causality test on each pair of standardized asset residuals, but the simple framework of equation (A.1) might lead to the detection of spurious causality. In fact, if we take three assets, i, j and z, and we have $i \to_G j$ (the series i causes the series j in the sense of Granger causality) and $j \to_G z$, then by using the standard Granger causality test we might also find that $i \to_G z$, but such a causality could be a by-product of the presence of j, and not a real direct causality impact. To control for such effects, typical of VAR models with a cross-sectional dimension larger than two, Billio et al. (2012) augment the model in (A.1) with a so-called background series and/or a set of common factors, leading to the following model

$$\begin{bmatrix} \varepsilon_{i,t} \\ \varepsilon_{j,t} \end{bmatrix} = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix} \begin{bmatrix} \varepsilon_{i,t-1} \\ \varepsilon_{j,t-1} \end{bmatrix} + \begin{bmatrix} c_1 \\ c_2 \end{bmatrix} \varepsilon_{z,t-1} + \beta F_{t-1} + \begin{bmatrix} \varphi_{i,t} \\ \varphi_{j,t} \end{bmatrix}$$
(A.4)

where F_t might contain common factors, and with i, j, z = 1, 2, ..., N, and $i \neq j \neq z$. Obviously, we have N-2 possible background series for each pair of GARCH standardized residuals i, j.

The Granger causality test must therefore discriminate between the various choices of z, conditional to the possible presence of common factors F_t . The approach of Billio et al. (2012) made use of information criteria to specify the VAR lag structure and select the background series. Once the background series had been selected, the authors ran a set of Granger causality tests including/excluding the background series and/or the common factors, and selected the test outcome with the highest p-value (of the Granger's causality test) to produce more robust results.

We point out that the approach is computationally intensive because detecting the causality from i to j, for a specific choice of i and j, involves estimating N-2 bivariate systems to select the background series, and then further estimates to perform the causality testing.

For each pair of GARCH standardized residuals i and j, the test enables us to check for the presence of causality from i to j, with a decision rule associated with the above-described procedure. The decision rule provides for just two outcomes, a 0 or a 1: if causality is present, the approach of Billio et al. (2012) leads to the creation of a connection from node i to node j. The adjacency matrix gives rise to a directed and unweighted network.

Following the intuition of Billio et al. (2012), we interpret the adjacency matrix based on Granger causality as a proxy for a channel of shocks spreading and we take it as exogenous. Furthermore, before estimating our model, we proceed to a normalization of the network.

A.2 Quantile causality: a baseline case

Granger causality tests focus on the possible presence of causality relations affecting the mean dynamic evolution of x_t and y_t . As noted earlier, this approach might easily be extended to testing causality among higher-order moments.

Despite the usefulness and simplicity of this approach, focusing on the moments alone might prove too restrictive, since it could preclude or limit the possibility of studying the causal relations between x_t and y_t in specific regions of their distributions, or the risk measures associated with the distributions of the two variables, such as quantiles or conditional moments.

For instance, there might be situations where x_t and y_t are not linked by causal relations in the center of their distributions, or in the mean, but they might still be strongly dependent on one another in the tails of their distributions. This phenomenon could be crucial in the context of financial returns (given that their distributions are typically characterized by fat tails, Cont (2001)), or when we are interested in the existence of causal relations during periods of market turmoil (when both the variables analyzed take values in their tails), or during a negative period for the target variable (when only the dependent variable takes values in its tail). There might also be more interesting cases in which the causal relations are asymmetrical between the left and right tails. The classical Granger's test and its baseline variants are obviously unable to detect such structural features.

This shortcoming can be overcome by building causality tests based on the quantile regression method introduced by Koenker and Bassett (1978). In such a framework, we postulate the existence of a linear relationship between the quantile of a target variable and the values taken by an explanatory variable. If we deal with two variables, we thus have two equations, and the bivariate system becomes:

$$Q_{y_t}(\tau) = \beta_{0,1}(\tau) + \beta_{1,1}(\tau)y_{t-1} + \beta_{2,1}(\tau)x_{t-1}$$

$$Q_{x_t}(\theta) = \beta_{0,2}(\theta) + \beta_{1,2}(\theta)y_{t-1} + \beta_{2,2}(\theta)x_{t-1}$$
(A.5)

where $Q_{y_t}(\tau)$ and $Q_{x_t}(\theta)$ denote the τ -th and θ -th conditional quantiles of y_t and x_t , respectively. As in the idea behind Granger causality, testing for causality from x_t to the τ -th conditional quantile of y_t involves testing the significance of the coefficient $\beta_{2,1}(\tau)$. Likewise, testing for causality from y_t to the θ -th conditional quantile of x_t involves testing the significance of $\beta_{1,2}(\theta)$.

Unlike the Granger causality testing approach, the estimation of the conditional quantiles is based on the marginal model for each target variable. We therefore test for causality from x_t to y_t , and from y_t to x_t using two different models (i.e. two different conditional quantile linear models, one for each target variables). A recently-developed system approach is also viable, as discussed later on.

Financial variables are often highly correlated, due mainly to their sensitivity to market trends and shocks. The influence of these common factors can be isolated by including one or more control variables reproducing the market movements in the bivariate system (A.5). If we let \mathbf{F}_t be the vector of these common factor variables observed in t, we rewrite the (A.5) as

$$Q_{y_t}(\tau) = \beta_{0,1}(\tau) + \beta_{1,1}(\tau)y_{t-1} + \beta_{2,1}(\tau)x_{t-1} + \gamma_1 F'_{t-1}$$

$$Q_{x_t}(\theta) = \beta_{0,2}(\theta) + \beta_{1,2}(\theta)y_{t-1} + \beta_{2,2}(\theta)x_{t-1} + \gamma_2 F'_{t-1}$$
(A.6)

The latter conditional quantiles thus mimic the presence of common factors used in the Billio et al. (2012) approach. Similarly, we can introduce a further background variable, and increase the lag structure within the conditional quantile specification. Note that we could introduce lags for both the conditioning and the modelled variables.

If we focus on the conditional quantile for x_t (similarly for y_t), with or without additional control or background variables, the detection of causality is associated with the presence of a significant impact of y_{t-1} (or all lags if the models structure is more complex) on the conditional quantile of x_t .

Within a quantile regression framework, we can also assess the causality impact of x_t by comparing the first equation in (A.5) with its restricted version:

$$Q_{y_t}^{(r)}(\tau) = \beta_{0,1}^{(r)}(\tau) + \beta_{1,1}^{(r)}(\tau)y_{t-1}, \tag{A.7}$$

on the basis of the testing procedure proposed by Koenker and Bassett (1982). The test verifies the null hypothesis that the additional variable in the first equation belonging to the system (A.5), i.e. x_{t-1} , does not improve on the goodness-of-fit achieved with the restricted model given in (A.7). If the null hypothesis of the test, which reads like the F-test, is rejected at a given significance level, then there is evidence of causality from x_t to the τ -th conditional quantile of y_t . The same method could likewise be applied to test the causality relations from y_t to the θ -th conditional quantile of x_t .

Finally, we note that the quantile causality network is specific to the τ quantile used to test the link between x_t and y_t . The above-outlined method might therefore generate a collection of quantile causality networks. When the focus is on the risk, however, our interest lies in the left tail of the distribution, and therefore on small values of τ , ideally between 1% and 10%.

Using conditional quantile causality testing based on an assessment of the significance of the coefficients, we thus obtain the quantile causality network at quantile τ . We stress that the estimated network depends on the chosen quantile, so we can estimate a collection of quantile-based causality networks.

A.3 Quantile-on-quantile causality

So far, we have considered the causal impact of one variable on another, focusing either on moments (within a Granger causality framework) or on conditional quantiles. We can take the analysis further by directly linking the quantiles of both the causal and the dependent variables. In other words, our aim is to check whether the θ -th quantile of x_t causes the τ -th quantile of y_t , and vice versa, for θ equaling or differing from τ . In this way, we can test whether the power of x_t in causing the τ -th quantile of y_t changes over $\theta \in (0,1)$.

From a different viewpoint, while in traditional quantile regression we analyze the possible impact of the values taken by x_{t-1} (across its whole density) on the conditional quantile of y_t , for instance, we are interested here in testing the existence of causality when we restrict our attention to a neighborhood of a quantile of x_{t-1} . This would enable us to seek causal relations when both the analyzed variables take values in the respective tails, for example.

Two different approaches can be used for this purpose. The first is based on the method

introduced by Sim and Zhou (2015), who proposed a modified version of quantile regression in which observations (and then the corresponding loss function) are weighted using a kernel function. In particular, Sim and Zhou (2015) use kernel-based weights to estimate the relations in quantiles between oil prices and stock returns. In our framework, following Sim and Zhou (2015), the conditional quantiles in (A.5) are rewritten, respectively, as follows:

$$Q_{y_t}(\tau,\theta) = \beta_{0,1}(\tau,\theta) + \beta_{1,1}(\tau,\theta)y_{t-1} + \beta_{2,1}(\tau,\theta)x_{t-1}$$

$$Q_{x_t}(\theta,\tau) = \beta_{0,2}(\theta,\tau) + \beta_{1,2}(\theta,\tau)y_{t-1} + \beta_{2,2}(\theta,\tau)x_{t-1}$$
(A.8)

where the only difference lies in the fact that the parameters have both τ and θ subscripts. In fact, the estimated parameters depend on the quantile levels of both y_t and x_t . Despite the similarities in the form, the equations in system (A.8) differ from those in (A.5) in the manner in which the unknown parameters are estimated. Notably, the coefficients belonging to the first equation of the system (A.8) come from the following minimization problem:

$$\min_{\beta_{0,1}(\tau,\theta),\beta_{1,1}(\tau,\theta),\beta_{2,1}(\tau,\theta)} \sum_{t=1}^{T} \rho_{\tau} \left[y_{t} - \beta_{0,1}(\tau,\theta) - \beta_{1,1}(\tau,\theta) y_{t-1} - \beta_{2,1}(\tau,\theta) x_{t-1} \right] * \\
*K \left(\frac{F_{T}(x_{t}) - \theta}{h} \right), \tag{A.9}$$

where $\rho_{\tau}(e) = e(\tau - \mathbf{I}_{\{e<0\}})$ is the asymmetric loss function on the basis of the quantile regression method introduced by Koenker and Bassett (1978), and $\mathbf{I}_{\{\cdot\}}$ is the indicator function, taking a value of 1 if the condition in $\{\cdot\}$ is true, and a value of 0 otherwise. The difference with respect to the classical quantile regression of Koenker and Bassett (1978) lies in $K(\cdot)$, i.e. the kernel function, with bandwidth h, where $F_T(x_t) = T^{-1} \sum_{k=1}^T \mathbf{I}_{\{x_k < x_t\}}$, to focus on the impact of x_t in the neighborhood of its θ -th quantile. The parameters in the second equation of the system (A.8) are likewise estimated from

$$\min_{\beta_{0,2}(\theta,\tau),\beta_{1,2}(\theta,\tau),\beta_{2,2}(\theta,\tau)} \sum_{t=1}^{T} \rho_{\theta} \left[x_{t} - \beta_{0,2}(\theta,\tau) - \beta_{1,2}(\theta,\tau) y_{t-1} - \beta_{2,2}(\theta,\tau) x_{t-1} \right] *$$

$$*K \left(\frac{F_{T}(y_{t}) - \tau}{h} \right).$$
(A.10)

If we remove the kernel function, the parameters no longer depend on θ , and we return to the traditional quantile regression estimator. As discussed in the previous section, we can generalize the conditional quantiles in (A.8) by adding common factors and background variables. Notably, what Sim and Zhou (2015) call the *quantile-on-quantile* approach corresponds to a non-parametric quantile regression where the knots used to obtain the local quantiles are fixed at specific quantiles of the dependent variable, Koenker (2005).

The construction of a quantile-causality network by following the quantile-on-quantile approach of Sim and Zhou (2015) builds on the significance of the coefficients associated with the impact of the covariates quantiles. In particular, focusing on x_t , the existence of causality from the θ quantile of y_t to the τ quantile of x_t depends on the coefficient $\beta_{1,2}(\theta,\tau)$. If the conditional quantiles have a more complex structure, including several lags, the number of coefficients to be tested naturally increases. When accounting for the presence of common factors and background

variables, we can again follow a procedure similar to that of Billio et al. (2012).

When estimating the quantile causality network by means of the quantile-on-quantile method, the estimated network depends on two quantile levels: one referring to the dependent variable and the other to the explanatory variable. As in the previous section, if we are interested in monitoring the causality networks during periods of financial turmoil, both quantiles would be placed in the range of 1% - 10%.

We note that such an approach has some similarities with the VAR for VaR model introduced by White et al. (2015), which, in a multivariate framework, estimates the dependence across quantiles for a collection of series, also accounting for the possible presence of an auto-regressive structure, in the spirit of the CAViaR by Engle and Manganelli (2004). Clearly, the quantile-on-quantile dependence might be seen as a special case.

A.4 Quantile causality: a nonparametric test

The previous quantile regression-based approaches for detecting causality have a relevant feature in common, that is their parametric nature. In this section, we refer to another approach that is non-parametric. Non-parametric techniques offer the important advantage of requiring no particular assumption concerning the distributions for the processes underlying the variables of interest. We refer here to the nonparametric testing procedure proposed by Jeong et al. (2012).

First, we assume that the conditional variable is x_t , and we define the following vectors: $\mathcal{Y}_{t-1} \equiv (y_{t-1}, ..., y_{t-p}), \ \mathcal{X}_{t-1} \equiv (x_{t-1}, ..., x_{t-q}), \ \mathcal{Z}_{t-1} \equiv (y_{t-1}, ..., y_{t-p}, x_{t-1}, ..., x_{t-q}), \ \text{for } (p,q) > 1$. Note that the vectors refer to the lags of one or both the variables of interest. Further, $F_{y_t|\mathcal{Z}_{t-1}}(y_t|\mathcal{Z}_{t-1})$ and $F_{y_t|\mathcal{Y}_{t-1}}(y_t|\mathcal{Y}_{t-1})$ denote the distributions of y_t , conditional on \mathcal{Z}_{t-1} and \mathcal{Y}_{t-1} , respectively; the distribution of y_t is assumed to be absolutely continuous in y for almost all $\nu = (\mathcal{Y}, \mathcal{Z})$. For the sake of simplicity, we denote the τ -th quantile of y_t conditional on \mathcal{Z}_{t-1} as $Q_{\tau}(\mathcal{Z}_{t-1})$ and the τ -th quantile of y_t conditional on \mathcal{Y}_{t-1} as $Q_{\tau}(\mathcal{Y}_{t-1})$.

The definition of causality in quantiles in Jeong et al. (2012) focuses on the conditional quantiles of the series densities: x_t does not cause y_t in its τ -th quantile, with respect to \mathcal{Z}_{t-1} , if $Q_{\tau}(\mathcal{Z}_{t-1}) = Q_{\tau}(\mathcal{Y}_{t-1})$; conversely, x_t causes y_t in its τ -th quantile, with respect to \mathcal{Z}_{t-1} , if $Q_{\tau}(\mathcal{Z}_{t-1}) \neq Q_{\tau}(\mathcal{Y}_{t-1})$. From the previous definition, the hypotheses to be tested read as follows:

$$\begin{cases}
H_0: P[F_{y_t|\mathcal{Z}_{t-1}}(Q_{\tau}(\mathcal{Y}_{t-1})|\mathcal{Z}_{t-1}) = \tau] = 1 \\
H_1: P[F_{y_t|\mathcal{Z}_{t-1}}(Q_{\tau}(\mathcal{Y}_{t-1})|\mathcal{Z}_{t-1}) = \tau] < 1
\end{cases}$$
(A.11)

The non-parametric test developed by Jeong et al. (2012) is based on a measure of distance defined as:

$$J_T = \mathbb{E}\left[[F_{y_t|\mathcal{Z}_{t-1}}(Q_\tau(\mathcal{Y}_{t-1})|\mathcal{Z}_{t-1}) - \tau]^2 g_{\mathcal{Z}_{t-1}}(\mathcal{Z}_{t-1}) \right], \tag{A.12}$$

where $g_{\mathcal{Z}_{t-1}}(\mathcal{Z}_{t-1})$ is the marginal density function of \mathcal{Z}_{t-1} . J_T is estimated using the feasible kernel-based estimator:

$$\widehat{J}_T = \frac{1}{T(T-1)h^m} \sum_{t=1}^T \sum_{s \neq t} K\left(\frac{\mathcal{Z}_{t-1} - \mathcal{Z}_{s-1}}{h}\right) \widetilde{\epsilon}_t \widetilde{\epsilon}_s, \tag{A.13}$$

where m = p + q, $K(\cdot)$ is the kernel function with bandwidth h, $\tilde{\epsilon}_t = \mathbf{I}_{\{y_t \leq \widetilde{Q}_\tau(\mathcal{Y}_{t-1})\}} - \tau$, $\mathbf{I}_{\{\cdot\}}$ being the indicator function that takes a value of 1 if the condition in $\{\cdot\}$ is true, and zero otherwise. In Jeong et al. (2012), $\widetilde{Q}_\tau(\mathcal{Y}_{t-1}) \equiv \widetilde{F}_{y_t|\mathcal{Y}_{t-1}}^{-1}(\tau|\mathcal{Y}_{t-1})$, where

$$\widetilde{F}_{y_t|\mathcal{Y}_{t-1}}(y_t|\mathcal{Y}_{t-1}) = \frac{\sum_{s \neq t} C_{t-1,s-1} \mathbf{1}_{\{y_s \leq y_t\}}}{\sum_{s \neq t} C_{t-1,s-1}}$$
(A.14)

is the Nadaraya-Watson kernel estimator of $F_{y_t|\mathcal{Y}_{t-1}}(y_t|\mathcal{Y}_{t-1})$, with the kernel function $C_{t-1,s-1} = C(\mathcal{Y}_{t-1} - \mathcal{Y}_{s-1})/a$, and a is the bandwidth.

Under a set of precise assumptions, Jeong et al. (2012) derived the limiting distribution for the test statistic

$$Th^{m/2}\widehat{J}_T \stackrel{L}{\to} \mathcal{N}(0, \sigma_0^2),$$
 (A.15)

where

$$\sigma_0^2 = 2\mathbb{E}\left[\sigma_{\epsilon}^4(\mathcal{Z}_{t-1})g_{\mathcal{Z}_{t-1}}(\mathcal{Z}_{t-1})\right] \left(\int K^2(u)du\right),\tag{A.16}$$

and $\sigma_{\epsilon}^2(\mathcal{Z}_{t-1}) = \tau(1-\tau)$.

The unknown parameter σ_0^2 is estimated as

$$\widehat{\sigma}_0^2 = \frac{2\tau^2 (1-\tau)^2}{T(T-1)h^m} \sum_{s \neq t} K^2 \left(\frac{\mathcal{Z}_{t-1} - \mathcal{Z}_{s-1}}{h} \right). \tag{A.17}$$

Then, after estimating all the quantities of interest, the standardized statistic can be computed:

$$\widehat{J}_T^* = \frac{Th^{m/2}\widehat{J}_T}{\widehat{\sigma}_0}. (A.18)$$

The above-described method could likewise be applied to test the causality impact of y_t on the quantiles of x_t , simply by inverting the roles of x_t and y_t .

If we detect quantile causality using the non-parametric test of Jeong et al. (2012), then we have a dependence on the chosen quantile, as in the previous quantile-based causality cases. To keep the focus on the risk side, we suggest using quantile levels in the range 1%-10%.

B Banks and Insurance companies

In the empirical analyses we use the bank or the insurance companies as in the following lists. Within parentheses we report the ticker. A single star * identifies companies included only in the first sample - 2006 to 2008 - while a double star identifies companies included only in the second sample - 2011-2015. When stars are not present, the companies are included in both samples.

Banks:

Astoria Financial Corporation (AF*), Associated Banc-Corp (ASB), Bank of America (BAC), Bancfirst Corporation (BANF*), Credicorp (BAP), BB&T (BBT), Bbx Capital Corporation

(BBX*), Bank Mutual Corporation (BKMU*), Bank of Hawaii (BOH), Bok Financial Corporation (BOKF), Boston Private Financial Holdings (BPFH*), Brookline Bancorp (BRKL*), Bancorpsouth (BXS*), Citigroup (C), Cathay General Bancorp (CATY*), Commerce Bancshares (CBSH), Capitol Federal Financial (CFFN), Cullen/Frost Bankers (CFR), Chemical Financial Corporation (CHFC*), Comerica (CMA), City National (CN), Central Pacific Financial Corporation (CPF*), CVB Financial Corporation (CVBF*), Doral Financial Corporation (DRLCQ*), East West Bancorp (EWBC), First Horizon National (FHN**), Fifth Third Bancorp (FITB**), Firstmerit (FMER**), First Niagara Financial Group (FNFG**), First Republic Bank (FRC**), Fulton Financial Corporation (FULT**), Huntington Bancshares Incorporated (HBAN**), Hudson City Bancorp (HCB**), Investors Bancorp (ISBC**), Jp Morgan Chase & Co. (JPM**), Keycorp (KEY**), M&T Bank (MTB**).

Insurances:

American Financial Group (AFG**), Aflac (AFL), American International Group (AIG), Assurant (AIZ), The Allstate Corporation (ALL), Berkshire Hathaway (BRKA), Brown & Brown (BRO), Cincinnati Financial Corporation (CINF), Cna Financial Corporation (CNA), Cno Financial Group (CNO*), Erie Indemnity Company (ERIE**), Genworth Financial (GNW), The Hartford Financial Services Group (HIG), Kemper (KMPR*), Loews (L), Lincoln National (LNC), Mbia (MBI*), Metlife (MET), Markel (MKL**), Marsh & Mclennan (MMC), Old Republic International Corporation (ORI*), Principal Financial Group (PFG), Progressive Ohio (PGR), Prudential Financial (PRU), Reinsurance Group of America (RGA**), Torchmark (TMK), The Travelers Companies (TRV), Unum Group (UNM), W.R. Berkley Corporation (WRB).

C Descriptive statistics

STOCK	MEAN	MED	ST DEV	MIN	MAX	$1^{st} Q$	$3^{rd} Q$	SKEW	KURT	LB_1	LB_5	LB_{10}	SLB_1	SLB_5	SLB_{10}	MV
\mathbf{C}	-0.262	-0.127	4.296	-30.661	45.632	-1.113	0.711	0.673	30.033	5.716	0.000	0.002	0.000	0.000	0.000	236630.90
BAC	-0.157	-0.039	3.765	-30.416	24.060	-0.977	0.688	-0.468	19.555	72.686	0.295	0.126	0.000	0.000	0.000	212543.10
BBT	-0.056	-0.070	3.108	-26.608	21.198	-1.167	0.849	-0.094	18.219	0.326	1.159	0.000	0.000	0.000	0.000	21015.19
CMA	-0.139	-0.102	3.361	-20.720	18.805	-1.266	0.885	0.049	10.555	23.288	6.167	0.032	0.000	0.000	0.000	9392.46
ASB	-0.058	-0.030	2.905	-18.999	19.354	-0.916	0.705	0.076	14.517	94.558	0.339	0.009	0.107	0.000	0.000	4619.78
$^{\mathrm{CN}}$	-0.053	0.000	2.730	-17.379	20.213	-0.964	0.827	0.048	13.119	0.080	0.035	0.003	0.000	0.000	0.000	3763.18
CBSH	-0.003	-0.040	2.113	-12.894	15.113	-0.808	0.651	0.696	13.581	0.000	0.000	0.000	0.000	0.000	0.000	3428.85
AF	-0.077	-0.096	2.676	-18.972	13.698	-1.172	0.820	-0.251	11.170	0.054	0.032	0.011	0.000	0.000	0.000	3162.41
BOKF	-0.016	-0.018	2.228	-13.426	12.315	-0.848	0.764	0.055	12.242	63.004	0.283	0.052	0.000	0.000	0.000	3135.55
CFR	-0.008	-0.051	2.302	-13.183	16.306	-0.915	0.852	0.539	13.505	0.007	0.000	0.000	0.000	0.000	0.000	2853.67
BOH	-0.017	-0.019	2.458	-25.508	12.946	-0.859	0.834	-1.279	23.778	0.009	0.002	0.006	0.000	0.000	0.000	2695.89
BAP	0.104	0.089	2.572	-17.439	11.226	-1.017	1.368	-0.507	8.735	76.155	6.593	1.739	0.000	0.000	0.000	2572.86
CFFN	0.043	0.052	1.636	-7.664	6.606	-0.619	0.755	-0.127	6.254	10.379	14.081	0.059	0.000	0.000	0.000	2366.73
EWBC	-0.109	-0.155	3.730	-23.669	24.166	-1.255	0.869	0.263	12.930	47.310	11.160	19.360	0.000	0.000	0.000	2089.00
BXS	0.008	-0.077	2.854	-15.116	19.711	-1.165	1.088	0.592	11.463	6.615	0.065	0.173	0.000	0.000	0.000	1895.44
CATY	-0.055	-0.121	3.631	-23.886	21.314	-1.504	1.227	0.167	11.957	39.151	1.648	1.971	0.000	0.000	0.000	1865.30
CVBF	-0.029	-0.145	3.257	-18.965	20.210	-1.419	1.101	0.702	10.139	1.833	1.097	3.680	0.000	0.000	0.000	1241.16
BPFH	-0.198	-0.138	3.671	-30.902	17.640	-1.403	1.161	-1.226	17.170	13.155	2.258	2.606	0.007	0.001	0.000	1171.49
DRLCQ	-0.443	-0.522	6.085	-36.564	36.891	-2.885	1.968	-0.166	9.962	84.586	85.266	69.825	1.543	0.144	0.000	1097.65
CPF	-0.169	-0.199	3.925	-22.119	29.376	-1.668	1.259	0.553	12.079	99.950	0.257	1.170	0.000	0.000	0.000	1042.41
BRKL	-0.038	-0.076	2.461	-14.973	11.878	-1.119	0.990	0.158	7.816	0.088	0.175	0.259	0.000	0.000	0.000	918.22
BBX	-0.330	-0.235	6.086	-48.290	35.188	-2.062	1.308	-0.797	16.628	15.046	30.970	3.020	0.000	0.000	0.000	828.74
CHFC	-0.017	-0.034	2.904	-19.297	17.823	-1.368	1.131	0.267	10.009	0.173	1.038	6.583	0.035	0.000	0.000	775.59
BKMU	0.011	-0.080	1.970	-14.235	9.307	-0.844	0.917	-0.048	9.507	0.000	0.000	0.000	0.076	0.000	0.000	718.23
BANF	0.039	0.000	2.709	-24.878	21.243	-1.325	1.233	0.039	19.231	0.158	0.044	0.667	0.000	0.000	0.000	675.88

Table C.9: Descriptive statistics of the daily returns generated by the 25 U.S. banks with the largest market value, as recorded at the beginning of 2006, in the period between January 3, 2006 and December 31, 2008. From left to right the table reports, for each stock, the mean (%), the median (%), the standard deviation (%), the minimum (%), the maximum (%), the first and the third quartiles (%), the skewness, the kurtosis, the p-value of the Ljung-Box test (%), at the significance level of the 5%, applied for both the returns (LB) and the squared returns (SLB) with lags equal to 1, 5 and 10, respectively, and the market value recorded in January 2006.

STOCK	MEAN	MED	ST DEV	MIN	MAX	1^{st} Q	3^{rd} Q	SKEW	KURT	LB_1	LB_5	LB_{10}	SLB_1	SLB_5	SLB_{10}	MV
JPM	0.035	0.048	1.698	-9.888	8.101	-0.789	0.901	-0.206	7.043	0.338	0.628	0.272	0.000	0.000	0.000	186018.60
BAC	0.018	0.000	2.295	-22.713	15.481	-1.050	1.103	-0.508	14.427	1.340	0.000	0.000	0.000	0.000	0.000	136331.90
$^{\mathrm{C}}$	0.007	0.000	2.129	-17.934	12.968	-0.944	0.981	-0.533	10.310	8.722	0.001	0.001	0.000	0.000	0.000	132889.30
BBT	0.029	0.065	1.497	-11.274	6.693	-0.716	0.890	-0.685	8.638	0.001	0.000	0.000	0.000	0.000	0.000	18978.31
FITB	0.025	0.106	1.736	-12.067	8.709	-0.817	0.908	-0.430	8.619	0.001	0.000	0.000	0.000	0.000	0.000	12549.82
MTB	0.026	0.034	1.355	-8.059	6.554	-0.624	0.732	-0.222	6.892	0.008	0.000	0.001	0.000	0.000	0.000	10488.55
KEY	0.032	0.074	1.807	-10.987	8.319	-0.906	1.031	-0.228	7.187	0.000	0.000	0.000	0.000	0.000	0.000	8339.36
BAP	-0.016	-0.009	1.711	-20.843	6.231	-0.918	0.894	-1.487	21.228	41.290	15.174	10.460	9.994	0.071	1.654	7814.11
CMA	-0.001	0.041	1.772	-11.118	6.100	-0.879	0.942	-0.701	7.149	18.082	0.247	0.554	0.000	0.000	0.000	6795.13
HBAN	0.038	0.102	1.794	-10.472	8.542	-0.931	1.059	-0.301	6.869	0.000	0.000	0.000	0.000	0.000	0.000	5750.23
HCB	-0.018	0.000	1.805	-11.304	14.569	-0.748	0.793	-0.325	11.772	0.010	0.178	1.702	0.001	0.000	0.000	5161.57
FRC	0.065	0.064	1.546	-16.359	8.376	-0.715	0.921	-0.901	15.518	0.004	0.016	0.081	0.064	0.013	0.047	3821.80
CFR	-0.001	0.066	1.396	-8.121	7.493	-0.677	0.758	-0.102	6.964	0.004	0.000	0.001	0.000	0.000	0.000	3667.12
BOKF	0.009	0.059	1.403	-9.594	6.062	-0.709	0.720	-0.390	7.708	0.005	0.003	0.097	0.000	0.000	0.000	3553.32
CBSH	0.025	0.090	1.316	-6.635	6.047	-0.697	0.776	-0.307	5.971	0.002	0.001	0.004	0.000	0.000	0.000	3552.96
EWBC	0.060	0.091	1.747	-9.353	8.752	-0.867	0.992	-0.158	6.443	0.412	0.009	0.000	0.000	0.000	0.000	3321.22
$^{\mathrm{CN}}$	0.030	0.000	1.577	-8.200	17.330	-0.687	0.821	0.850	17.467	6.143	0.012	0.009	8.723	0.002	0.000	3083.03
FHN	0.017	0.032	1.884	-11.185	7.559	-1.003	1.034	-0.415	6.255	6.398	0.799	2.454	0.000	0.000	0.000	2999.39
FNFG	-0.020	0.017	1.708	-14.511	13.548	-0.878	0.922	-0.556	13.534	3.662	2.318	0.807	4.568	2.718	1.949	2907.62
ASB	0.017	0.059	1.666	-12.420	7.165	-0.858	1.005	-0.669	8.529	0.003	0.000	0.000	0.000	0.000	0.000	2538.15
BOH	0.023	0.085	1.323	-8.229	5.795	-0.671	0.799	-0.296	6.328	0.027	0.005	0.031	0.000	0.000	0.000	2272.40
FULT	0.018	0.085	1.678	-10.064	8.031	-0.806	0.913	-0.389	7.399	0.006	0.000	0.001	0.000	0.000	0.000	2216.42
CFFN	0.004	0.000	0.915	-5.721	5.001	-0.504	0.498	-0.252	7.127	0.000	0.000	0.001	0.000	0.000	0.000	1880.95
FMER	-0.005	0.058	1.765	-11.931	9.523	-0.951	0.999	-0.422	8.063	2.032	0.284	0.080	0.000	0.000	0.000	1855.60
ISBC	0.070	0.056	1.326	-8.360	7.474	-0.570	0.734	0.004	7.640	0.000	0.000	0.000	0.000	0.000	0.000	1645.87

Table C.10: Descriptive statistics of the daily returns generated by the 25 U.S. banks with the largest market value, as recorded at the beginning of 2011, in the period between January 3, 2011 and December 31, 2015. From left to right the table reports, for each stock, the mean (%), the median (%), the standard deviation (%), the minimum (%), the maximum (%), the first and the third quartiles (%), the skewness, the kurtosis, the p-value of the Ljung-Box test (%), at the significance level of the 5%, applied for both the returns (LB) and the squared returns (SLB) with lags equal to 1, 5 and 10, respectively, and the market value recorded in January 2011.

STOCK	MEAN	MED	ST DEV	MIN	MAX	1^{st} Q	3^{rd} Q	SKEW	KURT	LB_1	LB_5	LB_{10}	SLB_1	SLB_5	SLB_{10}	MV
AIG	-0.500	-0.055	6.469	-93.626	35.853	-0.853	0.681	-5.387	75.716	0.000	0.000	0.000	0.000	0.000	0.000	164653.50
BRKA	0.011	-0.001	1.717	-12.883	14.953	-0.523	0.536	0.622	24.924	93.137	0.610	0.965	0.000	0.000	0.000	110164.70
MET	-0.045	0.000	3.687	-31.156	24.686	-0.991	0.915	-0.396	23.343	40.365	1.162	3.180	0.000	0.000	0.000	37890.41
PRU	-0.117	-0.055	3.973	-26.327	32.390	-0.989	0.964	0.207	22.244	20.640	0.081	0.203	0.000	0.000	0.000	36860.64
ALL	-0.066	0.000	2.761	-23.799	19.628	-0.735	0.621	-0.579	28.043	93.526	0.017	0.000	0.000	0.000	0.000	32610.33
TRV	0.002	0.000	2.655	-20.067	22.758	-0.917	0.850	0.438	20.729	0.000	0.000	0.000	0.000	0.000	0.000	28563.55
$_{ m HIG}$	-0.219	-0.034	6.060	-72.486	70.487	-1.020	0.935	-0.386	64.131	15.143	0.000	0.000	36.736	0.000	0.000	24387.18
AFL	-0.002	0.021	2.601	-18.902	14.961	-0.801	0.700	-0.530	18.381	0.100	0.001	0.000	0.000	0.000	0.000	22472.57
PGR	-0.090	-0.082	2.382	-13.947	21.490	-1.010	0.888	0.536	17.712	6.330	0.000	0.000	0.006	0.000	0.000	20556.40
${ m L}$	-0.015	0.091	2.745	-19.939	21.220	-0.906	0.946	-0.820	20.944	0.010	0.001	0.011	0.000	0.000	0.000	18655.31
LNC	-0.137	0.018	4.884	-50.891	36.235	-0.981	0.931	-1.672	39.635	3.661	0.073	0.093	0.000	0.000	0.000	16040.52
MMC	-0.036	-0.038	2.033	-13.075	11.690	-0.924	0.829	0.078	10.357	0.000	0.000	0.000	0.000	0.000	0.000	15753.62
PFG	-0.098	0.000	4.252	-31.978	34.190	-0.938	1.053	0.098	20.411	61.640	0.000	0.001	0.000	0.000	0.000	13717.85
GNW	-0.332	-0.088	7.528	-78.552	63.599	-1.295	0.939	-0.014	44.676	0.255	0.000	0.000	0.000	0.000	0.000	13158.24
CNA	-0.091	0.000	3.326	-39.476	24.106	-1.058	0.935	-2.559	39.779	97.177	5.119	0.584	0.000	0.000	0.000	7953.97
MBI	-0.357	-0.087	6.493	-41.264	38.219	-1.786	1.198	-0.076	12.808	4.703	0.584	5.430	0.000	0.000	0.000	7842.20
WRB	-0.003	-0.095	2.284	-11.607	15.049	-1.062	0.903	0.931	12.003	0.007	0.003	0.008	0.000	0.000	0.000	7607.63
CINF	-0.057	-0.022	2.653	-22.399	16.765	-0.836	0.729	-0.559	18.647	0.001	0.000	0.000	0.000	0.000	0.000	7239.32
AIZ	-0.049	0.034	3.069	-28.679	19.721	-0.869	0.903	-0.962	24.650	29.164	0.012	0.000	0.000	0.000	0.000	6239.33
TMK	-0.029	0.017	2.208	-15.228	13.967	-0.633	0.691	-0.495	17.199	54.298	0.005	0.079	0.000	0.000	0.000	5853.66
UNM	-0.027	-0.049	3.428	-35.145	20.010	-1.051	0.984	-1.268	28.910	0.282	0.000	0.005	0.000	0.000	0.000	5751.81
ORI	-0.075	0.000	3.340	-29.214	31.979	-0.932	0.792	0.115	28.746	1.547	0.000	0.000	0.000	0.000	0.000	5033.62
BRO	-0.050	0.000	2.077	-17.218	11.196	-0.867	0.835	-0.630	13.710	18.359	15.353	23.086	0.378	0.001	0.000	4632.19
CNO	-0.198	0.000	6.043	-55.048	59.157	-1.009	0.851	-0.891	36.127	0.000	0.000	0.000	0.000	0.000	0.000	3741.03
KMPR	-0.138	-0.045	2.827	-16.981	20.799	-1.066	0.871	-0.261	16.077	36.300	0.347	0.002	0.000	0.000	0.000	3326.30

Table C.11: Descriptive statistics of the daily returns generated by the 25 U.S. insurance companies with the largest market value, as recorded at the beginning of 2006, in the period between January 3, 2006 and December 31, 2008. From left to right the table reports, for each stock, the mean (%), the median (%), the standard deviation (%), the minimum (%), the maximum (%), the first and the third quartiles (%), the skewness, the kurtosis, the p-value of the Ljung-Box test (%), at the significance level of the 5%, applied for both the returns (LB) and the squared returns (LB) with lags equal to 1, 5 and 10, respectively, and the market value recorded in January 2006.

STOCK	MEAN	MED	ST DEV	MIN	MAX	1^{st} Q	3^{rd} Q	SKEW	KURT	LB_1	LB_5	LB_{10}	SLB_1	SLB_5	SLB_{10}	MV
BRKA	0.039	0.004	1.089	-6.289	7.791	-0.558	0.601	0.454	9.381	0.000	0.000	0.000	0.000	0.000	0.000	115505.80
AIG	0.020	0.071	1.892	-10.580	9.819	-0.831	0.962	-0.272	6.851	18.211	0.000	0.001	0.000	0.000	0.000	61232.09
MET	0.006	0.018	1.897	-10.460	8.551	-0.923	1.067	-0.299	5.819	21.753	0.372	0.823	0.000	0.000	0.000	47239.25
PRU	0.026	0.037	1.864	-11.469	8.822	-0.929	1.134	-0.386	6.904	2.498	0.004	0.002	0.000	0.000	0.000	29888.47
TRV	0.056	0.085	1.154	-7.893	6.205	-0.554	0.685	-0.181	7.791	0.017	0.346	0.141	0.000	0.000	0.000	25156.82
AFL	0.005	0.037	1.588	-10.758	8.334	-0.751	0.772	-0.288	8.789	11.965	0.421	0.745	0.000	0.000	0.000	24977.64
${ m L}$	-0.001	0.021	1.130	-6.089	5.547	-0.598	0.620	-0.177	6.179	0.485	0.062	0.182	0.000	0.000	0.000	17509.66
ALL	0.053	0.045	1.294	-10.700	7.294	-0.602	0.725	-0.297	10.258	0.013	0.000	0.000	0.022	0.000	0.000	16607.85
MMC	0.056	0.034	1.166	-8.679	8.926	-0.561	0.706	0.128	9.663	0.001	0.000	0.002	0.000	0.000	0.000	16043.64
PGR	0.037	0.051	1.215	-7.178	6.798	-0.601	0.702	-0.294	7.486	0.147	0.173	1.376	0.000	0.000	0.000	14019.72
$_{ m HIG}$	0.039	0.104	2.068	-15.366	14.438	-0.912	1.020	-0.071	10.936	0.031	0.000	0.000	0.000	0.000	0.000	11897.86
PFG	0.026	0.136	1.835	-12.159	8.351	-0.849	1.004	-0.422	7.102	0.107	0.000	0.001	0.000	0.000	0.000	10108.24
LNC	0.047	0.096	2.169	-13.056	9.641	-1.007	1.132	-0.329	6.844	8.444	0.014	0.004	0.000	0.000	0.000	9350.80
UNM	0.025	0.057	1.527	-11.128	9.625	-0.762	0.945	-0.285	7.207	0.013	0.000	0.000	0.000	0.000	0.000	8137.82
CNA	0.021	0.035	1.348	-9.850	8.464	-0.648	0.666	-0.188	9.603	0.004	0.004	0.120	0.000	0.000	0.000	7966.17
GNW	-0.100	0.077	3.460	-48.533	15.461	-1.468	1.431	-2.801	39.098	20.702	0.816	8.069	46.063	83.411	99.480	6219.79
CINF	0.050	0.086	1.188	-7.681	6.446	-0.590	0.706	-0.183	7.535	0.000	0.000	0.000	0.000	0.000	0.000	5292.70
TMK	0.061	0.115	1.216	-9.837	8.233	-0.548	0.706	-0.303	10.318	0.000	0.000	0.000	0.000	0.000	0.000	5184.13
WRB	0.055	0.080	1.095	-6.691	5.396	-0.540	0.651	-0.018	6.488	0.140	0.853	0.484	0.000	0.000	0.000	4491.24
RGA	0.037	0.086	1.461	-11.464	10.052	-0.613	0.779	-0.880	12.999	0.024	0.000	0.004	0.000	0.000	0.000	4464.94
MKL	0.067	0.057	1.125	-10.812	6.529	-0.509	0.598	-0.560	14.716	0.004	0.008	0.093	0.000	0.000	0.000	4084.38
AIZ	0.059	0.097	1.395	-7.794	6.927	-0.736	0.884	-0.279	6.412	0.076	0.000	0.014	0.001	0.000	0.000	3730.86
BRO	0.023	0.030	1.271	-9.392	9.262	-0.624	0.688	-0.253	11.251	95.429	0.122	0.010	0.000	0.000	0.000	3726.44
AFG	0.064	0.102	1.092	-7.551	6.529	-0.529	0.659	-0.219	7.525	0.008	0.004	0.030	0.000	0.000	0.000	3648.48
ERIE	0.030	0.042	1.275	-9.423	8.906	-0.616	0.667	-0.089	10.218	0.014	0.272	1.656	0.000	0.000	0.000	3541.31

Table C.12: Descriptive statistics of the daily returns generated by the 25 U.S. insurance companies with the largest market value, as recorded at the beginning of 2011, in the period between January 3, 2011 and December 31, 2015. From left to right the table reports, for each stock, the mean (%), the median (%), the standard deviation (%), the minimum (%), the maximum (%), the first and the third quartiles (%), the skewness, the kurtosis, the p-value of the Ljung-Box test (%), at the significance level of the 5%, applied for both the returns (LB) and the squared returns (LB) with lags equal to 1, 5 and 10, respectively, and the market value recorded in January 2011.

							7								
POR	MEA	MED	STD	MIN	MAX	$1^{st} Q$	3^{rd} Q	SKE	KUR	LB_1	LB_5	LB_{10}	SLB_1	SLB_5	SLB_{10}
Agric	0.096	0.050	2.701	-15.270	18.240	-0.930	1.140	0.078	10.597	35.606	0.424	0.016	0.268	0.000	0.000
Food	0.010	0.030	1.165	-7.250	7.400	-0.485	0.580	-0.076	12.281	1.713	0.000	0.000	0.000	0.000	0.000
Soda	-0.012	0.070	1.629	-8.020	11.140	-0.675	0.715	0.054	10.820	93.185	3.668	0.045	0.000	0.000	0.000
Beer	0.031	0.040	1.187	-7.700	10.120	-0.480	0.530	0.811	18.398	0.018	0.000	0.000	0.000	0.000	0.000
Smoke	0.030	0.070	1.479	-7.400	13.310	-0.600	0.725	0.338	16.033	12.642	0.000	0.000	0.665	0.000	0.000
Toys	-0.040	0.010	1.791	-9.630	9.250	-0.855	0.780	-0.140	8.508	37.202	2.289	0.544	0.000	0.000	0.000
Fun	-0.083	0.050	2.423	-12.170	16.580	-0.900	0.830	0.023	11.578	0.003	0.000	0.002	0.000	0.000	0.000
Books	-0.124	-0.060	1.953	-11.240	11.180	-0.785	0.590	0.004	11.861	42.319	0.069	0.341	0.000	0.000	0.000
Hshld	0.010	0.030	1.260	-7.250	9.440	-0.455	0.520	0.077	13.616	0.002	0.000	0.000	0.000	0.000	0.000
Clths	-0.023	-0.010	2.012	-11.520	12.690	-0.905	0.895	0.099	8.974	48.150	0.212	0.680	0.000	0.000	0.000
Hlth	-0.038	0.020	1.348	-9.020	8.290	-0.610	0.545	-1.041	15.622	19.523	0.002	0.000	0.000	0.000	0.000
MedEq	-0.026	0.030	1.349	-7.210	11.690	-0.560	0.560	0.153	15.334	55.714	0.057	0.319	0.000	0.000	0.000
Drugs	0.004	0.020	1.258	-6.530	11.140	-0.485	0.580	0.459	16.120	0.017	0.000	0.000	0.000	0.000	0.000
Chems	-0.010	0.110	2.039	-11.380	13.060	-0.720	0.880	-0.416	11.100	4.333	0.002	0.010	0.000	0.000	0.000
Rubbr	-0.015	0.000	1.701	-10.140	7.800	-0.740	0.840	-0.384	7.555	0.178	1.160	11.422	0.000	0.000	0.000
Txtls	-0.052	-0.010	2.114	-12.230	10.030	-1.010	0.820	-0.143	8.827	14.373	2.701	0.045	0.002	0.000	0.000
BldMt	-0.050	0.020	1.889	-10.670	8.540	-0.835	0.795	-0.365	8.230	61.661	2.884	4.807	0.000	0.000	0.000
Cnstr	-0.057	-0.050	2.767	-12.400	15.560	-1.355	1.280	0.091	7.184	28.169	3.002	9.516	0.000	0.000	0.000
Steel	0.011	0.160	3.015	-15.930	20.060	-1.190	1.430	-0.033	10.453	54.236	0.540	0.193	0.000	0.000	0.000
FabPr	-0.023	0.100	2.126	-11.830	10.520	-0.965	0.900	-0.702	8.651	15.807	0.124	0.381	0.000	0.000	0.000
Mach	-0.016	0.150	2.144	-12.250	13.910	-0.895	1.025	-0.315	11.089	18.366	0.000	0.000	0.000	0.000	0.000
ElcEq	0.000	0.100	2.006	-13.100	14.080	-0.740	0.875	-0.039	12.589	0.492	0.000	0.000	0.000	0.000	0.000
Autos	-0.080	0.070	2.349	-11.230	11.700	-0.995	1.010	-0.062	8.630	99.063	1.838	3.530	0.000	0.000	0.000
Aero	-0.009	0.070	1.777	-7.650	13.570	-0.725	0.805	0.497	12.728	2.192	0.000	0.000	0.000	0.000	0.000
Ships	0.004	0.030	1.715	-9.400	10.620	-0.810	0.815	-0.110	9.046	0.011	0.001	0.001	0.000	0.000	0.000
Guns	0.050	0.040	1.669	-9.000	10.220	-0.670	0.855	-0.316	9.248	0.000	0.000	0.000	0.000	0.000	0.000
Gold	0.017	-0.050	3.120	-14.160	25.560	-1.505	1.530	1.087	14.187	3.891	0.245	0.399	0.000	0.000	0.000
Mines	0.059	0.210	3.299	-16.990	19.850	-1.425	1.690	0.090	8.463	63.781	4.439	19.951	0.000	0.000	0.000
Coal	0.014	0.000	4.170	-19.340	21.360	-1.805	1.885	-0.204	7.777	37.757	0.120	0.013	0.000	0.000	0.000
Oil	0.039	0.120	2.417	-15.380	19.270	-1.030	1.230	0.181	14.674	0.004	0.000	0.000	0.000	0.000	0.000
Util	0.016	0.100	1.608	-8.920	14.430	-0.545	0.660	0.934	18.802	0.001	0.000	0.000	0.000	0.000	0.000
Telcm	-0.001	0.070	1.748	-9.670	14.510	-0.645	0.665	0.698	17.768	3.763	0.000	0.000	0.007	0.000	0.000
PerSv	-0.004	0.000	1.749	-8.690	8.950	-0.740	0.795	-0.139	7.585	9.509	0.008	0.003	0.000	0.000	0.000
BusSv	-0.022	0.040	1.616	-7.890	11.940	-0.670	0.660	0.301	11.959	0.398	0.000	0.000	0.000	0.000	0.000
Comps	-0.011	0.040	1.830	-9.890	11.870	-0.830	0.795	0.166	9.166	0.426	0.011	0.110	0.000	0.000	0.000
Chips	-0.051	0.020	1.841	-8.900	10.700	-0.905	0.860	0.002	8.591	0.486	0.001	0.004	0.000	0.000	0.000
LabEq	-0.014	0.070	1.655	-9.000	12.700	-0.700	0.760	-0.023	11.949	60.608	0.006	0.020	0.000	0.000	0.000
Paper	-0.043	0.050	1.535	-9.610	8.630	-0.610	0.630	-0.275	10.620	0.123	0.000	0.000	0.002	0.000	0.000

Boxes	0.028	0.110	1.881	-9.040	10.920	-0.855	0.800	-0.016	7.738	78.496	4.496	8.536	0.000	0.000	0.000
Trans	-0.006	0.020	1.785	-8.760	9.330	-0.870	0.890	-0.229	7.122	5.101	0.082	0.289	0.003	0.000	0.000
Whlsl	-0.028	0.030	1.476	-8.500	9.740	-0.585	0.650	-0.210	11.386	6.327	0.135	0.228	0.000	0.000	0.000
Rtail	-0.016	-0.040	1.612	-8.310	11.750	-0.745	0.710	0.375	9.746	7.334	0.006	0.011	0.000	0.000	0.000
Meals	0.015	0.060	1.580	-8.290	8.910	-0.805	0.800	0.048	7.214	92.449	0.753	0.790	0.000	0.000	0.000
Banks	-0.060	-0.040	2.735	-16.280	16.750	-0.860	0.665	0.314	12.215	1.838	0.093	0.102	0.000	0.000	0.000
Insur	-0.062	0.000	2.124	-11.530	17.840	-0.650	0.600	0.362	16.953	18.539	0.000	0.001	0.000	0.000	0.000
RlEst	-0.119	-0.080	2.740	-15.500	18.650	-1.180	1.065	0.125	10.400	45.850	0.253	1.581	0.000	0.000	0.000
Fin	-0.046	0.030	2.824	-16.270	17.940	-1.035	1.045	0.232	11.047	33.562	0.005	0.046	0.000	0.000	0.000
Other	-0.033	0.020	1.570	-9.880	9.990	-0.540	0.550	-0.270	12.982	44.794	1.394	0.069	0.000	0.000	0.000

Table C.13: Descriptive statistics of the daily returns generated by the 48 industry portfolios in the period between January 3, 2006 and December 31, 2008. From left to right the table reports, for each stock, the mean (%), the median (%), the standard deviation (%), the minimum (%), the maximum (%), the first and the third quartiles (%), the skewness, the kurtosis, and the p-value of the Ljung-Box test (%), at the significance level of the 5%, applied for both the returns (LB) and the squared returns (SLB) with lags equal to 1, 5 and 10, respectively.

POR	MEA	MED	STD	MIN	MAX	$1^{st} Q$	3^{rd} Q	SKE	KUR	LB_1	LB_5	LB_{10}	SLB_1	SLB_5	SLB_{10}
Agric	0.045	0.050	1.399	-6.940	7.650	-0.700	0.840	0.175	6.106	74.914	11.260	12.269	0.000	0.000	0.000
Food	0.063	0.100	0.855	-4.620	3.670	-0.420	0.550	-0.274	5.070	8.731	0.316	0.581	0.000	0.000	0.000
Soda	0.062	0.060	1.066	-6.570	5.150	-0.490	0.650	-0.202	6.850	0.007	0.000	0.000	0.000	0.000	0.000
Beer	0.064	0.075	0.833	-4.280	4.020	-0.430	0.520	-0.139	4.934	14.505	0.010	0.023	0.000	0.000	0.000
Smoke	0.075	0.090	0.966	-4.780	4.390	-0.488	0.640	-0.165	4.698	35.988	17.203	21.629	0.000	0.000	0.000
Toys	0.043	0.070	1.300	-8.290	6.680	-0.600	0.790	-0.270	6.759	34.771	50.545	67.397	0.000	0.000	0.000
Fun	0.060	0.120	1.624	-8.960	6.410	-0.840	0.968	-0.127	5.209	26.336	1.394	3.720	0.000	0.000	0.000
Books	0.055	0.090	1.332	-8.720	6.910	-0.650	0.820	-0.567	7.973	94.914	0.490	1.377	0.000	0.000	0.000
Hshld	0.039	0.050	0.851	-4.090	4.010	-0.430	0.540	-0.300	5.228	39.275	1.251	6.061	0.000	0.000	0.000
Clths	0.066	0.070	1.332	-8.330	6.790	-0.610	0.818	-0.286	6.471	81.731	0.000	0.000	0.000	0.000	0.000
$_{ m Hlth}$	0.058	0.130	1.264	-10.100	5.040	-0.580	0.770	-0.971	9.741	91.847	9.331	0.267	0.001	0.000	0.000
MedEq	0.062	0.100	1.075	-6.910	4.780	-0.480	0.708	-0.528	6.520	17.852	1.430	0.098	0.000	0.000	0.000
Drugs	0.083	0.110	1.034	-4.720	4.550	-0.430	0.670	-0.312	5.204	68.920	7.524	1.500	0.000	0.000	0.000
Chems	0.044	0.080	1.299	-8.210	6.620	-0.600	0.748	-0.383	7.219	40.159	0.001	0.001	0.000	0.000	0.000
Rubbr	0.069	0.100	1.204	-5.960	6.710	-0.620	0.730	-0.177	5.742	34.775	0.009	0.137	0.000	0.000	0.000
Txtls	0.107	0.130	1.534	-8.130	6.830	-0.700	0.988	-0.262	5.268	94.660	0.008	0.048	0.000	0.000	0.000
BldMt	0.055	0.100	1.434	-8.410	6.330	-0.720	0.850	-0.204	5.708	49.169	0.003	0.017	0.000	0.000	0.000
Cnstr	0.039	0.080	1.646	-10.030	6.020	-0.860	0.978	-0.398	5.522	98.631	0.011	0.139	0.000	0.000	0.000
Steel	-0.016	0.000	1.730	-10.760	8.600	-0.945	0.960	-0.163	6.304	88.503	0.044	0.281	0.000	0.000	0.000
FabPr	-0.009	0.060	2.007	-15.450	8.170	-1.108	1.070	-0.266	7.224	94.000	0.034	0.211	0.000	0.000	0.000
Mach	0.022	0.060	1.392	-8.790	6.790	-0.690	0.750	-0.169	6.720	42.870	0.001	0.003	0.000	0.000	0.000
ElcEq	0.025	0.040	1.351	-7.740	7.560	-0.718	0.740	-0.069	6.608	57.250	0.049	0.335	0.000	0.000	0.000
Autos	0.034	0.080	1.456	-8.530	6.380	-0.660	0.850	-0.374	6.056	12.214	0.070	0.295	0.000	0.000	0.000
Aero	0.059	0.095	1.167	-7.030	5.050	-0.568	0.740	-0.457	6.455	85.949	0.328	3.428	0.000	0.000	0.000
Ships	0.091	0.195	1.733	-8.900	7.180	-0.880	1.070	-0.173	5.049	25.610	2.883	11.182	0.000	0.000	0.000
Guns	0.106	0.120	1.069	-5.730	3.740	-0.488	0.770	-0.345	4.931	32.729	6.839	18.811	0.000	0.000	0.000
Gold	-0.079	-0.150	2.375	-11.760	8.760	-1.480	1.298	0.074	4.495	33.125	70.504	68.383	0.586	0.000	0.000
Mines	-0.043	-0.045	1.882	-8.500	10.040	-1.010	0.970	0.060	5.541	6.054	0.108	0.554	0.000	0.000	0.000
Coal	-0.206	-0.135	2.886	-18.440	18.080	-1.795	1.288	-0.012	7.057	20.566	0.015	0.176	0.001	0.000	0.000
Oil	0.009	0.020	1.422	-8.350	5.480	-0.720	0.780	-0.342	6.057	54.006	0.005	0.001	0.003	0.000	0.000
Util	0.039	0.070	0.886	-6.050	4.550	-0.470	0.558	-0.311	6.626	8.517	0.007	0.020	0.000	0.000	0.000
Telcm	0.058	0.110	0.960	-6.600	4.650	-0.440	0.628	-0.521	6.756	52.461	0.002	0.013	0.000	0.000	0.000
PerSv	0.033	0.080	1.345	-8.650	4.810	-0.700	0.888	-0.457	5.261	25.112	29.973	29.929	0.000	0.000	0.000
BusSv	0.060	0.090	1.106	-6.310	4.870	-0.490	0.678	-0.290	5.843	91.176	0.049	0.141	0.000	0.000	0.000
Comps	0.027	0.100	1.251	-6.260	5.500	-0.660	0.770	-0.190	5.481	74.150	10.162	7.555	0.001	0.000	0.000
Chips	0.057	0.090	1.245	-5.760	5.070	-0.600	0.760	-0.117	4.863	99.802	1.614	0.298	0.000	0.000	0.000
LabEq	0.062	0.100	1.318	-7.460	5.740	-0.620	0.780	-0.283	6.327	73.759	0.251	0.216	0.000	0.000	0.000
Paper	0.056	0.100	1.084	-6.330	5.170	-0.460	0.630	-0.356	6.220	5.275	0.090	0.533	0.000	0.000	0.000

Boxes	0.045	0.090	1.218	-8.280	7.420	-0.570	0.720	-0.340	6.851	11.874	0.035	0.526	0.000	0.000	0.000
Trans	0.053	0.130	1.187	-6.010	4.760	-0.588	0.720	-0.321	5.009	90.781	0.800	1.182	0.000	0.000	0.000
Whlsl	0.057	0.090	1.039	-6.710	5.440	-0.470	0.650	-0.239	6.954	42.801	0.029	0.089	0.000	0.000	0.000
Rtail	0.067	0.095	0.936	-5.990	4.790	-0.428	0.620	-0.369	6.227	11.008	0.213	0.118	0.000	0.000	0.000
Meals	0.063	0.090	0.982	-5.440	4.870	-0.468	0.638	-0.313	6.208	4.122	0.084	0.000	0.000	0.000	0.000
Banks	0.055	0.070	1.384	-10.670	7.950	-0.638	0.770	-0.316	9.177	0.001	0.000	0.000	0.000	0.000	0.000
Insur	0.066	0.115	1.149	-8.520	6.940	-0.510	0.690	-0.342	9.017	0.021	0.000	0.000	0.000	0.000	0.000
RlEst	0.048	0.070	1.427	-10.310	6.600	-0.680	0.820	-0.367	7.586	54.196	0.036	0.024	0.000	0.000	0.000
Fin	0.044	0.100	1.421	-8.930	8.900	-0.668	0.790	-0.207	7.838	0.456	0.000	0.000	0.000	0.000	0.000
Other	0.049	0.050	1.070	-6.750	5.890	-0.510	0.618	-0.078	7.505	0.098	0.000	0.000	0.000	0.000	0.000

Table C.14: Descriptive statistics of the daily returns generated by the 48 industry portfolios in the period between January 3, 2011 and December 31, 2015. From left to right the table reports, for each stock, the mean (%), the median (%), the standard deviation (%), the minimum (%), the maximum (%), the first and the third quartiles (%), the skewness, the kurtosis, and the p-value of the Ljung-Box test (%), at the significance level of the 5%, applied for both the returns (LB) and the squared returns (SLB) with lags equal to 1, 5 and 10, respectively.

D Figures of primitive causality networks




Figure D.9: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 banks ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 10% quantile q_{10} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 10% quantile q_{10} . Panel d) reports the network extracted by a not parametric methodologie on the 10% quantile q_{10} . All the causality regression are computed without the market factor.




Figure D.10: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 banks ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 50% quantile q_{50} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 50% quantile q_{50} . Panel d) reports the network extracted by a not parametric methodology on the 50% quantile q_{50} . All the causality regression are computed without the market factor.




Figure D.11: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 banks ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 90% quantile q_{90} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 90% quantile q_{90} . Panel d) reports the network extracted by a not parametric methodology on the 90% quantile q_{90} . All the causality regression are computed without the market factor.




Figure D.12: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 Insurers ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 10% quantile q_{10} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 10% quantile q_{10} . Panel d) reports the network extracted by a not parametric methodologie on the 10% quantile q_{10} . All the causality regression are computed without the market factor.




Figure D.13: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 Insurers ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 50% quantile q_{50} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 50% quantile q_{50} . Panel d) reports the network extracted by a not parametric methodology on the 50% quantile q_{50} . All the causality regression are computed without the market factor.




Figure D.14: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 Insurers ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 90% quantile q_{90} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 90% quantile q_{90} . Panel d) reports the network extracted by a not parametric methodology on the 90% quantile q_{90} . All the causality regression are computed without the market factor.




Figure D.15: This figure visualizes 4 different networks for the period 2006-2008 relative to the 48 Fama and French industry portofolios. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 10% quantile q_{10} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 10% quantile q_{10} . Panel d) reports the network extracted by a not parametric methodologie on the 10% quantile q_{10} . All the causality regression are computed without the market factor.




Figure D.16: This figure visualizes 4 different networks for the period 2006-2008 relative to the 48 Fama and French industry portofolios. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 50% quantile q_{50} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 50% quantile q_{50} . Panel d) reports the network extracted by a not parametric methodology on the 50% quantile q_{50} . All the causality regression are computed without the market factor.




Figure D.17: This figure visualizes 4 different networks for the period 2006-2008 relative to the 48 Fama and French industry portofolios. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by the standard granger causality. Panel b) indicates the network extracted by a quantile baseline regression methodology on the 90% quantile q_{90} . Panel c) indicates the network extracted by a quantile on quantile methodology on the 90% quantile q_{90} . Panel d) reports the network extracted by a not parametric methodology on the 90% quantile q_{90} . All the causality regression are computed without the market factor.




Figure D.18: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 banks ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by a Not parametric methodologie on the 10% quantile q_{10} . Panel b) indicates the network extracted by a quantile on quantile methodology on the median q_{50} . Panel c) displays the network extracted by a Not parametric methodologie on the 10% quantile q_{10} . Panel b) indicates the network extracted by a quantile on quantile methodology on the 90% quantile q_{90} . Panel d) reports the network extracted by the standard granger causality. All the causality regression are computed without the market factor.




Figure D.19: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 25 Insurers ordered for market capitalization. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by a Not parametric methodologie on the 10% quantile q_{10} . Panel b) indicates the network extracted by a quantile on quantile methodology on the median q_{50} . Panel c) displays the network extracted by a Not parametric methodologie on the 10% quantile q_{10} . Panel b) indicates the network extracted by a quantile on quantile methodology on the 90% quantile q_{90} . Panel d) reports the network extracted by the standard granger causality. All the causality regression are computed without the market factor.




Figure D.20: This figure visualizes 4 different networks for the period 2006-2008 relative to the first 48 Fama and French industry portfolios. In this case the networks visualized are some of the primitive networks used for the resulting network computations. Panel a) reports the network extracted by a Not parametric methodology on the 10% quantile q_{10} . Panel b) indicates the network extracted by a quantile on quantile methodology on the median q_{50} . Panel c) displays the network extracted by a Not parametric methodology on the 10% quantile q_{10} . Panel b) indicates the network extracted by a quantile on quantile methodology on the 90% quantile q_{90} . Panel d) reports the network extracted by the standard granger causality. All the causality regression are computed without the market factor.

E Networks summary measures

The network *Density* monitors the number of edges of the network relative to the maximum number of edges that the network might present. The *Density* equals $D = \frac{E}{V(V-1)}$ where E is the total number of edges observed in a given network and V is the number of nodes of the network. For further details we refer to Wasserman and Faust (1994).

The second measure we consider is Assortativity, denote by r, also called homophily, which captures the nodes tendency to connect with nodes having similar properties. In this work, we use a special type of assortativity, called assortativity by degree, see Newman (2010) and Newman (2002). This measure monitors the node willingness to create links with nodes having similar degree. In this case, the degree k_i , i = 1, 2, ... V, has a double role, the first (the standard one) represents the number of links ending at node i, the second one is the value assigned to that node i for computing the assortativity. In the latter case, the degree might be a measure of the number of edges connecting the node to other nodes of the network, but could also be any continuous variable, for instance associated with the relevance of the edges. In order to distinguish between these two roles, we denote the degree by k_i in the first case, and x_i in the second case. Note that, the two values might be identical. We define the mean value μ of x_i as:

$$\mu = \frac{\sum_{i}^{K} \sum_{j}^{K} W_{ij} x_{i}}{2E} = \frac{\sum_{i} k_{i} x_{i}}{2E}$$
 (E.19)

where 2E are the ends for all edges across the network, and W_{ij} is the element of the unnormalized and un-weighted adjacency matrix W. The covariance between x_i and x_j is a way to measure the co-variation between node i and node j with respect to the variable x, which is, in the simplest case, the degree. Across the edges, this covariance can be formally defined as:

$$cov(x_{i}, x_{j}) = \frac{\sum_{ij} W_{ij}(x_{i} - \mu)(x_{j} - \mu)}{\sum_{ij} W_{ij}} = \frac{1}{2E} \sum_{ij} W_{ij}(x_{i} - \mu)(x_{j} - \mu)$$

$$= \frac{1}{2E} \sum_{ij} W_{ij}(x_{i}x_{j} - \mu x_{i} - \mu x_{j} + \mu^{2})$$

$$= \frac{1}{2E} \sum_{ij} W_{ij}x_{i}x_{j} - \frac{\sum_{ij} W_{ij}x_{i}\mu}{2E} - \frac{\sum_{ij} W_{ij}x_{j}\mu}{2E} - \frac{\sum_{ij} W_{ij}\mu^{2}}{2E}$$

$$= \frac{1}{2E} \sum_{ij} W_{ij}x_{i}x_{j} - \mu^{2}$$

$$= \frac{1}{2E} \sum_{ij} W_{ij}x_{i}x_{j} - \frac{\sum_{ij} k_{i}k_{j}x_{i}x_{j}}{(2E)^{2}} = \frac{1}{2E} \sum_{ij} \left(W_{ij} - \frac{k_{i}k_{j}}{2E}\right) x_{i}x_{j}$$
(E.20)

In order to bound the assortativity coefficient in $-1 \le r \le 1$, the covariance is normalized by the maximum covariance value that can be reached by the network. The latter corresponds to the case where all the nodes share edges with nodes having the same degree (or $x_i = x_j$).

$$cov(x_i, x_j)_{max} = cov(x_i, x_i) = \frac{1}{2E} \sum_{ij} \left(W_{ij} - \frac{k_i k_j}{2E} \right) x_i^2$$
 (E.21)

Finally, as out interest lies in the evaluation of the assortativity by degree, we substitute x_i and x_j with k_i and k_j , and obtain:

$$r = \frac{\sum_{ij} (W_{ij} - k_i k_j / 2E) k_i k_j}{\sum_{ij} (W_{ij} - k_i k_j / 2E) k_i^2}$$
 (E.22)

If the W matrix is not symmetric, then the network associated with this matrix is directed, and equation (E.22) must be modified.⁷ In that case, 2E becomes $\sum_{ij} W_{ij} = M$ and we distinguish the degree in indegree k_i^{in} and outdegree k_i^{out} , respectively, for the i-th node.

Formula E.23 must be slightly modified. We are not using directly k^{in} and k^{out} but the excess degree, labeled with the symbol e. For an un-directed network, the excess degree or remaining degree is the number of edges leaving a given vertex minus one. As an example, if the l-th edge links the vertex i with the vertex j, and the two vertexes have degree k_i and k_j , respectively, then the excess degree for the i_{th} and j_{th} node is $e_i = k_i - 1$ and $e_j = k_j - 1$, respectively. If we have a directed network, each node has an excess outdegree e^{out} and an excess indegree e^{in} . Thus, if the l-th edge starts from vertex i and goes to vertex j, then the excess outdegree for the i-th node is $e_i^{out} = k_i^{out} - 1$ and excess indegree for the j-th node is $e_j^{in} = k_j^{in} - 1$.

In the case of a directed network, to compute the assortativity by degree we use:

$$r = \frac{\sum_{ij} e_i^{in} e_j^{out} - \frac{1}{M} \sum_i e_i^{in} \sum_j e_j^{out}}{\sqrt{\left(\sum_i (e_i^{in})^2 - \frac{1}{M} (\sum_i e_i^{in})^2\right) \left(\sum_j (e_j^{out})^2 - \frac{1}{M} (\sum_j e_j^{out})^2\right)}}.$$
 (E.23)

Since the assortativity measure is a ratio, indeterminate forms are also possible. The probability having indeterminate forms $\frac{0}{0}$ increases with a limited number of nodes and high number of edges. In this circumstances the actual number of links among the nodes is almost equal to that we would expect if the links were random, the effect is therefore the reduction of the numerator to zero. In addition, the almost completeness of the network makes null denominator because all the nodes have the same degree and consequently they behave as in a perfect assortativity mixing pattern (nodes with same degree are connected with node having exactly the same degree) where there is only one category given by the degree. Therefore, the indeterminateness arises for the impossibility to capture the network homophily tendency, because of the coexistence of two scenarios: a fully assortative pattern from one side and completely random from the other.

The last two measures we consider are closely related. The first, is the Eigenvector Centrality introduced by Bonacich (1987) This measure captures the nodes relevance (or centrality) as a function of the relevance of neighbor nodes. In other words, we make the prestige of the node proportional to the prestige of its neighbors. Formally, if we define with x_i the prestige of the i_{th} node and W the adjacency matrix, we assume that

$$x_i = \sum_j W_{ij} x_j. \tag{E.24}$$

We can rewrite the previous expression E.24 by using a matrix notation. We denote by \mathbf{x}

⁷For details see Newman (2002) and Newman (2003).

the V-dimensional vector collecting the centrality score. The vector must satisfy the following equality:

$$\mathbf{x} = W\mathbf{x} \tag{E.25}$$

The estimation of the eigenvector centrality requires the use of an iterative procedure. In particular, starting from an initial guess for the centrality scores, \mathbf{x}_0 , which we set equal to a vector of ones, the centrality scores are updated following

$$\mathbf{x}_t = W\mathbf{x}_{t-1} = W^t\mathbf{x}_0 \tag{E.26}$$

where t denotes th t - th iteration. The vector \mathbf{x}_0 can also be seen as a decomposition of linear independent vectors, where c_i is the scalar associated with the i - th component.

$$\mathbf{x}_0 = \sum_{i}^{N} c_i \mathbf{v}_i \tag{E.27}$$

If we plug equation E.27 into equation E.26, by using the equivalence that $W\mathbf{x} = \lambda \mathbf{x}$ we obtain:

$$\mathbf{x}_{t} = W^{t} \mathbf{x}_{0} = W^{t} \sum_{i}^{N} c_{i} \mathbf{v}_{i} = \sum_{i}^{N} c_{i} W^{t} \mathbf{v}_{i} = \sum_{i}^{N} c_{i} W^{t-1} \lambda_{i} \mathbf{v}_{i} = \sum_{i}^{N} c_{i} \lambda_{i}^{t} \mathbf{v}_{i}$$

$$= \left(\frac{\lambda_{1}}{\lambda_{1}}\right)^{t} \sum_{i}^{N} c_{i} \lambda_{i}^{t} \mathbf{v}_{i} = \lambda_{1}^{t} \sum_{i}^{N} c_{i} \left(\frac{\lambda_{i}}{\lambda_{1}}\right)^{t} \mathbf{v}_{i}$$
(E.28)

where λ_i is the i-th eigenvalue associated to the i-th eigenvector, and λ_1 is the maximum eigenvalue.⁸ What we learn from the expression in E.29 is the following:

since λ_1 is the maximum eigenvalue, as soon $t \to \infty$ the $\sum_{i=1}^{N} c_i \left(\frac{\lambda_i}{\lambda_1}\right)^t \mathbf{v}_i$ tends to $c_1 \mathbf{v}_1$ and thus the eigenvector centrality is proportional to the first eigenvalue as in the equation E.29 we have

$$\mathbf{x}_t = \lambda_1^t c_1 \mathbf{v}_1 \tag{E.29}$$

In other words, combining equation E.29 with E.24, we can observe that the centrality score is a function of the first eigenvalue

$$x_i = \frac{1}{\lambda_1} \sum_j W_{ij} x_j. \tag{E.30}$$

The eigenvector centrality can be computed not only for adjacency matrix W but also for the normalized adjacency matrix.

⁸The existence of the maximum eigenvalue is guaranteed by the Perron- Frobenius theorem.

⁹The eigenvector centrality does not converge when the maximum eigenvalue λ_1 tends to zero. The maximum eigenvalue coefficient decreases with the sparsity of the adjacency matrix, Van Mieghem (2010)

F Figures of Combined Network




Figure F.21: This figure visualizes the 3 different resulting networks for the period 2006-2008 relative to the first 25 banks ordered for market capitalization. Panel a) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method. Panel b) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 50% quantile, and the standard granger causality method. Panel c) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 90% quantile, and the standard granger causality method. All the causality regression are computed without the market factor.




Figure F.22: This figure visualizes the 3 different resulting networks for the period 2011-2015 relative to the first 25 banks ordered for market capitalization. Panel a) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method. Panel b) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 50% quantile, and the standard granger causality method. Panel c) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 90% quantile, and the standard granger causality method. All the causality regression are computed without the market factor.




Figure F.23: This figure visualizes the 3 different resulting networks for the period 2006-2008 relative to the first 25 Insurance companies ordered for market capitalization. Panel a) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method. Panel b) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 50% quantile, and the standard granger causality method. Panel c) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 90% quantile, and the standard granger causality method. All the causality regression are computed without the market factor.




Figure F.24: This figure visualizes the 3 different resulting networks for the period 2011-2015 relative to the first 25 Insurance companies ordered for market capitalization. Panel a) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method. Panel b) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 50% quantile, and the standard granger causality method. Panel c) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 90% quantile, and the standard granger causality method. All the causality regression are computed without the market factor.




Figure F.25: This figure visualizes the 3 different resulting networks for the period 2006-2008 relative to 48 Industry portfolios. Panel a) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method. Panel b) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 50% quantile, and the standard granger causality method. Panel c) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 90% quantile, and the standard granger causality method. All the causality regression are computed without the market factor.




Figure F.26: This figure visualizes the 3 different resulting networks for the period 2011-2015 relative to 48 Industry portfolios. Panel a) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 10% quantile, and the standard granger causality method. Panel b) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 50% quantile, and the standard granger causality method. Panel c) reports the network extracted by combining causality network by using quantile regression (QB, Qo and QN) at the 90% quantile, and the standard granger causality method. All the causality regression are computed without the market factor.

G Sensitivity analysis for weekly Returns

Table G.15: The table reports the δ of model (5) that represent the weights for networks combination. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based network, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks. Columns 3 to 5 refer to the crisis sample while columns 6 to 8 to the most recent sample. The second row identifies the three different networks which are optimally combined: baseline quantile causality - QB; quantile-on-quantile causality Qo; non-parametric quantile causality - QN. Parameters are, by construction, positive and sum up to one (within each row and within each period). A star identifies parameters significant at the 5% confidence level.

		2	006-200	8	2	010-201	5
Quantile	Factor	QB	Qo	QN	QB	Qo	QN
			25 Ba	nks			
10%	N	0.060	0.054*	0.885*	0.001	0.625	0.374
50%	N	0.142	0.108*	0.750*	0.094	0.067	0.838*
90%	N	0.028	0.026	0.946*	0.000	0.725*	0.275*
10%	Y	0.024	0.109	0.868*	0.000*	0.526*	0.474*
50%	Y	0.054	0.062*	0.885*	0.201	0.066*	0.733*
90%	Y	0.123	0.020	0.857*	0.000	0.877*	0.123*
		25 In	surance	Compa	nies		
10%	N	0.000	0.034	0.966*	0.058	0.831*	0.111*
50%	N	0.842*	0.125	0.034	0.059	0.045	0.896*
90%	N	0.000	0.028	0.972*	0.000	0.456*	0.544*
10%	Y	0.001	0.028	0.972*	0.000	0.488*	0.512*
50%	Y	0.678*	0.106	0.216*	0.053	0.041	0.907*
90%	Y	0.000	0.046	0.954	0.000	0.554	0.446
		48 I	Industry	Portfol	lio		
10%	N	0.000	1.000*	0.000	0.032	0.968*	0.000
50%	N	0.202	0.000	0.798	0.214	0.023	0.763
90%	N	0.108	0.892*	0.000	0.016	0.984*	0.000
10%	Y	0.390	0.540	0.071	0.038	0.755*	0.207
50%	Y	0.237	0.613	0.150	0.089	0.024	0.888*
90%	Y	0.000	1.000*	0.000	0.111	0.857*	0.032

Table G.16: The table reports residual correlation descriptive analyses for the Banks dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

2006-2008

Model	N Networks	Factor	Min	Max	q10	q50	q90	% elements ≤ -0.1
${}$ Q(10%)	4	N	-0.279	0.584	-0.108	0.071	0.279	11.3%
Q(50%)	4	N	-0.344	0.597	-0.106	0.060	0.316	11.0%
Q(90%)	4	N	-0.297	0.597	-0.127	0.062	0.299	14.0%
Q(10%)	4	Y	-0.291	0.567	-0.135	0.046	0.253	16.0%
$\mathrm{Q}(50\%)$	4	Y	-0.346	0.479	-0.118	0.055	0.251	11.7%
Q(90%)	4	Y	-0.301	0.533	-0.137	0.038	0.235	16.3%
G	1	N	-0.381	0.670	-0.128	0.111	0.372	13.3%
G	1	Y	-0.394	0.641	-0.128	0.089	0.378	11.7%
4-F-CAPM	0	N.A	-0.358	0.678	-0.108	0.257	0.502	10.7%
			201	1-2015				
Model	${f N}$	Factor	Min	Max	q10	$\mathbf{q50}$	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.265	0.424	-0.121	-0.015	0.137	14.7%
$\mathrm{Q}(50\%)$	4	N	-0.390	0.643	-0.107	0.053	0.282	13.3%
Q(90%)	4	N	-0.248	0.432	-0.125	0.010	0.136	15.3%
Q(10%)	4	Y	-0.281	0.505	-0.131	-0.011	0.145	18.7%
$\mathrm{Q}(50\%)$	4	Y	-0.316	0.609	-0.127	0.054	0.245	15.0%
Q(90%)	4	Y	-0.304	0.508	-0.139	0.001	0.132	17.7%
G	1	N	-0.211	0.598	-0.027	0.151	0.362	3.0%
G	1	Y	-0.211	0.598	-0.027	0.146	0.361	3.3%
4-F-CAPM	0	NA	-0.122	0.669	0.054	0.266	0.487	0.3%

Table G.17: The table reports residual correlation descriptive analyses for the Insurance Companies dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

		·	200	6-2008				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.440	0.682	-0.222	0.002	0.248	27.0%
Q(50%)	4	N	-0.423	0.694	-0.161	0.044	0.260	16.7%
Q(90%)	4	N	-0.448	0.676	-0.207	0.009	0.255	25.7%
Q(10%)	4	Y	-0.455	0.686	-0.212	0.002	0.257	26.7%
Q(50%)	4	Y	-0.369	0.515	-0.181	0.018	0.225	20.7%
Q(90%)	4	Y	-0.464	0.671	-0.214	0.001	0.253	27.0%
G	1	N	-0.466	0.831	-0.166	0.044	0.316	17.7%
G	1	Y	-0.460	0.714	-0.166	0.048	0.323	15.0%
4-F-CAPM	0	N.A	-0.370	0.847	-0.149	0.089	0.422	14.0%
			201	1-2015				
Model	N	Factor	Min	Max	q10	$\mathbf{q50}$	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.290	0.504	-0.121	0.009	0.157	14.7%
Q(50%)	4	N	-0.258	0.547	-0.149	0.019	0.202	16.7%
Q(90%)	4	N	-0.289	0.345	-0.140	-0.004	0.168	18.3%
Q(10%)	4	Y	-0.333	0.503	-0.153	-0.012	0.189	20.7%
Q(50%)	4	Y	-0.268	0.590	-0.157	0.019	0.191	18.3%
Q(90%)	4	Y	-0.299	0.398	-0.152	-0.001	0.199	21.3%
\mathbf{G}	1	N	-0.136	0.658	-0.005	0.121	0.317	1.7%
G	1	Y	-0.136	0.658	-0.014	0.116	0.315	1.7%
4-F-CAPM	0	N.A	-0.103	0.658	0.017	0.138	0.332	0.3%

Table G.18: The table reports residual correlation descriptive analyses for the Industry portfolios dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

		·	200	6-2008				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.465	0.611	-0.191	-0.001	0.201	25.6%
$\mathrm{Q}(50\%)$	4	N	-0.380	0.661	-0.176	0.011	0.247	22.9%
Q(90%)	4	N	-0.443	0.608	-0.195	-0.006	0.206	25.9%
Q(10%)	4	Y	-0.517	0.624	-0.217	-0.011	0.211	27.1%
Q(50%)	4	Y	-0.467	0.690	-0.199	0.006	0.237	25.2%
Q(90%)	4	Y	-0.533	0.599	-0.195	-0.014	0.216	25.2%
G	1	N	-0.384	0.687	-0.146	0.025	0.257	18.6%
G	1	Y	-0.401	0.687	-0.157	0.011	0.239	19.8%
4-F-CAPM	0	N.A	-0.486	0.733	-0.219	0.003	0.267	27.1%
			201	1-2015				
Model	N	Factor	Min	Max	q10	$\mathbf{q50}$	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.463	0.567	-0.140	-0.008	0.135	19.7%
Q(50%)	4	N	-0.447	0.519	-0.138	0.007	0.176	16.6%
Q(90%)	4	N	-0.470	0.572	-0.144	-0.013	0.138	19.6%
Q(10%)	4	Y	-0.442	0.501	-0.148	-0.003	0.145	19.3%
Q(50%)	4	Y	-0.454	0.508	-0.136	0.003	0.160	16.7%
Q(90%)	4	Y	-0.443	0.567	-0.144	-0.005	0.153	18.4%
G	1	N	-0.470	0.577	-0.140	0.007	0.199	18.0%
G	1	Y	-0.503	0.577	-0.139	0.006	0.177	17.7%
4-F-CAPM	0	N.A	-0.456	0.592	-0.147	0.007	0.208	18.4%

Table G.19: The table report the δ of model (5) that represent the weights for networks combination. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based network, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks. Columns 3 to 5 refer to the crisis sample while columns 6 to 8 to the most recent sample. The second row identifies the three different networks which are optimally combined: quantile-on-quantile causality Qo; non-parametric quantile causality - QN and Granger Causality. Parameters are, by construction, positive and sum up to one (within each row and within each period). A star identifies parameters significant at the 5% confidence level.

		20	006-200	8	2	010-201	5
Quantile	Factor	Qo	QN	$\mathbf{G}\mathbf{R}$	Qo	QN	GR
			25 Ba	nks			
10%	N	0.071	0.915*	0.014	0.630	0.354	0.017
50%	N	0.128	0.848*	0.024	0.132	0.714*	0.154
90%	N	0.008	0.884*	0.108	0.706*	0.243*	0.051*
10%	Y	0.082	0.843*	0.076	0.538	0.428*	0.034*
50%	Y	0.094	0.726	0.180	0.092	0.745*	0.163
90%	Y	0.002	0.893*	0.105	0.876*	0.121*	0.003*
		25 Ins	surance	Compa	nies		
10%	N	0.034	0.966*	0.000	0.804	0.141	0.055*
50%	N	0.000	0.942*	0.058	0.054	0.615	0.331
90%	N	0.028	0.972	0.000	0.364	0.436	0.200
10%	Y	0.006	0.949*	0.045	0.489	0.511	0.000
50%	Y	0.516	0.474	0.009	0.051	0.846*	0.102
90%	Y	0.046	0.954*	0.000	0.554	0.446	0.000*
		48 I	ndustry	Portfo	lio		
10%	N	0.173	0.057	0.770	1.000*	0.000	0.000*
50%	N	0.079	0.248	0.673	0.030	0.868*	0.102
90%	N	1.000*	0.000	0.000	1.000*	0.000	0.000*
10%	Y	0.619	0.052	0.329	0.640	0.320	0.039
50%	Y	0.000	0.676	0.324	0.029	0.876*	0.095
90%	Y	0.810	0.000*	0.190	0.918	0.032	0.050

Table G.20: The table reports residual correlation descriptive analyses for the Banks dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (Qo and QN) at the 10% quantile, combined with the Granger Causality Network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

		200	6-2008						
N	Factor	Min	Max	q10	q50	q90	% elements		
Networks							≤ -0.1		
4	N	-0.304	0.584	-0.117	0.067	0.273	12.0%		
4	N	-0.319	0.587	-0.105	0.072	0.342	11.0%		
4	N	-0.269	0.600	-0.147	0.050	0.278	16.0%		
4	Y	-0.295	0.571	-0.136	0.035	0.236	15.3%		
4	Y	-0.346	0.490	-0.119	0.048	0.288	12.3%		
4	Y	-0.313	0.485	-0.142	0.031	0.244	18.0%		
1	N	-0.381	0.670	-0.128	0.111	0.372	13.3%		
1	Y	-0.394	0.641	-0.128	0.089	0.378	11.7%		
0	N.A	-0.358	0.678	-0.108	0.257	0.502	10.7%		
2011-2015									
N	Factor	Min	Max	q10	q50	q90	% elements		
Networks							≤ -0.1		
4	N	-0.263	0.422	-0.122	-0.013	0.135	15.3%		
4	N	-0.332	0.528	-0.116	0.052	0.243	12.7%		
4	N	-0.253	0.431	-0.127	0.010	0.138	15.7%		
4	Y	-0.283	0.503	-0.132	-0.008	0.140	19.0%		
4	Y	-0.311	0.522	-0.122	0.052	0.237	13.0%		
4	Y	-0.304	0.508	-0.140	0.001	0.132	17.7%		
1	N	-0.211	0.598	-0.027	0.151	0.362	3.0%		
1	Y	-0.211	0.598	-0.027	0.146	0.361	3.3%		
0	NA	-0.122	0.669	0.054	0.266	0.487	0.3%		
	## A	Networks N 4 N 4 N 4 Y 4 Y 4 Y 1 N 1 Y 0 N.A Factor Networks A N 4 N 4 N 4 Y 4 Y 4 Y 1 N 1 N 1 N 1 N 1 Y	Networks Factor Min 4 N -0.304 4 N -0.319 4 N -0.269 4 Y -0.295 4 Y -0.346 4 Y -0.346 4 Y -0.381 1 Y -0.394 0 N.A -0.358 Networks Factor Min Networks Min -0.263 4 N -0.263 4 N -0.263 4 N -0.253 4 Y -0.283 4 Y -0.311 4 Y -0.304 1 Y -0.304 1 N -0.211 1 Y -0.211	Networks Image: Content of the con	Networks Factor Min Max q10 4 N -0.304 0.584 -0.117 4 N -0.319 0.587 -0.105 4 N -0.269 0.600 -0.147 4 Y -0.295 0.571 -0.136 4 Y -0.346 0.490 -0.119 4 Y -0.313 0.485 -0.142 1 N -0.381 0.670 -0.128 1 Y -0.394 0.641 -0.128 0 N.A -0.358 0.678 -0.108 Networks Factor Min Max q10 Networks Y -0.263 0.422 -0.122 4 N -0.263 0.422 -0.122 4 N -0.332 0.528 -0.116 4 N -0.253 0.431 -0.127 4 Y -0.283 0.503 -	Networks Factor Min Max q10 q50 1 N -0.304 0.584 -0.117 0.067 4 N -0.319 0.587 -0.105 0.072 4 N -0.269 0.600 -0.147 0.050 4 Y -0.295 0.571 -0.136 0.035 4 Y -0.346 0.490 -0.119 0.048 4 Y -0.346 0.490 -0.119 0.048 4 Y -0.313 0.485 -0.142 0.031 1 N -0.381 0.670 -0.128 0.111 1 Y -0.394 0.641 -0.128 0.089 0 N.A -0.358 0.678 -0.108 0.257 Protor Min Max q10 q50 Networks T T -0.122 -0.013 4 N -0.263 0.422 -0.	Networks Factor Min Max q10 q50 q90 Networks -0.304 0.584 -0.117 0.067 0.273 4 N -0.319 0.587 -0.105 0.072 0.342 4 N -0.269 0.600 -0.147 0.050 0.278 4 Y -0.295 0.571 -0.136 0.035 0.236 4 Y -0.346 0.490 -0.119 0.048 0.288 4 Y -0.313 0.485 -0.142 0.031 0.244 1 N -0.381 0.670 -0.128 0.111 0.372 1 Y -0.394 0.641 -0.128 0.011 0.372 1 Y -0.358 0.678 -0.108 0.257 0.502 Networks Factor Min Max q10 q50 q90 Networks - - - - -		

Table G.21: The table reports residual correlation descriptive analyses for the Insurance Companies dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (Qo and QN) at the 10% quantile, combined with the Granger Causality Network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

		·	200	6-2008				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.441	0.682	-0.222	0.002	0.248	27.0%
$\mathrm{Q}(50\%)$	4	N	-0.435	0.756	-0.169	0.027	0.335	20.7%
Q(90%)	4	N	-0.448	0.676	-0.207	0.009	0.255	25.7%
Q(10%)	4	Y	-0.436	0.684	-0.203	0.005	0.246	28.0%
Q(50%)	4	Y	-0.468	0.574	-0.185	0.016	0.238	22.7%
Q(90%)	4	Y	-0.464	0.671	-0.214	0.001	0.253	27.0%
G	1	N	-0.466	0.831	-0.166	0.044	0.316	17.7%
G	1	Y	-0.460	0.714	-0.166	0.048	0.323	15.0%
4-F-CAPM	0	N.A	-0.370	0.847	-0.149	0.089	0.422	14.0%
			201	1-2015				
Model	${f N}$	Factor	Min	Max	q10	$\mathbf{q50}$	$\mathbf{q90}$	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.293	0.482	-0.123	0.009	0.158	14.7%
Q(50%)	4	N	-0.238	0.593	-0.132	0.025	0.217	15.0%
Q(90%)	4	N	-0.289	0.348	-0.137	-0.006	0.167	19.3%
Q(10%)	4	Y	-0.333	0.503	-0.153	-0.012	0.189	20.7%
Q(50%)	4	Y	-0.276	0.590	-0.140	0.020	0.210	17.0%
Q(90%)	4	Y	-0.299	0.398	-0.152	-0.001	0.199	21.3%
G	1	N	-0.136	0.658	-0.005	0.121	0.317	1.7%
G	1	Y	-0.136	0.658	-0.014	0.116	0.315	1.7%
4-F-CAPM	0	N.A	-0.103	0.658	0.017	0.138	0.332	0.3%

Table G.22: The table reports residual correlation descriptive analyses for the Industry portfolios dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (Qo and QN) at the 10% quantile, combined with the Granger Causality Network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

20	ገበ	6.	-2	N	08

\mathbf{Model}	${f N}$	Factor	Min	Max	q10	$\mathbf{q50}$	$\mathbf{q90}$	% elements
	Networks							≤ -0.1
${Q(10\%)}$	4	N	-0.370	0.592	-0.153	0.013	0.237	20.0%
Q(50%)	4	N	-0.357	0.604	-0.151	0.020	0.251	19.3%
Q(90%)	4	N	-0.465	0.660	-0.196	-0.002	0.212	25.8%
Q(10%)	4	Y	-0.431	0.595	-0.182	0.007	0.221	22.5%
Q(50%)	4	Y	-0.415	0.687	-0.169	0.006	0.231	21.5%
Q(90%)	4	Y	-0.508	0.592	-0.187	-0.007	0.215	23.8%
G	1	N	-0.384	0.687	-0.146	0.025	0.257	18.6%
G	1	Y	-0.401	0.687	-0.157	0.011	0.239	19.8%
4-F-CAPM	0	N.A	-0.486	0.733	-0.219	0.003	0.267	27.1%
			201	1-2015				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
-Q(10%)	4	N	-0.463	0.567	-0.138	-0.008	0.133	19.3%
Q(50%)	4	N	-0.464	0.519	-0.137	0.003	0.172	16.8%
Q(90%)	4	N	-0.471	0.572	-0.144	-0.013	0.139	20.0%
Q(10%)	4	Y	-0.434	0.495	-0.148	-0.005	0.145	19.1%
$\mathrm{Q}(50\%)$	4	Y	-0.509	0.513	-0.141	0.004	0.163	16.5%
Q(90%)	4	Y	-0.424	0.563	-0.144	-0.004	0.150	19.3%
G	1	N	-0.470	0.577	-0.140	0.007	0.199	18.0%
G	1	Y	-0.503	0.577	-0.139	0.006	0.177	17.7%

H Tables for Monthly returns

Table H.23: The table reports the δ of model (5) that represent the weights for networks combination. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based network, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks. Columns 3 to 6 refer to the crisis sample while columns 7 to 10 to the most recent sample. The second row identifies the four different networks which are optimally combined: baseline quantile causality - QB; quantile-on-quantile causality Qo; non-parametric quantile causality - QN; Granger causality. Parameters are, by construction, positive and sum up to one (within each row and within each period). A star identifies parameters significant at the 5% confidence level.

			2006	S-2008			2010	-2015	
Quantile	Factor	QB	Qo	QN	QR	QB	Qo	QN	QR
				25 Ba	nks				
10%	N	0.031	0.091	0.878	0.000	0.000	0.692*	0.271	0.037
50%	N	0.495	0.000*	0.345	0.159	0.083	0.123*	0.689*	0.105
90%	N	0.002	0.007	0.872*	0.119	0.000	0.739*	0.142	0.119
10%	Y	0.061	0.182	0.681*	0.077	0.000	0.604	0.375	0.022
50%	Y	0.131	0.109*	0.451*	0.310*	0.075	0.046*	0.746*	0.132
90%	Y	0.131	0.022	0.639*	0.207	0.000	0.954*	0.000	0.046
			25 Ins	surance	Compar	nies			
10%	N	0.248	0.000*	0.638*	0.115	0.000	0.651	0.137	0.213
50%	N	0.689	0.000	0.175*	0.136	0.052	0.063	0.637	0.248
90%	N	0.000	0.042*	0.829*	0.129	0.000	0.449	0.453	0.098
10%	Y	0.000	0.005*	0.916*	0.079	0.000	1.000*	0.000	0.000
50%	Y	0.000	0.071*	0.831*	0.098	0.082	0.054	0.862	0.003
90%	Y	0.000	0.000	0.907	0.093	0.000	0.186	0.814*	0.000
			48 I	ndustry	Portfoli	io			
10%	N	0.149	0.253	0.599	0.000	0.020	0.980	0.000	0.000
50%	N	0.085	0.119	0.000	0.797	0.309	0.031	0.520	0.141
90%	N	0.140	0.000	0.860	0.000	0.014	0.203	0.633*	0.150
10%	Y	0.304	0.226	0.356*	0.114	0.012	0.576	0.294	0.118
50%	Y	0.426	0.137	0.000	0.437	0.093	0.027	0.767*	0.113
90%	Y	0.166	0.030	0.749	0.055	0.103	0.668	0.106	0.123

Table H.24: The table reports residual correlation descriptive analyses for the Banks dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

Table H.25: Add caption

			200	6-2008				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.485	0.748	-0.229	0.062	0.358	23.7%
Q(50%)	4	N	-0.401	0.736	-0.193	0.095	0.412	20.7%
Q(90%)	4	N	-0.496	0.696	-0.236	0.071	0.367	26.3%
Q(10%)	4	Y	-0.499	0.797	-0.248	0.034	0.325	30.3%
Q(50%)	4	Y	-0.495	0.755	-0.226	0.055	0.351	26.3%
Q(90%)	4	Y	-0.504	0.737	-0.257	0.050	0.323	28.3%
\mathbf{G}	1	N	-0.605	0.756	-0.172	0.190	0.528	17.0%
${ m G}$	1	Y	-0.599	0.826	-0.190	0.141	0.505	18.0%
4-F-CAPM	0	N.A	-0.404	0.826	0.040	0.417	0.699	7.0%
			201	1-2015				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.415	0.527	-0.209	-0.007	0.209	28.0%
Q(50%)	4	N	-0.536	0.602	-0.189	0.033	0.288	21.3%
Q(90%)	4	N	-0.455	0.512	-0.200	0.007	0.227	27.7%
Q(10%)	4	Y	-0.382	0.563	-0.226	-0.011	0.228	28.0%
Q(50%)	4	Y	-0.449	0.582	-0.201	0.038	0.280	22.7%

27.7%

8.3%

8.3%

0.7%

Y

Ν

Y

NA

-0.423

-0.314

-0.388

-0.155

0.564

0.677

0.677

0.793

-0.223

-0.083

-0.083

0.098

-0.003

0.165

0.162

0.385

0.224

0.460

0.455

0.604

4

1

1

0

 $\hat{Q(90\%)}$

 \mathbf{G}

G

4-F-CAPM

Table H.26: The table reports residual correlation descriptive analyses for the Insurance Companies dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

			2006	6-2008				
Model	N Networks	Factor	Min	Max	q10	q50	q90	% elements ≤ -0.1
Q(10%)	4	N	-0.684	0.693	-0.264	0.040	0.372	26.3%
Q(50%)	4	N	-0.674	0.760	-0.261	0.039	0.366	26.0%
Q(90%)	4	N	-0.694	0.744	-0.276	0.044	0.367	27.3%
Q(10%)	4	Y	-0.674	0.734	-0.277	0.054	0.383	29.0%
Q(50%)	4	Y	-0.694	0.755	-0.271	0.041	0.370	28.7%
Q(90%)	4	Y	-0.679	0.734	-0.270	0.055	0.383	28.3%
G	1	N	-0.665	0.771	-0.267	0.055	0.399	27.7%
G	1	Y	-0.629	0.771	-0.240	0.065	0.398	24.7%
4-F-CAPM	0	N.A	-0.639	0.886	-0.190	0.157	0.519	16.7%
			201	L-2015				
Model	N	Factor	Min	Max	q10	$\mathbf{q50}$	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.543	0.610	-0.206	0.010	0.236	26.0%
Q(50%)	4	N	-0.428	0.648	-0.224	0.037	0.283	24.3%
Q(90%)	4	N	-0.450	0.577	-0.229	0.001	0.247	27.7%
Q(10%)	4	Y	-0.533	0.629	-0.239	0.008	0.264	31.7%
Q(50%)	4	Y	-0.442	0.697	-0.236	0.036	0.287	26.7%
Q(90%)	4	Y	-0.446	0.626	-0.242	0.002	0.280	31.0%
G	1	N	-0.230	0.826	-0.015	0.221	0.473	2.3%
\mathbf{G}	1	Y	-0.230	0.826	-0.023	0.218	0.473	2.7%
4-F-CAPM	0	N.A	-0.204	0.826	0.045	0.246	0.475	0.7%

Table H.27: The table reports residual correlation descriptive analyses for the Industry portfolios dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger causality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

	2006-2008										
Model	N	Factor	Min	Max	q10	q50	q90	% elements			
	Networks							$\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ $			
${}$ Q(10%)	4	N	-0.702	0.653	-0.338	-0.029	0.267	39.0%			
Q(50%)	4	N	-0.654	0.680	-0.353	-0.032	0.296	39.7%			
Q(90%)	4	N	-0.619	0.658	-0.347	-0.027	0.289	38.7%			
Q(10%)	4	Y	-0.749	0.638	-0.319	-0.022	0.265	37.3%			
Q(50%)	4	Y	-0.675	0.675	-0.371	-0.031	0.318	40.8%			
Q(90%)	4	Y	-0.690	0.689	-0.348	-0.030	0.277	39.5%			
G	1	N	-0.654	0.685	-0.368	-0.023	0.293	39.2%			
G	1	Y	-0.695	0.705	-0.373	-0.020	0.301	40.3%			
4-F-CAPM	0	N.A	-0.667	0.713	-0.378	-0.025	0.309	39.5%			
			201	1-2015							
Model	N	Factor	Min	Max	q10	q50	q90	% elements			
	Networks							≤ -0.1			
${}$ Q(10%)	4	N	-0.583	0.623	-0.234	-0.008	0.210	30.4%			
Q(50%)	4	N	-0.499	0.617	-0.206	0.014	0.248	26.1%			
Q(90%)	4	N	-0.551	0.589	-0.224	-0.006	0.244	29.1%			
Q(10%)	4	Y	-0.534	0.543	-0.228	-0.002	0.244	29.5%			
Q(50%)	4	Y	-0.443	0.529	-0.216	0.008	0.241	25.3%			
Q(90%)	4	Y	-0.446	0.585	-0.230	-0.002	0.242	29.5%			
\mathbf{G}	1	N	-0.502	0.579	-0.228	0.009	0.259	28.0%			
G	1	Y	-0.478	0.594	-0.220	0.010	0.251	27.7%			
4-F-CAPM	0	N.A	-0.502	0.626	-0.231	0.021	0.283	28.4%			

I Homogeneity

Table I.28: The table reports the δ of model (5), that represent the weights for networks combination. The model replaces the matrix \mathcal{R} with a scalar coefficient, and thus we have, across assets, a homogeneous impact of the network. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based network, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks. Columns 3 to 6 refer to the crisis sample while columns 7 to 10 to the most recent sample. The second row identifies the four different networks which are optimally combined: baseline quantile causality - QB; quantile-on-quantile causality Qo; non-parametric quantile causality - QN; Granger causality. Parameters are, by construction, positive and sum up to one (within each row and within each period). A star identifies parameters significant at the 5% confidence level.

			2010-2015								
Quantile	Factor	QB	Qo	QN	QG	QB	Qo	QN	QG		
25 Banks											
10%	N	0.000	0.159	0.841*	0.000	0.028	0.656*	0.264*	0.052		
50%	N	0.199	0.146*	0.589*	0.066*	0.140*	0.273*	0.349*	0.238*		
90%	N	0.001	0.000	0.984*	0.000	0.048	0.565*	0.281*	0.106		
10%	Y	0.000*	0.398*	0.403*	0.199*	0.036	0.771*	0.160	0.033*		
50%	Y	0.140*	0.204*	0.444*	0.212*	0.198*	0.187*	0.409*	0.206*		
90%	Y	0.247*	0.000*	0.471*	0.282*	0.000	0.939*	0.061*	0.000*		
	25 Insurance Companies										
10%	N	0.061	0.151*	0.789*	0.000*	0.078	0.922*	0.000	0.000		
50%	N	0.193	0.099*	0.495*	0.212*	0.158	0.111*	0.659*	0.072*		
90%	N	0.000*	0.000*	1.000*	0.000*	0.000*	0.492*	0.508*	0.000*		
10%	Y	0.040	0.082	0.856*	0.022	0.007	0.645*	0.347*	0.000		
50%	Y	0.090	0.218*	0.631*	0.060	0.038	0.072	0.831*	0.058		
90%	Y	0.000	0.000	0.973*	0.027	0.000*	0.566*	0.434*	0.000*		
	48 Industry Portfolio										
10%	N	0.000*	0.000*	0.000*	1.000*	0.074	0.926*	0.000*	0.000*		
50%	N	0.000	0.213	0.177	0.610	0.000*	1.000*	0.000*	0.000*		
90%	N	0.000*	0.000*	0.000*	1.000*	0.000*	1.000*	0.000*	0.000*		
10%	Y	1.000*	0.000*	0.000*	0.000*	0.000*	1.000*	0.000*	0.000*		
50%	Y	1.000*	0.000*	0.000*	0.000*	0.705	0.001*	0.294*	0.000*		
90%	Y	0.782*	0.218*	0.000*	0.000*	0.000*	1.000*	0.000*	0.000*		

Table I.29: The table reports scalar coefficient ρ replacing matrix \mathcal{R} in model (5), to impose homogeneous reaction of the assets to network exposure. The top panel focused on the banks dataset, the middle panel on the insurance companies dataset and the bottom panel on the industry portfolios dataset. The first column identifies the quantiles used to estimate the quantile-based network, and the second column indicates if a common factor was used (Y) or not used (N) in the estimation of the causality networks. The results identify the four different networks which are optimally combined: baseline quantile causality; quantile-on-quantile causality; non-parametric quantile causality; Granger causality. A star identifies parameters significant at the 5% confidence level.

Quantile	Factor	2006-2008	2011-2015							
25 Banks										
10%	N	0.687*	0.727*							
50%	N	0.756*	0.851*							
90%	N	0.604*	0.774*							
10%	Y	0.656*	0.713*							
50%	Y	0.894*	0.886*							
90%	Y	0.639*	0.673*							
25 Insurance Companies										
10%	N	0.497*	0.554*							
50%	N	0.301*	0.613*							
90%	N	0.469*	0.618*							
10%	Y	0.491*	0.581*							
50%	Y	0.472*	0.638*							
90%	Y	0.477*	0.564*							
	48 Industry companies									
10%	N	-0.077*	0.460*							
50%	N	-0.117*	0.025*							
90%	N	-0.077*	0.324*							
10%	Y	0.175*	0.282*							
50%	Y	0.129*	0.123*							
90%	Y	0.100*	0.252*							

Table I.30: The table reports residual correlation descriptive analyses for the Banks dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

2006-2008										
Model	N Networks	Factor	Min	Max	q10	q50	q90	% elements ≤ -0.1		
10%	4	N	-0.443	0.578	-0.207	0.082	0.364	23.0%		
50%	4	N	-0.469	0.640	-0.205	0.114	0.425	19.3%		
90%	4	N	-0.461	0.602	-0.193	0.031	0.371	23.3%		
10%	4	Y	-0.377	0.573	-0.198	0.061	0.322	21.3%		
50%	4	Y	-0.526	0.558	-0.184	0.057	0.329	18.0%		
90%	4	Y	-0.588	0.546	-0.206	0.061	0.347	19.3%		
Granger	1	N	-0.366	0.662	-0.120	0.221	0.475	12.3%		
Granger	1	Y	-0.591	0.626	-0.143	0.155	0.428	12.3%		
Multifactor	-	-	-0.358	0.678	-0.108	0.257	0.502	10.7%		
			201	1-2015						
Model	N	Factor	Min	Max	q10	q50	q90	% elements		
	Networks				_	_	_	≤ -0.1		
10%	4	N	-0.450	0.394	-0.146	-0.019	0.186	20.0%		
50%	4	N	-0.416	0.594	-0.145	0.091	0.308	15.7%		
90%	4	N	-0.639	0.444	-0.136	0.005	0.169	17.7%		
10%	4	Y	-0.320	0.462	-0.154	-0.013	0.181	22.0%		
50%	4	Y	-0.423	0.560	-0.168	0.071	0.290	17.7%		
90%	4	Y	-0.369	0.482	-0.156	-0.005	0.166	23.0%		
Granger	1	N	-0.468	0.598	-0.052	0.169	0.369	6.7%		
Granger	1	Y	-0.472	0.598	-0.053	0.167	0.369	6.7%		
Multifactor	_	_	-0.122	0.669	0.054	0.266	0.487	0.3%		


Table I.31: The table reports residual correlation descriptive analyses for the Insurance Companies dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

			200	6-2008				
Model	N	Factor	Min	Max	q10	$\mathbf{q50}$	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.499	0.809	-0.261	-0.004	0.294	31.7%
Q(50%)	4	N	-0.468	0.812	-0.213	0.018	0.379	23.3%
Q(90%)	4	N	-0.477	0.804	-0.257	-0.008	0.315	30.7%
Q(10%)	4	Y	-0.475	0.806	-0.259	-0.008	0.325	30.7%
Q(50%)	4	Y	-0.461	0.810	-0.243	0.009	0.346	26.3%
Q(90%)	4	Y	-0.485	0.802	-0.266	-0.011	0.314	30.7%
\mathbf{G}	1	N	-0.425	0.831	-0.190	0.047	0.398	20.0%
G	1	Y	-0.420	0.836	-0.188	0.046	0.384	20.0%
4-F-CAPM	-	-	-0.370	0.847	-0.149	0.089	0.422	14.0%
			201	1-2015				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.302	0.554	-0.115	0.002	0.182	14.7%
Q(50%)	4	N	-0.293	0.636	-0.135	0.032	0.235	15.3%
Q(90%)	4	N	-0.313	0.510	-0.136	0.001	0.181	18.0%
Q(10%)	4	Y	-0.314	0.593	-0.152	-0.002	0.193	21.0%
Q(50%)	4	Y	-0.249	0.650	-0.142	0.031	0.220	19.0%
Q(90%)	4	Y	-0.292	0.545	-0.149	0.006	0.194	22.0%
G	1	N	-0.109	0.658	0.001	0.116	0.322	1.0%
G	1	Y	-0.103	0.658	-0.011	0.120	0.318	1.0%
4-F-CAPM	_	_	-0.103	0.658	0.017	0.138	0.332	0.3%


Table I.32: The table reports residual correlation descriptive analyses for the Industry portfolios dataset. The first column identify the various models, while the second column indicates the number of networks used in the model. In the first column Q(10%) identifies the use of a combination of causality networks from quantile regression (QB, Qo and QN) at the 10% quantile, combined with the Granger causality network. Similarly, when the reference quantile is 50% or 90%. With G we denote the model using just the Granger casuality network, while the last line refers to the 4-factor CAPM. The table reports statistics for the residuals correlations: the minimum, maximum, the 10% quantile q_{10} , the median q_{50} , the 90% quantile and the number of elements of the correlation matrix lower than -0.1.

		·	200	6-2008				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.444	0.756	-0.204	0.007	0.274	25.6%
$\mathrm{Q}(50\%)$	4	N	-0.449	0.757	-0.198	0.014	0.276	24.9%
Q(90%)	4	N	-0.444	0.756	-0.204	0.007	0.274	25.6%
Q(10%)	4	Y	-0.476	0.731	-0.226	-0.013	0.253	28.2%
Q(50%)	4	Y	-0.494	0.733	-0.228	-0.007	0.265	28.2%
Q(90%)	4	Y	-0.463	0.728	-0.227	-0.005	0.261	28.5%
G	1	N	-0.444	0.756	-0.204	0.007	0.274	25.6%
G	1	Y	-0.415	0.752	-0.203	0.009	0.273	25.4%
4-F-CAPM	-	-	-0.486	0.733	-0.219	0.003	0.267	27.1%
			201	1-2015				
Model	N	Factor	Min	Max	q10	q50	q90	% elements
	Networks							≤ -0.1
Q(10%)	4	N	-0.434	0.560	-0.161	-0.015	0.154	24.3%
Q(50%)	4	N	-0.450	0.588	-0.148	0.007	0.204	18.8%
Q(90%)	4	N	-0.424	0.551	-0.159	-0.010	0.170	23.5%
Q(10%)	4	Y	-0.431	0.564	-0.166	-0.010	0.179	23.0%
Q(50%)	4	Y	-0.461	0.589	-0.152	0.007	0.191	19.1%
Q(90%)	4	Y	-0.432	0.567	-0.166	-0.005	0.183	22.6%
G	1	N	-0.467	0.589	-0.144	0.009	0.208	18.4%
G	1	Y	-0.445	0.592	-0.148	0.008	0.208	18.4%
4-F-CAPM	_	_	-0.456	0.592	-0.147	0.007	0.208	18.4%

J Estimated Parameters




(a) 2006 - 2008




(b) 2011 - 2015

Figure J.27: This figure exhibits the $\rho's$ coefficients signficant (blue asterisks '*') and not significant (green circle 'o') by using different resulting networks obtained combining different causality methodologies. The $\rho's$ parameters are relative to first 25 banks monthly returns ordered for market capitalization. Panel a) reports the coefficients relative to the period 2006 – 2008. Panel b) reports the coefficients relative to the period 2011 – 2015.




(a) 2006 - 2008




(b) 2011 - 2015

Figure J.28: This figure exhibits the $\rho's$ coefficients signficant (blue asterisks '*') and not significant (green circle 'o') by using different resulting networks obtained combining different causality methodologies. The $\rho's$ parameters are relative to first 25 Insurers monthly returns ordered for market capitalization. Panel a) reports the coefficients relative to the period 2006 – 2008. Panel b) reports the coefficients relative to the period 2011 – 2015.



(a) 2006 - 2008



(b) 2011 - 2015

Figure J.29: This figure exhibits the $\rho's$ coefficients significant (blue asterisks '*') and not significant (green circle 'o') by using different resulting networks obtained combining different causality methodologies. The $\rho's$ parameters are relative to 48 Fama and French industry portfolios monthly returns ordered for market capitalization. Panel a) reports the coefficients relative to the period 2006 – 2008. Panel b) reports the coefficients relative to the period 2011 – 2015.