

Call for contributions

Climate action

Managing soils functions to mitigate and adapt to climate change (PICO)

EUROSOIL2020CONT-2246

STRATEGIES TO REDUCE AMMONIA VOLATILIZATION FROM N MINERAL FERTILIZERS IN THE PO RIVER BASIN, NORTHERN ITALY

Marta Mencaroni¹, Nicola Dal Ferro¹, Jacopo Furlanetto¹, Matteo Longo¹, Barbara Lazzaro¹, Brian Grant², Ward Smith², Francesco Morari¹

¹DAFNAE, University of Padova, Legnaro (Padova), Italy, ²Ottawa Research and Development Centre, Ottawa, Canada

Presentation Preference: Oral communication

Would you like to receive more information about applying for a grant once this becomes available?: Yes

Content: Ammonia (NH₃) losses from nitrogen mineral fertilizers is a debated issue in Europe due to their contribution to fine particulate matter formation (PM_{2.5} and PM₁₀ fractions). In particular, up to 70% NH₃ can be lost from urea-based fertilizers under broadcast application. A number of Best Available Techniques (BATs) for preventing and reducing emissions have been recommended although uncertainties in their effectiveness at the site-specific level still occur. The Life PREPAIR project aims to study reduction methods for NH₃ emissions from urban to agricultural sources in the Po river basin, northern Italy, one of the most air-polluted areas in Europe. Within the PREPAIR framework, the present study aims to identify site-specific based BATs which meet the criteria for the UNECE benchmark for mineral fertilizers, namely a NH₃ emission reduction of at least 30% compared to urea surface broadcast.

The study was conducted across the Veneto region, northeastern Italy. Firstly, a potential risk map of NH₃ volatilization was created by overlapping soil properties (e.g. pH, CEC) and climatic conditions which are well-known to affect NH₃ losses. In high risk hot spots, N losses (e.g. NH₃ and N₂O volatilization, N leaching, etc..) were estimated using a modified version of DNDC v. CAN (Dutta et al., 2016). Inputs for DNDC v. CAN were collected via a questionnaire conducted among farmers to identify business-as-usual practices for different crops. Alternative BATs, such as NH₄NO₃ (AN), organic fertilizers (e.g. beef digestate, liquid manure) and different incorporation methods (injection, deep incorporation), were compared to urea broadcast application for maize (*Zea mais* L.) and winter wheat (*Triticum aestivum* L.) production.

The most promising BATs were further tested in a field experiment on bare soil using a wind tunnel combined with a FTIR gas analyzer (Gasmeter DX4015) for continuous gas measurements. Wind speed, soil water content, and both air and soil temperature were also continuously monitored and used to improve prediction of NH₃ volatilization dynamics.

Modeling results showed that AN closed-slot injection and deep incorporation of organic fertilizers significantly reduced ammonia loss both for maize (-75% and -96% respectively) and winter wheat (-87% and -98%). Nevertheless, some increase in nitrate leaching was observed, mostly in case of winter wheat (+24% for injection of AN and +89% for organic fertilizers). Field results on bare soil confirmed the effectiveness of closed-slot injection, being able to bring a reduction up to 68% compared to NH₃ emissions after surface broadcast. Contrasting results were observed with urea incorporation or mixing only, that showed enhanced NH₃ emissions (+50%).

Our study showed that NH₃ loss can be highly reduced, and that N injection practices are among the most effective solutions in Veneto region. Furthermore, a site-specific approach is highly recommended to reduce N losses in air and water, which are strongly related to soil and climatic properties and their interactions with agricultural management.

Disclosure of Interest: None Declared

Keywords: ammonia volatilization