

UNIVERSITÀ DEGLI STUDI DI PADOVA

FACOLTÀ DI SCIENZE MM.FF.NN.

DIPARTIMENTO DI MATEMATICA PURA ED APPLICATA

TESI DI DOTTORATO IN MATEMATICA

The regularity of the minimum time  
function via nonsmooth analysis and  
geometric measure theory

**COORDINATORE:** Ch.mo Prof. Franco Cardin

**SUPERVISORE:** Ch.mo Prof. Giovanni Colombo

**DOTTORANDO:** Nguyen Tien Khai



# Acknowledgments



# Sommario



# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Preliminary</b>	<b>7</b>
2.1	Nonsmooth analysis and geometric measure theory . . . . .	7
2.1.1	Nonsmooth analysis . . . . .	7
2.1.2	Geometric measure theory . . . . .	9
2.2	Positive reach and external sphere condition . . . . .	10
2.2.1	Positive reach . . . . .	10
2.2.2	External sphere condition . . . . .	14
2.3	Control theory . . . . .	15
2.3.1	Control systems . . . . .	15
2.3.2	Minimum time function . . . . .	16
2.3.3	Controllability and continuity . . . . .	16
2.4	Differential inclusions . . . . .	18
<b>I</b>	<b>On the structure of the minimum time function</b>	<b>21</b>
<b>3</b>	<b>Semiconcavity type results</b>	<b>23</b>
3.1	Nonlinear control system . . . . .	24
3.2	Statement of the main results . . . . .	26
3.3	Preparatory Lemmas . . . . .	27
3.3.1	Transporting proximal normals . . . . .	27
3.3.2	Type of singularities of the minimum time function . . . . .	37
3.3.3	Wedgedness and exposed rays . . . . .	40
3.4	Proof of the main results of Chapter 3 . . . . .	45
3.4.1	Proof of Theorem 3.2.1 . . . . .	45
3.4.2	Proof of Theorem 3.2.3 . . . . .	46
3.5	The case of optimal points . . . . .	49
3.6	Examples . . . . .	53
3.7	The case of differential inclusions . . . . .	60
3.7.1	Hypotheses and some consequences . . . . .	60

3.7.2	The hypograph of the minimum time function satisfies an external sphere condition . . . . .	63
3.7.3	The inner ball property of attainable sets . . . . .	72
<b>4</b>	<b>Semiconvexity type results</b>	<b>77</b>
4.1	Linear control system and normality . . . . .	78
4.2	Quantitative strict convexity of reachable sets . . . . .	79
4.2.1	Linear autonomous control systems . . . . .	79
4.2.2	Linear nonautonomous control system in the case $N = 2$	88
4.3	Nonlinear control system in the case $N = 2$ . . . . .	90
4.3.1	Strict convexity of the reachable set . . . . .	90
4.3.2	Optimal points . . . . .	93
4.3.3	The epi-graph of the minimum time function has positive reach . . . . .	95
<b>II</b>	<b>Regularity of a class of continuous functions</b>	<b>99</b>
<b>5</b>	<b>External sphere condition and continuous functions</b>	<b>101</b>
5.1	Notation . . . . .	102
5.2	Main results . . . . .	103
5.2.1	Statement of main results . . . . .	103
5.2.2	Outline of proof of Theorem 5.2.2 . . . . .	104
5.3	Preparatory Lemmas . . . . .	105
5.3.1	Closedness of the set of bad points . . . . .	105
5.3.2	Zero Lebesgue measure of some special subsets of the set of bad points . . . . .	111
5.4	Proof of our main results of Chapter 5 . . . . .	114
5.4.1	Proof of Theorem 5.2.2 . . . . .	114
5.4.2	Proof of Theorem 5.2.1 . . . . .	119
5.4.3	Proof of Colloraries . . . . .	120
<b>6</b>	<b>Rectifiability of the set of bad points</b>	<b>121</b>
6.1	Notations . . . . .	121
6.2	Rectifiability results for the set of bad points . . . . .	123
6.2.1	Preparatory Lemmas . . . . .	123
6.2.2	Proof of main results . . . . .	126
6.3	A counterexample . . . . .	128
<b>7</b>	<b>Appendix</b>	<b>131</b>
7.1	Appendix A . . . . .	131
7.2	Appendix B . . . . .	133
7.3	Appendix C . . . . .	136



# Chapter 1

## Introduction

The minimum time problem is classical in control theory. Given a nonempty closed target  $\mathcal{S}$  and a control system

$$\begin{cases} \dot{y}(t) = f(t, y(t), u(t)) & a.e. \\ u(t) \in \mathcal{U} & a.e. \\ y(0) = x, \end{cases} \quad (1.0.1)$$

where the function  $f : \mathbb{R} \times \mathbb{R}^N \times \mathcal{U} \rightarrow \mathbb{R}^N$  is smooth enough and the control set  $\mathcal{U}$  is a compact nonempty subset of  $\mathbb{R}^M$ , for each admissible control  $u(\cdot) \in \mathcal{U}_{ad}$ , i.e.  $u(\cdot)$  is measurable and takes value in  $\mathcal{U}$ , there exists a unique solution  $y^{x,u}(\cdot)$  of (1.0.1) which is the trajectory starting from  $x$  under the control  $u(\cdot)$ . The minimum time needed to steer  $x$  to  $\mathcal{S}$ , regarded as a function of  $x$ , is called the minimum time function and is denoted by

$$T_{\mathcal{S}}(x) := \inf \{ \theta_{\mathcal{S}}(x, u) \mid u(\cdot) \in \mathcal{U}_{ad} \},$$

where  $\theta_{\mathcal{S}}(x, u) := \inf \{ t \geq 0 \mid y^{x,u}(t) \in \mathcal{S} \}$ . In general,  $T_{\mathcal{S}} \in [0, \infty]$ . The controllable set  $\mathcal{R}$  consists of all points  $x \in \mathbb{R}^N$  such that  $T_{\mathcal{S}}(x)$  is finite. The regularity of the minimum time function is related on one hand to the controllability properties of system (1.0.1), on the other one to the regularity of the target and of the dynamics, together with suitable relations between them.

Such topics were studied by several authors (see, e.g., [11, 12, 16, 17, 20, 21, 23, 29, 65] and reference therein) under different viewpoints. In particular, it is well known that in general the minimum time function  $T$  is not everywhere differentiable. It is also well known that suitable controllability conditions imply the Hölder continuity of  $T$  (see, e.g., [11, Chapter IV] and references therein). However, the latter fact does not provide information on differentiability. In a 1995 paper (see [20] and also Chapter 8 in the book [21]), Cannarsa and Sinestrari found a connection between the control system and the target which actually implies the semiconcavity

(or the semiconvexity) of  $T$ . Semiconcave functions are – essentially –  $\mathcal{C}^2$ -perturbations of concave functions and therefore inherit several regularity properties from convexity. Several features of semiconcavity were thoroughly studied (see Chapters 3, 4, 5 in [21] and references therein), thus providing a rich set of information on the structure of the minimum time function and suggesting semiconcavity/semiconvexity as a good regularity class for such value functions. The main result in [20] shows that if the target satisfies a *uniform internal ball condition* (see Definition 2.2.2 below) and the control system is smooth enough, then  $T$  is semiconcave, provided a strong *control-liability assumption*, called Petrov condition, holds. A partially symmetric result, contained in [20], states that if the target is convex and the control system is linear, then  $T$  is semiconvex, provided, again, Petrov condition holds. The latter requires that the minimized Hamiltonian at all boundary points of  $\mathcal{S}$ , computed along unit normal vectors, be bounded away from zero locally uniformly, i.e., for all  $R > 0$  there exists  $\mu > 0$  such that for all  $x \in \text{bdry}\mathcal{S} \cap B(0, R)$ ,

$$\min_{u \in \mathcal{U}} \langle f(x, u), \zeta \rangle < -\mu, \quad \text{for all } \zeta \in N_{\mathcal{S}}(x), \|\zeta\| = 1. \quad (1.0.2)$$

It is well known that Petrov condition is equivalent to the local Lipschitz continuity of  $T$  (see, e.g., [21, Section 8.2]).

In an entirely different setting, a class of sets which includes both convex and  $\mathcal{C}^2$ -sets was studied independently by several authors (including Federer [39], Canino [14], Clarke, Stern and Wolenski [26], Poliquin, Rockafellar and Thibault [53]) under different names, for example *sets with positive reach* [39],  *$\varphi$ -convex sets* [14], *proximally smooth sets* [26], and *prox-regular sets* [53]. Such sets, which in this thesis will be called sets with positive reach, are characterized by a strong external sphere condition (see Definition 2.2.1 below): every normal vector must be realized by a locally uniform ball. By observing that a convex set satisfies the same type of external sphere condition with an arbitrarily large radius, it is natural to expect that sets with positive reach enjoy locally several properties that convex sets enjoy globally. In particular, this holds for the metric projection, which is unique in a neighborhood of a set with positive reach  $K$ . This fact is used in proving all the regularity properties which are satisfied by sets with positive reach (see, e.g., [39, Section 4]). Semiconcave functions and sets with positive reach, through the hypograph, are linked together (see, e.g., Theorem 5.2 in [26], where semiconvex functions are called *lower- $\mathcal{C}^2$* ): a locally Lipschitz function is semiconcave if and only if its hypograph has positive reach. Of course an entirely symmetric characterization for semiconvex functions can be expressed using the epigraph. Trying to generalize to functions whose hypo/epigraph has positive reach some regularity properties enjoyed by semiconcave/convexity functions was therefore a natural challenge. Some results on this line were obtained in [27, 28], including the a.e. twice differ-

entiability (see Theorem 2.2.2 below) together some results on the structure of singularities.

In several control problems, controllability assumptions weaker than Petrov condition hold, and therefore the minimum time function is not locally Lipschitz. A natural question therefore is trying to understand whether the structure of the minimum time function remains unchanged if in the above setting the controllability assumptions are weakened. In other words it is natural to investigate whether the hypograph/epigraph of  $T$  has positive reach if  $T$  is supposed to be only continuous.

This thesis has been inspired by the above question. It is divided into two parts. The first one is devoted to the analysis of the minimum time function. The second one is motivated by the first part, and contains results on the regularity of merely continuous functions.

**Part I: On the structure of the minimum time function**

This part is dedicated to two types of regularity of the minimum time function  $T$ . More precisely, we will study in Chapter 3 semiconcavity type results for  $T$ . We first assume that the nonlinear control system is (essentially)  $\mathcal{C}^2$ , the target  $\mathcal{S}$  satisfies an internal sphere condition, and  $T$  is continuous, and study the hypograph of  $T$  in the complement of  $\mathcal{S}$ . Since the internal sphere property is closed with respect to the union operator, one can see intuitively that the reachable set  $\mathcal{R}^t$ , which is the set of points reachable from  $\mathcal{S}$  in time less than  $t$ , inherits such property from  $\mathcal{S}$ . By combining this fact and the Hamiltonian function, a regularity result on the hypograph of  $T$  can be obtained. The corresponding theorem is as follows:

**Theorem 1.0.1** *Under the above assumptions, the hypograph of  $T$  satisfies an external sphere condition.*

From this theorem, we obtain that if  $T$  is Lipschitz then  $T$  is semiconcave (see [50]). However, here the situation is more complicated than in the Lipschitz case: the main results depend on the pointedness of the normal cone to the hypograph. Indeed, from a representation of generalized supergradient of  $T$ , we prove that

**Theorem 1.0.2** *Together with the above assumptions, if the normal cone to the hypograph is pointed in the complement of  $\mathcal{S}$ , then the hypograph of  $T$  has positive reach.*

In the last section of this chapter, we also prove Theorem 1.0.1 for a class of differential inclusions taken from [22]. Moreover, in the spirit of [17] we finally extend this result to arbitrary target  $\mathcal{S}$ .

The next chapter is devoted to semiconvexity type results for  $T$ . For a linear control system and a convex target, the result is contained in [29]. However, for a nonlinear control system, one has to face the difficulty that the convexity (even the external sphere property) of the reachable set  $\mathcal{R}^t$

can be easily broken after any small time  $t$  (see example 4.3 in [20]). This is quite natural since the union of convex sets usually has inner corners or even cusps. Moreover, the "external normal regularity" of the reachable set is also related to the uniqueness of the optimal trajectory from a point to the target  $\mathcal{S}$ . Hence, finding a class of nonlinear control systems such that the convexity (or the external sphere property) of  $\mathcal{R}^t$  still holds up to small time  $t$  is a natural problem.

This chapter is first devoted to a result of this type. Indeed, we assume that the target is the origin and the linear part of the nonlinear control system at the origin is *normal* (see the definition in Theorem 1.0.3 below) and study the reachable set  $\mathcal{R}^t$ . More precisely, for  $t > 0$  small the normality of the linearization together with a further condition on the Taylor development at 0 of the nonlinear control system yields the strict convexity of the reachable set  $\mathcal{R}_L^t$  corresponding to the linear nonautonomous systems which are obtained by linearizing the nonlinear control system along the optimal trajectories. Therefore, it is reasonable to conjecture that the convexity of the reachable set, for  $t > 0$  small, still holds also for a suitable nonlinear control system. For this approach, the main preliminary result is as follows:

**Theorem 1.0.3** *Consider the linear control system*

$$\dot{x} = Ax + Bu, \quad (1.0.3)$$

where  $A \in \mathbb{M}_{N \times N}$ ,  $B \in \mathbb{M}_{N \times M}$ ,  $M \leq N$  and  $u = (u_1, u_2, \dots, u_M) \in \mathbb{R}^M$ ,  $|u_j| \leq 1$  for  $j = 1, 2, \dots, M$ .

Assume that (1.0.3) is normal, i.e., for every column  $b_j$ ,  $j = 1, 2, \dots, M$  of  $B$ ,

$$\text{rank} [b_j, Ab_j, \dots, A^{N-1}b_j] = N.$$

Then for all  $T > 0$  there exists a constant  $\gamma > 0$ , depending only on  $N, M, A, B, T$  such that for all  $x, y \in \mathcal{R}^T$ , for all  $\zeta \in N_{\mathcal{R}^T}(x)$ , it holds

$$\langle \zeta, y - x \rangle \leq -\gamma \|\zeta\| \|y - x\|^N. \quad (1.0.4)$$

However, the exponent of strict convexity of  $\mathcal{R}_L^t$  is  $N$  as in (1.0.4), while the exponent of the perturbative term appearing from the linearization is 2. Therefore, this approach is effective only in the two dimensional case (see Theorem 4.3.1). On the basis of the preceding result, we will prove that the epigraph of  $T$  in a neighborhood of 0 has positive reach (see Theorem 4.3.3). This will require proving that all points close enough to the origin are indeed optimal.

## Part II: The regularity of a class of continuous functions

Since verifying that a set has positive reach is often demanding, finding sufficient conditions for this property appears of some interest. In [49], a class of sets which are characterized by an external sphere condition (at each

point on the boundary, there exists *one* proximal normal vector realized by a locally uniform ball) is considered. The authors proved that if a set satisfies this condition and is *wedged* (this concept was introduced by Rockafellar in [54]) then it has positive reach. This results was later generalized in [50] by the same authors to investigate the relationships among functions whose hypograph satisfies an external sphere condition, the functions with positive reach hypograph and semiconcave functions. Wedgedness of a set  $C$  is equivalent to the pointedness of the Clarke normal cone to  $C$ , i.e. the normal cone does not contain lines (see [25] and [58]). Moreover, the pointedness assumption for the proximal normal cone to the hypograph of the minimum time function  $T$  appears pivotal in our result [19] (mentioned in the first part) for computing generalized gradients of  $T$  and then for proving that the hypograph of  $T$  has positive reach.

Several counterexamples (see. e.g, [49]), though, show that the external sphere condition is in general weaker than positive reach. In particular, in Example 2 in Section 6 of Chapter 3, we constructed a minimum time function with a constant dynamics and a  $C^{1,1}$  target such that its hypograph satisfies an external sphere condition but has not positive reach everywhere. On the other hand, the pointedness assumption for the normal cone to the hypograph of a continuous function is hard to verify since it is related to the representation formula for its generalized supergradient (this problem is studied in [30]). Therefore, the problem of understanding whether some concavity features are preserved under the external sphere condition appears natural. In Chapter 5 an answer to this question is provided. Our main result reads -essentially- as follows

**Theorem 1.0.4** *Let  $\Omega \subset \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be continuous. Assume that the hypograph of  $f$  satisfies the weak external sphere condition. Then there exists a closed set  $\Gamma$  with zero Lebesgue measure such that the hypograph of the restricted function  $f_{\Omega \setminus \Gamma}$  has positive reach.*

Consequently, a function satisfying the assumption of the above theorem enjoys several regularity properties inherited by functions whose hypograph has positive reach. Therefore, using Theorem 1.0.1 and Theorem 1.0.4 the pointedness assumption of the hypograph of  $T$  in Chapter 3 is removed and the a.e. twice differentiability of  $T$  for a class of differential inclusions is also obtained.

In general, however, sets with null Lebesgue measure can be very irregular and possess almost no structure. A natural question is then that of investigating the properties of the singular set  $\Sigma(f)$  for special classes of a.e. differentiable functions  $f$ .

When  $f$  is convex or concave, the properties of  $\Sigma(f)$  were first investigated in [36] and then developed in [64], [63], [61], [62], [2] and [3]. The basic approach in such papers is that of estimating the size of  $\Sigma(f)$ . We mention here a result which is essentially due to L. Zajíček and was later extended

to semiconcave functions by G. Alberti, L. Ambrosio and P. Cannarsa [1]. By  $\partial^F f(x)$  we denote here the Fréchet supergradient of  $f$  at  $x$ .

**Theorem 1.0.5 ([1])** *Let  $f$  be locally semiconcave. Then, for any  $k = 1, \dots, N$  the singular set  $\Sigma^k(f) := \{x \in \Omega \mid \dim \partial^F f(x) = k\}$  is countably  $(N - k)$ -rectifiable. In particular,  $\Sigma(f)$  is countably  $(N - 1)$ -rectifiable and  $\Sigma^N(u)$  is at most countable.*

The result also holds for the case of a continuous function  $f$  which has the hypograph with positive reach (see in [27]). Therefore, at the end we will study in Chapter 6 the rectifiability of the zero Lebesgue measure set  $\Gamma$ . The corresponding result is Theorem 1.0.4.

# Chapter 2

## Preliminary

### 2.1 Nonsmooth analysis and geometric measure theory

#### 2.1.1 Nonsmooth analysis

We quickly review in this subsection some basic concepts from nonsmooth analysis. Standard references are in [25, 44, 58].

Let  $x \in Q$  and  $v \in \mathbb{R}^N$ . We say that  $v$  is a *proximal normal* to  $Q$  at  $x$  (and we will denote this fact by  $v \in N_Q^P(x)$ ) if there exists  $\sigma = \sigma(v, x) \geq 0$  such that

$$\langle v, y - x \rangle \leq \sigma \|y - x\|^2 \quad \text{for all } y \in Q; \quad (2.1.1)$$

equivalently  $v \in N_Q^P(x)$  if and only if there exists  $\lambda > 0$  such that  $\pi_Q(x + \lambda v) = \{x\}$ . We say that the proximal normal  $v$  is *realized by a ball of radius*  $\rho > 0$  if  $\rho$  is the supremum of all  $\lambda$  such that  $\pi_Q(x + \lambda v) = \{x\}$ . In this case the best constant  $\sigma$  such that (2.1.1) holds true is  $\|v\|/(2\rho)$ . The following further concepts of normal vectors will be used (see [25, Chapter I] and [58, Chapter VI]). Let  $x \in Q$  and  $v \in \mathbb{R}^N$ . We say that:

1.  $v$  is a *Fréchet normal* (or Bouligand normal) to  $K$  at  $x$  ( $v \in N_Q^F(x)$ ) if

$$\limsup_{Q \ni y \rightarrow x} \langle v, \frac{y - x}{\|y - x\|} \rangle \leq 0;$$

2.  $v$  is a *limiting, or Mordukhovich, normal* to  $Q$  at  $x$  ( $v \in N_Q^L(x)$ ) if

$$v \in \{w \mid w = \lim w_n, w_n \in N_Q^P(x_n), x_n \rightarrow x\}$$

and is a *Clarke normal* ( $v \in N_Q^C(x)$ ) if  $v \in \overline{\text{co}}N_Q^L(x)$ . It is well known that  $N_Q^P(x)$  is convex.

Let  $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{+\infty\}$  be a lower semicontinuous function. By using  $\text{epi}(f) := \{(x, \xi) \mid \xi \geq f(x)\}$  and  $\text{hypo}(f) := \{(x, \xi) \mid \xi \leq f(x)\}$  one can

define some concepts of generalized differential for  $f$  at  $x \in \text{dom}(f) = \{x \in \mathbb{R}^N \mid f(x) < +\infty\}$ . Let  $x \in \text{dom}(f)$ ,  $v \in \mathbb{R}^N$ . We say that:

1.  $v$  is a *proximal subgradient* of  $f$  at  $x$  ( $v \in \partial_P f(x)$ ) if  $(v, -1) \in N_{\text{epi}(f)}^P(x, f(x))$ ; equivalently (see [25, Theorem 1.2.5]),  $v \in \partial_P f(x)$  iff there exist  $\sigma, \eta > 0$  such that for all  $y \in B(x, \eta) \cap \text{dom}(f)$ , it holds

$$f(y) \geq f(x) + \langle v, y - x \rangle - \sigma \|y - x\|^2; \quad (2.1.2)$$

2.  $v$  is a *proximal supergradient* of  $f$  at  $x$  ( $v \in \partial^P f(x)$ ) if  $(-v, 1) \in N_{\text{hypo}(f)}^P(x, f(x))$ ; equivalently  $v \in \partial^P f(x)$  iff  $-v \in \partial_P(-f)(x)$ , i.e., iff there exist  $\sigma, \eta > 0$  such that for all  $y \in B(x, \eta) \cap \text{dom}(f)$ , it holds

$$f(y) \leq f(x) + \langle v, y - x \rangle + \sigma \|y - x\|^2; \quad (2.1.3)$$

3.  $v$  is a *Fréchet subgradient* of  $f$  at  $x$  ( $v \in \partial_F f(x)$ ) if  $(v, -1) \in N_{\text{epi}(f)}^F(x, f(x))$ , i.e.,

$$\liminf_{y \rightarrow x} \frac{f(y) - f(x) - \langle v, y - x \rangle}{\|y - x\|} \geq 0;$$

4.  $v$  is a *Fréchet supergradient* of  $f$  at  $x$  ( $v \in \partial^F f(x)$ ) if  $(-v, 1) \in N_{\text{hypo}(f)}^F(x, f(x))$ ;

5.  $v$  is a *limiting subgradient* of  $f$  at  $x$  ( $v \in \partial_L f(x)$ ) if  $(v, -1) \in N_{\text{epi}(f)}^L(x, f(x))$ .

6.  $v$  is a *limiting supergradient* of  $f$  at  $x$  ( $v \in \partial^L f(x)$ ) if  $(-v, 1) \in N_{\text{hypo}(f)}^L(x, f(x))$ .

7.  $v$  is a *Clarke generalized gradient* of  $f$  at  $x$  ( $v \in \partial f(x)$ ) if  $(v, -1) \in N_{\text{epi}(f)}^C(x, f(x))$ . We recall that if  $f$  is Lipschitz continuous in a neighborhood of  $x$ , then  $v \in \partial f(x)$  if and only if  $v \in \overline{\text{co}}\{\zeta \mid \zeta = \lim Df(x_i), x_i \in \text{dom}(Df), x_i \rightarrow x\}$  (see [25, Theorem 8.1]).

It follows readily from the definitions that the inclusions

$$N_Q^P(x) \subseteq N_Q^F(x) \subseteq N_Q^L(x) \subseteq N_Q^C(x)$$

hold, together with their analogues for the sub- and supergradient. Moreover, if a vector  $v$  belongs to both the Fréchet sub- and supergradient of  $f$  at  $x$ , then  $f$  is Fréchet differentiable at  $x$  and  $Df(x) = v$ .

For a not necessarily Lipschitz function  $f$ , the *horizon subgradient*  $\partial_\infty f$  plays an important role. This is defined as

$$\partial_\infty f(x) = \{v \in \mathbb{R}^N \mid (v, 0) \in N_{\text{epi}(f)}^C(x, f(x))\},$$



and is clearly a closed convex cone. In particular, if  $f$  is not locally Lipschitz in a neighborhood of  $x$ , then  $\partial f(x)$  may be represented using  $\partial_\infty$ , namely (see [44, Prop. 2.6] or [58, Theorem 8.49])

$$\partial f(x) = \text{cl} (\text{co } \partial_L f(x) + \text{co } \partial_\infty f(x)). \quad (2.1.4)$$

Finally, we also consider a notion of *proximal* horizon supergradient, namely the convex cone

$$\partial^\infty f(x) = \{v \in \mathbb{R}^N \mid (-v, 0) \in N_{\text{hypo}(f)}^P(x, f(x))\}.$$

### 2.1.2 Geometric measure theory

We just introduce in this subsection some definitions needed our results. For basic concepts of geometric measure theory we refer to [5, 37].

Let  $k \geq 0$  and  $A \subset \mathbb{R}^N$  be fixed. The *k-dimensional Hausdorff measure* of  $A$  is defined as

$$\mathcal{H}^k(A) := \lim_{\delta \rightarrow 0^+} \mathcal{H}_\delta^k(A) = \sup_{\delta > 0} \mathcal{H}_\delta^k(A)$$

where for any  $\delta > 0$  we set

$$\mathcal{H}_\delta^k(A) := \inf \left\{ \sum_{i \in I} (\text{diam } A_i)^k \mid A \subset \bigcup_{i \in I} A_i, \text{diam } A_i < \delta \right\}.$$

The *Hausdorff dimension* of  $A$  is

$$\mathcal{H}\text{-dim}(A) := \inf\{k \geq 0 \mid \mathcal{H}^k(A) = 0\} = \sup\{k \geq 0 \mid \mathcal{H}^k(A) = \infty\}.$$

It is well known (see e.g. [40, 46]) that  $\mathcal{H}^k$  is a Borel measure on  $\mathbb{R}^N$ ;  $\mathcal{H}^0$  is the counting measure. Moreover, if  $k \in \mathbb{N}$  and  $S$  is a  $k$ -dimensional Lipschitz surface, then the surface measure of  $S$  coincides with  $\frac{2^k}{\omega_k} \mathcal{H}^k \llcorner S$ .

Let  $k \in \mathbb{N}$ ; we say that  $A \subset \mathbb{R}^N$  is *countably k-rectifiable* if

$$A \subset \mathcal{N} \cup \bigcup_{i=1}^{\infty} S_i$$

where  $S_i$  are suitable Lipschitz  $k$ -dimensional surfaces and  $\mathcal{N}$  is a  $\mathcal{H}^k$ -negligible set. We say that  $A$  is *k-rectifiable* if it is countably  $k$ -rectifiable and  $\mathcal{H}^k(A) < \infty$ .

Any countably  $k$ -rectifiable set  $A$  satisfies  $\mathcal{H}\text{-dim}(A) = k$ . It is well known that, if  $f : A \subset \mathbb{R}^k \rightarrow \mathbb{R}^N$  is Lipschitz continuous, then  $f(A)$  is countably  $k$ -rectifiable; if  $A$  is bounded, then  $f(A)$  is  $k$ -rectifiable.

In what follows, given  $A \subset \mathbb{R}^N$  we define its  $\epsilon$ -neighborhood  $(A)_\epsilon$  by

$$(A)_\epsilon := \{x \in \mathbb{R}^N \mid \text{there exists } y \in A \text{ such that } \|x - y\| < \epsilon\}.$$

Let  $\mathcal{K}$  denote the set of closed subsets of  $S^{N-1} \subset \mathbb{R}^N$ ; for  $A, B \in \mathcal{K}$  we introduce the *Hausdorff distance*  $d_{\mathcal{H}}(A, B)$  by

$$d_{\mathcal{H}}(A, B) = \inf\{\epsilon > 0 \mid A \subset (B)_{\epsilon} \text{ and } B \subset (A)_{\epsilon}\}.$$

It turns out (see e.g. [6]) that  $(\mathcal{K}, d_{\mathcal{H}})$  is a complete compact metric space.

## 2.2 Positive reach and external sphere condition

### 2.2.1 Positive reach

The concept of reach originates from the unique nearest point property. More precisely, the reach of a subset  $Q$  of  $\mathbb{R}^N$  is the largest  $r$  (possibly  $\infty$ ) such that if  $x \in \mathbb{R}^N$  and the distance,  $d_Q(x)$ , from  $x$  to  $Q$  smaller than  $r$ , then  $Q$  contains a unique point,  $\pi_Q(x)$ , nearest to  $x$ .

Let  $Q \subseteq \mathbb{R}^N$  be closed. We denote by  $\partial Q$  the topological boundary of  $Q$ , and, for  $x \in \mathbb{R}^N$ ,

$$\begin{aligned} d_Q(x) &= \inf\{\|y - x\| \mid y \in Q\} && \text{(the distance of } x \text{ from } Q) \\ \pi_Q(x) &= \{y \in Q \mid \|y - x\| = d_Q(x)\} && \text{(the metric projections of } x \text{ into } Q). \end{aligned}$$

Moreover, we set

$$\text{Unp}(Q) = \{x \in \mathbb{R}^N : \pi_Q(x) \text{ is a singleton}\}.$$

If  $x \in Q$  then

$$\text{reach}(x, Q) = \sup\{r \mid B(x, r) \subset \text{Unp}(Q)\}$$

where  $B(x, r) = \{y \mid \|y - x\| < r\}$ .

Also

$$\text{reach}(Q) = \inf\{\text{reach}(Q, x) \mid x \in Q\}.$$

**Remark 2.2.1** *The function  $\text{reach}(Q, \cdot)$  is continuous in  $Q$ . Moreover if  $\text{reach}(Q) > 0$  then  $Q$  is closed.*

If  $\text{reach}(Q) > 0$  we say that  $Q$  has positive reach. Moreover, a set with positive reach can be alternatively defined as follow (see in [14, 35]).

**Definition 2.2.1** *Let  $Q \subset \mathbb{R}^N$  be closed. We say that  $Q$  has positive reach if there exists a continuous function  $\varphi : Q \rightarrow [0, \infty)$  be continuous such that for all  $x, y \in Q, v \in N_Q^P(x)$ , the inequality*

$$\langle v, y - x \rangle \leq \varphi(x) \|v\| \|x - y\|^2$$

*holds, i.e.  $v \in N_Q^P(x)$  is realized by a ball of radius  $\frac{1}{2\varphi(x)}$ .*

**Remark 2.2.2** If a set  $Q$  has positive reach then the function  $\varphi(\cdot)$  in the definition (2.2.1) can be replaced by  $\frac{1}{2 \operatorname{reach}(\cdot, Q)}$ .

**Proof.** The proof can be found in [39, Theorem 4.8 (7)].  $\square$

It is therefore clear that every closed and convex set has positive reach, with  $\operatorname{reach}(Q) = \infty$ , and every closed set with a  $\mathcal{C}^{1,1}$ -boundary has positive reach, with  $\operatorname{reach}(Q) = L/2$ , where  $L$  is the Lipschitz constant of a suitable parametrization of  $\partial Q$ . Some properties of the distance from a set with positive reach  $Q$  and the metric projection onto  $Q$  are important features of this class of sets.

**Theorem 2.2.1** Let  $Q \subset \mathbb{R}^N$  be a set with positive reach. Then there exists an open set  $U \supset K$  such that

- (1)  $d_Q \in \mathcal{C}^{1,1}(U \setminus Q)$  and  $Dd_Q(y) = \frac{y - \pi_Q(y)}{d_K(y)}$  for every  $y \in U \setminus Q$ ;
- (2)  $\pi_Q : U \rightarrow Q$  is a locally Lipschitz single-valued map. In particular, the function  $\pi_Q : \{x \in \mathbb{R}^N \mid d(x, Q) < \frac{\operatorname{reach}(Q)}{2}\} \rightarrow Q$  is Lipschitz with Lipschitz ratio 2.

Moreover,

- (3)  $Q$  has finite perimeter in  $\mathbb{R}^N$  (provided it is compact);
- (4) for every  $x \in Q$ ,  $N_Q^P(x) = N_Q^C(x)$ ;
- (5) the set valued map  $N_Q^P(\cdot)$  has closed graph in  $\partial Q \times \mathbb{R}^N$ .

**Proof.** The proof of (1) and (2) can be found in [14, Proposition 2.6, 2.9, Remark 2.10] or in [39, §4]. The proof of (3) is in [27, §5], while (4) and (5) can be found in several papers, including [53].  $\square$

**Remark 2.2.3** Conditions (1) and (2) in Theorem 2.2.1 are actually equivalent to positive reach, as it is proved, e.g., in [39, §4]. Examples of finite dimensional sets with positive reach can be found, e.g., in [39].

We also give here Lemma 3.1 in [32] which concerns an estimate of the excess of the convex hull of a set with positive reach  $Q$  over  $Q$ .

**Lemma 2.2.1** Let  $Q$  be a set with positive reach and let  $x \in \operatorname{co}Q$  and  $d_Q(x) < \operatorname{reach}(Q)$ . Then

$$\|x - \pi_Q(x)\| \leq \frac{1}{2 \operatorname{reach}(Q)} \sum_{i,j=1}^{N+1} t_i t_j \|x_i - x_j\|^2,$$

where  $t_i \geq 0$ ,  $\sum_{i=1}^{N+1} t_i = 1$ ,  $x_i \in Q$ , and  $x = \sum_{i=1}^{N+1} t_i x_i$ .

**Proof.** From Remark 2.2.2, we have for each  $i = 1, \dots, N+1$ ,

$$\langle x - \pi_Q(x), x_i - \pi_Q(x) \rangle \leq \frac{1}{2 \operatorname{reach}(Q)} \|x - \pi_Q(x)\| \|x_i - \pi_Q(x)\|^2,$$

so that

$$\langle x - \pi_Q(x), \sum_{i=1}^{N+1} t_i x_i - \pi_Q(x) \rangle \leq \frac{\|x - \pi_Q(x)\|}{2 \operatorname{reach}(Q)} \sum_{i=1}^{N+1} t_i \|x_i - \pi_Q(x)\|^2.$$

Recalling that  $x = \sum_{i=1}^{N+1} t_i x_i$ , we thus obtain

$$\|x - \pi_Q(x)\| \leq \frac{1}{2 \operatorname{reach}(Q)} \sum_{i=1}^{N+1} t_i \|x_i - \pi_Q(x)\|^2. \quad (2.2.1)$$

Putting  $I = \sum_{i=1}^{N+1} t_i \|x_i - x\|^2$ , from an elementary computation taking into account the condition  $\sum_{i=1}^{N+1} t_i (x - x_i) = 0$ , we obtain, for all  $v \in \mathbb{R}^N$ ,

$$\sum_{i=1}^{N+1} t_i \|x_i - v\|^2 = \|x - v\|^2 + I. \quad (2.2.2)$$

Now we compute  $I$ . Taking  $v = x_j$  in (2.2.2), we have

$$\sum_{i=1}^{N+1} t_i \|x_i - x_j\|^2 = \|x - x_j\|^2 + I.$$

Thus we obtain both

$$t_j \sum_{i=1}^{N+1} t_i \|x_i - x_j\|^2 = t_j \|x - x_j\|^2 + t_j I$$

and

$$\sum_{j=1}^{N+1} \sum_{i=1}^{N+1} t_j t_i \|x_i - x_j\|^2 = \sum_{j=1}^{N+1} t_j \|x - x_j\|^2 + \sum_{j=1}^{N+1} t_j I.$$

From  $\sum_{j=1}^{N+1} t_j = 1$  and  $I = \sum_{j=1}^{N+1} t_j \|x - x_j\|^2$ , we obtain

$$I = \frac{1}{2} \sum_{j=1}^{N+1} \sum_{i=1}^{N+1} t_j t_i \|x_i - x_j\|^2.$$

Using this expression in (2.2.2) with  $\pi_Q(x)$  in place of  $v$ , we obtain

$$\sum_{i=1}^{N+1} t_i \|x_i - \pi_Q(x)\|^2 = \|x - \pi_Q(x)\|^2 + \frac{1}{2} \sum_{j=1}^{N+1} \sum_{i=1}^{N+1} t_j t_i \|x_i - x_j\|^2.$$

Thus

$$\|x - \pi_Q(x)\| \leq \frac{1}{2 \operatorname{reach}(Q)} \left( \|x - \pi_Q(x)\|^2 + \frac{1}{2} \sum_{j=1}^{N+1} \sum_{i=1}^{N+1} t_j t_i \|x_i - x_j\|^2 \right).$$

Since  $\|x - \pi_Q(x)\| = d_Q(x) < \operatorname{reach}(Q)$ , the proof is concluded.  $\square$

In both optimal control and partial differential equations theory, *semi-concave functions* play an important role (see, e.g., [11, 21]). Let  $\Omega \subset \mathbb{R}^N$  be open: a function  $f : \Omega \rightarrow \mathbb{R}$  is said to be semiconcave if for every  $x \in \Omega$  and every  $\delta > 0$  there exists a constant  $C > 0$  such that

$$f(x) - C \|x\|^2 \text{ is concave in } B(x, \delta).$$

Semiconcave functions are therefore locally Lipschitz. Moreover, thanks to Theorem 5.2 in [26], the hypograph of such functions has positive reach. More in general, upper semicontinuous functions which have hypograph with positive reach (or l.s.c. functions which have epigraph with positive reach) enjoy several of the regularity properties, except Lipschitz continuity, that semiconcave functions satisfy. Such functions identify the class which we want to show that our minimum time belongs to. To this aim, we state a result which collects the main properties.

**Theorem 2.2.2** *Let  $\Omega \subset \mathbb{R}^N$  be open, and let  $f : \overline{\Omega} \rightarrow \mathbb{R} \cup \{+\infty\}$  be proper, upper semicontinuous, and such that  $\operatorname{hypo}(f)$  has positive reach. Then there exists a sequence of sets  $\Omega_h \subseteq \Omega$  such that  $\Omega_h$  is compact in  $\operatorname{dom}(f)$  and*

- (1) *the union of  $\Omega_h$  covers  $\mathcal{L}^N$ -almost all  $\operatorname{dom}(f)$ ;*
- (2) *for all  $x \in \bigcup_h \Omega_h$  there exist  $\delta = \delta(x) > 0$ ,  $L = L(x) > 0$  such that  $f$  is Lipschitz on  $B(x, \delta)$  with ratio  $L$ , and hence semiconcave on  $B(x, \delta)$ .*

Consequently,

- (3)  *$f$  is a.e. Fréchet differentiable and admits a second order Taylor expansion around a.e. point of its domain.*

Moreover, the set of points where the graph of  $f$  is nonsmooth has small Hausdorff dimension. More precisely, for every  $k = 1, \dots, N$ , the set

$$\{x \in \operatorname{int} \operatorname{dom}(f) \mid \operatorname{Dim}(\partial^P f(x)) \text{ is } \geq k\}$$

is countably  $\mathcal{H}^{N-k}$ -rectifiable.

This result is essentially Theorem 5.1 in [27].

### 2.2.2 External sphere condition

**Definition 2.2.2** Let  $Q \subset \mathbb{R}^N$  be closed and let  $\theta(\cdot) : \partial Q \rightarrow (0, \infty]$ . We say that  $Q$  satisfies the  $\theta(\cdot)$ -external sphere condition if and only if for every  $x \in \partial Q$ , there exists a vector  $v_x \neq 0$  such that  $v_x \in N_Q^P(x)$  is realized by a ball of radius  $\theta(x)$ , i.e.,

$$\left\langle \frac{v_x}{\|v_x\|}, y - x \right\rangle \leq \frac{1}{2\theta(x)} \|y - x\|^2.$$

for all  $y \in Q$ .

Moreover, denote  $Q'$  by the closure of the complement of  $Q$ , we also say that the set  $Q$  satisfies the  $\theta(\cdot)$ -internal sphere condition if  $Q'$  satisfies the  $\theta(\cdot)$ -external sphere condition.

In general, a set which satisfies an  $\theta(\cdot)$ -external sphere condition doesn't have positive reach (see. e.g, [49]). However, under a wedgedness assumption these two concepts are equivalent in [49]. The wedgedness assumption was first introduced by Rockafellar in [54].

Let  $C \subset \mathbb{R}^N$  be a cone (i.e., if  $x \in C$  and  $\lambda \geq 0$ , then  $\lambda x \in C$ ). We say that  $C$  is *wedged* if  $C \cap (-C) = \{0\}$ . In [55, Corollary 18.7.1, p. 169] it is proved that

$$\begin{aligned} &\text{if } C \text{ is closed, convex, and wedged,} \\ &\text{then it is the closed convex hull of its exposed rays.} \end{aligned} \tag{2.2.3}$$

We recall (see [55, p.163]) that an exposed ray  $\mathbb{R}^+v$  of a convex cone  $C$  is defined by the property that there exists a linear functional  $h$  which is zero on it and is such that if  $h(p) = 0$  and  $p \in C$  then  $p \in \mathbb{R}^+v$ .

**Theorem 2.2.3** Let  $Q$  satisfy the  $\theta(\cdot)$ -external sphere condition. Assume that the Clarke normal cone  $N_Q^C(x)$  is wedged at every  $x \in \partial Q$  then  $Q$  has positive reach.

To end up this subsection, we expose here the relationships among functions whose hypograph satisfies an external sphere condition, the functions with positive reach hypograph and semiconcave functions (see in [50]).

**Theorem 2.2.4** Let  $f : \Omega \subset \mathbb{R}^N \rightarrow \mathbb{R}$  be Lipschitz. Then  $f$  is semiconcave if and only if the hypograph of  $f$  satisfies a  $\theta(\cdot)$ -external sphere condition.

**Proof.** The proof is based on Theorem 2.2.3 under remark that Clarke normal cones to  $\text{hypo}(f)$  at every point on  $\text{graph}(f)$  are wedged.  $\square$

## 2.3 Control theory

### 2.3.1 Control systems

We just consider here autonomous control systems, namely,  $f(x, u)$  does not depend on  $t$ ; this is done for the sake of simplicity since the results we present can be easily extended to the nonautonomous case. Standard references are in [11, 21, 13].

**Definition 2.3.1** *A control system consists of a pair  $(f, \mathcal{U})$ , where  $\mathcal{U} \subset \mathbb{R}^m$  is a closed set and  $f : \mathbb{R}^N \times \mathcal{U} \rightarrow \mathbb{R}^N$  is a continuous function. The set  $\mathcal{U}$  is called the control set, while  $f$  is called the dynamics of the system. The state equation associated with the system is*

$$\begin{cases} \dot{y}(t) = f(y(t), u(t)), & t \in [0, +\infty) \text{ a.e.} \\ u(\cdot) \in \mathcal{U}_{ad}, \\ y(0) = x, \end{cases} \quad (2.3.1)$$

where  $\mathcal{U}_{ad}$  the set of admissible controls, i.e., the measurable functions  $u : \mathbb{R} \rightarrow \mathbb{R}^m$ , such that  $u(t) \in \mathcal{U}$  a.e.

Two basic assumptions of the control systems are

(H1) The control set  $\mathcal{U}$  is nonempty and compact.

(H2) The function  $f$  satisfies:

$$\|f(y, u) - f(x, u)\| \leq L\|y - x\| \quad \forall x, y \in \mathbb{R}^N, \forall u \in \mathcal{U},$$

for a positive constant  $L$ .

Under the assumption (H2), for any  $u(\cdot) \in \mathcal{U}_{ad}$ , there is a unique Carathéodory solution of (2.3.1) denoted by  $y^{x, u(\cdot)}$ . The solution  $y^{x, u(\cdot)}$  is called the trajectory starting from  $x$  associated with the control  $u(\cdot)$ . The attainable set  $\mathcal{A}(T)$  from  $x$  in time  $T$  is thus defined by

$$\mathcal{A}^T(x) = \{y^{(u, x)}(t) \mid t \leq T, u(\cdot) \in \mathcal{U}_{ad}\} \quad (2.3.2)$$

Observe that assumption (H1) and (H2), together with the continuity of  $f$ , imply

$$\|f(x, u)\| \leq C + L\|x\|, \quad x \in \mathbb{R}^N, u \in \mathcal{U},$$

where  $C = \max\{\|f(0, u)\| \mid u \in \mathcal{U}\}$ . Therefore, the set  $\mathcal{A}^T(x)$  is bounded for all  $x \in \mathbb{R}^N$  and  $T < \infty$ . An upper bound of norm of points in the attainable set  $\mathcal{A}^T(x)$  can be found in the Appendix.

The set  $\mathcal{A}^T(x)$  is in general not closed. However, by standard results,

**Remark 2.3.1**  $\mathcal{A}^T(x)$  is compact if  $f(z, \mathcal{U})$  is convex for every  $z \in \mathbb{R}^N$ .

**Proof.** The proof is based on Filippov's Lemma and the compactness property for the trajectories for the control system (see Theorem 7.1.5 and Theorem 7.1.6 in [21]).  $\square$

### 2.3.2 Minimum time function

Together with the system 2.3.1, we consider a closed nonempty set  $\mathcal{S} \subset \mathbb{R}^N$ , which is called the target.

We first set  $T(x) = 0$  for all  $x \in \mathcal{S}$ . For a fixed  $x \in \mathbb{R}^N \setminus \mathcal{S}$ , we define

$$\theta(x, u) := \min \{t \geq 0 \mid y^{x,u}(t) \in \mathcal{S}\}.$$

Of course,  $\theta(x, u) \in (0, +\infty]$ , and  $\theta(x, u)$  is the time taken for the trajectory  $y^{x,u}(\cdot)$  to reach  $\mathcal{S}$ , provided  $\theta(x, u) < +\infty$ . The *minimum time*  $T(x)$  to reach  $\mathcal{S}$  from  $x$  is defined by

$$T(x) := \inf \{\theta(x, u) \mid u(\cdot) \in \mathcal{U}_{\text{ad}}\}. \quad (2.3.3)$$

Equivalently,

$$T(x) := \inf \{t \geq 0 \mid \mathcal{A}^t(x) \cap \mathcal{S} \neq \emptyset\}.$$

**Remark 2.3.2** *If  $f(z, \mathcal{U})$  is convex for every  $z \in \mathbb{R}^N$  then*

$$T(x) := \min \{\theta(x, u) \mid u(\cdot) \in \mathcal{U}_{\text{ad}}\}. \quad (2.3.4)$$

**Proof.** This is a consequence of Remark 2.3.1. □

A minimizing control in (2.3.4), say  $\bar{u}(\cdot)$ , is called an *optimal control*. The trajectory  $y^{x, \bar{u}}(\cdot)$  associated with  $\bar{u}(\cdot)$  is called an *optimal trajectory*.

We finally give in this subsection a result which one can see intuitively from the definition of the minimum time function:

**Theorem 2.3.1** (*Dynamic Programming Principle*) *Assume that the control system satisfies (H1) and (H2). For  $x \in \mathbb{R}^N \setminus \mathcal{S}$ , and for  $0 < t \leq T(x)$ , we have*

$$T(x) = t + \inf \{T(y) \mid y \in \mathcal{A}^t(x)\}. \quad (2.3.5)$$

*Equivalently, for all  $u(\cdot)$  if we set  $x(\cdot) = y^{x,u}(\cdot)$  then the function  $t \mapsto t + T(x(t))$  is increasing in  $[0, T(x)]$ .*

*Moreover, if  $x(\cdot)$  is an optimal trajectory then  $t \mapsto t + T(x(t))$  is constant in  $[0, T(x)]$ , i.e.,*

$$T(x(t)) = t - s + T(x(s)) \quad \text{for } 0 \leq s \leq t \leq T(x).$$

**Proof.** The proof can be found in several books, e.g., [11, 21, 13]. □

### 2.3.3 Controllability and continuity

Continuity properties of the minimal time function is a widely studied topic, mainly in connection with controllability. In this subsection, we will give shortly some definitions and some basic results which are concerned with



our works. For the references we prefer to quote [11, 21].

We first introduce the notations

$$\begin{aligned}\mathcal{R}(t) &= \{x \in \mathbb{R}^N \mid T(x) < t\}, \quad t > 0, \\ \mathcal{R} &= \bigcup_{t>0} \mathcal{R}(t) = \{x \in \mathbb{R}^N \mid T(x) < \infty\},\end{aligned}$$

where the letter  $\mathcal{R}$  stands for *reachable*:  $\mathcal{R}(t)$  is the set of points which can reach to the target  $\mathcal{S}$  with the control dynamics in time less than  $t$ , namely, for all  $x \in \mathcal{R}(t)$

$$\mathcal{A}^t(x) \cap \mathcal{S} \neq \emptyset.$$

The set  $\mathcal{R}$  is also called the *controllable set*.

**Definition 2.3.2** *The system  $(f, \mathcal{U})$  is small-time controllable on  $\mathcal{S}$  (briefly STCS) if  $\mathcal{S} \subseteq \text{int}\mathcal{R}(t)$  for all  $t > 0$ . If  $\mathcal{S} = \{0\}$  this property is called small-time local controllability (STLC).*

Note that STLC is equivalent to the continuity of  $T$  in 0 if  $\mathcal{S} = \{0\}$ , since  $T(0) = 0$  by definition. The next Proposition extends this observation to the general case.

**Proposition 2.3.1** *Assume that the control system  $(f, \mathcal{U})$  satisfies (H1), (H2) and the target  $\mathcal{S}$  is compact. Then the following statements are equivalent:*

- (i) *the system  $(f, \mathcal{U})$  is STCS;*
- (ii)  *$T$  is continuous in  $x$  for all  $x \in \partial\mathcal{S}$ ;*
- (iii) *there exists  $\delta > 0$  and  $\omega_T : [0, \delta] \rightarrow [0, \infty[$  such that  $\lim_{s \rightarrow 0} \omega_T(s) = 0$  and  $T(x) \leq \omega_T(d_{\mathcal{S}}(x))$  for all  $x \in B(\mathcal{S}, \delta) = \{z \in \mathbb{R}^N \mid d_{\mathcal{S}}(z) < \delta\}$ .*

**Proof.** The proof is in [11, Proposition 1.2, Chapter IV]. □

**Remark 2.3.3** *Under assumption in Proposition 2.3.1,  $T(x) > 0$  if and only if  $x \notin \mathcal{S}$ .*

Some consequence of STCS which is in [11, Chapter IV].

**Proposition 2.3.2** *Under assumptions in Proposition 2.3.1, if the system  $(f, \mathcal{U})$  is STCS then:*

- (i) *the controllable set  $\mathcal{R}$  is open;*
- (ii)  *$T$  is continuous in  $\mathcal{R}$ ;*
- (iii)  *$\lim_{x \rightarrow x_0} T(x) = +\infty$  for any  $x_0 \in \partial\mathcal{R}$ .*

We now introduce a special controllability condition which implies the Lipschitz continuity of the minimum time function.

**Definition 2.3.3** *We say that the control system  $(f, \mathcal{U})$  and the target  $\mathcal{S}$  satisfy the Petrov condition if, for any  $R > 0$ , there exists  $\mu > 0$  such that*

$$\min_{u \in \mathcal{U}} \langle f(x, u), \nu \rangle \leq -\mu \|\nu\|, \quad \forall x \in \partial \mathcal{S} \cap B(0, R), \nu \in N_{\mathcal{S}}^P(x). \quad (2.3.6)$$

**Theorem 2.3.2** *Under assumptions (H0), (H1) and the compactness of  $\mathcal{S}$ , if the control system  $(f, \mathcal{U})$  then:*

(i) *for any  $R > 0$ , there exist  $\delta, k > 0$  such that*

$$B(\mathcal{S}, \delta) \cap B(0, R) \subset \mathcal{R} \quad \text{and} \quad T(x) \leq kd_{\mathcal{K}}(x), \quad x \in \mathcal{S}, \delta) \cap B(0, R);$$

(ii) *the minimum time function  $T$  is locally Lipschitz continuous on  $\mathcal{R}$ .*

**Proof.** Standard reference for the proof is in [21, Chapter VIII].  $\square$

We finally give in the subsection a theorem in which the minimum time function is just continuous under the weak Petrov condition.

**Theorem 2.3.3** *Under the assumptions (H1), (H2) and the compactness of the target  $\mathcal{S}$ , if the control system satisfies the weak Petrov condition, i.e., there exist  $\delta > 0$  and a continuous nondecreasing function  $\mu : [0, \infty) \rightarrow [0, +\infty)$  with the properties*

(a)  $\mu(0) = 0$ ,  $\mu(\rho) > 0$  for  $\rho > 0$  and  $\int_0^\delta \frac{d\rho}{\mu(\rho)} d\rho < +\infty$ ;

(b) *for all  $x \in B(\mathcal{S}, \delta) \setminus \mathcal{S}$ , there exists  $\bar{s} \in \pi_{\mathcal{S}}(x)$  such that*

$$\min_{u \in \mathcal{U}} \langle f(x, u), x - \bar{s} \rangle \leq -\mu(d_{\mathcal{S}}(x))d_{\mathcal{S}}(x).$$

*Then the control system  $(f, \mathcal{S})$  is SCTS and  $T$  is continuous in  $\mathcal{R}$ .*

**Proof.** There are various versions, obtained with different methods, which can be found, e.g., in [20, 42, 45, 51, 60].  $\square$

## 2.4 Differential inclusions

We shall give here some basic definitions and theorems which are needed in Subsection 3.7. For the reference, we refer to [9].

Consider differential inclusions of the form

$$\begin{cases} \dot{x}(t) \in F(x(t)), \\ x(0) = x_0, \end{cases} \quad (2.4.1)$$

where  $F : \mathbb{R}^N \rightrightarrows \mathbb{R}^N$  is a multifunction, and  $x_0 \in \mathbb{R}^N$  is the starting point.

**Definition 2.4.1** Let  $y^{x_0}(\cdot) : (a, b) \rightarrow \mathbb{R}^N$  where  $a < 0 < b$  be such that  $y^{x_0}(0) = x_0$ . We say that  $y^{x_0}(\cdot)$  is a solution of (2.4.1) if  $y^{x_0}(\cdot)$  is absolutely continuous and

$$\dot{y}^{x_0}(t) \in F(y^{x_0}(t)) \quad \text{a.e. } t \in (a, b).$$

The solution  $y^{x_0}(\cdot)$  is usually called a trajectory starting from  $x_0$  and associated with (2.4.1).

There are several results about the existence of solution to the differential inclusion (2.4.1). For the reference, we refer to Chapter 2 and Chapter 3 in [9].

The attainable  $\mathcal{A}^T(x_0)$  from  $x_0$  in time  $T$  is now denoted by

$$\mathcal{A}^T(x_0) = \{y^{x_0}(t) \mid 0 \leq t \leq T\}.$$

We will end this subsection with a theorem which gives the existence of a minimal time  $T$ . In the following statement, we prefer to consider assumptions which will be used in Section 3.7.

The multifunction  $F : \mathbb{R}^N \rightrightarrows \mathbb{R}^N$  is Lipschitz with respect to the Hausdorff distance if there exists a constant  $M > 0$  such that

$$d_{\mathcal{H}}(F(x), F(y)) \leq M\|y - x\|.$$

**Theorem 2.4.1** Assume that  $F$  is Lipschitz with respect to the Hausdorff distance and  $F(x)$  is nonempty, convex, and compact for each  $x \in \mathbb{R}^n$ . If there exists a constant  $M_2 > 0$  so that  $\max\{\|v\| \mid v \in F(x)\} \leq M_2(1 + \|x\|)$  then for all  $T > 0$  the attainable set  $\mathcal{A}^T(x_0)$  is compact.

**Proof.** One can find an original version and the proof in Section 2, Chapter 2 [9].  $\square$



## Part I

# On the structure of the minimum time function



## Chapter 3

# Semiconcavity type results

We will first study in this Chapter a minimum time problem with a nonlinear smooth dynamics and a target satisfying an internal sphere condition. Under the assumptions that the minimum time  $T$  be continuous and the proximal normal cone to the hypograph of  $T$  be wedged, we show that  $\text{hypo}(T)$  has positive reach. Consequently,  $T$  satisfies the list of properties in Theorem (2.2.2). In particular,  $T$  is a.e. twice differentiable.

The result is based on an analysis of how proximal normals (to the complement of the target) are transported by the adjoint flow, which in turn permits a representation of the generalized gradient of  $T$  in terms of suitable adjoint vectors (Theorems 3.2.1 and 3.2.2). Here the wedgedness assumption plays a major role: actually *exposed rays* of the normal cone to the hypograph are special normals, as they can be approximated by normals at differentiability points of  $T$  (Lemma 3.3.7). Moreover, wedgedness is used in Theorem 3.2.3 in order to obtain a uniform estimate for radii of the balls realizing proximal normals to the hypograph. We show also through an example (Example 2 in Section 3.6) that if the normal cone is not wedged, then Theorem 3.2.3 may fail. However, an external sphere condition to the hypograph of  $T$  still holds (see Proposition 3.2.1). An analysis of this general case will be discussed in Chapter 5 and Chapter 6, where topological and measure theoretic results on the set where the normal cone is not wedged are given.

Moreover, on the basis in Chapter 5 and Chapter 6, we also study the minimum time function for a class of differential inclusions. For such class, under an internal sphere condition on the target  $\mathcal{S}$  the hypograph of  $T$  still satisfies an external sphere condition. The proof will be based on the Hamiltonian function and Pontryagin's maximum principle. At the end of this chapter, we will partially extend our result to an arbitrary target  $\mathcal{S}$ .

The chapter is structured as follows: Section 3.1 is devoted to some notations, while Section 3.2 contains assumptions and statement of the main results. Detailed arguments begin in Section 3.3, which contains several

lemmas whose geometrical meaning is illustrated, and ends with a result (Theorem 3.3.1) giving a representation of the normal cone to the hypograph of  $(T)$ , under the wedgedness assumption. Section 3.4 is devoted to the conclusion of the proof of the main theorems, which is now only a simple use of the lemmas contained in Section 3.4. Section 3.5 is dedicated to an improvement of Theorems 3.2.1 and 3.2.2 for an *optimal point*, i.e. a point which is crossed by a time-optimal trajectory and Section 3.6 contains examples. Finally, our results will be extended for a class of differential inclusion in Section 3.7.

### 3.1 Nonlinear control system

We consider throughout the chapter a nonlinear control system of the form

$$\begin{cases} \dot{y}(t) = f(y(t), u(t)) & a.e. \\ u(t) \in \mathcal{U} & a.e. \\ y(0) = x, \end{cases} \quad (3.1.1)$$

where the Lipschitz function  $f : \mathbb{R}^N \times \mathcal{U} \rightarrow \mathbb{R}^N$  and the control set  $\mathcal{U}$ , a compact nonempty subset of  $\mathbb{R}^m$ , are given. We recall that  $\mathcal{U}_{\text{ad}}$  the set of admissible controls, i.e., the measurable functions  $u : \mathbb{R} \rightarrow \mathbb{R}^m$ , such that  $u(t) \in \mathcal{U}$  a.e. For any  $u(\cdot) \in \mathcal{U}_{\text{ad}}$ , the unique Carathéodory solution of (3.1.1) is denoted by  $y^{x,u}(\cdot)$ .

The adjoint vectors associated with a trajectory  $y^{x,u}(\cdot)$  can be represented using the fundamental solution matrix  $M(\cdot, x, u)$  of the linear equation

$$\dot{p}(t) = D_x f(y^{x,u}(t), u(t)) p(t), \quad p(0) = \mathbb{I}^{N \times N}. \quad (3.1.2)$$

We also define  $M^{-1}(\cdot, x, u)$  to be the fundamental solution matrix of the time reversed adjoint equation

$$\dot{q}(t) = -q(t) D_x f(y^{x,u}(t), u(t)), \quad q(0) = \mathbb{I}^{N \times N}. \quad (3.1.3)$$

Suppose we are now given a closed nonempty set  $\mathcal{S} \subset \mathbb{R}^N$ , which is called the target. For a fixed  $x \in \mathbb{R}^N \setminus \mathcal{S}$ , we define

$$\theta(x, u) := \min \{t \geq 0 \mid y^{x,u}(t) \in \mathcal{S}\}.$$

Of course,  $\theta(x, u) \in (0, +\infty]$ , and  $\theta(x, u)$  is the time taken for the trajectory  $y^{x,u}(\cdot)$  to reach  $\mathcal{S}$ , provided  $\theta(x, u) < +\infty$ . The *minimum time*  $T(x)$  to reach  $\mathcal{S}$  from  $x$  is defined by

$$T(x) := \inf \{\theta(x, u) \mid u(\cdot) \in \mathcal{U}_{\text{ad}}\}. \quad (3.1.4)$$

In general, an *optimal trajectory*, i.e., a trajectory which attains the infimum in (3.1.4) does not exist. Therefore, we need also to consider *minimizing*



sequences and limiting optimal trajectories steering  $x$  to the target  $\mathcal{S}$ . In particular, we will consider the limits of end-points (thus belonging to  $\mathcal{S}$ ) of minimizing sequences of trajectories. More precisely,

$$\begin{aligned} \mathcal{S}_x &= \{ \bar{x} \in \mathcal{S} \mid \text{there exist sequences } \{x_n\} \subset \mathcal{S}^c \text{ and } \{\bar{u}_n(\cdot)\} \subset \mathcal{U}_{\text{ad}} \\ &\quad \text{such that } x_n \rightarrow x, \theta(x_n, \bar{u}_n) \rightarrow T(x), y^{x_n, \bar{u}_n}(\theta(x_n, \bar{u}_n)) \rightarrow \bar{x} \}. \end{aligned}$$

Observe that if  $T(x) < +\infty$ , then  $\emptyset \neq \mathcal{S}_x \subseteq \text{bdry}\mathcal{S}$ .

For any  $\bar{x} \in \mathcal{S}_x$  we define also

$$\begin{aligned} \mathcal{U}_{\bar{x}} &= \{ \{\bar{u}_n(\cdot)\} \subset \mathcal{U}_{\text{ad}} \mid \text{there exists a sequence } \{x_n\} \text{ satisfying} \\ &\quad x_n \rightarrow x, \theta(x_n, \bar{u}_n) \rightarrow T(x), \text{ and } y^{x_n, \bar{u}_n}(\theta(x_n, \bar{u}_n)) \rightarrow \bar{x} \}, \end{aligned}$$

i. e., the set of *minimizing sequences of controls* steering  $x$  to  $\bar{x}$ . Together with  $\mathcal{U}_{\bar{x}}$  we define also

$$\begin{aligned} \mathcal{T}_{\bar{x}} &= \{ \{y^{x_n, \bar{u}_n}(\cdot)\} \mid x_n \rightarrow x, \bar{u}_n \in \mathcal{U}_{\text{ad}}, \\ &\quad \theta(x_n, \bar{u}_n) \rightarrow T(x), \text{ and } y^{x_n, \bar{u}_n}(\theta(x_n, \bar{u}_n)) \rightarrow \bar{x} \}, \end{aligned}$$

i. e., the set of trajectories corresponding to minimizing sequences of controls steering  $x$  to  $\bar{x}$ .

Correspondingly, the *limiting adjoint trajectories* related to *minimizing sequences of controls* are defined by the following

$$\begin{aligned} \mathcal{M}_{\bar{x}} &= \{ M : [0, T(x)] \rightarrow \mathbb{M}^{N \times N} \mid \exists \{y^{x_n, \bar{u}_n}(\cdot)\} \subset \mathcal{T}_{\bar{x}} \text{ such that} \\ &\quad M(\cdot) \text{ is uniform limit on } [0, T(x)] \text{ of } M(\cdot, x_n, \bar{u}_n) \}. \end{aligned} \quad (3.1.5)$$

**Remark 3.1.1** *If  $T(\cdot)$  is everywhere finite, both  $\mathcal{S}_x, \mathcal{T}_{\bar{x}}$  are nonempty. By compactness,  $\mathcal{M}_{\bar{x}}$  is nonempty as well for all  $\bar{x} \in \mathcal{S}_x$ . Moreover, if  $F(x) := \{f(x, u) \mid u \in \mathcal{U}\}$  is convex for all  $x$ , then the infimum is attained and the sets  $\mathcal{S}_x, \mathcal{U}_{\bar{x}}$ , and  $\mathcal{T}_{\bar{x}}$  can be substituted by the simpler sets*

$$\begin{aligned} \mathcal{S}_x &= \{ \bar{x} \in \mathcal{S} \mid \text{there exists } \bar{u} \in \mathcal{U}_{\text{ad}} \text{ such that} \\ &\quad \theta(x, \bar{u}) = T(x), \bar{x} = y^{x, \bar{u}}(T(x)) \} \\ \mathcal{U}_{\bar{x}} &= \{ \bar{u} \in \mathcal{U}_{\text{ad}} \mid \theta(x, \bar{u}) = T(x), y^{x, \bar{u}}(T(x)) = \bar{x} \} \\ \mathcal{T}_{\bar{x}} &= \{ y^{x, \bar{u}} \mid \bar{u} \in \mathcal{U}_{\bar{x}} \}. \end{aligned}$$

Finally, the *Maximized Hamiltonian*, namely the function

$$H : \mathbb{R}^N \times \mathbb{R}^N \longrightarrow \mathbb{R}, \quad H(x, p) = \max_{u \in \mathcal{U}} \langle f(x, u), p \rangle,$$

will be important in our analysis.

### 3.2 Statement of the main results

We repeat first the setting we are concerned with and specify our assumptions.

We consider the nonlinear system (3.1.1) under the following assumptions:

(H1)  $\mathcal{U} \subset \mathbb{R}^N$  is compact.

(H2)  $f : \mathbb{R}^N \times \mathcal{U} \rightarrow \mathbb{R}^N$  is continuous and satisfies:

$$\|f(x, u) - f(y, u)\| \leq L \|x - y\| \quad \forall x, y \in \mathbb{R}^N, u \in \mathcal{U},$$

for a positive constant  $L$ . Moreover, the differential of  $f$  with respect to the  $x$  variable,  $D_x f$ , exists everywhere, is continuous with respect to both  $x$  and  $u$  and satisfies the following Lipschitz condition:

$$\|D_x f(x, u) - D_x f(y, u)\| \leq L_1 \|x - y\| \quad \forall x, y \in \mathbb{R}^N, u \in \mathcal{U},$$

for a positive constant  $L_1$ .

(H3) The minimum time function  $T : \mathbb{R}^N \rightarrow [0, +\infty)$  is everywhere finite and continuous, (i.e. controllability and small time controllability hold).

(H4) The target  $\mathcal{S}$  is nonempty, closed, and satisfies the internal sphere condition of radius  $\rho > 0$ .

**Remark 3.2.1** *Conditions ensuring small time controllability when the target is not necessarily a singleton can be found in the previous chapter.*

Our analysis will be based on the transportation of certain vectors, normal to the closure of the complement of the target  $\mathcal{S}$ , by means of the (limiting) adjoint flow. More precisely, two sets of transported normals will be considered, according with the Hamiltonian:

$$N_0(x) = \{M^T(r)v \mid M(\cdot) \in \mathcal{M}_{\bar{x}}, v \in N_{\mathcal{S}^c}^P(\bar{x}), \bar{x} \in \mathcal{S}_x \text{ and } H(M^T(r)v, x) = 0\}$$

$$N_1(x) = \{M^T(r)v \mid M(\cdot) \in \mathcal{M}_{\bar{x}}, v \in N_{\mathcal{S}^c}^P(\bar{x}), \bar{x} \in \mathcal{S}_x \text{ and } H(M^T(r)v, x) = 1\}$$

Our main results are the following three theorems, together with the corollary.

**Theorem 3.2.1** *Let  $x \in \mathcal{S}^c$  and  $r = T(x)$ . Under the conditions (H1), (H2), (H3), and (H4), together with the further assumption*

$$N_{\text{hypo}(T)}^P(x, T(x)) \text{ is wedged,} \tag{3.2.1}$$

*the (proximal) horizontal supergradient of the minimum time function  $T(\cdot)$  at the point  $x$  can be computed as follows:*

$$\partial^\infty T(x) = -\text{co}(N_0(x)). \tag{3.2.2}$$

**Theorem 3.2.2** *Let  $x \in \mathcal{S}^c$  and  $r = T(x)$ . Under the same assumptions of Theorem 3.2.1, the proximal supergradient of the minimum time function at the point  $x$  can be computed as follows:*

$$\partial^P T(x) = -[\text{co}(N_1(x)) + \text{co}(N_0(x))]. \quad (3.2.3)$$

**Theorem 3.2.3** *Let the assumptions of Theorem 3.2.1 hold for all  $x \in \mathcal{S}^c$ . Then for every closed set  $\mathcal{S}' \subset \mathcal{S}^c$ ,  $\text{hypo}(T) \cap (\mathcal{S}' \times \mathbb{R})$  has positive reach.*

**Corollary 3.2.1** *Let the assumptions of Theorem 3.2.1 hold. Then the minimum time function  $T$  satisfies all the properties listed in Theorem 2.2.2.*

The last result is concerned with the case where the wedgedness assumption (3.2.1) does not hold. We will present here, for the sake of brevity, only a partial result together with two examples, a thorough analysis being postponed to the next chapter.

**Proposition 3.2.1** *Let the assumptions (H1), (H2), (H3), and (H4) hold. Then the hypograph of the minimum time function  $T$  satisfies the external sphere condition with a locally uniform radius, namely for every  $x \in \mathcal{S}^c$  there exists a unit proximal normal  $v$  to  $\text{hypo}(T)$  at  $(x, T(x))$  which is realized by a sphere with a locally constant radius  $\sigma > 0$ .*

The proof of the previous Proposition is a straightforward consequence of Lemmas 3.3.2 and 3.3.3.

**Remark 3.2.2** *The constant  $\sigma$  can be explicitly computed, and depend only on  $x$ , on  $f$  and  $\mathcal{U}$ , and on the constants  $L$ ,  $L_1$  and  $\rho$  appearing in the assumptions (H2) and (H4).*

### 3.3 Preparatory Lemmas

This section is devoted to several partial results which are needed to prove Theorem 3.2.2 and Theorem 3.2.2. In particular, the proof of “ $\supseteq$ ” inclusions in (3.2.1) and (3.2.2) will be based on Lemma 3.3.2 and Lemma 3.3.3 below.

#### 3.3.1 Transporting proximal normals

In this subsection we do not assume that  $\mathcal{S}$  satisfies the internal sphere condition, nor that the normal cone to the hypograph of  $T(\cdot)$  at  $(x, T(x))$  is wedged.

The following notation for sublevels of the minimum time function will be used: for  $r > 0$  we set

$$\begin{aligned} \mathcal{S}(r) &:= \{x \in \mathbb{R}^N \mid T(x) < r\} \\ \mathcal{S}^c(r) &:= \{x \in \mathbb{R}^N \mid T(x) \geq r\} \end{aligned}$$

We state first a technical lemma, showing that the limiting adjoint flow transports proximal normals to the complement of the target to proximal normals to the complement of sublevels of  $T$ . Moreover, the radius of the ball which realizes the transported normal can be explicitly estimated.

**Lemma 3.3.1** *Assume that  $\mathcal{S}$  is closed and let the assumptions (H1), (H2), and (H3) hold. Let  $x \in \mathcal{S}^c$  and set  $r = T(x) > 0$ . Fix  $\bar{x} \in \mathcal{S}_x$ ,  $v \in N_{\overline{\mathcal{S}^c}}^P(\bar{x})$  and  $M(\cdot) \in \mathcal{M}_{\bar{x}}$ . Then*

$$M^T(r)v \in N_{\mathcal{S}^c(r)}^P(x).$$

*More precisely, assume that  $v$  is realized by a ball of radius  $\rho > 0$ . Then there exists an explicitly computable continuous function  $K$  depending only on  $r$ ,  $\|x\|$ ,  $\rho$  such that for all  $z \in \mathcal{S}^c(r)$  we have*

$$\langle M^T(r)v, z - x \rangle \leq K(r, \|x\|, \rho) \|M^T(r)v\| \|z - x\|^2. \quad (3.3.1)$$

**Proof.** Let  $x_n \rightarrow x$ ,  $\bar{x} \in \mathcal{S}_x$ , and  $\{\bar{u}_n\} \subset \mathcal{U}_{\text{ad}}$  be such that  $\{y^{x_n, \bar{u}_n}(\cdot)\} \in \mathcal{T}_{\bar{x}}$  and  $M(\cdot, x_n, \bar{u}_n)$  converges to  $M(\cdot)$  uniformly on  $[0, T(x)]$ . By definition of proximal normal realized by a  $\rho$ -ball,

$$\langle v, \bar{z} - \bar{x} \rangle \leq \frac{\|v\|}{2\rho} \|\bar{z} - \bar{x}\|^2 \quad \text{for all } \bar{z} \in \overline{\mathcal{S}^c}.$$

Fix  $z \in \mathcal{S}^c(r)$ . We define

$$\bar{x}_n = y^{x_n, \bar{u}_n}(\theta(x_n, \bar{u}_n)), \quad \bar{z}_n = y^{z, \bar{u}_n}(\theta(x_n, \bar{u}_n)),$$

and observe that  $\bar{x}_n \in \mathcal{S}$ ,  $\bar{x}_n \rightarrow \bar{x}$  and we can assume without loss of generality that  $\bar{z}_n$  converges to a point  $\bar{z}$  which belongs to  $\overline{\mathcal{S}^c}$  since  $\theta(x_n, \bar{u}_n) \rightarrow r \leq T(z)$ .

We set for simplicity  $\alpha_n(\cdot) = y^{x_n, \bar{u}_n}(\cdot)$ ,  $\beta_n(\cdot) = y^{z, \bar{u}_n}(\cdot)$ ,  $t_n = \theta(x_n, \bar{u}_n)$ , so that

$$\bar{x}_n = x_n + \int_0^{t_n} f(\alpha_n(s), \bar{u}_n(s)) ds, \quad \bar{z}_n = z + \int_0^{t_n} f(\beta_n(s), \bar{u}_n(s)) ds,$$

whence

$$\begin{aligned} \bar{z}_n - \bar{x}_n &= z - x_n + \\ &\int_0^{t_n} \left( \int_0^1 D_x f(\alpha_n(s) + \tau(\beta_n(s) - \alpha_n(s)), \bar{u}_n(s)) d\tau \right) (\beta_n(s) - \alpha_n(s)) ds. \end{aligned}$$

We define now

$$\begin{aligned} A_n^1(s) &= D_x f(\alpha_n(s), \bar{u}_n(s)), \\ A_n^2(s) &= \int_0^1 D_x f(\alpha_n(s) + \tau(\beta_n(s) - \alpha_n(s)), \bar{u}_n(s)) d\tau, \end{aligned}$$

and observe that, thanks to (H2), for all  $s \in [0, t_n]$  we have

$$\|A_n^2(s) - A_n^1(s)\| \leq \frac{L_1}{2} \|\beta_n(s) - \alpha_n(s)\|. \quad (3.3.2)$$

Using (iv) in Lemma 7.1.1 and the definition of  $L_2$  in (7.1.1), we obtain

$$\|A_n^1(s)\| \leq L_2(s, \|x_n\|) \quad (3.3.3)$$

for all  $s \in [0, t_n]$ . Thus

$$\|A_n^2(s)\| \leq L_2(s, \|x_n\|) + \frac{L_1}{2} \|\beta_n(s) - \alpha_n(s)\| \quad (3.3.4)$$

for all  $s \in [0, t_n]$ . Now, Gronwall's Lemma yields

$$\|\beta_n(s) - \alpha_n(s)\| \leq e^{Ls} \|z - x_n\|, \quad (3.3.5)$$

so that combining (3.3.4) and (3.3.5) we obtain

$$\|A_n^2(s)\| \leq L_2(s, \|x_n\|) + \frac{L_1}{2} e^{Ls} \|z - x_n\|. \quad (3.3.6)$$

Define  $M_n^2(\cdot)$  to be the solution of the problem

$$\dot{p}(s) = A_n^2(t)p(s), \quad p(0) = \mathbb{I}^{N \times N}.$$

Recalling that  $M(\cdot, x, u)$  is the fundamental solution of (3.1.2) set  $M_n^1(\cdot) = M(\cdot, x_n, \bar{u}_n)$ ,  $z_n^1(s) = M_n^1(s)(z - x_n)$  and  $z_n^2(s) = M_n^2(s)(z - x_n)$ , for all  $s \in [0, t_n]$ . Using these notations, we can write

$$\begin{aligned} \langle v, \bar{z}_n - \bar{x}_n \rangle &= \langle v, z_n^2(t_n) \rangle \\ &= \langle v, z_n^1(t_n) \rangle + \langle v, z_n^2(t_n) - z_n^1(t_n) \rangle \\ &= \langle v, M_n^1(t_n)(z - x_n) \rangle + \langle v, (M_n^2(t_n) - M_n^1(t_n))(z - x_n) \rangle \\ &\geq \langle v, M_n^1(t_n)(z - x_n) \rangle - \|v\| \|(M_n^2(t_n) - M_n^1(t_n))(z - x_n)\|. \end{aligned} \quad (3.3.7)$$

To simplify our writing, we set, for all  $s \geq 0$  and  $y, z \in \mathbb{R}^N$ ,  $L_3(s, y, z) = \frac{L_1}{2} e^{Ls} \|z - y\|$ . By (3.3.3), (3.3.6), Lemma 7.1.3, and (3.3.2), we have

$$\begin{aligned} &\|(M_n^2(t_n) - M_n^1(t_n))(z - x_n)\| \\ &\leq e^{[2L_2(t_n, \|x_n\|) + L_3(t_n, x_n, z)]t_n} \int_0^{t_n} \|A_n^2(s) - A_n^1(s)\| ds \|z - x_n\| \\ &\leq \frac{L_1}{2} e^{[2L_2(t_n, \|x_n\|) + L_3(t_n, x_n, z)]t_n} \int_0^{t_n} \|\beta_n(s) - \alpha_n(s)\| ds \|z - x_n\|. \end{aligned}$$

Recalling (3.3.5), we obtain

$$\begin{aligned} & \| (M_n^2(t_n) - M_n^1(t_n))(z - x_n) \| \\ & \leq \frac{L_1}{2} e^{[2L_2(t_n, \|x_n\|) + L_3(t_n, x_n, z) + L]t_n} \|z - x_n\|^2. \end{aligned} \quad (3.3.8)$$

Therefore, by passing to the limit in (3.3.7) and (3.3.8) (recall that  $M_n^1(\cdot) \rightarrow M(\cdot)$  uniformly), we have

$$\begin{aligned} & \langle M^T(r)v, z - x \rangle \\ & \leq \langle v, \bar{z} - \bar{x} \rangle + \|v\| \frac{L_1}{2} e^{[2L_2(r, \|x\|) + L_3(r, x, z) + L]r} \|z - x\|^2 \\ & \leq \frac{\|v\|}{2\rho} \|\bar{z} - \bar{x}\|^2 + \|v\| \frac{L_1}{2} e^{[2L_2(r, \|x\|) + L_3(r, x, z) + L]r} \|z - x\|^2. \end{aligned}$$

Moreover, from (3.3.5) we have  $\|\bar{z} - \bar{x}\| \leq e^{Lr} \|z - x\|$ . Therefore,

$$\langle M^T(r)v, z - x \rangle \leq \left( \frac{L_1}{2} e^{[2L_2(r, \|x\|) + L_3(r, x, z) + L]r} + \frac{e^{2Lr}}{2\rho} \right) \|v\| \|z - x\|^2 \quad (3.3.9)$$

for all  $z \in \mathcal{S}^c(r)$ .

Observe that

$$\begin{aligned} \|v\| & = \|(M^T(r))^{-1}M^T(r)v\| \\ & \leq \|M(r)^{-1}\| \|M^T(r)v\|. \end{aligned}$$

By (ii) in Lemma 7.1.2 we obtain

$$\|M(r)^{-1}\| \leq e^{L_2(r, \|x\|)r}.$$

Combining the above inequalities with (3.3.9) we thus have

$$\begin{aligned} & \langle M^T(r)v, z - x \rangle \\ & \leq \left( \frac{L_1}{2} e^{[3L_2(r, \|x\|) + L_3(r, x, z) + L]r} + \frac{e^{2Lr + L_2(r, \|x\|)}}{2\rho} \right) \|M^T(r)v\| \|z - x\|^2. \end{aligned} \quad (3.3.10)$$

In order to complete the proof, we consider two cases.

If  $\|z - x\| < 1$ , then  $L_3(r, x, z) \leq \frac{L_1}{2} e^{Lr}$ . Thus, by (3.3.10) we have

$$\begin{aligned} & \langle M^T(r)v, z - x \rangle \\ & \leq \left( \frac{L_1}{2} e^{[3L_2(r, \|x\|) + \frac{L_1}{2} e^{Lr} + L]r} + \frac{e^{2Lr + L_2(r, \|x\|)}}{2\rho} \right) \|M^T(r)v\| \|z - x\|^2. \end{aligned} \quad (3.3.11)$$

If instead  $\|z - x\| \geq 1$ , then  $\langle M^T(r)v, z - x \rangle \leq \|M^T(r)v\| \|z - x\|^2$ .  
Therefore, in both cases we have that

$$\langle M^T(r)v, z - x \rangle \leq K(r, \|x\|, \rho) \|M^T(r)v\| \|z - x\|^2 \quad \text{for all } z \in \mathcal{S}^c(r), \quad (3.3.12)$$

where the continuous function  $K$ , defined for  $r, \delta \geq 0$  and  $\rho > 0$  as

$$K(r, \delta, \rho) := \max \left\{ 1, \frac{L_1}{2} e^{[3L_2(r, \delta) + \frac{L_1}{2} e^{Lr} + L]r} + \frac{e^{2Lr + L_2(r, \delta)}}{2\rho} \right\}, \quad (3.3.13)$$

depends only on the variables  $r, \delta, \rho$  and on the constants  $L, L_1, K_1, K_2$ .  
The proof is complete.  $\square$

**Remark 3.3.1** *It follows from (3.3.13) that  $K(r, \delta, \rho)$  is nondecreasing with respect to both  $r$  and  $\delta$ .*

The next lemma establishes that normals transported along the *limiting adjoint flow* generate horizontal proximal normals to the hypograph of  $T(\cdot)$ , provided their Hamiltonian is zero. Moreover, the radius of the ball realizing them can be explicitly estimated.

**Lemma 3.3.2** *Let  $\mathcal{S}$  be closed and let the assumptions (H1), (H2), and (H3) hold. Let  $x \in \mathcal{S}^c$ , set  $r := T(x) > 0$ , and let  $\xi \in N_0(x)$ . Then  $-\xi \in \partial^\infty T(x)$ , or, equivalently,  $(\xi, 0) \in N_{\text{hypo}(T(x))}^P(x, T(x))$ .*

*More precisely, let  $\bar{x} \in \mathcal{S}_x$  and let  $v \in N_{\overline{\mathcal{S}^c}}^P(\bar{x})$ ,  $M(\cdot) \in \mathcal{M}_{\bar{x}}$  be such that  $H(M^T(r)v, x) = 0$ . Assume that  $v$  is realized by a ball of radius  $\rho$ . Then there exists an explicitly computable continuous function  $K_3(r, x, \rho)$ , depending only on  $r, x, \rho$ , such that for all  $z \in \overline{\mathcal{S}^c}$  and all  $\beta \leq T(z)$  we have*

$$\langle M^T(r)v, z - x \rangle \leq K_3(r, x, \rho) \|M^T(r)v\| \left( \|z - x\|^2 + |\beta - T(x)|^2 \right). \quad (3.3.14)$$

**Proof.** Let  $v \in N_{\overline{\mathcal{S}^c}}^P(\bar{x})$  be such that

$$\langle v, \bar{z} - \bar{x} \rangle \leq \frac{\|v\|}{2\rho} \|\bar{z} - \bar{x}\|^2 \quad \forall \bar{z} \in \overline{\mathcal{S}^c}. \quad (3.3.15)$$

Recalling Lemma 3.3.1, for all  $z \in \mathcal{S}^c(r)$  we have

$$\langle M^T(r)v, z - x \rangle \leq K(r, \|x\|, \rho) \|M^T(r)v\| \|z - x\|^2. \quad (3.3.16)$$

Let  $z \in \overline{\mathcal{S}^c}$ . Two cases may occur:

- (i)  $T(z) \geq T(x)$ ,
- (ii)  $T(z) < T(x)$ .

In the first case, (3.3.14) follows immediately from (3.3.16).

In the second case, define  $r_1 = T(z)$  and take sequences  $\{x_n\}$ , with  $x_n \rightarrow x$ ,

$\{\bar{u}_n\} \subset \mathcal{U}_{\text{ad}}$  and  $\{\alpha_n(\cdot) := y^{x_n, \bar{u}_n}(\cdot)\}$  corresponding to  $M(\cdot)$ , according to the definition given in (3.1.5). For all  $n$  large enough there exists  $r_n^1 < r$  for which

$$\bar{x}_n^1 := \alpha_n(r - r_n^1) = x_n + \int_0^{r-r_n^1} f(\alpha_n(s), \bar{u}_n(s)) ds$$

is such that  $T(\bar{x}_n^1) = r_1$ . We can assume without loss of generality that  $\alpha_n(\cdot)$  converges uniformly to some  $\alpha(\cdot)$  and that  $r_n^1 \rightarrow \bar{r}_1$ . Observe that  $\bar{r}_1 < r$ . Setting  $\bar{x}^1 = \alpha(r - \bar{r}_1) (= \lim \bar{x}_n^1)$ , one can easily see that  $T(\bar{x}^1) = r_1$  by the continuity of  $T(x)$ . Then, by Lemma 3.3.1 we obtain that

$$\langle M^T(r_1)v, z - \bar{x}^1 \rangle \leq K(r_1, \|\bar{x}^1\|, \rho) \|M^T(r_1)v\| \|z - \bar{x}^1\|^2. \quad (3.3.17)$$

We write

$$\langle M^T(r)v, z - x \rangle = \langle M^T(r)v, z - \bar{x}^1 \rangle + \langle M^T(r)v, \bar{x}^1 - x \rangle$$

and perform some estimates.

First, we consider

$$\langle M^T(r)v, z - \bar{x}^1 \rangle = \langle M^T(r_1)v, z - \bar{x}^1 \rangle + \langle (M^T(r) - M^T(r_1))v, z - \bar{x}^1 \rangle.$$

By (3.3.17) we have

$$\begin{aligned} \langle M^T(r)v, z - \bar{x}^1 \rangle &\leq K(r_1, \|\bar{x}^1\|, \rho) \|M^T(r_1)v\| \|z - \bar{x}^1\|^2 \\ &\quad + \|(M^T(r) - M^T(r_1))v\| \|z - \bar{x}^1\|. \end{aligned}$$

Moreover from (ii) in Lemma 7.1.2 we have

$$\begin{aligned} \|M^T(r_1)v\| &\leq \|(M^T(r - r_1))^{-1}\| \|M^T(r)v\| \\ &\leq e^{L_2(r-r_1, \|x\|)(r-r_1)} \|M^T(r)v\| \\ &\leq e^{L_2(r, \|x\|)r} \|M^T(r)v\|. \end{aligned}$$

Also, using (iv) in Lemma 7.1.1 we obtain

$$\begin{aligned} \|(M^T(r) - M^T(r_1))v\| &\leq \int_{r_1}^r \|\dot{M}^T(s)v\| ds \\ &\leq \int_{r_1}^r e^{L_2(r, \|x\|)r} \|M^T(r)v\| ds \\ &= e^{L_2(r, \|x\|)r} \|M^T(r)v\| |r - r_1|. \end{aligned}$$

Therefore,

$$\begin{aligned} \langle M^T(r)v, z - \bar{x}^1 \rangle &\leq K(r_1, \|\bar{x}^1\|, \rho) e^{L_2(r, \|x\|)r} \|M^T(r)v\| \|z - \bar{x}^1\|^2 \\ &\quad + e^{L_2(r, \|x\|)r} \|(M^T(r)v)\| |r - r_1| \|z - \bar{x}^1\|. \end{aligned}$$



Recalling (i) in Lemma 7.1.1 for  $\alpha(\cdot) = y^{x_n, \bar{u}_n(\cdot)}$ ,  $t = r - r_1$ , and then taking  $n \rightarrow \infty$ , we obtain

$$\|\bar{x}^1 - x\| \leq \frac{(L\|x\| + K_1)(e^{L(r-r_1)} - 1)}{L} \leq \frac{(L\|x\| + K_1)(e^{Lr} - 1)}{L}, \quad (3.3.18)$$

from which it follows that  $\|\bar{x}^1\| \leq e^{Lr}\|x\| + \frac{(e^{Lr}-1)K_1}{L}$ . Hence,

$$\begin{aligned} \langle M^T(r)v, z - \bar{x}^1 \rangle &\leq R_1(r, \|x\|, \rho) e^{L_2(r, \|x\|)r} \|M^T(r)v\| \|z - \bar{x}^1\|^2 \\ &\quad + e^{L_2(r, \|x\|)r} \|M^T(r)v\| |r - r_1| \|z - \bar{x}^1\|, \end{aligned} \quad (3.3.19)$$

where

$$R_1(r, \delta, \rho) = K \left( r, e^{Lr}\delta + \frac{(e^{Lr}-1)K_1}{L}, \rho \right), \quad \text{for } r, \delta \geq 0, \rho > 0.$$

Observe also that we obtain from (iii) in Lemma 7.1.1 that

$$\begin{aligned} \|z - \bar{x}^1\| &\leq \lim_{n \rightarrow \infty} \left( \|z - x_n\| + \int_0^{r-r_1^n} \|f(\alpha_n(s), \bar{u}_n(s))\| ds \right) \\ &\leq \lim_{n \rightarrow \infty} \left( \|z - x_n\| + \int_0^{r-r_1^n} (Le^{Ls}\|x_n\| + e^{Ls}K_1) ds \right) \\ &\leq \|z - x\| + L_4(r, \|x\|) |r - r_1|, \end{aligned}$$

where  $L_4(s, \delta) = Le^{Ls}\delta + e^{Ls}K_1$  for  $s, \delta \geq 0$ .

Combining the above inequality and (3.3.19), we obtain

$$\langle M^T(r)v, z - \bar{x}^1 \rangle \leq R_2(r, \|x\|, \rho) \|M^T(r)v\| (\|z - x\|^2 + |r - r_1|^2), \quad (3.3.20)$$

where we have defined, for  $r, \delta \geq 0, \rho > 0$ ,

$$R_2(r, \delta, \rho) = e^{L_2(r, \delta)r} \left( 2R_1(r, \delta, \rho) \left( \frac{3}{2} + L_4^2(r, \delta) \right) + L_4(r, \delta) \right). \quad (3.3.21)$$

Second, we consider

$$\begin{aligned} &\langle M^T(r)v, \bar{x}_n^1 - x \rangle \\ &= \langle M^T(r)v, x_n - x \rangle + \left\langle M^T(r)v, \int_0^{r-r_1^n} f(\alpha_n(s), \bar{u}_n(s)) ds \right\rangle \\ &= \langle M^T(r)v, x_n - x \rangle + \left\langle M^T(r)v, \int_0^{r-r_1^n} f(x, \bar{u}_n(s)) ds \right\rangle \\ &\quad + \left\langle M^T(r)v, \int_0^{r-r_1^n} (f(\alpha_n(s), \bar{u}_n(s)) - f(x, \bar{u}_n(s))) ds \right\rangle. \end{aligned}$$

Recalling that  $H(M^T(r)v, x) = 0$ , we obtain from the above expression that

$$\begin{aligned}
\langle M^T(r)v, \bar{x}_n^1 - x \rangle &\leq \langle M^T(r)v, x_n - x \rangle + \\
&\quad \left\langle M^T(r)v, \int_0^{r-r_n^1} (f(\alpha_n(s), \bar{u}_n(s)) - f(x, \bar{u}_n(s))) ds \right\rangle \\
&\leq \|M^T(r)v\| \left( \|x_n - x\| + \int_0^{r-r_n^1} \|f(\alpha_n(s), \bar{u}_n(s)) - f(x, \bar{u}_n(s))\| ds \right) \\
&\leq \|M^T(r)v\| \left( \|x_n - x\| + L \int_0^{r-r_n^1} \|\alpha_n(s) - x\| ds \right) \\
&\leq \|M^T(r)v\| \left( (L+1) \|x_n - x\| + L \int_0^{r-r_n^1} \int_0^s \|f(\alpha_n(\tau), \bar{u}_n(\tau))\| d\tau ds \right).
\end{aligned}$$

By (iii) in Lemma 7.1.1, recalling that  $\bar{r}_1 < r$  we now obtain that

$$\begin{aligned}
\langle M^T(r)v, \bar{x}_n^1 - x \rangle &\leq \|M^T(r)v\| \left( (L+1) \|x_n - x\| \right. \\
&\quad \left. + L \int_0^{r-r_1} \int_0^s (Le^{Lr} \|x_n\| + e^{Lr} K_1) d\tau ds \right),
\end{aligned}$$

whence, taking  $n \rightarrow \infty$ ,

$$\langle M^T(r)v, \bar{x}^1 - x \rangle \leq \frac{L(Le^{Lr} \|x\| + e^{Lr} K_1)}{2} \|M^T(r)v\| |r - r_1|^2. \quad (3.3.22)$$

Set now, for  $r, \delta \geq 0, \rho > 0$ ,

$$K_3(r, \delta, \rho) = R_2(r, \delta, \rho) + \frac{L(Le^{Lr} \delta + e^{Lr} K_1)}{2}. \quad (3.3.23)$$

Recalling (3.3.20) and (3.3.22), the proof is complete.  $\square$

Now we prove a similar result for normals such that the Hamiltonian along the *limiting adjoint flow* is 1. Actually, if  $\xi$  is such a vector, we show that  $(\xi, 1)$  is a proximal normal to the hypograph of  $T(\cdot)$ , and again the radius of the sphere which realizes it can be explicitly estimated.

**Lemma 3.3.3** *Let  $\mathcal{S}$  be closed and let the assumptions (H1), (H2), and (H3) hold. Let  $x \in \mathcal{S}^c$ , set  $r := T(x) > 0$ , and let  $\xi \in N_1(x)$ . Then  $-\xi \in \partial^P T(x)$ , or, equivalently,  $(\xi, 1) \in N_{\text{hypo}(T(x))}^P(x, T(x))$ .*

*More precisely, let  $\bar{x} \in \mathcal{S}_x$  and let  $v \in N_{\overline{\mathcal{S}^c}}^P(\bar{x})$ ,  $M(\cdot) \in \mathcal{M}_{\bar{x}}$  be such that  $H(M^T(r)v, x) = 1$  and assume that  $v$  is realized by a ball of radius  $\rho > 0$ . Then there exists an explicitly computable continuous function  $K_6(r, \|x\|, \rho)$  depending only on  $r, \|x\|, \rho$  such that for all  $z \in \overline{\mathcal{S}^c}$  and all  $\beta \leq T(z)$  we have*

$$\langle M^T(r)v, z - x \rangle + \beta - r \leq K_6(r, \|x\|, \rho) \|(M^T(r)v, 1)\| (\|z - x\|^2 + |\beta - r|^2). \quad (3.3.24)$$

**Proof.** Let  $v \in N_{\mathcal{S}^c}^P(\bar{x})$  be such that

$$\langle v, \bar{z} - \bar{x} \rangle \leq \frac{\|v\|}{2\rho} \|\bar{z} - \bar{x}\|^2 \quad \forall \bar{z} \in \overline{\mathcal{S}^c}.$$

Let  $z \in \mathcal{S}^c$ . Two cases may occur:

- (i)  $T(z) \geq T(x)$ ,
- (ii)  $T(z) < T(x)$ .

**First case.** Recalling that  $H(M^T(r)v, x) = 1$ , one can find  $\bar{u} \in \mathcal{U}$  such that

$$\langle M^T(r)v, f(x, \bar{u}) \rangle = 1.$$

Set  $z_{\bar{u}}(\cdot) := y^{z, \bar{u}}(\cdot)$  to be the trajectory starting from  $z$  with the constant control  $\bar{u}$ , namely  $z_{\bar{u}}(t) = z + \int_0^t f(z_{\bar{u}}(s), \bar{u}) ds$ .

Taking  $T(x) \leq r_1 \leq T(z)$ , we have that  $z_{\bar{u}}(r_1 - r) \in \mathcal{S}^c(r)$ . Recalling Lemma 3.3.1, we obtain that

$$\langle M^T(r)v, z_{\bar{u}}(r_1 - r) - x \rangle \leq K(r, \|x\|, \rho) \|M^T(r)v\| \|z_{\bar{u}}(r_1 - r) - x\|^2. \quad (3.3.25)$$

We estimate

$$\begin{aligned} \langle M^T(r)v, z - z_{\bar{u}}(r_1 - r) \rangle &= \left\langle M^T(r)v, - \int_0^{r_1-r} f(z_{\bar{u}}(t), \bar{u}) dt \right\rangle \\ &= \left\langle M^T(r)v, - \int_0^{r_1-r} f(x, \bar{u}) dt \right\rangle \\ &\quad + \left\langle M^T(r)v, \int_0^{r_1-r} (f(x, \bar{u}) - f(z_{\bar{u}}(t), \bar{u})) dt \right\rangle \\ &\leq r - r_1 + L \|M^T(r)v\| \int_0^{r_1-r} \|z_{\bar{u}}(t) - x\| dt. \end{aligned}$$

Combining the above inequality with (3.3.25) we get

$$\begin{aligned} \langle M^T(r)v, z - x \rangle &\leq r - r_1 + L \|M^T(r)v\| \int_0^{r_1-r} \|z_{\bar{u}}(t) - x\| dt \\ &\quad + K(r, \|x\|, \rho) \|M^T(r)v\| \|z_{\bar{u}}(r_1 - r) - x\|^2. \end{aligned} \quad (3.3.26)$$

Moreover,

$$\begin{aligned} \|z_{\bar{u}}(s) - x\| &\leq \|z - x\| + \int_0^s \|f(z_{\bar{u}}(\tau), \bar{u})\| dt \\ &\leq \|z - x\| + \tilde{K}(\|x\|)s + L \int_0^s \|z_{\bar{u}}(\tau) - x\| d\tau, \end{aligned}$$

where we set, for  $\delta \geq 0$ ,  $\tilde{K}(\delta) := L\delta + K_1$ . Thus, Gronwall's inequality yields, for all  $0 \leq s \leq r_1 - r$ ,

$$\|z_{\bar{u}}(s) - x\| \leq e^{Ls} \|z - x\| + \tilde{K}(\|x\|) \left( s + \frac{e^{Ls} - Ls - 1}{L} \right). \quad (3.3.27)$$

Since  $e^{Ls} - Ls - 1 \leq L(e^L - 1)s$  for all  $s \in [0, 1]$ , we obtain from (3.3.27)

$$\|z_{\bar{u}}(s) - x\| \leq e^L \|z - x\| + \tilde{K}(\|x\|)e^{Ls} \quad \text{for all } s \in [0, 1]. \quad (3.3.28)$$

Now we consider two subcases.

*First subcase:*  $0 \leq r_1 - r \leq 1$ . Combining (3.3.28) with (3.3.26) we obtain

$$\langle M^T(r)v, z - x \rangle + r_1 - r \leq K_5(r, \|x\|, \rho) \|M^T(r)v\| (\|z - x\|^2 + |r_1 - r|^2), \quad (3.3.29)$$

where for  $r, \delta \geq 0, \rho > 0$  we set

$$K_5(r, \delta, \rho) = e^L \left( \frac{L}{2} + 2e^L K(r, \delta, \rho)(1 + \tilde{K}(\delta)^2) + \frac{\tilde{K}(\delta)}{2} \right). \quad (3.3.30)$$

*Second subcase:*  $r_1 - r > 1$ . Recalling Lemma 3.3.1, we obtain

$$\begin{aligned} \langle M^T(r)v, z - x \rangle + r_1 - r \\ \leq (K(r, \|x\|, \rho) + 1) \|(M^T(r)v, 1)\| (\|z - x\|^2 + |r_1 - r|^2). \end{aligned} \quad (3.3.31)$$

Observe now that, if  $\beta \leq T(x)$ , recalling Lemma 3.3.1 we have

$$\langle M^T(r)v, z - x \rangle + \beta - T(x) \leq K(r, \|x\|, \rho) \|M^T(r)v\| (\|z - x\|^2 + |\beta - T(x)|^2). \quad (3.3.32)$$

We are now ready to conclude the first case. Indeed, it suffices to combine (3.3.29), (3.3.32), and (3.3.31) and recall (3.3.30), obtaining, for all  $z \in \mathcal{S}^c(r)$  and  $\beta \leq T(z)$ ,

$$\begin{aligned} \langle M^T(r)v, z - x \rangle + \beta - T(x) \\ \leq (K_5(r, \|x\|, \rho) + 1) \|(M^T(r)v, 1)\| (\|z - x\|^2 + |\beta - T(x)|^2). \end{aligned} \quad (3.3.33)$$

**Second case.** It is entirely similar to the proof of the second case of Lemma (3.3.2). Indeed, by using the condition  $H(M^T(r)v, x) = 1$  we can replace (3.3.22) with

$$\langle M^T(r)v, \bar{x}^1 - x \rangle \leq T(x) - T(z) + \frac{L(Le^{Lr} \|x\| + e^{Lr} K_1)}{2} |r - r_1|^2. \quad (3.3.34)$$

Then, combining (3.3.20) and (3.3.34) we obtain

$$\begin{aligned} \langle M^T(r)v, z - x \rangle + \beta - T(x) \\ \leq K_3(r, \|x\|, \rho) \|M^T(r)v\| (\|z - x\|^2 + |\beta - T(x)|^2) \end{aligned} \quad (3.3.35)$$

for all  $\beta \leq T(z)$ ,  $z \in \overline{\mathcal{S}^c}$  and  $T(z) \leq T(x)$ .

To conclude the proof of the Lemma we recall (3.3.35), (3.3.33), (3.3.30) and set, for  $r, \delta \geq 0, \rho > 0$ ,

$$K_6(r, \delta, \rho) = \max\{K_5(r, \delta, \rho) + 1, K_3(r, \delta, \rho)\}. \quad (3.3.36)$$

□

The next subsection is to show that singularities of  $T$  may be only of “upwards type”. Assuming that the target satisfies the internal sphere condition of radius  $\rho$ , we show that if  $\xi$  belongs to the proximal subgradient of  $T(\cdot)$  at  $x$ , then it belongs also to the proximal supergradient. Moreover  $-\xi$  is the transported vector by the limiting adjoint flow of a normal to  $\mathcal{S}^c$ , which is realized by  $\rho$ , and the radius of the sphere realizing  $(-\xi, 1)$  as a proximal normal to the hypograph of  $T(\cdot)$  can be explicitly estimated. In this lemma, the internal sphere condition (H4) is used for the first time.

### 3.3.2 Type of singularities of the minimum time function

In order to simplify our writing, we will replace the functions  $K$ ,  $K_3$ , and  $K_6$  appearing respectively in Lemma 3.3.1, Lemma 3.3.2, and Lemma 3.3.3 by the explicit (continuous) function

$$k(r, \|x\|, \rho) = \max\{K_6(r, \|x\|, \rho), K(r, \|x\|, \rho)\}. \quad (3.3.37)$$

**Lemma 3.3.4** *Let the assumptions (H1) – (H4) hold and let  $x \in \mathcal{S}^c$  and let  $\xi \in \partial_P T(x)$ . Then*

- (i)  $\xi \in \partial^P T(x)$  and therefore  $T$  is differentiable at  $x$ ;
- (ii)  $-\xi \in N_1(x)$ .

Moreover, for all  $z \in \overline{\mathcal{S}^c}$  and for all  $\beta \leq T(z)$ ,

$$\langle -\xi, z - x \rangle + \beta - T(x) \leq k(T(x), \|x\|, \rho) \|(-\xi, 1)\| (\|z - x\|^2 + |\beta - T(x)|^2). \quad (3.3.38)$$

**Proof.** Set  $r = T(x)$  and let  $\xi \in \partial_P T(x)$ . By Proposition IV.2.3 in [11],  $H(x, -\xi) \geq 1$ , so that  $\xi \neq 0$ . It follows from the definition of proximal subgradient that there exists  $\sigma \geq 0$  such that

$$\langle \xi, z - x \rangle \leq \sigma \|z - x\|^2, \quad \forall z \in \mathcal{S}(r). \quad (3.3.39)$$

Let  $\bar{x} \in \mathcal{S}_x$  and  $M(\cdot) \in \mathcal{M}_{\bar{x}}$ , and take a sequence  $\{y^{x_n, \bar{u}_n}(\cdot)\} \subset \mathcal{T}_{\bar{x}}$  such that  $M(\cdot)$  is the uniform limit of  $M(\cdot, x_n, \bar{u}_n)$ . We claim that  $(M^T(r))^{-1}\xi \in N_{\mathcal{S}}^P(\bar{x})$ .

Indeed, take  $\bar{z} \in \mathcal{S}$  and set  $\bar{z}_n^-(\cdot) = y^-(\cdot, \bar{z}, \bar{u}_n)$  where  $y^-(\cdot, \bar{z}, \bar{u}_n)$  is the solution of

$$\begin{cases} \dot{y}(t) &= -f(y(t), \bar{u}_n(r-t)) \text{ a.e.} \\ y(0) &= \bar{z}. \end{cases}$$

We set  $z_n = z_n^-(\theta(x_n, \bar{u}_n))$  and consider  $\bar{z}_n = y^{z_n, \bar{u}_n}(\theta(x_n, \bar{u}_n))$ . We can assume without loss of generality that  $\{z_n\}$  converges to some  $z$ , which is easily seen belonging to  $\mathcal{S}(r)$ .

To simplify our writing, we set  $t_n = \theta(x_n, \bar{u}_n)$ ,  $\alpha_n(\cdot) = y^{x_n, \bar{u}_n}(\cdot)$ ,  $\bar{x}_n = \alpha_n(t_n)$ , and  $M_n^1(\cdot) = M(\cdot, x_n, \bar{u}_n)$ . Let also  $\beta_n(\cdot) = y^{z_n, \bar{u}_n}(\cdot)$ ,  $A_n(t) =$

$\int_0^1 D_x f(\alpha_n(t) + \tau(\beta_n(t) - \alpha_n(t)), \bar{u}_n(t)) d\tau$  and let  $M_n^2(\cdot)$  be the fundamental solution of  $\dot{p}(t) = A_n(t)p(t)$ ,  $p(0) = \mathbb{I}^{N \times N}$ . Finally, we set  $w_n^i(t) = M_n^i(z_n - x_n)$  for  $i \in \{1, 2\}$ .

Using Lemma 7.1.2 and the same argument leading to (3.3.8) we can perform the following estimate:

$$\begin{aligned} \langle M^T(r)^{-1}\xi, \bar{z}_n - \bar{x}_n \rangle &= \langle M^T(r)^{-1}\xi, w_n^2(t_n) \rangle \\ &= \langle M^T(r)^{-1}\xi, w_n^1(t_n) \rangle + \langle M^T(r)^{-1}\xi, w_n^2(t_n) - w_n^1(t_n) \rangle \\ &\leq \langle M^T(r)^{-1}\xi, w_n^1(t_n) \rangle + \|M^T(r)^{-1}\| \|\xi\| \|w_n^2(t_n) - w_n^1(t_n)\| \\ &\leq \langle M^T(r)^{-1}\xi, w_n^1(t_n) \rangle + \tilde{K}_0 \|z_n - x_n\|^2 \\ &\leq \langle M^T(r)^{-1}\xi, w_n^1(t_n) \rangle + \tilde{K}_1 \|\bar{z} - \bar{x}_n\|^2, \end{aligned}$$

where  $\tilde{K}_0$  and  $\tilde{K}_1$  are suitable constants. Taking  $n \rightarrow \infty$  in the above inequalities, we obtain

$$\begin{aligned} \langle M^T(r)^{-1}\xi, \bar{z} - \bar{x} \rangle &\leq \langle M^T(r)^{-1}\xi, M^T(r)(z - x) \rangle + \tilde{K}_1 \|\bar{z} - \bar{x}\|^2 \\ &= \langle \xi, z - x \rangle + \tilde{K}_1 \|\bar{z} - \bar{x}\|^2. \end{aligned}$$

Recalling (3.3.39) and Lemma 7.1.2, we thus obtain

$$\begin{aligned} \langle M^T(r)^{-1}\xi, \bar{z} - \bar{x} \rangle &\leq \sigma \|\xi\| \|z - x\|^2 + \tilde{K}_1 \|\bar{z} - \bar{x}\|^2 \\ &\leq \tilde{K}_2 \|\bar{z} - \bar{x}\|^2, \end{aligned}$$

for a suitable constant  $\tilde{K}_2$ . The above inequality in turn implies that

$$(M^T(r))^{-1}\xi \in N_{\mathcal{S}}^P(\bar{x}). \quad (3.3.40)$$

Thanks to (H4), there exists  $0 \neq \zeta \in N_{\overline{\mathcal{S}^c}}^P(\bar{x})$ . Therefore, both  $\mathcal{S}$  and  $\overline{\mathcal{S}^c}$  admit at  $\bar{x}$  an external nonzero proximal normal. This means that  $\mathcal{S}$  is smooth at  $\bar{x}$ , and so, by (H4), the unique external normal to  $\overline{\mathcal{S}^c}$  at  $\bar{x}$ , namely  $-M^T(r)^{-1}\xi$ , must be realized by a ball of radius  $\rho$ .

Using Proposition IV.2.3 in [11] we see that  $H(x, -\xi) \geq 1$ , and so we can choose  $\lambda \in (0, 1)$  such that  $H(-\lambda\xi, x) = 1$ . Applying Lemma 3.3.3 for  $v = \lambda M^T(r)^{-1}\xi$ , we obtain that  $\lambda\xi \in \partial^P T(x)$ . Therefore,  $T$  is differentiable at  $x$  and so  $\lambda\xi = \xi$ . Thus both (i) and (ii) are proved.

In order to complete the proof, we apply the last statement of Lemma 3.3.3.

□

The next lemma classifies limiting normals, and shows that limiting subgradients generate proximal normals to the hypograph which are *horizontal/non-horizontal* according to the *unboundedness/boundedness* of the corresponding sequence of proximal subgradients. Also, the radius of the sphere realizing the limiting vector can be explicitly estimated.

**Lemma 3.3.5** *Let the assumptions (H1) – (H4) hold, and let  $\{x_n\}$  be a sequence converging to  $x \in \mathcal{S}^c$ . Assume that there exists a sequence  $\{\xi_n\}$  satisfying  $\xi_n \in \partial_P T(x_n)$ .*

*Then the following alternatives hold true:*

(i) *If  $\limsup_{n \rightarrow \infty} \|\xi_n\| < +\infty$  then there exists a subsequence  $\{\xi_{n_k}\}$  converging to a vector  $\xi$  such that  $-\xi \in N_1(x)$ . Moreover,  $(-\xi, 1) \in N_{\text{hypo}(T)}^P(x, T(x))$  and, for all  $z \in \mathcal{S}^c$  and all  $\beta \leq T(z)$  the inequality*

$$\langle -\xi, z - x \rangle + \beta - T(x) \leq k(T(x), \|x\|, \rho) \|(-\xi, 1)\| (\|z - x\|^2 + |\beta - T(x)|^2) \quad (3.3.41)$$

*holds.*

(ii) *If  $\limsup_{n \rightarrow \infty} \|\xi_n\| = +\infty$  then there exists a subsequence of  $\{\frac{\xi_n}{\|\xi_n\|}\}$  converging to a vector  $\xi$  such that  $-\xi \in N_0(x)$ . Moreover,  $(-\xi, 0)$  is in  $N_{\text{hypo}(T)}^P(x, T(x))$  and for all  $z \in \mathcal{S}^c$  and all  $\beta \leq T(z)$ , the inequality*

$$\langle -\xi, z - x \rangle \leq k(T(x), \|x\|, \rho) (\|z - x\|^2 + |\beta - T(x)|^2) \quad (3.3.42)$$

*holds.*

**Proof.** Set  $r = T(x)$ . Recalling Lemma (3.3.4), the function  $T(\cdot)$  is differentiable at  $x_n$ . Taking  $\bar{x}_n \in \mathcal{S}_{x_n}$  and  $M_n(\cdot) \in \mathcal{M}_{\bar{x}_n}$ , it follows from Lemma 3.3.4 that for all  $n \in \mathbb{N}$

a)  $-M_n^T(T(x_n))^{-1}\xi_n \in N_{\mathcal{S}^c}^P(\bar{x}_n)$  and each  $-M_n^T(T(x_n))^{-1}\xi_n$  is realized by a ball of radius  $\rho$ , namely

$$\langle -M_n^T(T(x_n))^{-1}\xi_n, \bar{z} - \bar{x}_n \rangle \leq \frac{\|M_n^T(T(x_n))^{-1}\xi_n\|}{2\rho} \|\bar{z} - \bar{x}_n\|^2, \quad \forall \bar{z} \in \overline{\mathcal{S}^c}. \quad (3.3.43)$$

b)  $H(-\xi_n, x_n) = 1$ .

If  $\limsup_{n \rightarrow \infty} \|\xi_n\| < +\infty$ , we choose subsequences  $\{\bar{x}_{n_k}\}$  and  $\{\xi_{n_k}\}$  converging respectively to  $\bar{x} \in \mathcal{S}$  and  $\bar{\xi}$ . By compactness, without loss of generality we can assume that  $\{M_{n_k}(\cdot)\}$  converges uniformly to  $M(\cdot)$ . We now take  $n_k \rightarrow \infty$  in (3.3.43) and obtain

$$\langle -M^T(r)^{-1}\bar{\xi}, \bar{z} - \bar{x} \rangle \leq \frac{\|M^T(r)^{-1}\bar{\xi}\|}{2\rho} \|\bar{z} - \bar{x}\|^2. \quad (3.3.44)$$

Thus  $-M^T(r)^{-1}\bar{\xi} \in N_{\mathcal{S}^c}^P(\bar{x})$  and  $-M^T(r)^{-1}\bar{\xi}$  is realized by a ball of radius  $\rho$ .

On the other hand, we also take  $n_k \rightarrow \infty$  in b) and obtain  $H(-\bar{\xi}, x) = 1$ .

One can also easily show that  $M^T(\cdot) \in \mathcal{M}_{\bar{x}}$ , so that  $-\bar{\xi} \in N_1(x)$ . Recalling Lemma 3.3.3 and setting  $\xi := \bar{\xi}$  the proof of (i) is concluded.

Analogously, if  $\limsup_{n \rightarrow \infty} \|\xi_n\| = +\infty$ , we can assume that  $-\bar{\xi} = -\lim_{n_k \rightarrow \infty} \frac{\xi_{n_k}}{\|\xi_{n_k}\|}$ , together with  $-M^T(r)^{-1}\bar{\xi} \in N_{\mathcal{S}^c}^P(\bar{x})$  and  $H(-\bar{\xi}, x) = 0$ . Thus  $-\bar{\xi} \in N_0(x)$ . Finally, recalling Lemma 3.3.2 and setting  $\xi := \bar{\xi}$  we conclude the proof of (ii).  $\square$

### 3.3.3 Wedgedness and exposed rays

The final results of this section use for the first time the wedgedness assumption for the normal cone  $N_{\text{hypo}(T)}^P(x, T(x))$ . They show essentially that  $N_{\text{hypo}(T)}^P(x, T(x))$  is a closed cone, and that horizontal (resp. non-horizontal) exposed rays of  $N_{\text{hypo}(T)}^P(x, T(x))$  belong to  $N_0(x)$  (resp.  $N_1(x)$ ). As a byproduct of our argument we obtain a representation of  $N_{\text{hypo}(T)}^P(x, T(x))$  through  $N_0(x)$  and  $N_1(x)$  (see Theorem 3.3.1).

**Lemma 3.3.6** *Let  $s \in \mathcal{S}^c$  and let the assumptions (H1) - H(4) hold. Assume that  $N_{\text{hypo}(T)}^P(x, T(x))$  is wedged and set*

$$\begin{aligned}\tilde{N}_0(x) &= \{(\xi, 0) \mid \xi \in N_0(x)\}, \\ \tilde{N}_1(x) &= \{\lambda(\xi, 1) \mid \xi \in N_1(x), \lambda \geq 0\}, \\ N(x) &= \text{co}\tilde{N}_0(x) + \text{co}\tilde{N}_1(x).\end{aligned}$$

*Then  $N(x) \subseteq N_{\text{hypo}(T)}^P(x, T(x))$  is a closed, convex, and wedged cone.*

**Proof.** Thanks to Lemmas 3.3.2 and 3.3.3 and the definition of  $k$  in (3.3.37), every  $\zeta \in \tilde{N}_0(x) \cup \tilde{N}_1(x)$  satisfies the following property: for every  $y \in \mathcal{S}^c$  and every  $\beta \leq T(y)$ , the inequality

$$\langle \zeta, (y - x, \beta - T(x)) \rangle \leq k(T(x), \|x\|, \rho) \|\zeta\| \left( \|y - x\|^2 + |\beta - T(x)|^2 \right) \quad (3.3.45)$$

holds. It follows immediately from the above property that both  $\tilde{N}_0(x)$  and  $\tilde{N}_1(x)$  are cones contained in  $N_{\text{hypo}(T)}^P(x, T(x))$ . Thus  $\text{co}\tilde{N}_0(x)$  and  $\text{co}\tilde{N}_1(x)$  are contained in  $N_{\text{hypo}(T)}^P(x, T(x))$ , and so they are wedged. Set  $N_0^1 = \{\xi \in \mathbb{R}^N \mid \xi \in N_0(x), \|\xi\| = 1\}$ , and observe that on one hand  $\tilde{N}_0(x) = \{\lambda(\xi, 0) \mid \xi \in N_0^1, \lambda \geq 0\}$ , on the other  $N_0^1$  (by the continuity of the Hamiltonian) is compact and  $0 \notin N_0^1$ . Analogously, observe that  $N_1(x)$  is compact and does not contain zero. Therefore, using Lemma 7.2.1, we obtain that both  $\text{co}\tilde{N}_0(x)$  and  $\text{co}\tilde{N}_1(x)$  are closed, and the proof is concluded.  $\square$

**Lemma 3.3.7** *Let  $x \in \mathcal{S}^c$  and let the assumptions of Theorem 3.2.1 hold. Let  $\tilde{N}$  be a closed convex cone in  $\mathbb{R}^{N+1}$  with the property*

$$N(x) \subseteq \tilde{N} \subseteq N_{\text{hypo}(T)}^P(x, T(x)). \quad (3.3.46)$$

*Let  $\zeta$  belong to an exposed ray of  $\tilde{N}$ . The following statements hold true:*

- (i) *if  $\zeta = (\xi, 0)$ , with  $\xi \in \mathbb{R}^N$ , then  $\xi \in N_0(x)$ ;*
- (ii) *if  $\zeta = (\xi, \lambda)$ , with  $\xi \in \mathbb{R}^N$  and  $\lambda > 0$ , then  $\xi/\lambda \in N_1(x)$ .*



Moreover,  $\zeta$  satisfies (3.3.45) for all  $y \in \mathcal{S}^c$  and all  $\beta \leq T(y)$ .

**Proof.** By our assumption on  $\zeta$ , there exists  $\bar{v} = (v_0, \lambda_0)$  satisfying  $v_0 \in \mathbb{R}^N$ ,  $\|v_0\| = 1$ , and  $\lambda_0 \in \mathbb{R}$  such that

$$\begin{cases} \langle (v_0, \lambda_0), \zeta \rangle = 0 \\ \langle (v_0, \lambda_0), w \rangle \leq 0 \quad \forall w \in \tilde{N} \\ \langle (v_0, \lambda_0), w \rangle = 0, \text{ and } 0 \neq w \in \tilde{N} \Rightarrow \frac{w}{\|w\|} = \frac{\zeta}{\|\zeta\|}. \end{cases} \quad (3.3.47)$$

We now begin proving (i). Since  $\zeta = (\xi, 0) \in N_{\text{hypo}(T)}^P(x, T(x))$ , there exists a constant  $\sigma \geq 0$  such that, for all  $z \in \overline{\mathcal{S}^c}$  and all  $\beta \leq T(z)$ , the inequality

$$\langle \xi, z - x \rangle \leq \sigma (\|z - x\|^2 + |\beta - T(x)|^2) \quad (3.3.48)$$

holds. Set now  $x_n = x + \frac{v_0}{n} + \frac{\xi}{n\sqrt{n}}$ . Then, by the Density Theorem (see [25, Theorem 1.3.1]), for each  $n$  there exists  $z_n$  such that

$$\partial_P T(z_n) \neq \emptyset, \quad (3.3.49)$$

$$\|z_n - x_n\| \leq \frac{1}{n^2}. \quad (3.3.50)$$

First, we show that

$$T(z_n) \leq T(x) \quad \text{for all } n \text{ large enough.} \quad (3.3.51)$$

Indeed, assume by contradiction that  $T(z_n) > T(x)$ . Taking  $z = z_n$  and  $\beta = T(x)$  in (3.3.48), we obtain

$$\langle \xi, z_n - x \rangle \leq \sigma \|z_n - x\|^2.$$

It follows from the above inequality, (3.3.47), and (3.3.50) that there exists a suitable constant  $\sigma_1$  for which

$$\frac{\|\xi\|^2}{n\sqrt{n}} \leq \frac{\sigma_1}{n^2}$$

for all  $n$  large enough, a contradiction.

Second, we claim that there exists  $\sigma_2$  such that

$$|T(z_n) - T(x)| > \sigma_2 n^{-\frac{3}{4}} \quad \text{for all } n \text{ large enough.} \quad (3.3.52)$$

Indeed, taking  $z = z_n$  and  $\beta = T(z_n)$  in (3.3.48) we obtain

$$\langle \xi, z_n - x \rangle \leq \sigma (\|z_n - x\|^2 + |T(z_n) - T(x)|^2).$$

From the above inequality, (3.3.47), and (3.3.50), one can easily see that (3.3.52) holds.

On the other hand, by (3.3.49) and Lemma 3.3.4 we know that  $T$  is differentiable at  $z_n$  and we write  $\xi_n = DT(z_n)$ . Recalling (3.3.38), for all  $z \in \mathcal{S}^c$  and all  $\beta \leq T(z)$  the inequality

$$\begin{aligned} \langle -\xi_n, z - z_n \rangle + \beta - T(z_n) \\ \leq k(T(z_n), \|z_n\|, \rho) \|(-\xi_n, 1)\| (\|z - z_n\|^2 + |\beta - T(z_n)|^2) \end{aligned} \quad (3.3.53)$$

holds.

We claim that  $\|\xi_n\| \rightarrow +\infty$ .

Assume by contradiction that there exists a constant  $Q$  such that  $\|\xi_n\| \leq Q$  for all  $n$ . Taking  $z = x$ ,  $\beta = T(x)$  in (3.3.53) and recalling (3.3.51), we obtain that

$$\begin{aligned} (T(x) - T(z_n)) \left(1 - k(T(z_n), \|z_n\|, \rho) \sqrt{Q^2 + 1} |T(x) - T(z_n)|\right) \\ \leq \|x - z_n\| \left(Q + k(T(z_n), \|z_n\|, \rho) \sqrt{Q^2 + 1} \|x - z_n\|\right). \end{aligned}$$

By the continuity of  $T(\cdot)$  and  $k(\cdot)$  and by (3.3.51), (3.3.50), and (3.3.52), there exists a constant  $Q_1 > 0$  such that

$$\frac{Q_1}{n^{\frac{3}{4}}} \leq \frac{1}{n} \quad \text{for all } n \text{ large enough,}$$

a contradiction.

Now, recalling (ii) in Lemma 3.3.5 and assuming without loss of generality that  $\lim_{n \rightarrow \infty} -\frac{\xi_n}{\|\xi_n\|} = -\bar{\xi}$ , we see that  $(-\bar{\xi}, 0) \in \tilde{N}_0(x) \subseteq \tilde{N}$ . By (3.3.51) we can take  $z = x$  and  $\beta = T(z_n)$  in (3.3.53), obtaining

$$\left\langle -\frac{\xi_n}{\|(-\xi_n, 1)\|}, \frac{x - z_n}{\|x - z_n\|} \right\rangle \leq k(T(z_n), \|z_n\|, \rho) \|x - z_n\|.$$

Taking  $n \rightarrow \infty$  in the above inequality and recalling (3.3.50) we obtain

$$\langle -\bar{\xi}, -v_0 \rangle \leq 0,$$

or, equivalently,  $\langle (-\bar{\xi}, 0), (v_0, \lambda_0) \rangle \geq 0$ . Therefore, we obtain from (3.3.47) that  $(-\bar{\xi}, 0) = \frac{(\xi, 0)}{\|\xi\|}$ . Thus  $\xi = -\bar{\xi}$  and the proof of claim (i) is concluded.

*Ad* (ii). We now take  $\zeta = (\xi, 1)$  and take  $\bar{v} = (v_0, \lambda_0)$  satisfying (3.3.47). Set  $x_n = x + \frac{v_0}{n}$ . Then by the Density Theorem (see Theorem 1.3.1 in [25]) for each  $n$  there exists  $z_n$  such that

$$\partial_P T(z_n) \neq \emptyset, \quad (3.3.54)$$

$$\|z_n - x_n\| \leq \frac{1}{n^2}. \quad (3.3.55)$$

Recalling Lemma 3.3.4, (3.3.54) implies that  $T(\cdot)$  is differentiable at  $z_n$ . Moreover, if we set  $\xi_n = DT(z_n)$ , then  $-\xi_n \in N_1(z_n)$  and for all  $z \in \mathcal{S}^c$  and  $\beta \leq T(z)$  we have

$$\begin{aligned} & \langle -\xi_n, z - z_n \rangle + \beta - T(z_n) \\ & \leq k(T(z_n), \|z_n\|, \rho) \|(-\xi_n, 1)\| (\|z - z_n\|^2 + |\beta - T(z_n)|^2). \end{aligned} \quad (3.3.56)$$

We claim that the sequence  $\{\xi_n\}$  is bounded.

Suppose by contradiction that  $\limsup_{n \rightarrow \infty} \|\xi_n\| = +\infty$ . Then assuming without loss of generality that  $-\frac{\xi_n}{\|\xi_n\|} \rightarrow -\bar{\xi}$ , (ii) of Lemma 3.3.5 yields that  $-\bar{\xi} \in N_0(x)$  and  $(-\bar{\xi}, 0) \in \tilde{N}_0(x)$ .

In order to obtain a contradiction, we consider two cases:

- a)  $T(x) \geq T(z_n)$  for infinitely many  $n$ ;
- b)  $T(x) < T(z_n)$  for infinitely many  $n$ .

In the first case, we can choose  $z = x$ ,  $\beta = T(z_n)$  in (3.3.56), obtaining

$$\left\langle -\frac{\xi_n}{\|(-\xi_n, 1)\|}, \frac{x - z_n}{\|x - z_n\|} \right\rangle \leq k(T(z_n), z_n, \rho) \|x - z_n\|.$$

Taking  $n \rightarrow \infty$  and recalling (3.3.55) we get

$$\langle -\bar{\xi}, -v_0 \rangle \leq 0, \quad (3.3.57)$$

which implies  $\langle (-\bar{\xi}, 0), (v_0, \lambda_0) \rangle \geq 0$ . Thus, combining  $(-\bar{\xi}, 0) \in \tilde{N}_0(x)$  with (3.3.47) we obtain  $\frac{(-\bar{\xi}, 0)}{\|-\bar{\xi}\|} = \frac{(\xi, 1)}{\|(\xi, 1)\|}$ , a contradiction.

In the second case, since  $(\xi, 1) \in N_{\text{hypo}(T)}^P(x, T(x))$  there exists  $\sigma \geq 0$  such that

$$\langle \xi, z_n - x \rangle + T(z_n) - T(x) \leq \sigma (\|z_n - x\|^2 + |T(z_n) - T(x)|^2) \quad \text{for all } n. \quad (3.3.58)$$

The above inequality implies that there exists  $\sigma_1$  such that, for all  $n$  large enough,

$$T(z_n) - T(x) = |T(z_n) - T(x)| \leq \sigma_1 \|z_n - x\|. \quad (3.3.59)$$

Recalling (3.3.56) and taking  $z = x$ ,  $\beta = T(x)$ , we have, for all  $n$  large enough,

$$\begin{aligned} & \left\langle \frac{-\xi_n}{\|(-\xi_n, 1)\|}, \frac{x - z_n}{\|z_n - x\|} \right\rangle + \frac{T(x) - T(z_n)}{\|(-\xi_n, 1)\| \|z_n - x\|} \leq \\ & \leq k(T(z_n), \|z_n\|, \rho) \left( \|x - z_n\| + \frac{|T(x) - T(z_n)|^2}{\|x - z_n\|} \right). \end{aligned} \quad (3.3.60)$$

Taking  $n \rightarrow \infty$  in both (3.3.59) and (3.3.60) we obtain

$$\langle -\bar{\xi}, v_0 \rangle \geq 0, \quad (3.3.61)$$

which implies in turn that  $\langle (-\bar{\xi}, 0), (v_0, \lambda_0) \rangle \geq 0$ . Thus, combining the condition  $(-\bar{\xi}, 0) \in \tilde{N}_0(x)$  with (3.3.47), we obtain  $\frac{(-\bar{\xi}, 0)}{\|-\bar{\xi}\|} = \frac{(\xi, 1)}{\|(\xi, 1)\|}$ , a contradiction.

We can now assume that

$$\|\xi_n\| \leq Q \quad \text{for all } n, \quad (3.3.62)$$

for a suitable constant  $Q$ , and without loss of generality that

$$\lim_{n \rightarrow \infty} \xi_n = \bar{\xi}. \quad (3.3.63)$$

From (i) of Lemma 3.3.5 we have that  $-\bar{\xi} \in N_1(x)$ ,  $(-\bar{\xi}, 1) \in \tilde{N}_1(x)$ , and (3.3.41) with  $\bar{\xi}$  in place of  $\xi$  holds.

We claim that there exists a constant  $\sigma_2$  such that

$$|T(z_n) - T(x)| \leq \sigma_2 \|z_n - x\| \quad \forall n. \quad (3.3.64)$$

In the case  $T(x) < T(z_n)$ , this was already proved (see (3.3.59)).

Assume now  $T(x) \geq T(z_n)$ . Then, using (3.3.56) with  $z = x$  and  $\beta = T(x)$ , we obtain, for all  $n$  large enough,

$$\begin{aligned} & \langle -\xi_n, x - z_n \rangle + T(x) - T(z_n) \\ & \leq k(T(z_n), \|z_n\|, \rho) \|(-\xi_n, 1)\| (\|x - z_n\|^2 + |T(x) - T(z_n)|^2). \end{aligned} \quad (3.3.65)$$

The above inequality and (3.3.62) imply, for all  $n$  large enough,

$$\begin{aligned} T(x) - T(z_n) & \leq k(T(z_n), \|z_n\|, \rho) \sqrt{Q^2 + 1} (\|x - z_n\|^2 \\ & \quad + |T(x) - T(z_n)|^2) + Q \|z_n - x\|, \end{aligned}$$

from which, by the local boundedness of  $k$ , the inequality (3.3.64) follows. Summing (3.3.58) and (3.3.65) we obtain, for a suitable constant  $\sigma_3 \geq 0$  that for all  $n$  large enough

$$\left\langle \xi_n + \xi, \frac{z_n - x}{\|z_n - x\|} \right\rangle \leq \sigma_3 \left( \|z_n - x\| + \frac{|T(z_n) - T(x)|^2}{\|z_n - x\|} \right).$$

Taking  $n \rightarrow \infty$  in the above inequality and using (3.3.64), (3.3.55) we obtain

$$\langle \bar{\xi} + \xi, v_0 \rangle \leq 0,$$

or, equivalently,

$$\langle (\xi, 1), (v_0, \lambda_0) \rangle \leq \langle (-\bar{\xi}, 1), (v_0, \lambda_0) \rangle.$$

Recalling (3.3.47), we have  $\langle (\xi, 1), (v_0, \lambda_0) \rangle = 0$ , whence  $\langle (-\bar{\xi}, 1), (v_0, \lambda_0) \rangle \geq 0$ . Note that  $(-\bar{\xi}, 1) \in \tilde{N}_1(x)$ , so that  $\langle (-\bar{\xi}, 1), (v_0, \lambda_0) \rangle = 0$  by (3.3.47). Moreover, using again (3.3.47), we finally arrive to

$$\frac{(-\bar{\xi}, 1)}{\|(-\bar{\xi}, 1)\|} = \frac{(\xi, 1)}{\|(-\xi, 1)\|}.$$

Therefore we see that  $\xi = -\bar{\xi} \in N_1(x)$  and the proof is concluded.  $\square$

The lemmas contained in this section yield immediately the following result.

**Theorem 3.3.1** *Let  $x \in \mathcal{S}^c$  and let the assumptions of Theorem 3.2.1 hold. Then*

$$N_{\text{hypo}(T)}^P(x, T(x)) = N(x),$$

where  $N(x)$  was defined in the statement of Lemma 3.3.6, so that  $N_{\text{hypo}(T)}^P(x, T(x))$  is a closed (convex) cone.

**Proof.** Assume by contradiction that there exists  $\zeta \in N_{\text{hypo}(T)}^P(x, T(x)) \setminus N(x)$ . Set

$$\tilde{N} = \text{co}(N(x) \cup \{\lambda\zeta \mid \lambda \geq 0\})$$

and observe that  $\tilde{N}$  is a closed convex cone which satisfies (3.3.46). Clearly,  $\zeta$  belongs to an exposed ray of  $\tilde{N}$ , so that, by Lemma 3.3.7,  $\zeta \in \tilde{N}_0(x) \cup \tilde{N}_1(x)$ , a contradiction.  $\square$

## 3.4 Proof of the main results of Chapter 3

### 3.4.1 Proof of Theorem 3.2.1

It is clear that the “ $\supseteq$ ” inclusion in (3.2.2) follows from Lemma 3.3.2 and the convexity of  $\partial^\infty T(x)$ .

In order to prove the “ $\subseteq$ ” inclusion, take  $\xi \in \partial^\infty T(x)$ , i.e.,  $(-\xi, 0) \in N_{\text{hypo}(T)}^P(x, T(x))$ . Since  $N_{\text{hypo}(T)}^P(x, T(x))$  is wedged and closed (see Theorem 3.3.1), recalling (2.2.3) we can find numbers  $\alpha_i, \beta_i \geq 0$  and vectors  $\xi_i, \zeta_i \in \mathbb{R}^N, i \in \{1, \dots, N+2\}$ , such that

$$\begin{cases} (-\xi_i, 1) & \text{belongs to an exposed ray of } N_{\text{hypo}(T)}^P(x, T(x)) \\ (-\zeta_i, 0) & \text{belongs to an exposed ray of } N_{\text{hypo}(T)}^P(x, T(x)) \\ (-\xi, 0) & = \sum_{i=1}^{N+2} \alpha_i (-\xi_i, 1) + \sum_{i=1}^{N+2} \beta_i (-\zeta_i, 0). \end{cases} \quad (3.4.1)$$

From the above equality we deduce that  $\alpha_i = 0$  for all  $i \in \{1, \dots, N+2\}$ . Thus, we have

$$(-\xi, 0) = \sum_{i=1}^{N+2} \beta_i (-\zeta_i, 0). \quad (3.4.2)$$

Recalling (i) in Lemma 3.3.7 we obtain  $-\zeta_i \in N_0(x)$ . Setting  $\bar{\zeta}_i = (\sum_{j=1}^{N+2} \beta_j) \zeta_i$  and  $\bar{\beta}_i = \frac{\beta_i}{\sum_{i=1}^{N+2} \beta_i}$ , one can easily see  $-\bar{\zeta}_i \in N_0(x)$  and  $\sum_{i=1}^{N+2} \bar{\beta}_i = 1$ .

From (3.4.2), we obtain

$$\xi = - \sum_{i=1}^{N+2} \bar{\beta}_i (-\bar{\zeta}_i).$$

The proof is concluded.  $\square$

**Proof of Theorem 3.2.2.** Observe that from the very definition it follows that if  $\xi \in \partial^P T(x)$  and  $\zeta \in \partial^\infty T(x)$  then  $\xi + \zeta \in \partial^P T(x)$ . Thus the “ $\supseteq$ ” inclusion in (3.2.3) follows from Lemma 3.3.3, Lemma 3.3.2 and the above observation.

In order to prove the “ $\subseteq$ ” inclusion, take  $\xi \in \partial^P T(x)$ , i.e.  $(-\xi, 1) \in N_{\text{hypo}(T)}^P(x, T(x))$ . Since  $N_{\text{hypo}(T)}^P(x, T(x))$  is wedged and closed (see Theorem 3.3.1), recalling (2.2.3) we can find numbers  $\alpha_i, \beta_i \geq 0$  and vectors  $\xi_i, \zeta_i \in \mathbb{R}^N, i \in \{1, \dots, N+2\}$ , such that

$$\begin{cases} (-\xi_i, 1) & \text{belongs to an exposed ray of } N_{\text{hypo}(T)}^P(x, T(x)) \\ (-\zeta_i, 0) & \text{belongs to an exposed ray of } N_{\text{hypo}(T)}^P(x, T(x)) \\ (-\xi, 1) & = \sum_{i=1}^{N+2} \alpha_i (-\xi_i, 1) + \sum_{i=1}^{N+2} \beta_i (-\zeta_i, 0). \end{cases} \quad (3.4.3)$$

From the above equality we deduce that  $\sum_{i=1}^{N+2} \alpha_i = 1$ . Thus, recalling (ii) in Lemma 3.3.7 we obtain that  $\sum_{i=1}^{N+2} \alpha_i (-\xi_i) \in \text{co}(N_1(x))$ .

On the other hand, arguing similarly to the above proof we see that  $\sum_{i=1}^{N+2} \beta_i (-\zeta_i) \in \text{co}(N_0(x))$ . Therefore,

$$\xi = - \left( \sum_{i=1}^{N+2} \alpha_i (-\xi_i) + \sum_{i=1}^{N+2} \beta_i (-\zeta_i) \right) \in -[\text{co}(N_1(x)) + \text{co}(N_0(x))].$$

The proof is concluded.  $\square$

### 3.4.2 Proof of Theorem 3.2.3

We need the following technical lemma.

**Lemma 3.4.1** *Assume that  $N_{\text{hypo}(T)}^P(x, T(x))$  is wedged for all  $x \in \mathcal{S}^c$ . Then for each continuous function  $\theta : \mathcal{S}^c \rightarrow [0, \infty)$ , there exists a continuous function  $\psi_\theta : \mathcal{S}^c \rightarrow (0, 1]$  such that*

$$\langle \zeta_1, \zeta_2 \rangle \geq \psi_\theta(x) - 1 \quad (3.4.4)$$

for all  $x \in \mathcal{S}^c$  and for all  $\zeta_1, \zeta_2 \in N_{\text{hypo}(T)}^P(x, T(x))$  satisfying both  $\|\zeta_1\| = \|\zeta_2\| = 1$  and

$$\langle \zeta_j, (z - x, \beta - T(x)) \rangle \leq \theta(x) (\|z - x\|^2 + |\beta - T(x)|^2) \quad (3.4.5)$$

for all  $z \in \mathcal{S}^c, \beta \leq T(x)$ , and  $j = 1, 2$ .

**Proof.** We only need to show that for every  $n \in \mathbb{N}$  there exists a continuous function  $\psi_n : \overline{B(0, n)} \cap \mathcal{S}^c \rightarrow (0, 1]$  satisfying (3.4.4) with  $\psi_\theta(x)$  replaced by  $\psi_n(x)$ . It is easy to see that the following statement is sufficient to this aim.

Let, for all  $m, n \in \mathbb{N}$ ,  $\mathcal{K}_n^m = \overline{B(0, n)} \cap \mathcal{S}^c(\frac{1}{m})$ , and observe that, by the continuity of  $T(\cdot)$ ,  $\mathcal{K}_n^m$  is compact. Fix  $n$ . We claim that for each  $m \in \mathbb{N}$  there exists a constant  $k_m \in (0, 1]$  such that

$$\langle \zeta_1, \zeta_2 \rangle \geq k_m - 1, \quad (3.4.6)$$

for all  $x \in \mathcal{K}_n^m$ ,  $\zeta_1, \zeta_2 \in N_{\text{hypo}(T)}^P(x, T(x))$  satisfying  $\|\zeta_1\| = \|\zeta_2\| = 1$  and (3.4.5).

Indeed, assume by contradiction that there exists a sequence  $\{x_i\} \subset \mathcal{K}_n^m$  together with vectors  $\zeta_1^i, \zeta_2^i \in N_{\text{hypo}(T)}^P(x_i, T(x_i))$  satisfying  $\|\zeta_1^i\| = \|\zeta_2^i\| = 1$  and

$$\langle \zeta_j^i, (z - x_i, \beta - T(x_i)) \rangle \leq \theta(x_i)(\|z - x_i\|^2 + |\beta - T(x_i)|^2), \quad (3.4.7)$$

for all  $z \in \mathcal{S}^c$ ,  $\beta \leq T(x_i)$  and  $j \in \{1, 2\}$ , but such that

$$\lim_{i \rightarrow \infty} \langle \zeta_1^i, \zeta_2^i \rangle = -1. \quad (3.4.8)$$

We can assume without loss of generality that  $\{x_i\}$ ,  $\{\zeta_1^i\}$  and  $\{\zeta_2^i\}$  converge respectively to  $\bar{x} \in \mathcal{K}_n^m$ ,  $\bar{\zeta}_1$  and  $\bar{\zeta}_2$ . By the continuity of  $T(\cdot)$ ,  $\theta(\cdot)$  and (3.4.7) we obtain

$$\bar{\zeta}_i \in N_{\text{hypo}(T)}^P(\bar{x}, T(\bar{x})) \quad \text{for } i \in \{1, 2\}.$$

On the other hand, from  $\|\zeta_1^i\| = \|\zeta_2^i\| = 1$  and (3.4.8) we get

$$\bar{\zeta}_1 = -\bar{\zeta}_2.$$

But then the normal cone  $N_{\text{hypo}(T)}^P(\bar{x}, T(\bar{x}))$  contains a line, and this is a contradiction.  $\square$

*End of the proof of Theorem 3.2.3.*

We need to find a continuous function  $\varphi : \mathcal{S}^c \rightarrow [0, \infty)$  such that for all  $x \in \mathcal{S}^c$ ,  $\zeta \in N_{\text{hypo}(T)}^P(x, T(x))$  and for all  $z \in \mathcal{S}^c$ ,  $\beta \leq T(z)$  we have

$$\langle \zeta, (z - x, \beta - T(x)) \rangle \leq \varphi(x) \|\zeta\| (\|z - x\|^2 + |\beta - T(x)|^2). \quad (3.4.9)$$

Observe that for every  $\zeta \in N_{\text{hypo}(T)}^P(x, T(x))$ , by the wedgedness assumption and recalling Theorem 3.3.1, we have

$$\zeta = \sum_{i=1}^{N+2} \zeta_i, \quad (3.4.10)$$

where each  $\zeta_i$  belongs to an exposed ray of  $N_{\text{hypo}(T)}^P(x, T(x))$ . For  $k \in \{1, 2, \dots, N+2\}$ , we set

$$N_k^P(x) = \left\{ \zeta \mid \zeta = \sum_{i=1}^k \zeta_i, \right. \\ \left. \text{where } \zeta_i \text{ belongs to an exposed ray of } N_{\text{hypo}(T)}^P(x, T(x)) \right\}. \quad (3.4.11)$$

Of course  $N_k^P(x) \subseteq N_{\text{hypo}(T)}^P(x, T(x))$  and  $N_{N+2}^P(x) = N_{\text{hypo}(T)}^P(x, T(x))$ .

Now, we are going to construct by induction a continuous function  $\varphi_k(\cdot)$  such that

$$\left\langle \zeta^k, (z - x, \beta - T(x)) \right\rangle \leq \varphi_k(x) \left\| \zeta^k \right\| (\|z - x\|^2 + |\beta - T(x)|^2), \quad (3.4.12)$$

for all  $x \in \mathcal{S}^c$ ,  $\zeta^k \in N_k^P(x)$  and for all  $z \in \mathcal{S}^c$ ,  $\beta \leq T(z)$ .

For  $k = 1$  we choose  $\varphi_1(x) := k(T(x), \|x\|, \rho)$ . Recalling Lemma 3.3.7 and Lemma 3.3.3, 3.3.2, we obtain that for all  $\zeta^1 \in N_1^P(x)$  and for all  $z \in \mathcal{S}^c$ ,  $\beta \leq T(z)$

$$\left\langle \zeta^1, (z - x, \beta - T(x)) \right\rangle \leq \varphi_1(x) \left\| \zeta^1 \right\| (\|z - x\|^2 + |\beta - T(x)|^2). \quad (3.4.13)$$

Thus (3.4.12) holds.

Assume now that (3.4.12) is satisfied for  $k = h \geq 1$ . We want to show that (3.4.12) holds for  $k = h + 1$ , with

$$\varphi_{h+1}(x) = \sqrt{\frac{\varphi_h(x)^2 + \varphi_1(x)^2}{\psi_{\max\{\varphi_1, \varphi_h\}}(x)}}, \quad (3.4.14)$$

where the function  $\psi_{\max\{\varphi_1, \varphi_h\}}(\cdot)$  is given by Lemma 3.4.1 for  $\theta(\cdot) = \max\{\varphi_1(\cdot), \varphi_h(\cdot)\}$ . Indeed, given  $\zeta^{h+1} \in N_{h+1}^P(x)$ , one can write

$$\zeta^{h+1} = \zeta^h + \zeta^1,$$

where  $\zeta^h \in N_h^P(x)$  and  $\zeta^1 \in N_1^P(x)$ . From (3.4.13) and the inductive assumption, one can easily see that

$$\left\langle \zeta^{h+1}, (z - x, \beta - T(x)) \right\rangle \\ \leq \left( \varphi_1(x) \left\| \zeta^1 \right\| + \varphi_h(x) \left\| \zeta^h \right\| \right) (\|z - x\|^2 + |\beta - T(x)|^2), \quad (3.4.15)$$

for all  $z \in \mathcal{S}^c$ ,  $\beta \leq T(z)$ .

On the other hand, by inductive assumption, (3.4.13) and Lemma 3.4.1 applied for  $\theta(\cdot) = \max\{\varphi_1(\cdot), \varphi_h(\cdot)\}$ , we obtain

$$\left\langle \frac{\zeta^h}{\left\| \zeta^h \right\|}, \frac{\zeta^1}{\left\| \zeta^1 \right\|} \right\rangle \geq \psi_{\max\{\varphi_1, \varphi_h\}}(x) - 1.$$



Thus, since  $\psi(x) \in (0, 1]$ , we see that

$$\|\zeta^h + \zeta^1\|^2 \geq \psi_{\max\{\varphi_1, \varphi_h\}}(x) \left( \|\zeta^h\|^2 + \|\zeta^1\|^2 \right).$$

Therefore,

$$\|\zeta^h + \zeta^1\|^2 \geq \frac{\psi_{\max\{\varphi_1, \varphi_h\}}(x)}{\varphi_h(x)^2 + \varphi_1(x)^2} \left( \varphi_h(x) \|\zeta^h\| + \varphi_1(x) \|\zeta^1\| \right)^2.$$

Combining the above inequality, (3.4.14) and (3.4.15) we obtain that

$$\left\langle \zeta^{h+1}, (z - x, \beta - T(x)) \right\rangle \leq \varphi_{h+1}(x) \|\zeta^{h+1}\| \left( \|z - x\|^2 + |\beta - T(x)|^2 \right),$$

for all  $z \in \mathcal{S}^c$ ,  $\beta \leq T(z)$ .

To conclude the proof, we choose  $\varphi(\cdot) = \varphi_{N+2}(\cdot)$ .  $\square$

### 3.5 The case of optimal points

This section is devoted to the representation of *supergradient* and *horizontal gradient* at optimal points. The corresponding formulas are easier than in the general case and the structure of the Hamiltonian exhibits special properties.

The definition of optimal points here is based on the classical definition (see, e.g., Definition 2.24, p. 119 in [11]), but is adapted to *limiting optimal trajectories*, since optimal trajectories may not exist.

**Definition 3.5.1** *Let  $x \in \mathcal{S}^c$  and set  $r = T(x)$ . The point  $x$  is called an optimal point if there exist  $\tau > 0$  and  $x_\tau \in \mathcal{S}^c$  such that*

(i)  $T(x_\tau) = r + \tau$ ;

(ii) *there exist  $\bar{x}_\tau \in \mathcal{S}_{x_\tau}$  and  $\{\bar{u}_n\} \subset \mathcal{U}_{\bar{x}_\tau}$ , together with the corresponding sequence  $x_n \rightarrow x_\tau$ , such that  $y^{x_n, \bar{u}_n}(\tau) \rightarrow x$ .*

At optimal points, the Hamiltonian has a special behavior. More precisely, let  $x$  be an optimal point with  $T(x) = r > 0$ . Then the Hamiltonian  $H(x, \cdot)$  is additive on the proximal normal cone to  $\mathcal{S}^c(r)$ . It follows from this property that the *supergradient* and *horizontal supergradient* of  $T$  are contained, respectively, in the 1-level set and the 0-level set of the Hamiltonian.

**Theorem 3.5.1** *Let  $x \in \mathcal{S}^c$  be an optimal point. Under the same assumptions of Theorem 3.2.1, the (proximal) horizontal gradient and the supergradient of the minimum time function  $T(\cdot)$  at the point  $x$  can be computed as follows:*

(a)  $\partial^\infty T(x) = [-\text{co}(N(x))] \cap \{-\xi \mid H(\xi, x) = 0\},$

$$(b) \partial^P T(x) = [-\text{co}(N(x))] \cap \{-\xi \mid H(\xi, x) = 1\},$$

where

$$N(x) = \{M^T(r)v \mid M(\cdot) \in \mathcal{M}_{\bar{x}}, v \in N_{\mathcal{S}^c}^P(\bar{x}), \bar{x} \in \mathcal{S}_x\}. \quad (3.5.1)$$

The proof of Theorem 3.5.1 requires some preliminary lemmas. The first one gives an information on a lower bound of the Hamiltonian computed at a proximal normal of the sublevel of  $T$  at an optimal point.

**Lemma 3.5.1** *Let  $x \in \mathcal{S}^c$  be an optimal point, and let  $\xi \in N_{\mathcal{S}^c(T(x))}^P(x)$ . Then  $H(x, \xi) \geq 0$ .*

**Proof.** Set  $r = T(x)$ . Let  $\tau, x_\tau, \bar{x}_\tau, \bar{u}_n$  and  $x_n$  be with the properties stated in Definition 3.5.1. To simplify our writing, we set  $\gamma_n(\cdot) = y^{x_n, \bar{u}_n}(\cdot)$ . Assuming without loss of generality that  $\gamma_n(\cdot)$  converges uniformly to  $\gamma(\cdot)$ , one can easily check that  $\gamma(t) \in \mathcal{S}^c(r)$  for all  $t \in [0, \tau]$ . For, should  $\bar{t} \in (0, \tau]$  exist such that  $T(\gamma(\bar{t})) < r$ , then one would have  $T(x_\tau) < r + \tau$ , a contradiction. Now, since  $\xi \in N_{\mathcal{S}^c(r)}^P(x)$  there exists  $\sigma > 0$  such that for all  $t \in [0, \tau]$  we have

$$\langle \xi, \gamma(t) - x \rangle \leq \sigma \|\gamma(t) - x\|^2, \quad (3.5.2)$$

namely, for all  $t \in [0, \tau]$ ,

$$\lim_{n \rightarrow \infty} \langle \xi, \gamma_n(t) - x \rangle \leq \sigma \lim_{n \rightarrow \infty} \|\gamma_n(t) - x\|^2.$$

Equivalently, for all  $t \in [0, \tau]$

$$\begin{aligned} \lim_{n \rightarrow \infty} \left\langle \xi, \gamma_n(\tau) - \int_{\tau-t}^{\tau} f(\gamma_n(s), \bar{u}_n(s)) ds - x \right\rangle &\leq \\ &\leq \sigma \lim_{n \rightarrow \infty} \left\| \gamma_n(\tau) - \int_{\tau-t}^{\tau} f(\gamma_n(s), \bar{u}_n(s)) ds - x \right\|^2. \end{aligned}$$

Recalling (ii) in Definition 3.5.1, we obtain that for all  $t \in [0, \tau]$

$$\lim_{n \rightarrow \infty} \left\langle \xi, - \int_{\tau-t}^{\tau} f(\gamma_n(s), \bar{u}_n(s)) ds \right\rangle \leq \sigma \lim_{n \rightarrow \infty} \left\| \int_{\tau-t}^{\tau} f(\gamma_n(s), \bar{u}_n(s)) ds \right\|^2.$$

From (iii) of Lemma 7.1.1 and (i), (ii) in Definition 3.5.1, one can see that

$$\lim_{n \rightarrow \infty} \left\langle \xi, - \int_{\tau-t}^{\tau} f(\gamma_n(s), \bar{u}_n(s)) ds \right\rangle \leq O(t^2) \text{ for } t \rightarrow 0^+.$$

Thus, for  $t \rightarrow 0^+$ ,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \left\langle \xi, - \int_{\tau-t}^{\tau} f(x, \bar{u}_n(s)) ds \right\rangle \\ \leq O(t^2) + \limsup_{n \rightarrow \infty} \left\langle \xi, \int_{\tau-t}^{\tau} (f(\gamma_n(s), \bar{u}_n(s)) - f(x, \bar{u}_n(s))) ds \right\rangle. \end{aligned}$$

Applying the Lipschitz condition of the function  $f(\cdot, \cdot)$  and (iii) of Lemma 7.1.1 we easily obtain that

$$\limsup_{n \rightarrow \infty} \left\langle \xi, - \int_{\tau-t}^{\tau} f(x, \bar{u}_n(s)) ds \right\rangle \leq O(t^2) \quad \text{for } t \rightarrow 0^+.$$

Therefore, there exists a constant  $Q > 0$  such that for each  $t \in [0, \tau]$  one can find  $n_t \in \mathbb{N}$  with the property

$$\left\langle \xi, - \frac{\int_{\tau-t}^{\tau} f(x, \bar{u}_{n_t}(s)) ds}{t} \right\rangle \leq Qt.$$

Set  $\bar{f}_t = \frac{\int_{\tau-t}^{\tau} f(x, \bar{u}_{n_t}(s)) ds}{t}$ . Since  $\bar{f}_t \in \text{co}(f(x, \mathcal{U}))$ , by the compactness of  $\mathcal{U}$ , there exists a sequence  $\{t_n\} \subseteq [0, \tau]$  converging to 0 and  $\bar{f} \in \text{co}f(x, \mathcal{U})$  such that both

$$\bar{f} = \lim_{n \rightarrow \infty} \bar{f}_{t_n}$$

and

$$\langle \xi, \bar{f} \rangle \geq 0$$

hold. Since

$$H(x, \xi) = \max\{\langle \xi, f \rangle \mid f \in \text{co}f(x, \mathcal{U})\},$$

the proof is concluded.  $\square$

The next Lemma is the key point in order to obtain the additivity of the Hamiltonian.

**Lemma 3.5.2** *Let  $x \in \mathcal{S}^c$  be an optimal point, and set  $T(x) = r$ . Then there exists  $\bar{f} \in \text{co}f(x, \mathcal{U})$  such that, for all  $\xi \in N_{\mathcal{S}^c(r)}^P(x)$ ,*

$$H(x, \xi) = \langle \xi, \bar{f} \rangle.$$

**Proof.** Let  $\tau, x_\tau, \bar{x}_\tau, \bar{u}_n$  and  $x_n$  be with the properties stated in Definition 3.5.1.

To simplify our writing, we set  $\gamma_n(\cdot) = y^{x_n, \bar{u}_n}(\cdot)$ . Assuming without loss of generality that  $\gamma_n(\cdot)$  converges uniformly to  $\gamma(\cdot)$ , we see that  $\gamma(\tau) = x$  and  $T(\gamma(\tau - t)) = r + t$  for all  $t \in [0, \tau]$ . Pick  $v \in \mathcal{U}$ , and define, for each  $t \in [0, \tau]$ ,  $\beta_{v,t}(\cdot) = y^{\gamma(\tau-t), v}(\cdot)$ , where  $v(\cdot)$  is the constant control  $v(t) \equiv v$ . Observe that  $\beta_{v,t}(t) \in \mathcal{S}^c(r)$  for all  $t \in [0, \tau]$ .

Let now  $\xi \in N_{\mathcal{S}^c(r)}^P$ , together with a constant  $\sigma \geq 0$  such that for all  $t \in [0, \tau]$

$$\langle \xi, \beta_{v,t}(t) - x \rangle \leq \sigma \|\beta_{v,t}(t) - x\|^2.$$

Recalling (ii) in Definition 3.5.1, the latter is equivalent to

$$\begin{aligned} & \lim_{n \rightarrow \infty} \left\langle \xi, \int_0^t (f(\beta_{v,t}(s), v) - f(\gamma_n(\tau - t + s), \bar{u}_n(\tau - t + s))) ds \right\rangle \\ & \leq \sigma \lim_{n \rightarrow \infty} \left\| \int_0^t (f(\beta_{v,t}(s), v) - f(\gamma_n(\tau - t + s), \bar{u}_n(\tau - t + s))) ds \right\|^2 \end{aligned} \quad (3.5.3)$$

for all  $t \in [0, \tau]$ . Moreover, by (iii) of Lemma 7.1.1, there exists a constant  $M$  such that, for all  $n \in \mathbb{N}$ ,  $t \in [0, \tau]$  and  $s \in [0, t]$ ,

$$\|\gamma_n(\tau - t + s) - \gamma_n(\tau)\| \leq Mt,$$

so that for all  $t \in [0, \tau]$  and  $s \in [0, t]$

$$\lim_{n \rightarrow \infty} \|\gamma_n(\tau - t + s) - x\| \leq Mt.$$

Combining the above inequality with (3.5.3) and recalling the Lipschitz condition on  $f$ , we obtain that, for  $t \rightarrow 0^+$ ,

$$\limsup_{n \rightarrow \infty} \left\langle \xi, \int_0^t (f(x, v) - f(x, \bar{u}_n(r - t + s))) ds \right\rangle \leq O(t^2),$$

or, equivalently,

$$\limsup_{n \rightarrow \infty} \left\langle \xi, f(x, v) - \frac{\int_0^t f(x, \bar{u}_n(r - t + s)) ds}{t} \right\rangle \leq O(t).$$

By arguing as in the proof of Lemma 3.5.1, we can find  $\bar{f} \in \text{co}(f(x, U))$  independent of  $\xi$  and  $v$  such that

$$\langle \xi, f(x, v) \rangle \leq \langle \xi, \bar{f} \rangle.$$

The proof is therefore complete.  $\square$

The desired additivity property follows immediately from the above Lemma.

**Corollary 3.5.1** *Let  $x \in \mathcal{S}^c$  be an optimal point, and set  $T(x) = r$ . Then for all  $\xi_1, \xi_2 \in N_{\mathcal{S}^c(r)}^P(x)$ , the property*

$$H(x, \xi_1 + \xi_2) = H(x, \xi_1) + H(x, \xi_2)$$

*holds.*

We are now ready to prove Theorem 3.5.1.

**Proof of Theorem 3.5.1.**

*Proof of part a).* It is clear that the “ $\subseteq$ ” inclusion of the equality in (a) follows from Theorem 3.2.1 and Corollary 3.5.1.

To prove the “ $\supseteq$ ” inclusion, take  $\xi \in [-\text{co}(N(x))] \cap \{-\xi \mid H(x, \xi) = 0\}$ , namely,

$$\xi = - \sum_{i=1}^m M_i^T(r) v_i, \quad \text{where } M_i^T(r) v_i \in N(x) \quad (3.5.4)$$

and

$$H(x, \sum_{i=1}^m M_i^T(r) v_i) = 0. \quad (3.5.5)$$

Applying Lemma 3.3.1 we get that  $M_i^T(r)v_i \in N_{\mathcal{S}^c(r)}^P(x)$  for all  $i \in \{1, 2, \dots, m\}$ . Thus it follows from Lemma 3.5.1 that

$$H(x, M_i^T(r)v_i) \geq 0 \quad \text{for all } i \in \{1, 2, \dots, m\}. \quad (3.5.6)$$

Combining (3.5.5) and (3.5.6), we obtain from Corollary 3.5.1 that for all  $i \in \{1, 2, \dots, m\}$ ,  $H(x, M_i^T(r)v_i) = 0$ . Therefore  $M_i^T(r)v_i \in N_0(x)$  for all  $i \in \{1, 2, \dots, m\}$ . We conclude the proof using (3.5.4) and Theorem 3.2.1.

*Proof of part b).* Similarly to part (a), that the “ $\subseteq$ ” inclusion of the equality in (b) follows from Theorem 3.2.2 and Corollary 3.5.1.

To show the “ $\supseteq$ ” inclusion, let  $\xi \in [-\text{co}(N(x))] \cap \{-\xi \mid H(x, \xi) = 1\}$ . Recalling Lemma 3.5.1,  $\xi$  can be represented as

$$\xi = -\sum_{i=1}^m \alpha_i M_{0i}^T(r)v_i - \sum_{j=1}^m \beta_j M_{1j}^T(r)w_j, \quad (3.5.7)$$

where  $\alpha_i \geq 0$ ,  $\beta_j \geq 0$  and  $M_{0i}^T(r)v_i \in N_0(x)$ ,  $M_{1j}^T(r)w_j \in N_1(x)$ .

From  $M_{0i}^T(r)v_i \in N_{\mathcal{S}^c(r)}^P(x)$ ,  $M_{1j}^T(r)w_j \in N_{\mathcal{S}^c(r)}^P(x)$ , and Corollary 3.5.1, we have

$$H(x, \xi) = \sum_{i=1}^m \alpha_i H(x, M_{0i}^T(r)v_i) + \sum_{j=1}^m \beta_j H(x, M_{1j}^T(r)w_j) = \sum_{j=1}^m \beta_j, \quad (3.5.8)$$

so that  $\sum_{j=1}^m \beta_j = 1$ . The proof is concluded by using (3.5.8), (3.5.7), and Theorem 3.2.2.  $\square$

## 3.6 Examples

In this section we present some examples which illustrate our results. In particular, we provide an example showing that Theorem 3.2.3 is no longer valid if the wedgedness assumption (3.2.1) is dropped.

**Example 1.** Consider the dynamics  $x''(\cdot) \in [-1, 1] =: \mathcal{U}$ , i.e.

$$\begin{pmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{pmatrix} = A \begin{pmatrix} x_1(t) \\ x_2(t) \end{pmatrix} + \begin{pmatrix} 0 \\ u \end{pmatrix}, \quad u \in \mathcal{U}, \quad \text{where } A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad (3.6.1)$$

with the initial conditions  $x_1(0) = x_1^0$ ,  $x_2(0) = x_2^0$ . The target is the set (see Figure 1)

$$\begin{aligned} \mathcal{S} &= \{ (x_1, x_2) \in \mathbb{R}^2 \mid x_1 \leq 0 \} \cup \{ (x_1, x_2) \in \mathbb{R}^2 \mid x_1 \geq 0, x_2 \leq -x_1 \} \\ &\cup \{ (x_1, x_2) \in \mathbb{R}^2 \mid 0 \leq x_1 \leq 1, x_2 \geq x_1 \} \\ &\cup \{ (x_1, x_2) \in \mathbb{R}^2 \mid x_1 \geq 1, x_2 \geq 1 \}. \end{aligned}$$




Figure 1

Optimal trajectories are arcs of parabolas

$$x_1 = \frac{1}{2}(x_2)^2 - \frac{1}{2}(x_2^0)^2 + x_1^0 \quad (\text{corresponding to the control } u \equiv 1),$$

and

$$x_1 = -\frac{1}{2}(x_2)^2 + \frac{1}{2}(x_2^0)^2 + x_1^0 \quad (\text{corresponding to the control } u \equiv -1).$$

By direct computation, the minimum time function  $T$  is everywhere finite, continuous on the whole of  $\mathbb{R}^2$ , and the open set  $S^c$  can be divided into three regions, say  $H_1$ ,  $H_2$  and  $H_3$ , where  $T$  has a different explicit expression. More precisely, consider the curves

$$\begin{aligned} \gamma_1(t) &= \left( \sqrt{2t}(1-t), t \right), & 0 < t \leq 2 - \sqrt{3}, \\ \gamma_2(t) &= \left( \frac{1+t^2}{2}, t \right), & 2 - \sqrt{3} < t < 1, \\ \gamma_3(t) &= \left( \frac{3-8t+3t^2}{2}, t \right), & t \geq 2 - \sqrt{3}. \end{aligned}$$

Observe that  $\gamma_1(2 - \sqrt{3}) = \gamma_2(2 - \sqrt{3}) = \gamma_3(2 - \sqrt{3}) = 4 - 2\sqrt{3}$  and moreover all points  $\gamma_2(t)$ , with  $2 - \sqrt{3} < t < 1$ , are optimal (according to Definition

3.5.1), while all points  $\gamma_1(t)$ ,  $\gamma_2(t)$  are not optimal. Set

$$\begin{aligned} H_1 &= \{(x_1, x_2) \in \mathcal{S}^c \mid 0 \leq x_2 \leq 2 - \sqrt{3}, \gamma_1(x_2) \leq x_1 \leq \gamma_3(x_2)\} \\ &\cup \{(x_1, x_2) \in \mathcal{S}^c \mid x_2 \leq 0, -x_2 \leq \gamma_3(x_2)\}, \end{aligned}$$

$$\begin{aligned} H_2 &= \{(x_1, x_2) \in \mathcal{S}^c \mid 0 \leq x_2 \leq 2 - \sqrt{3}, x_2 \leq x_1 \leq \gamma_1(x_2)\} \\ &\cup \{(x_1, x_2) \in \mathcal{S}^c \mid 2 - \sqrt{3} \leq x_2 \leq 1, x_2 \leq x_1 \leq \gamma_2(x_2)\}, \end{aligned}$$

$$\begin{aligned} H_3 &= \{(x_1, x_2) \in \mathcal{S}^c \mid 2 - \sqrt{3} \leq x_2 \leq 1, x_1 \geq \gamma_2(x_2)\} \\ &\cup \{(x_1, x_2) \in \mathcal{S}^c \mid x_2 \leq 2 - \sqrt{3}, x_1 \geq \gamma_3(x_2)\}. \end{aligned}$$

The minimum time function  $T : \mathcal{S}^c \rightarrow \mathbb{R}$  can be explicitly computed as

$$T(x_1, x_2) = \begin{cases} x_2 - 1 + \sqrt{1 + 2x_1 + (x_2)^2} & := \theta_1(x_1, x_2), & (x_1, x_2) \in H_1 \\ 1 - x_2 - \sqrt{1 - 2x_1 + (x_2)^2} & := \theta_2(x_1, x_2), & (x_1, x_2) \in H_2 \\ 1 - x_2 & := \theta_3(x_1, x_2), & (x_1, x_2) \in H_3. \end{cases}$$

In the interior of each region  $H_i$ ,  $i = 1, 2, 3$ ,  $T$  is differentiable. Singularities appear of each point of the curves  $\gamma_i$ ,  $i = 1, 2, 3$ . Moreover  $T$  is Hölder continuous with exponent  $\frac{1}{2}$ .

In order to appreciate the role of nonsmoothness of the target, as well as optimality/non optimality of a point and failure of Petrov's condition, we compute the generalized differential of  $T$  at the three points

$$P_1 = \left(\frac{7}{16}, \frac{1}{8}\right), \quad P_2 = \left(\frac{5}{8}, \frac{1}{2}\right), \quad P_3 = \left(4 - 2\sqrt{3}, 2 - \sqrt{3}\right).$$

Observe that  $T(P_1) = \frac{1}{2}$ ,  $T(P_2) = \frac{1}{2}$ ,  $T(P_3) = \sqrt{3} - 1$ .

To this aim we compute the adjoint flow:

$$e^{A^T t} = \begin{pmatrix} 1 & 0 \\ t & 1 \end{pmatrix},$$

and the Hamiltonian

$$H((x_1, x_2), (\xi_1, \xi_2)) = x_2 \xi_1 + |\xi_2|.$$

The point  $P_1$  belongs to the curve  $\gamma_1$ , and is steered optimally in time  $\frac{1}{2}$  to both  $(\frac{5}{8}, \frac{5}{8})$  and  $(\frac{3}{8}, -\frac{3}{8})$ , where the normal cones to  $\mathcal{S}^c$  are respectively  $\mathbb{R}^+(-1, 1)$  and  $\mathbb{R}^+(-1, -1)$ , while  $P_2$  belongs to the curve  $\gamma_2$ , and is steered optimally to  $(1, 1)$  in time  $\frac{1}{2}$ , where the normal cone to  $\overline{\mathcal{S}^c}$  is  $\mathbb{R}^+ \text{co}\{(-1, 1), (0, 1)\}$ .  $P_2$  is an optimal point. Finally,  $P_3$  is steered optimally to both  $(2\sqrt{3} - 3, 3 - 2\sqrt{3})$  and  $(1, 1)$  in time  $\sqrt{3} - 1$ . Observe that  $H((1, 1), (-1, 1)) = 0$ , i.e., Petrov's condition fails, while at all other (nonzero) points  $P$  of the boundary of  $\mathcal{S}$  we have  $H(P, \zeta) > 0$  for all

$\zeta \in N_{\mathcal{S}^c}^P(P)$ ,  $\zeta \neq 0$ .

According to Theorem 3.2.2, and, of course, also to explicit computations from the expression of  $T$ , we have

$$\begin{aligned} \partial^c T(P_1) &= \partial^P T(P_1) \\ &= -\text{co} \left\{ \begin{array}{l} e^{A^T \frac{1}{2} v} \mid v = \begin{pmatrix} -\lambda \\ \lambda \end{pmatrix} \\ \text{or } v = \begin{pmatrix} -\lambda \\ -\lambda \end{pmatrix}, H(P_1, e^{A^T \frac{1}{2} v}) = 1 \end{array} \right\} \\ &= -\text{co} \left\{ \begin{pmatrix} \frac{8}{3} \\ -\frac{4}{3} \end{pmatrix}, \begin{pmatrix} \frac{8}{11} \\ -\frac{12}{11} \end{pmatrix} \right\}, \\ \partial^\infty T(P_1) &= \{0\}; \end{aligned}$$

$$\begin{aligned} \partial^c T(P_2) &= \partial^P T(P_2) \\ &= -\text{co} \left\{ e^{A^T \frac{1}{2} v} \mid v \in N_{\mathcal{S}^c}^P(1, 1), H(P_2, e^{A^T \frac{1}{2} v}) = 1 \right\} \\ &\quad -\text{co} \left\{ e^{A^T \frac{1}{2} v} \mid v \in N_{\mathcal{S}^c}^P(1, 1), H(P_2, e^{A^T \frac{1}{2} v}) = 0 \right\} \\ &= \begin{pmatrix} 0 \\ -1 \end{pmatrix} + \left\{ \lambda \begin{pmatrix} 1 \\ -\frac{1}{2} \end{pmatrix} \mid \lambda \geq 0 \right\} \\ &= \left\{ \begin{pmatrix} \lambda \\ -1 - \frac{\lambda}{2} \end{pmatrix} \mid \lambda \geq 0 \right\} \\ &= [-N(P_2)] \cap \left\{ \zeta \mid H(P_2, -\zeta) = 1 \right\} \\ &\quad (\text{where } N(P_2) \text{ was defined in (3.5.1)}), \\ \partial^\infty T(P_2) &= \left\{ \lambda \begin{pmatrix} 1 \\ -\frac{1}{2} \end{pmatrix} \mid \lambda \geq 0 \right\}; \end{aligned}$$

$$\begin{aligned} \partial^c T(P_3) &= \partial^P T(P_3) \\ &= -\text{co} \left\{ e^{A^T (\sqrt{3}-1)v} \mid v \in N_{\mathcal{S}^c}^P(1, 1) \right. \\ &\quad \left. \text{or } v \in N_{\mathcal{S}^c}^P(2\sqrt{3}-3, 3-2\sqrt{3}), \text{ and } H(P_3, v) = 1 \right\} \\ &= -\text{co} \left\{ \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} \frac{-1}{2(\sqrt{3}-1)} \\ \frac{-\sqrt{3}}{2(\sqrt{3}-1)} \end{pmatrix} \right\} \\ &\quad - \left\{ \begin{pmatrix} \lambda \\ (2-\sqrt{3})\lambda \end{pmatrix} \mid \lambda \geq 0 \right\}, \\ \partial^\infty T(P_3) &= \left\{ \begin{pmatrix} \lambda \\ (2-\sqrt{3})\lambda \end{pmatrix} \mid \lambda \geq 0 \right\}. \end{aligned}$$

Observe that the vector  $\bar{f} \in \text{co}(f(P_2, \mathcal{U}))$  appearing in the statement of Lemma 3.5.2 is given by  $\bar{f} = (1/2, -1)$ .

If the target is modified to become  $\mathcal{S}' := \mathcal{S} \setminus \{(x_1, x_2) \in \mathbb{R}^2 : x_2 \geq$



$1, x_1 \geq (x_2)^2/2 + 1/2 + (x_2 - 1)^4$  (note that the boundary of  $\mathcal{S}'$  is  $\mathcal{C}^2$  at  $(1, 1)$ , see Figure 2), then the graph of the new minimum time function  $T'$  is smooth at all points of  $\gamma_2$ , but the unique normal is horizontal, so that  $T'$  is not differentiable at those points.




Figure 2

The next two examples deal with the case where the normal cone to the hypograph of  $T$  is not wedged. We show first that Theorem 3.2.3 does not hold in general. Next we provide an example where – although the normal cone is not wedged – the situation is entirely analogous to the case where the cone is wedged.

**Example 2.** Set

$$\gamma_1(y) = \begin{cases} (1 - \sqrt{-y^2 - 2y}, y) & -2 \leq y \leq -1 \\ (-1 + \sqrt{-y^2 - 2y}, y) & -1 \leq y \leq 0 \\ (-1 - \sqrt{-y^2 + 4y}, y) & 0 \leq y \leq 3, \end{cases}$$

and

$$\gamma_2(y) = \begin{cases} (1 + \sqrt{-y^2 - 2y}, y) & -2 \leq y \leq 0 \\ (1 - \sqrt{-y^2 + 2y}, y) & 0 \leq y \leq 1 \\ (0, y) & 1 \leq y \leq 2 \\ (-1 + \sqrt{-y^2 + 4y}, y) & 2 \leq y \leq 3. \end{cases}$$

Observe now that the concatenation of  $\gamma_1$  with  $\gamma_2$  defines a  $\mathcal{C}^{1,1}$ -curve  $\gamma$ . We set the target  $\mathcal{S}$  to be the unbounded component of  $\mathbb{R}^2 \setminus \{\gamma\}$  (see Figure 3) and the dynamics to be

$$\begin{cases} \dot{x}(t) = u \\ \dot{y}(t) = 0 \\ u \in \mathcal{U} = [-1, 1]. \end{cases}$$




Figure 3

It is readily verified that the minimum time function is everywhere defined and continuous. Observe furthermore that Petrov's condition holds at no points of the segment  $[-1, 1] \times \{0\}$ .

Consider now the curve

$$\Gamma(t) = \frac{\gamma_1(t) + \gamma_2(t)}{2}, \quad t \in [-1, 1],$$

together with the function

$$T(t) = \gamma_2(t) - \Gamma(t) (= \Gamma(t) - \gamma_1(t)), \quad t \in [-1, 1],$$

which is the minimum time to reach  $\mathcal{S}$  from the point  $\Gamma(t)$ .

Observe that  $T(t)$  is constantly equal to 1 for  $-1 \leq t \leq 0$  and in this interval all points of  $\Gamma$  are maximum points for  $T$ . Therefore  $(0, 0, 1)$  is a unit limiting normal vector to the hypograph of  $T$  at  $(0, 0, 1)$ .

On the other hand, it can be easily computed that a unit tangent vector to the graph of  $T$  at  $(0, 0, 1)$  is

$$\left( -\frac{2 + \sqrt{2}}{2\sqrt{3}}, 0, \frac{2 - \sqrt{2}}{2\sqrt{3}} \right).$$

Since the latter has positive scalar product with the limiting normal  $(0, 0, 1)$ , it is clear that the hypograph of  $T$  is not regular at  $(0, 0, 1)$ . In particular, the normal vector  $(0, 0, 1)$  is not proximal, thus showing that  $\text{hypo}(T)$  doesn't have positive reach (see (4) in Theorem 2.2.1).

Observe that both  $(1, 0, 0)$  and  $(-1, 0, 0)$  are unit proximal normals to  $\text{hypo}(T)$  at  $(0, 0, 1)$ , so that  $N_{\text{hypo}(T)}^C(0, 0, 1)$  contains a line. Therefore, the assumption (3.2.1) in Theorem 3.2.3 cannot be dropped in general.

Observe finally that the hypograph of  $T$  satisfies the external sphere condition with radius  $\rho$  for a suitable  $\rho > 0$ . Therefore this is a simple example showing that this condition is weaker than positive reach.  $\square$

**Example 3.** We consider again the dynamics (3.6.1) appearing in Example 1 and modify the target in order to allow lines in the normal cone to the hypograph of  $T$ .

The target is the set (see Figure 4)

$$\begin{aligned} \mathcal{S} = & \{ (x_1, x_2) \in \mathbb{R}^2 \mid x_1 \leq 0 \} \cup \{ (x_1, x_2) \in \mathbb{R}^2 \mid x_1 \geq 1 \} \\ & \cup \{ (x_1, x_2) \in \mathbb{R}^2 \mid 0 \leq x_1 \leq 1, x_2 \leq x_1 - 1 \} \\ & \cup \{ (x_1, x_2) \in \mathbb{R}^2 \mid 0 \leq x_1 \leq 1, x_2 \geq x_1 \}. \end{aligned}$$




Figure 4

The minimum time function is everywhere finite and continuous, but Petrov's condition does not hold. Computations of the same type of Example 1 show that the normal cone to the hypograph of  $T$  at  $(1/2, 0, 1)$  is not wedged, however  $N_{\text{hypo}(T)}^C(1/2, 0, 1)$  can be represented exactly as in (3.2.3) and the hypograph of  $T$  has positive reach. More precisely,

$$\begin{aligned} N_{\text{hypo}(T)}^P(1/2, 0, 1) &= N_{\text{hypo}(T)}^C(1/2, 0, 1) \\ &= \mathbb{R} \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \mathbb{R}^+ \text{co} \left\{ \begin{pmatrix} \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{pmatrix}, \begin{pmatrix} -\frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} \end{pmatrix} \right\} \end{aligned}$$

and

$$\partial^P T(1/2, 0) = \partial^C T(1/2, 0) = \mathbb{R} \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \text{co} \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ -1 \end{pmatrix} \right\}.$$

Observe that  $H((1/2, 0), (1, 0)) = 0$ , while

$$H((1/2, 0), (1, 1)) = H((1/2, 0), (-1, -1)) = 1,$$

so that the conclusion of Theorem 3.2.2 holds. An explicit computation of the minimum time function shows also that the conclusion of Theorem 3.2.3 holds as well.  $\square$

### 3.7 The case of differential inclusions

We study in this section the minimum time problems in the case of differential inclusions

$$\begin{cases} \dot{x}(t) \in F(x(t)) & a.e. \\ x(0) = x_0, \end{cases} \quad (3.7.1)$$

with the closed target  $\mathcal{S}$ .

Here  $F : \mathbb{R}^N \rightrightarrows \mathbb{R}^N$  is a multifunction that describes the dynamics, and  $x_0 \in \mathbb{R}^N$  is the starting point. For each trajectory  $y^{x_0}(\cdot)$  satisfying (3.7.1), we denote

$$\theta(y^{x_0}(\cdot)) := \min \{t \geq 0 \mid y^{x_0}(t) \in \mathcal{S}\}.$$

Of course  $\theta(y^{x_0}(\cdot)) \in [0, \infty]$ , and  $\theta(y^{x_0}(\cdot))$  is the taken time for the trajectory  $y^{x_0}(\cdot)$  to reach  $\mathcal{S}$ . The minimum time  $T(x_0)$  to reach  $\mathcal{S}$  for  $x_0$  is defined by

$$T(x_0) := \inf \{\theta(y^{x_0}(\cdot)) \mid \theta(y^{x_0}(\cdot)) \text{ is a trajectory satisfying (3.7.1)}\}. \quad (3.7.2)$$

Under standing hypotheses on  $F$  which we are now going to introduce, (3.7.2) will admit a minimal value.

#### 3.7.1 Hypotheses and some consequences

**Hypothesis (F):**

- (F1)  $F(x)$  is nonempty, convex, and compact for each  $x \in \mathbb{R}^n$ .
- (F2)  $F$  is Lipschitz continuous with respect to the Hausdorff distance. Thus, if  $K$  is the Lipschitz constant of  $F$ , then  $K|p|$  is the Lipschitz constant of  $H(\cdot, p)$ , i.e.,

$$|H(y, p) - H(x, p)| \leq K|p||y - x| \quad \forall x, y \in \mathbb{R}^n, \forall p \in \mathbb{R}^n. \quad (3.7.3)$$

- (F3) There exists a constant  $K_2 > 0$  so that  $\max\{|v| \mid v \in F(x)\} \leq K_2(1 + |x|)$ .

**Hypothesis (H):**

- (H1) There exists a constant  $c_0 \geq 0$  so that  $x \mapsto H(x, p)$  is semiconvex with semiconvexity constant  $c_0|p|$ .

(H2) For all  $p \neq 0$ , the gradient  $\nabla_p H(\cdot, p)$  exists and is globally Lipschitz, i.e.,

$$|\nabla_p H(x, p) - \nabla_p H(y, p)| \leq K_1 |y - x| \quad \forall x, y \in \mathbb{R}^n, \forall p \in \mathbb{R}^n \setminus \{0\}, \quad (3.7.4)$$

for some constant  $K_1 \geq 0$ .

**Remark 3.7.1** Global Lipschitz continuity in both (F2) and (H2) was assumed just to simplify computations. Indeed, our results still hold if  $F$  is locally Lipschitz with respect to the Hausdorff distance, and  $\nabla_p H(\cdot, p)$  is locally Lipschitz in  $x$ , uniformly so over  $p$  in  $\mathbb{R}^n \setminus \{0\}$  (as was supposed in [22]).

The following lemma is proved in [22].

**Lemma 3.7.1** *Suppose  $f : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}$  is Lipschitz in  $(x, y)$  on a boxed neighborhood  $U_x \times U_y = \{(x, y) \mid \max\{|x - \bar{x}|, |y - \bar{y}|\} < \delta\}$ , and that for each  $y \in U_y$ , the function  $x \rightarrow f(x, y)$  is semiconvex on  $U_x$  with constant independent of  $y$ . Then, for any  $\xi = (\xi_x, \xi_y) \in \partial f(\bar{x}, \bar{y})$ , one has  $\xi_x \in \partial_x f(\bar{x}, \bar{y})$ .*

**Corollary 3.7.1** *Suppose  $H$  satisfies assumption (H1). Then*

$$\partial H(x, p) \subseteq \partial_x H(x, p) \times \partial_p H(x, p) \quad \forall p \neq 0.$$

The following proposition is a consequence of [22, Proposition 1].

**Proposition 3.7.1** *Suppose  $F$  satisfies (F) and (H1). Then*

(1) *for each  $x, z \in \mathbb{R}^n$  with  $x_{\pm} = x \pm z$ , we have*

$$H(x_+, p) + H(x_-, p) - 2H(x, p) \geq -c_0 |p| |z|^2; \text{ and}$$

(2) *for each  $x, y \in \mathbb{R}^n$ , and  $\xi \in \partial_x H(x, p)$ , we have*

$$H(y, p) - H(x, p) - \langle \xi, y - x \rangle \geq -c_0 |p| |y - x|^2.$$

The differentiability statement in the assumption (H2) is equivalent to the argmax set of  $v \mapsto \langle v, p \rangle$  being a singleton, which equals  $\nabla_p H(x, p)$  and will be also denoted by  $F_p(x)$ . In view of (H2), for all  $p \neq 0$  the function  $F_p(\cdot)$  is globally Lipschitz with constant  $K_1$ .

The main use of (H2) is given by the following result whose proof is straightforward.

**Proposition 3.7.2** *Assume (F) and (H), and let  $p(\cdot)$  be an absolutely continuous arc on  $[0, T]$ , with  $p(t) \neq 0$  for all  $t \in [0, T]$ . Then, for each  $x \in \mathbb{R}^n$ , the initial value problem*

$$\begin{cases} \dot{x}(t) = F_{p(t)}(x(t)) & \text{a.e. } t \in [0, T] \\ x(0) = x. \end{cases} \quad (3.7.5)$$

has a unique solution  $y(\cdot, x)$ . Moreover,  $x \mapsto y(t, x)$  is Lipschitz on  $\mathbb{R}^n$  and

$$|y(t, z) - y(t, x)| \leq e^{K_1 t} |z - x|. \quad (3.7.6)$$

We conclude this section with some simple consequences of Gronwall's lemma.

**Lemma 3.7.2** *Let  $G : [0, T] \times \mathbb{R}^n \rightrightarrows \mathbb{R}^n$  be an upper semicontinuous multifunction. Assume  $G(t, \cdot)$  satisfies hypotheses (F1), (F2) uniformly in  $t \in [0, T]$ , and is such that for some  $K_0 > 0$ ,*

$$|v| \leq K_0 |p| \quad \forall v \in G(t, p), \forall (t, p) \in [0, T] \times \mathbb{R}^n.$$

Let  $\bar{p}(\cdot)$  a solution of the differential inclusion

$$\begin{cases} \dot{p}(t) \in G(t, p(t)) & \text{a.e. } t \in [0, T] \\ p(0) = p_0. \end{cases} \quad (3.7.7)$$

Then

$$e^{-K_0 t} |p(0)| \leq |p(t)| \leq e^{K_0 t} |p(0)| \quad \forall t \in [0, T].$$

Moreover, for all  $0 \leq t_1 \leq t_2 \leq T$ ,

$$e^{-K_0(t_2-t_1)} |p(t_2)| \leq |p(t_1)| \leq e^{K_0(t_2-t_1)} |p(t_2)|$$

and

$$|p(t_2) - p(t_1)| \leq K_0 e^{K_0(t_2-t_1)} (t_2 - t_1) |p(t_2)|.$$

*Proof.* Since  $p(t) = p(0) + \int_0^t \dot{p}(s) ds$ , we have

$$|p(t)| \leq |p(0)| + \int_0^t |\dot{p}(s)| ds \leq |p(0)| + K_0 \int_0^t |p(s)| ds.$$

So, using Gronwall's inequality, we get:  $|p(t)| \leq e^{K_0 t} |p(0)|$ .

We are now going to prove that  $e^{-K_0 t} |p(0)| \leq |p(t)|$  for all  $t > 0$ . Fixing  $t > 0$ , we define  $g(s) := p(t - s)$  for all  $s \in [0, t]$ . Since  $\dot{g}(s) = -\dot{p}(t - s)$  for almost  $s \in [0, t]$ , we have  $g(s) = g(0) + \int_0^s \dot{g}(\tau) d\tau$  for all  $s \in [0, t]$ . Thus,

$$\begin{aligned} |g(s)| &\leq |g(0)| + \int_0^s |\dot{g}(\tau)| d\tau = |g(0)| + \int_0^s |\dot{p}(t - \tau)| d\tau \\ &\leq |g(0)| + K_0 \int_0^s |p(t - \tau)| d\tau = |g(0)| + K_0 \int_0^s |g(\tau)| d\tau \end{aligned}$$

Again by Gronwall's inequality, we obtain  $|g(s)| \leq e^{K_0 s} |g(0)|$  for all  $s \in [0, t]$ . In particular,  $|g(t)| \leq e^{K_0 t} |g(0)|$ . The proof is completed noting that  $g(t) = p(0)$  and  $g(0) = p(t)$ .  $\square$

**Corollary 3.7.2** *Let  $p(\cdot)$  be a solution of (3.7.7). Then either  $\bar{p}(t) = 0$  for all  $t \in [0, T]$  or  $p(t) \neq 0$  for all  $t \in [0, T]$ .*

**Lemma 3.7.3** *Let  $y(\cdot, x_0)$  be a solution of (3.7.1). Then, for all  $t > 0$ , the following holds:*

- i)  $|y(t, x_0)| \leq (|x_0| + 1)e^{K_2 t} - 1,$
- ii)  $|y(t, x_0) - x_0| \leq (|x_0| + 1)(e^{K_2 t} - 1) \leq K_2(|x_0| + 1)e^{K_2 t}.$

*Proof.* Since

$$y(t, x_0) = x_0 + \int_0^t \dot{y}(s, x_0) ds,$$

recalling (F3) we have

$$|y(t, x_0)| \leq |x_0| + K_2 t + K_2 \int_0^t |y(s, x_0)| ds.$$

Hence, Gronwall's inequality yields (i). Then, observing that

$$|y(t, x_0) - x_0| \leq K_2 \int_0^t (1 + |y(s, x_0)|) ds,$$

(ii) follows using (i) in the above estimate.  $\square$

### 3.7.2 The hypograph of the minimum time function satisfies an external sphere condition

In this part, we will assume that  $\mathcal{S}$  is nonempty, closed and has the inner ball property with balls of radius  $\rho_0 > 0$ . Moreover, assumptions (F) and (H) are also assumed throughout. Recall that  $c_0, K, K_1, K_2$  are the constants in (H1), (F2), (H2), (F3). Let us define, for any  $r > 0$ ,

$$\mathcal{S}'(r) = \{x \mid T(x) \geq r\}, \quad \mathcal{S}' = \{x \mid T(x) \geq 0\} \quad \text{and} \quad \mathcal{S}^c = \{x \in \mathbb{R}^n \mid x \notin \mathcal{S}\}.$$

Our main results are the following theorem, together with the corollary.

**Theorem 3.7.1** *Assume (F) and (H). Suppose further that  $\mathcal{S}$  is nonempty, closed and has the inner ball property balls of radius  $\rho_0 > 0$  and  $T(\cdot)$  is continuous in  $\mathcal{S}^c$ . Then, the hypograph of  $T(\cdot)$  satisfies a  $\rho_T(\cdot)$ -exterior sphere condition for some continuous function  $\rho_T(\cdot) : \mathcal{S}^c \rightarrow (0, \infty)$ .*

**Remark 3.7.2** *The function  $\rho_T(\cdot)$  can be explicitly computed and depends only on  $x, T(x)$ , and on  $c_0, K, K_1, K_2, \rho_0$ .*

Consequently, under the assumptions of Theorem 3.7.1,  $T(\cdot)$  enjoys the regularity properties will be described in Corollary 5.2.2. Moreover, the following corollary follows from Theorem 3.7.1 and [50, theorem 21].

**Corollary 3.7.3** *Under the assumptions of Theorem 3.7.1, if  $T(\cdot)$  is locally Lipschitz, then  $T(\cdot)$  is locally semiconcave.*

The main part of the proof of Theorem 3.7.1 is divided into three lemmas.

**Lemma 3.7.4** *Suppose  $T(\cdot)$  has not a local maximum at the point  $\bar{x} \in \mathcal{S}^c$ . Let  $r = T(\bar{x})$  and let  $\bar{x}^+(\cdot)$  be an optimal trajectory steering  $\bar{x}$  to  $\mathcal{S}$  in time  $r$ , and set  $\bar{x}^-(s) = \bar{x}^+(r-s)$ . Then, there exists an arc  $\bar{p}(\cdot)$  defined on  $[0, r]$ , with  $\bar{p}(s) \neq 0$  for all  $s \in [0, r]$ , such that*

$$\begin{cases} -\dot{\bar{p}}(s) \in \partial_x H(\bar{x}^-(s), -\bar{p}(s)) & \text{a.e. } s \in [0, r], \\ \dot{\bar{x}}^-(s) = -F_{-\bar{p}(s)}(\bar{x}^-(s)) & \text{a.e. } s \in [0, r]. \end{cases} \quad (3.7.8)$$

Moreover,  $-\bar{p}(r-t) \in N_{\mathcal{S}'(r-t)}^P(\bar{x}^+(t))$  is realized by a ball of radius  $\rho(r-t)$  for all  $t \in (0, r]$ , i.e.,

$$\left\langle \frac{-\bar{p}(r-t)}{|\bar{p}(r-t)|}, \bar{y} - \bar{x}^+(t) \right\rangle \leq \frac{1}{2\rho(r-t)} |\bar{y} - \bar{x}^+(t)|^2, \quad \forall \bar{y} \in \mathcal{S}'(r-t), \quad (3.7.9)$$

where

$$\rho(s) = \frac{\rho_0}{1 + 2c_0\rho_0 s} e^{-(K+2K_1)s}. \quad (3.7.10)$$

*Proof.* Set  $\bar{x}_1 = \bar{x}^+(r)$ . Of course,  $\bar{x}_1 \in \partial\mathcal{S}$ . Since  $\mathcal{S}$  satisfies the  $\rho_0$ -internal sphere condition, there exists a proximal normal vector  $v \neq 0$  to  $\mathcal{S}'$  at  $\bar{x}_1$  such that  $\overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0) \subseteq \mathcal{S}$ , i.e.,

$$\left\langle \frac{v}{|v|}, z - \bar{x}_1 \right\rangle \leq \frac{1}{2\rho_0} |z - \bar{x}_1|^2 \quad \forall z \in \mathcal{S}'. \quad (3.7.11)$$

Now, consider the reversed differential inclusion with initial data

$$\begin{cases} \dot{y}(s) \in -F(y(s)) & \text{a.e. } s \in [0, r], \\ y(0) \in \overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0) \subseteq \mathcal{S}. \end{cases} \quad (3.7.12)$$

The Hamiltonian associated with  $-F$  is defined by

$$H^-(x, p) := \sup_{v \in -F(x)} \langle v, p \rangle = \sup_{w \in F(x)} \langle w, -p \rangle = H(x, -p). \quad (3.7.13)$$

Let us recall that the attainable set from  $\overline{B}(\bar{x}_1 + \rho v, \rho)$ , denoted  $\mathcal{A}^-(r)$ , is defined to be the set of all points  $y(r)$  where  $y(\cdot)$  is a trajectory satisfying (3.7.12). Since  $\bar{x}^-(\cdot)$  is a solution of (3.7.12) with initial point  $y(0) = \bar{x}_1$ , and  $T(\cdot)$  has not a local maximum at the point  $\bar{x}$ , one has that  $\bar{x}^-(r) = \bar{x}$  is on the boundary of  $\mathcal{A}^-(r)$ . Indeed, suppose  $\bar{x}$  is not on the boundary of  $\mathcal{A}^-(r)$ , then there exists  $\epsilon > 0$  such that  $B(\bar{x}, \epsilon) \subset \mathcal{A}^-(r)$ . Thus,  $T(y) \leq r = T(\bar{x})$  for all  $y \in B(\bar{x}, \epsilon)$ , and we get a contradiction since  $T(\cdot)$  has not a local



maximum at  $\bar{x}$ . Now, since  $\bar{x}$  is on the boundary of  $\mathcal{A}^-(r)$ , by [24, Theorem 3.4.5], there is an arc  $\bar{p}(\cdot)$ , such that  $\|\bar{p}(\cdot)\|_\infty > 0$ , satisfying

$$(-\dot{\bar{p}}(s), \dot{\bar{x}}^-(s)) \in \partial H^-(\bar{x}^-(s), \bar{p}(s)) \quad a.e. \ s \in [0, r], \quad (3.7.14)$$

and

$$\bar{p}(0) \in N_{\overline{B}(\bar{x}_1 + \rho_0 \frac{v}{|v|}, \rho_0)}(\bar{x}_1). \quad (3.7.15)$$

From (3.7.14), (3.7.13) and Corollary 3.7.1, we have

$$-\dot{\bar{p}}(s) \in \partial_x H(\bar{x}^-(s), -\bar{p}(s)) \quad a.e. \ s \in [0, r]. \quad (3.7.16)$$

Moreover, owing to (3.7.3), for all  $v \in \partial_x H(x, p)$  we have  $|v| \leq K|p|$ . Therefore, applying Lemma 3.7.2 to  $G(s, -\bar{p}(s)) = \partial_x H(\bar{x}^-(s), -\bar{p}(s))$ , we get

$$e^{-Ks}|\bar{p}(0)| \leq |\bar{p}(s)| \leq e^{Ks}|\bar{p}(0)| \quad \forall s \in [0, r]. \quad (3.7.17)$$

Since  $\|\bar{p}\| > 0$ , we have  $\bar{p}(s) \neq 0$  for all  $s \in [0, r]$ . Therefore, from (3.7.14) and Corollary 3.7.1 we get

$$\dot{\bar{x}}^-(s) = -\partial_p H(\bar{x}^-(s), -\bar{p}(s)) = -F_{-\bar{p}(s)}(\bar{x}^-(s)) \quad a.e. \ s \in [0, r]. \quad (3.7.18)$$

We are now going to prove (3.7.9). Fix  $t \in (0, r]$  and let  $\bar{y} \in \mathcal{S}'(r-t)$ , i.e.,  $T(\bar{y}) \geq r-t$ . Let  $\bar{y}^+(\cdot)$  be the solution of the Cauchy problem

$$\begin{cases} \dot{\bar{y}}^+(s) = F_{-\bar{p}(r-t-s)}(\bar{y}^+(s)) & a.e. \ s \in [0, r-t], \\ \bar{y}^+(0) = \bar{y}. \end{cases} \quad (3.7.19)$$

Note that  $\bar{y}_1 := \bar{y}^+(r-t) \in \mathcal{S}'$ . Then,  $\bar{y}^-(s) := \bar{y}^+(r-t-s)$  satisfies  $\bar{y}^-(r-t) = \bar{y}$  and

$$\begin{cases} \dot{\bar{y}}^-(s) = -F_{-\bar{p}(s)}(\bar{y}^-(s)) & a.e. \ s \in [0, r-t], \\ \bar{y}^-(0) = \bar{y}_1. \end{cases} \quad (3.7.20)$$

From (3.7.18), (3.7.20) and (3.7.6), we have

$$|\bar{y}^-(s) - \bar{x}^-(s)| \leq e^{K_1(r-t)}|\bar{y}_1 - \bar{x}_1| \quad \forall s \in [0, r-t]. \quad (3.7.21)$$

In order to prove (3.7.9), observe that

$$\begin{aligned} \langle -\bar{p}(r-t), \bar{y}^-(r-t) - \bar{x}^-(r-t) \rangle &= \\ \langle -\bar{p}(0), \bar{y}^-(0) - \bar{x}^-(0) \rangle &+ \int_0^{r-t} \frac{d}{ds} \langle -\bar{p}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle ds. \end{aligned} \quad (3.7.22)$$

Moreover,

$$\begin{aligned} \frac{d}{ds} \langle -\bar{p}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle &= \\ &= \langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle + \langle -\bar{p}(s), \dot{\bar{y}}^-(s) - \dot{\bar{x}}^-(s) \rangle \\ &= \langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle + \langle -\bar{p}(s), -F_{-\bar{p}(s)}(\bar{y}^-(s)) + F_{-\bar{p}(s)}(\bar{x}^-(s)) \rangle \\ &= \langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle - H(\bar{y}^-(s), -\bar{p}(s)) + H(\bar{x}^-(s), -\bar{p}(s)). \end{aligned}$$

Recalling (3.7.16) and Proposition 3.7.1 it follows that

$$\langle -\dot{\bar{p}}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle - H(\bar{y}^-(s), -\bar{p}(s)) + H(\bar{x}^-(s), -\bar{p}(s)) \leq c_0 |\bar{p}(s)| |\bar{y}^-(s) - \bar{x}^-(s)|^2.$$

Therefore,

$$\frac{d}{ds} \langle -\bar{p}(s), \bar{y}^-(s) - \bar{x}^-(s) \rangle \leq c_0 |\bar{p}(s)| |\bar{y}^-(s) - \bar{x}^-(s)|^2. \quad (3.7.23)$$

Owing to (3.7.15) and the fact that  $\bar{p}(0) \neq 0$ , we have  $-\frac{\bar{p}(0)}{|\bar{p}(0)|} = \frac{v}{|v|}$ . Thus, by (3.7.11) and the fact that  $\bar{y}_1 \in \mathcal{S}'$  we obtain

$$\left\langle -\frac{\bar{p}(0)}{|\bar{p}(0)|}, \bar{y}_1 - \bar{x}_1 \right\rangle \leq \frac{1}{2\rho_0} |\bar{y}_1 - \bar{x}_1|^2. \quad (3.7.24)$$

Combining (3.7.22), (3.7.23), (3.7.24) and noting that  $\bar{x}^-(0) = \bar{x}_1$ ,  $\bar{y}^-(0) = \bar{y}_1$ ,  $\bar{x}^-(r-t) = \bar{x}^+(t)$ ,  $\bar{y}^-(r-t) = \bar{y}$ , we conclude that

$$\langle -\bar{p}(r-t), \bar{y} - \bar{x}^+(t) \rangle \leq \frac{|\bar{p}(0)|}{2\rho_0} |\bar{y}_1 - \bar{x}_1|^2 + c_0 \int_0^{r-t} |\bar{p}(s)| |\bar{y}^-(s) - \bar{x}^-(s)|^2 ds.$$

Thus, by (3.7.17) and (3.7.21),

$$\left\langle \frac{-\bar{p}(r-t)}{|-\bar{p}(r-t)|}, \bar{y} - \bar{x}^+(t) \right\rangle \leq \left( \frac{1}{2\rho_0} + c_0(r-t) \right) e^{(K+2K_1)(r-t)} |\bar{y} - \bar{x}^+(t)|^2.$$

So, (3.7.9) follows (3.7.10), and the proof is complete.  $\square$

**Lemma 3.7.5** *Suppose  $T(\cdot)$  has not a local maximum at the point  $\bar{x} \in \mathcal{S}^c$ . Let  $r = T(\bar{x})$  and let  $\bar{x}^+(\cdot)$  be an optimal trajectory steering  $\bar{x}$  to  $\mathcal{S}$  in time  $r$ . If  $\bar{p}(\cdot)$  is the arc in Lemma 3.7.4, then  $H(\bar{x}, -\bar{p}(r)) \geq 0$ .*

*Proof.* Fixing  $t \in (0, r)$ , by (3.7.9) and the fact that  $\bar{x} \in \mathcal{S}'(r-t)$ , we have

$$\langle -\bar{p}(r-t), \bar{x} - \bar{x}^+(t) \rangle \leq \frac{1}{2\rho(r-t)} |\bar{p}(r-t)| |\bar{x} - \bar{x}^+(t)|^2.$$

Equivalently,

$$\left\langle -\bar{p}(r-t), \int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right\rangle \leq \frac{1}{2\rho(r-t)} |\bar{p}(r-t)| \left| \int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right|^2.$$

Dividing by  $t$  both sides of the above inequality, we get

$$\left\langle -\bar{p}(r-t), \frac{\int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds}{t} \right\rangle \leq \frac{1}{2\rho(r-t)} |\bar{p}(r-t)| \frac{\left| \int_0^t -F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \right|^2}{t}.$$

As  $t \rightarrow 0$ , we obtain

$$\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{x}) \rangle \leq 0.$$

This implies that  $H(\bar{x}, -\bar{p}(r)) \geq 0$ .  $\square$

**Lemma 3.7.6** *Suppose  $T(\cdot)$  has not a local maximum at the point  $\bar{x} \in \mathcal{S}^c$ . Let  $r = T(\bar{x})$  and let  $\bar{x}^+(\cdot)$  be an optimal trajectory steering  $\bar{x}$  to  $\mathcal{S}$  in time  $r$ . If  $\bar{p}(\cdot)$  is the arc in Lemma 3.7.4, then there exists a positive constant  $\rho_T$  such that  $(-\bar{p}(r), \lambda) \in N_{\text{hypo}(T)}^P(\bar{x}, T(\bar{x}))$  is realized by a ball of radius  $\rho_T$ , i.e., for all  $\bar{y} \in \mathcal{S}^c$  and  $\beta \leq T(\bar{y})$*

$$\left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq \frac{1}{2\rho_T} (|\bar{y} - \bar{x}|^2 + |\beta - r|^2) \quad (3.7.25)$$

where  $\lambda = H(\bar{x}, -\bar{p}(r))$ . Moreover,  $\rho_T = \rho_T(\bar{x})$  where  $\rho_T(\cdot) : \mathcal{S}^c \rightarrow (0, \infty)$  is a continuous function that can be computed explicitly.

*Proof.* Let  $\bar{y} \in \mathcal{S}'$ . Two cases may occur:

- (i)  $T(\bar{y}) < T(\bar{x})$ ,
- (ii)  $T(\bar{y}) \geq T(\bar{x})$ .

*First case:*  $T(\bar{y}) =: r_1 < r = T(\bar{x})$ . Let  $\bar{x}_1 = \bar{x}^+(r - r_1)$  and write

$$\langle -\bar{p}(r), \bar{y} - \bar{x} \rangle = \langle -\bar{p}(r), \bar{y} - \bar{x}_1 \rangle + \langle -\bar{p}(r), \bar{x}_1 - \bar{x} \rangle. \quad (3.7.26)$$

Recalling Lemma 3.7.4 and noting that  $\bar{y} \in \mathcal{S}'(r_1)$ , we can estimate the first term in the right-hand side of the above identity as follows

$$\begin{aligned} \langle -\bar{p}(r), \bar{y} - \bar{x}_1 \rangle &= \langle -\bar{p}(r_1), \bar{y} - \bar{x}_1 \rangle + \langle -\bar{p}(r) + \bar{p}(r_1), \bar{y} - \bar{x}_1 \rangle \\ &\leq \frac{1}{2\rho(r_1)} |\bar{p}(r_1)| |\bar{y} - \bar{x}_1|^2 + |\bar{p}(r) - \bar{p}(r_1)| |\bar{y} - \bar{x}_1|. \end{aligned}$$

From Lemmas 3.7.2 and 3.7.3, we have that

$$|\bar{p}(r_1)| \leq e^{K(r-r_1)} |\bar{p}(r)|, \quad |\bar{p}(r) - \bar{p}(r_1)| \leq K e^{K(r-r_1)} (r - r_1) |\bar{p}(r)|$$

and

$$|\bar{y} - \bar{x}_1| \leq |\bar{y} - \bar{x}| + |\bar{x}_1 - \bar{x}| \leq |\bar{y} - \bar{x}| + K_2(|\bar{x}| + 1) e^{K_2(r-r_1)} (r - r_1).$$

Thus, observing that  $\rho(r_1) \geq \rho(r)$ , one can get the estimate

$$\langle -\bar{p}(r), \bar{y} - \bar{x}_1 \rangle \leq L_1(|x|, r) |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |r - r_1|^2) \quad (3.7.27)$$

where

$$L_1(|x|, r) = \frac{1 + K_2^2(|x| + 1)^2 e^{2K_2 r}}{2\rho(r)} e^{Kr} + K K_2 (|x| + 1) e^{(K+K_2)r} + 2K e^{Kr}.$$

We rewrite the right-most term of (3.7.26) as follows

$$\begin{aligned} \langle -\bar{p}(r), \bar{x}_1 - \bar{x} \rangle &= \langle -\bar{p}(r), \int_0^{r-r_1} F_{-\bar{p}(r-s)}(\bar{x}^+(s)) ds \rangle \\ &= \int_0^{r-r_1} \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle ds \end{aligned}$$

and observe that

$$\begin{aligned} \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle &= \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}^+(s)) \rangle \\ &+ \langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}) \rangle + \langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}) \rangle. \end{aligned}$$

Moreover, recalling that  $\lambda = H(\bar{x}, -\bar{p}(r))$ , we have

$$\langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}) \rangle = H(\bar{x}, -\bar{p}(r)) = \lambda,$$

$$\begin{aligned} \langle -\bar{p}(r), F_{-\bar{p}(r)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}) \rangle &\leq K |\bar{p}(r)| |\bar{x}^+(s) - \bar{x}| \\ &\leq KK_2(|\bar{x}| + 1)e^{K_2r} |\bar{p}(r)| s \end{aligned}$$

and

$$\begin{aligned} &\langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) - F_{-\bar{p}(r)}(\bar{x}^+(s)) \rangle \\ &= \langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle - H(\bar{x}^+(s), -\bar{p}(r)) \\ &= \langle -\bar{p}(r) + \bar{p}(r-s), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle + H(\bar{x}^+(s), -\bar{p}(r-s)) - H(\bar{x}^+(s), -\bar{p}(r)) \\ &\leq 2K_2(|\bar{x}^+(s)| + 1) |\bar{p}(r) - \bar{p}(r-s)| \leq 2KK_2(|\bar{x}| + 1)e^{(K_2+K)r} |\bar{p}(r)| s. \end{aligned}$$

Therefore,

$$\langle -\bar{p}(r), F_{-\bar{p}(r-s)}(\bar{x}^+(s)) \rangle \leq \lambda + L_2(|\bar{x}|, r) |\bar{p}(r)| s$$

where  $L_2(|\bar{x}|, r) = KK_2(|\bar{x}| + 1)(2e^{Kr} + 1)e^{K_2r}$ . Thus, in view of the above estimates,

$$\langle -\bar{p}(r), \bar{x}_1 - \bar{x} \rangle \leq \lambda(r - r_1) + \frac{L_2(|\bar{x}|, r)}{2} |\bar{p}(r)| |r - r_1|^2. \quad (3.7.28)$$

Combining (3.7.26), (3.7.27) and (3.7.28), we get

$$\langle -\bar{p}(r), \bar{y} - \bar{x} \rangle + \lambda(r_1 - r) \leq \frac{2L_1(|\bar{x}|, r) + L_2(|\bar{x}|, r)}{2} |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |r_1 - r|^2).$$

From Lemma 3.7.4, we have that  $\lambda \geq 0$ . Therefore, since  $r_1 < r$ , we conclude that

$$\left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq \frac{2L_1(|\bar{x}|, r) + L_2(|\bar{x}|, r)}{2} (|\bar{y} - \bar{x}|^2 + |\beta - r|^2) \quad (3.7.29)$$

for all  $\beta \leq r_1$ . So, if  $T(\bar{y}) < T(\bar{x})$ , then (3.7.25) holds true provided  $\rho_T(\bar{x})$  is such that

$$\rho_T(\bar{x}) \leq \rho(r) \text{ and } \rho_T(\bar{x}) \leq \frac{2}{2L_1(|\bar{x}|, r) + L_2(|\bar{x}|, r)}. \quad (3.7.30)$$

*Second case:*  $T(\bar{y}) = r_1 \geq r = T(\bar{x})$ .

In view of Lemma 3.7.4, we already know that (3.7.25) holds for all  $\beta \leq r$ . So let's prove (3.7.25) for  $r < \beta \leq r_1$ . Let  $\bar{y}^+(\cdot)$  be the solution of

$$\begin{cases} \dot{\bar{y}}(s) \in F_{-\bar{p}(r)}(\bar{y}(s)) & \text{a.e. } s \in [0, r_1 - \beta] \\ \bar{y}(0) = \bar{y}. \end{cases} \quad (3.7.31)$$

Set  $\bar{y}_1 = \bar{y}^+(\beta - r)$  and compute

$$\langle -\bar{p}(r), \bar{y} - \bar{x} \rangle = \langle -\bar{p}(r), \bar{y} - \bar{y}_1 \rangle + \langle -\bar{p}(r), \bar{y}_1 - \bar{x} \rangle. \quad (3.7.32)$$

Since  $r < \beta \leq r_1$ , one can see that  $T(\bar{y}_1) > r$ . Thus,  $\bar{y}_1 \in \mathcal{S}'(r)$ . Then, recalling Lemma 3.7.4 we get

$$\langle -\bar{p}(r), \bar{y}_1 - \bar{x} \rangle \leq \frac{1}{2\rho(r)} |\bar{p}(r)| |\bar{y}_1 - \bar{x}|^2.$$

Using Lemma 3.7.3, we also have

$$\begin{aligned} |\bar{y}_1 - \bar{x}| &\leq |\bar{y}_1 - \bar{y}| + |\bar{y} - \bar{x}| \\ &\leq K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}|\beta - r| + |\bar{y} - \bar{x}|. \end{aligned}$$

So,

$$\langle -\bar{p}(r), \bar{y}_1 - \bar{x} \rangle \leq \frac{K_2^2(|\bar{y}| + 1)^2 e^{2K_2(\beta-r)} + 1}{2\rho(r)} |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |\beta - r|^2). \quad (3.7.33)$$

On the other hand, recalling (3.7.4) we have

$$\begin{aligned} \langle -\bar{p}(r), \bar{y} - \bar{y}_1 \rangle &= \left\langle -\bar{p}(r), \int_0^{\beta-r} -F_{-\bar{p}(r)}(\bar{y}^+(s)) ds \right\rangle = \int_0^{\beta-r} \left\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{y}^+(s)) \right\rangle ds \\ &= \int_0^{\beta-r} \left\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{y}^+(s)) + F_{-\bar{p}(r)}(\bar{x}) \right\rangle ds + \int_0^{\beta-r} \left\langle -\bar{p}(r), -F_{-\bar{p}(r)}(\bar{x}) \right\rangle ds \\ &\leq K_1 |\bar{p}(r)| \int_0^{\beta-r} |\bar{y}^+(s) - \bar{x}| ds + \int_0^{\beta-r} -H(x, -\bar{p}(r)) ds \\ &= K_1 |\bar{p}(r)| \int_0^{\beta-r} |\bar{y}^+(s) - \bar{x}| ds + \lambda(r - \beta). \end{aligned}$$

Owing to Lemma 3.7.3, for all  $s \in [0, \beta - r]$

$$\begin{aligned} |\bar{y}^+(s) - \bar{x}| &\leq |\bar{y}^+(s) - \bar{y}| + |\bar{y} - \bar{x}| \\ &\leq K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}|\beta - r| + |\bar{y} - \bar{x}|. \end{aligned}$$

Therefore,

$$\langle -\bar{p}(r), \bar{y} - \bar{y}_1 \rangle \leq \lambda(r - \beta) + K_1 [1 + K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}] |\bar{p}(r)| (|\bar{y} - \bar{x}|^2 + |\beta - r|^2). \quad (3.7.34)$$

Combining (3.7.32), (3.7.33) and (3.7.34), we get

$$\left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq L_3 (|\bar{y} - \bar{x}|^2 + |\beta - r|^2)$$

where  $L_3 = \frac{K_2^2(|\bar{y}|+1)^2 e^{2K_2(\beta-r)} + 1}{2\rho(r)} + K_1[1 + K_2(|\bar{y}| + 1)e^{K_2(\beta-r)}]$ . The dependence of  $L_3$  on  $|\bar{y}|$  can be easily disposed of taking

$$L_4(|\bar{x}|, r) = \frac{K_2^2(|\bar{x}| + 2)^2 e^{2K_2} + 1}{2\rho(r)} + K_1[1 + K_2(|\bar{x}| + 2)e^{K_2}] + 1.$$

Then, the above inequality yields

$$\left\langle \frac{(-\bar{p}(r), \lambda)}{|(-\bar{p}(r), \lambda)|}, (\bar{y} - \bar{x}, \beta - r) \right\rangle \leq L_4(|\bar{x}|, r) (|\bar{y} - \bar{x}|^2 + |\beta - r|^2). \quad (3.7.35)$$

Recalling (3.7.30) and (3.7.35), and taking

$$\rho_T(\bar{x}) := \left( 2 \max \left\{ \frac{2L_1(|\bar{x}|, T(\bar{x})) + L_2(|\bar{x}|, T(\bar{x}))}{2}, L_4(|\bar{x}|, T(\bar{x})) \right\} \right)^{-1} \quad (3.7.36)$$

we obtain (3.7.25). Finally, since  $T(\cdot)$  is continuous on  $\mathcal{S}^c$ , one can easily see that  $\rho_T(\cdot)$  is also continuous on  $\mathcal{S}^c$ . The proof is complete.  $\square$

*Proof of Theorem 3.7.1.* Let  $\bar{x} \in \mathcal{S}^c$ . Let  $r = T(\bar{x})$  and let  $\bar{x}^+(\cdot)$  be an optimal trajectory steering  $\bar{x}$  to  $\mathcal{S}$  in time  $r$ . By the dynamic programming principle,  $T(\bar{x}^+(t)) = r - t$  for all  $t \in (0, r)$ . This implies that  $T(\cdot)$  has a not local maximum at  $\bar{x}^+(t)$  for all  $t \in (0, r)$ . Therefore, by applying Lemma 3.7.6, we obtain that for all  $t \in (0, r)$ , there exists a unit vector  $\bar{q}(t) \in N_{\text{hypo}(T)}^P(\bar{x}^+(t), T(\bar{x}^+(t)))$  realized by a ball of radius  $\rho_T(\bar{x}^+(t))$  where  $\rho_T(\cdot)$  is given by (3.7.36), i.e., for all  $\bar{y} \in \mathcal{S}^c$  and  $\beta \leq T(\bar{y})$

$$\begin{aligned} & \left\langle \bar{q}(t), (\bar{y} - \bar{x}^+(t), \beta - T(\bar{x}^+(t))) \right\rangle \\ & \leq \frac{1}{2\rho_T(\bar{x}^+(t))} (|\bar{y} - \bar{x}^+(t)|^2 + |\beta - T(\bar{x}^+(t))|^2). \end{aligned} \quad (3.7.37)$$

Since  $\bar{q}(t)$  is a unit vector in  $\mathbb{R}^{n+1}$  for all  $t \in (0, r)$ , there exists a sequence  $\{t_k\}$  which converges to  $0^+$  such that the sequence  $\{\bar{q}(t_k)\}$  converges to a unit vector  $\bar{q}$  in  $\mathbb{R}^{n+1}$ . Taking  $t = t_k$  and then letting  $k \rightarrow \infty$  in (3.7.37), by the continuity of  $T(\cdot)$  and  $\rho_T(\cdot)$ , we obtain that for all  $\bar{y} \in \mathcal{S}^c$  and  $\beta \leq T(\bar{y})$ ,

$$\left\langle \bar{q}, (\bar{y} - \bar{x}, \beta - T(\bar{x})) \right\rangle \leq \frac{1}{2\rho_T(\bar{x})} (|\bar{y} - \bar{x}|^2 + |\beta - T(\bar{x})|^2)$$

where  $\rho_T(\cdot)$  is given by (3.7.36). Therefore,  $\bar{q} \in N_{\text{hypo}(T)}^P(\bar{x}, T(\bar{x}))$  is realized by a ball of radius  $\rho_T(T(\bar{x}))$ . The proof is complete.  $\square$

We conclude this part with an example in which the Petrov's controllability condition does not hold and the minimum time function  $T$  is just continuous. Moreover, in this example, the multifunction  $F$  does not admit a  $C^1$  parameterization but the multifunction  $F$  and the Hamiltonian  $H$  satisfy the assumptions (F) and (H).

**Example 1.** Set

$$\gamma(t) = \begin{cases} (1, t) & t \leq 0 \\ (1 - \sqrt{-t^2 + 2t}, t) & 0 \leq t \leq 1 \\ (0, t) & t \geq 1. \end{cases}$$

We set the target  $\mathcal{S}$  to be the right part of  $\mathbb{R}^2 \setminus \{\gamma\}$  (see Figure 1) and the differential inclusion to be

$$(\dot{x}_1(t), \dot{x}_2(t)) \in F(x_1(t), x_2(t)) = \left\{ (u_1, h(x_2(t))u_2) \mid u_1, u_2 \in [0, 1] \right\},$$

where

$$h(x_2) = \begin{cases} 0 & \text{if } x_2 \leq 1 \\ x_2 - 1 & \text{if } x_2 \geq 1. \end{cases}$$




Figure 1

Observe first that  $\mathcal{S}$  has the inner ball property. Observe furthermore that for  $0 < t \leq 1$ , the point  $z_t = (1 - \sqrt{-t^2 + 2t}, t)$  is on the boundary of  $\mathcal{S}$ , and

$$\min_{v \in F(z_t)} \langle v, \nu \rangle = -\sqrt{-t^2 + 2t} |\nu|,$$

where  $\nu$  is the proximal vector to  $\mathcal{S}$ . Therefore, since  $\lim_{t \rightarrow 0^+} \sqrt{-t^2 + 2t} = 0$ , one can see that the Petrov's controllability condition does not hold in a neighborhood of  $(1, 0)$ . Moreover, by computing explicitly the minimum time function  $T$ ,

$$T(x_1, x_2) = \begin{cases} 1 - x_1 & \text{if } x_1 \leq 1, x_2 \leq 0 \\ 1 - \sqrt{-x_2^2 + 2x_2} - x_1 & \text{if } x_1 \leq 1 - \sqrt{-x_2^2 + 2x_2}, 0 < x_2 \leq 1 \\ -x_1 & \text{if } x_1 \leq 0, x_2 > 1, \end{cases}$$

one can prove that  $T$  is continuous, but is not Lipschitz at points  $(x_1, 0)$  for  $x_1 \leq 1$ .

We next show that  $F$  does not admit a  $C^1$  parameterization. We first recall a criterion in [22, p3] that if  $F$  admit a  $C^1$  parameterization, then the Hamiltonian  $H$  necessarily has the property

$$H(x, p) = -H(x, -p) \implies \partial_x H(x, p) = -\partial_x H(x, -p), \quad (3.7.38)$$

where  $\partial_x$  denotes the Clarke partial subgradient in  $x$ . In this example, the Hamiltonian  $H$  is computed as

$$H((x_1, x_2), (p_1, p_2)) = \begin{cases} 0 & p_1 < 0, p_2 < 0, \\ p_1 & p_1 \geq 0, p_2 < 0, \\ h(x_2)p_2 & p_1 < 0, p_2 \geq 0, \\ p_1 + h(x_2)p_2 & p_1 \geq 0, p_2 \geq 0. \end{cases}$$

At the point  $(x_1, x_2) = (1, 1)$ , one has that  $H((1, 1), (0, -1)) = H((1, 1), (0, 1)) = 0$ . However,

$$\partial_x H((1, 1), (0, -1)) = (0, 0) \quad \text{and} \quad \partial_x H((1, 1), (0, 1)) = (0, [0, 1]),$$

and so (3.7.38) is violated at the point  $(1, 1)$ . Thus,  $F$  does not admit a  $C^1$  parameterization.

Finally, since  $h$  is a convex function, one can also prove that  $F$  and  $H$  satisfies the assumptions (F) and (H). Therefore, by applying Theorem 3.7.1, the hypograph of  $T$  satisfies a  $\rho_T(\cdot)$ -exterior sphere condition.

### 3.7.3 The inner ball property of attainable sets

In this second part, we will study the attainable set  $\mathcal{A}(T)$  from 0 for the reversed differential inclusion

$$\begin{cases} \dot{x}(t) \in -F(x(t)) & a.e. \\ x(0) = 0. \end{cases} \quad (3.7.39)$$

For any  $T > 0$ , such set is defined by

$$\mathcal{A}(T) := \{y(t) \mid t \in [0, T] \text{ and } y(\cdot) \text{ is a solution of (3.7.39)}\}.$$

Let us recall that  $c_0$ ,  $K$  and  $K_1$  are the constants appearing in  $(H_1)$ , (3.7.3) and (3.7.4), respectively.

**Theorem 3.7.2** *Assume  $F$  satisfies (F) and (H). In addition, suppose that, for some  $R > 0$ ,  $F(x)$  satisfies the  $R$ -interior sphere condition for all  $x \in \mathbb{R}^n$ . If  $T > 0$  and  $e^{-3KT} > 2c_0RT^2$ , then the attainable set  $\mathcal{A}(T)$  satisfies the  $R(T)$ -internal sphere condition with*

$$R(T) = R \frac{(e^{-3KT} - 2c_0RT^2)}{(1 + KT + K_1T)^2}. \quad (3.7.40)$$



*Proof.* Let  $\bar{x} \in \partial\mathcal{A}(T)$  and let  $\bar{x}^-(\cdot)$  be a trajectory of (3.7.39) steering 0 to  $\bar{x}$  in time  $T$ . By the Pontryagin maximum principle, there exists an arc  $\bar{p}(\cdot)$  defined in  $[0, T]$ , with  $\bar{p}(s) \neq 0$  for all  $s \in [0, T]$ , such that

$$\begin{cases} -\dot{\bar{p}}(s) \in \partial_x H(\bar{x}^-(s), -\bar{p}(s)) & \text{a.e. } s \in [0, T] \\ \dot{\bar{x}}^-(s) = -F_{-\bar{p}(s)}(\bar{x}^-(s)) & \text{a.e. } s \in [0, T]. \end{cases} \quad (3.7.41)$$

We want to prove that, for  $r_0 := R(T)$  (where  $R(T)$  is defined in (3.7.40)),

$$B\left(\bar{x} + r_0 T \frac{-\bar{p}(T)}{|\bar{p}(T)|}, r_0 T\right) \subseteq \mathcal{A}(T). \quad (3.7.42)$$

Let  $\theta \in B(0, 1)$ . Considering the adjoint equation associated with  $\bar{p}(\cdot)$ , that is,

$$\begin{cases} \dot{\bar{z}}(s) = -\frac{\langle \dot{\bar{p}}(s), \bar{z}(s) \rangle}{|\bar{p}(s)|^2} \bar{p}(s) & \text{a.e.} \\ \bar{z}(T) = \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta, \end{cases} \quad (3.7.43)$$

one can see that

$$\langle \dot{\bar{z}}(s), \bar{p}(s) \rangle = -\langle \dot{\bar{p}}(s), \bar{z}(s) \rangle \text{ for a.e. } s \in [0, T]. \quad (3.7.44)$$

This implies that  $\frac{d}{dt} \langle \bar{z}(s), \bar{p}(s) \rangle = 0$  for a.e.  $s \in [0, T]$ . Therefore,  $\langle \bar{z}(s), \bar{p}(s) \rangle$  is constant for all  $s \in [0, T]$ . In particular,

$$\langle \bar{z}(s), \bar{p}(s) \rangle = \langle \bar{z}(T), \bar{p}(T) \rangle. \quad (3.7.45)$$

On the other hand, from (3.7.43) we have  $|\dot{\bar{z}}(s)| \leq K|\bar{z}(s)|$ . Thus, recalling Lemma 3.7.2 we obtain

$$e^{-K(t_2-t_1)} |\bar{z}(t_2)| \leq |\bar{z}(t_1)| \leq e^{K(t_2-t_1)} |\bar{z}(t_2)| \quad \text{for all } 0 \leq t_1 \leq t_2 \leq T. \quad (3.7.46)$$

We now set

$$\bar{y}_\theta(s) = \bar{x}(s) - r_0 s \bar{z}(s). \quad (3.7.47)$$

Our aim is to prove that  $\bar{y}_\theta(T) \in \mathcal{A}(T)$ . Since  $\bar{y}_\theta(0) = \bar{x}(0) = 0$  and  $\bar{y}_\theta(T) = \bar{x}(T) - r_0 T \left( \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right)$ , we need to show

$$\dot{\bar{y}}_\theta(s) \in -F(\bar{y}_\theta(s)) \quad \text{for a.e. } s \in [0, T].$$

Observe that  $F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \in \partial F(\bar{y}_\theta(s))$ . Since  $F(\bar{y}_\theta(s))$  is convex and satisfies the  $R$ -internal sphere condition, we have that  $\bar{p}(s)$  is an inner normal vector to  $\partial F(\bar{y}_\theta(s))$  at the point  $F_{-\bar{p}(s)}(\bar{y}_\theta(s))$ . Thus,  $-\dot{\bar{y}}_\theta(s) \in F(\bar{y}_\theta(s))$  (equivalently to  $\dot{\bar{y}}_\theta(s) \in -F(\bar{y}_\theta(s))$ ) if  $-\dot{\bar{y}}_\theta(s) \in B(F_{-\bar{p}(s)}(\bar{y}_\theta(s)) + r \frac{\bar{p}(s)}{|\bar{p}(s)|}, r)$ . Therefore,<sup>1</sup> our conclusion will follow from

$$\left\langle \frac{\bar{p}(s)}{|\bar{p}(s)|}, -\dot{\bar{y}}_\theta(s) - F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{1}{2r} |\dot{\bar{y}}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))|^2. \quad (3.7.48)$$

<sup>1</sup>Observe that for all  $r > 0$  and  $x \in \mathbb{R}^N$ ,  $y \in \overline{B}(x + rv, r) \Leftrightarrow \langle v, y - x \rangle \geq \frac{1}{2r} |y - x|^2$  where  $v \in \mathbb{R}^N$  is any unit vector.

Equivalently,

$$\left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{1}{2r} |\dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))|^2. \quad (3.7.49)$$

We are now going to prove (3.7.49). On account of (3.7.47), we have

$$\dot{y}_\theta(s) = -F_{-\bar{p}(s)}(\bar{x}(s)) - r_0\bar{z}(s) - r_0s\dot{\bar{z}}(s).$$

Thus,

$$\begin{aligned} & \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \\ &= \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, F_{-\bar{p}(s)}(\bar{y}_\theta(s)) - F_{-\bar{p}(s)}(\bar{x}(s)) - r_0\bar{z}(s) - r_0s\dot{\bar{z}}(s) \right\rangle \\ &= \frac{1}{|\bar{p}(s)|} (H(\bar{y}_\theta(s), -\bar{p}(s)) - H(\bar{x}(s), -\bar{p}(s))) \\ & \quad + \frac{r_0}{|\bar{p}(s)|} \langle \bar{p}(s), \bar{z}(s) \rangle + r_0s \frac{1}{|\bar{p}(s)|} \langle \bar{p}(s), \dot{\bar{z}}(s) \rangle. \end{aligned}$$

Recalling (3.7.44), (3.7.47) and (3.7.45), we conclude that

$$\begin{aligned} & \left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \\ &= \frac{1}{|\bar{p}(s)|} (H(\bar{y}_\theta(s), -\bar{p}(s)) - H(\bar{x}(s), -\bar{p}(s)) - \langle -\dot{\bar{p}}(s), \bar{y}_\theta(s) - \bar{x}(s) \rangle) + \frac{r_0}{|\bar{p}(s)|} \langle \bar{p}(T), \bar{z}(T) \rangle \\ & \geq -c_0 |\bar{y}_\theta(s) - \bar{x}(s)|^2 + r_0 \frac{|\bar{p}(T)|}{|\bar{p}(s)|} \left\langle \frac{\bar{p}(T)}{|\bar{p}(T)|}, \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right\rangle \\ & \geq -c_0 r_0^2 s^2 |\bar{z}(s)|^2 + \frac{r_0}{2} \frac{|\bar{p}(T)|}{|\bar{p}(s)|} \left| \frac{\bar{p}(T)}{|\bar{p}(T)|} - \theta \right|^2 \\ & = -c_0 r_0^2 s^2 |\bar{z}(s)|^2 + \frac{r_0}{2} \frac{|\bar{p}(T)|}{|\bar{p}(s)|} |\bar{z}(T)|^2. \end{aligned}$$

Recalling Lemma 3.7.2 and (3.7.46), we obtain

$$\left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{r_0}{2} (-2c_0 r_0 T^2 + e^{-3KT}) |\bar{z}(s)|^2. \quad (3.7.50)$$

Observe that  $0 < r_0 = R(T) = R \frac{(e^{-3KT} - 2c_0 RT^2)}{(1 + KT + K_1 T)^2} \leq R$ . Then,

$$\left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{r_0}{2} (-2c_0 RT^2 + e^{-3KT}) |\bar{z}(s)|^2. \quad (3.7.51)$$

On the other hand,

$$\begin{aligned} |\dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))| &\leq |F_{-\bar{p}(s)}(\bar{y}_\theta(s)) - F_{-\bar{p}(s)}(\bar{x}(s))| + r_0 |\bar{z}(s)| + r_0 s |\dot{\bar{z}}(s)| \\ &\leq K_1 r_0 s |\bar{z}(s)| + r_0 |\bar{z}(s)| + K r_0 s |\bar{z}(s)| \\ &\leq r_0 (K_1 T + KT + 1) |\bar{z}(s)|. \end{aligned}$$

Thus,

$$\left\langle -\frac{\bar{p}(s)}{|\bar{p}(s)|}, \dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s)) \right\rangle \geq \frac{e^{3KT} - 2c_0RT^2}{2r_0(K_1T + KT + 1)^2} |\dot{y}_\theta(s) + F_{-\bar{p}(s)}(\bar{y}_\theta(s))|^2,$$

and (3.7.49) follows. The proof is complete.  $\square$

Finally, let us denote by  $\mathcal{A}(x, T)$  the attainable set from  $x$  in time  $T$  for the differential inclusion in (3.7.39). One can see from Theorem 3.7.2 that there exists a time  $T_0 > 0$  such that for all  $0 < T < T_0$ , the set  $\mathcal{A}(x, T)$  satisfies the  $R(T)$ -exterior sphere condition with  $R(T)$  given by (3.7.40). Moreover, for any closed set  $\mathcal{S} \subset \mathbb{R}^N$ , let us set

$$\mathcal{A}(\mathcal{S}, T) = \bigcup_{x \in \mathcal{S}} \mathcal{A}(x, T).$$

**Corollary 3.7.4** *Suppose that  $\mathcal{S}$  is nonempty and closed. Under the assumptions in Theorem 3.7.2, there exists  $T_0 > 0$  such that, for all  $0 < T < T_0$ , then the set  $\mathcal{A}(\mathcal{S}, T)$  satisfies the  $R(T)$ -exterior sphere condition with  $R(T)$  given by (3.7.40).*

Consequences on minimum time function for a general target.

**Theorem 3.7.3** *Assume (F), (H) and for some  $R > 0$ ,  $F(x)$  satisfies the  $R$ -interior sphere condition for all  $x \in \mathbb{R}^n$ . Suppose further that  $\mathcal{S}$  is nonempty, closed and  $T(\cdot)$  is continuous in  $\mathcal{S}^c$ . Then, the hypograph of  $T(\cdot)$  satisfies a  $\rho_T(\cdot)$ -exterior sphere condition for some continuous function  $\rho_T(\cdot) : \mathcal{S}^c \rightarrow (0, \infty)$ .*

**Corollary 3.7.5** *Under the assumptions of Theorem 3.7.3, if  $T(\cdot)$  is locally Lipschitz, then  $T(\cdot)$  is locally semiconcave.*



## Chapter 4

# Semiconvexity type results

In this chapter we will be concerned with the minimum time function  $T$  around the origin. Under some suitable assumptions on the nonlinear control system, there exists an open neighborhood  $\mathcal{U}$  of 0 such that the *epigraph* of  $T|_{\mathcal{U}}$  has *positive reach*. Therefore,  $T$  enjoys several regularity properties of a semiconvex function (see Theorem 2.2.2). In particular,  $T$  is a.e twice differentiable in  $\mathcal{U}$ .

Our approach is based on the linear nonautonomous control systems which are obtained by linearizing the nonlinear control system along the optimal trajectories. From the normality assumption at 0 of such linear systems, the corresponding reachable sets up to small time will be strictly convex. Furthermore, the reachable set of the nonlinear control systems in  $2D$  can inherit the strict convexity from linear systems. Consequently, every point near 0 is optimal and is steered to the origin by a unique optimal trajectory.

A fundamental analysis for this approach is the study on the reachable set  $\mathcal{R}^T$  of the *normal* linear control system (see Definition 4.1.1). For such systems, the Pontryagin maximum principle gives complete information on optimal controls. More precisely, every optimal control  $\bar{u}$  on  $[0, T]$  is a sign function of a switching function  $g$  on  $[0, T]$  which has finitely many zero points, i.e.,  $\bar{u}$  is a bang-bang control with finitely many switching points on  $[0, T]$ . Therefore,  $\mathcal{R}^T$  is strictly convex (see other proof in [41]). However, to our aim, we need to have an optimal computation of the modulus of the strict convexity of  $\mathcal{R}^T$  (Theorem 4.2.1). The crucial point of the proof is classifying the set of zero points of  $g$  according to their multiplicity.

The chapter is organized as follows: in Section 4.1 recall some definitions of linear control systems and some basic facts, while Section 4.2 contains our results on the reachable set in the case of linear control system. The last Section 4.3 is devoted a study on the minimum time function for a nonlinear control system in  $2D$ .

## 4.1 Linear control system and normality

Consider the linear control system

$$\begin{cases} \dot{y}(t) = Ay(t) + Bu(t) & a.e. \\ u(t) \in \mathcal{U} & a.e. \\ y(0) = 0, \end{cases} \quad (4.1.1)$$

where  $A \in \mathbb{M}_{N \times N}$ ,  $B \in \mathbb{M}_{N \times M}$  and the control set  $\mathcal{U} = [-1, 1]^M$ . For any  $t > 0$ , we denote by  $\mathcal{U}_{ad}^t$ , the set of admissible controls on the interval  $[0, t]$ , i.e., the measurable function  $u : [0, t] \rightarrow \mathbb{R}^M$ , such that  $u(s) \in \mathcal{U}$  a.e.  $s \in [0, t]$ . For any  $u(\cdot) \in \mathcal{U}_{ad}^t$ , the unique Carathéodory solution of (4.1.1) is denoted by  $y^u(\cdot)$ . Moreover,

$$y^u(t) = \int_0^t e^{A(t-s)} Bu(s) ds.$$

Therefore, the reachable set from 0 in time  $T$  can be explicitly computed, and reads as:

$$\mathcal{R}^T = \left\{ \int_0^T e^{A(T-s)} Bu(s) ds \mid u(\cdot) \in \mathcal{U}_{ad}^T \right\}.$$

One can see that the set  $\mathcal{R}^T$  is convex and compact.

**Proposition 4.1.1 (Pontryagin maximum principle)** *Suppose  $\bar{x} \in \mathcal{R}^T$  is realized by the control  $\bar{u}(\cdot) \in \mathcal{U}_{ad}^T$ . Then  $\bar{x} \in \partial \mathcal{R}^T$  if and only if for all  $\bar{\zeta} \in N_{\mathcal{R}^T}$ , it holds*

$$\bar{u}_i(t) = \text{sign} \langle \bar{\zeta}, e^{A(T-t)} b_i \rangle, \quad a.e. \ t \in [0, T],$$

for all  $i = 1, 2, \dots, M$  where  $u = (u_1, u_2, \dots, u_M)$  and  $B = (b_1, b_2, \dots, b_M)$ .

A standard reference for the proof is in [41].

**Definition 4.1.1** *The system (4.1.1) is normal if and only if*

$$\text{Rank}[b_i, Ab_i, \dots, A^{N-1}b_i] = N$$

for  $i = 1, 2, \dots, M$  where  $B = (b_1, b_2, \dots, b_M)$ .

**Remark 4.1.1** *If the system (4.1.1) is normal then  $(A, B)$  satisfies Kalman rank condition. Therefore (4.1.1) is small time locally controllable.*

We state here a classical result for such systems.

**Theorem 4.1.1** *The linear control system (4.1.1) is normal if and only if the reachable set  $\mathcal{R}^T$  is strictly convex for any  $T > 0$ .*

**Proof.** One can find a proof in [41]. □

## 4.2 Quantitative strict convexity of reachable sets

### 4.2.1 Linear autonomous control systems

We will study in this subsection the modulus of strict convexity of the reachable sets in the case of normal linear autonomous control systems (4.1.1). The first Lemma clarifies the role of the normality assumption in the behavior of the switching functions  $g$ . The case of a single control is the pivotal one.

**Lemma 4.2.1** *Let  $A \in \mathbb{M}_{N \times N}$  and  $b \in \mathbb{R}^N$  be such that*

$$\text{Rank}[b, Ab, \dots, A^{N-1}b] = N. \quad (4.2.1)$$

Take  $\zeta \in \mathbb{R}^N$ , with  $\|\zeta\| = 1$ , and define, for  $s \in [0, \infty)$

$$g(s) = \langle e^{As}b, \zeta \rangle. \quad (4.2.2)$$

Then there exists a constant  $\mathcal{L}$ , depending only on  $A, b, N$  such that

$$\sum_{i=0}^{N-1} |g^{(i)}(s)| \geq \mathcal{L} e^{-\|A\|s}. \quad (4.2.3)$$

**Proof.** Set

$$K = \begin{pmatrix} b \\ Ab \\ \vdots \\ A^{N-1}b \end{pmatrix}$$

and observe that, by (4.2.1)

$$\mathcal{L} = \min_{\|\zeta\|=1} \|K\zeta\| > 0. \quad (4.2.4)$$

Fix  $\zeta \in \mathbb{R}^N$  with  $\|\zeta\| = 1$  and write  $\zeta_1(s) = e^{sA^T}\zeta$ . Observe that  $\zeta = e^{-A^T s}\zeta_1(s)$  and  $\|\zeta_1(s)\| \geq e^{-s\|A\|}$ . We compute now, for  $i = 0, 1, \dots, N-1$ ,

$$g^{(i)}(s) = \langle e^{As}A^i b, \zeta \rangle = \langle A^i b, \zeta_1(s) \rangle.$$

Therefore,

$$K\zeta_1(s) = \begin{pmatrix} b \\ Ab \\ \vdots \\ A^{N-1}b \end{pmatrix} \zeta_1(s) = \begin{pmatrix} g^{(0)}(s) \\ g^{(1)}(s) \\ \vdots \\ g^{(N-1)}(s) \end{pmatrix}$$

Using (4.2.4) we have that

$$\|K\zeta_1(s)\| \geq \mathcal{L}e^{-s\|A\|}.$$

On the other hand,

$$\|K\zeta_1(s)\| \leq \sum_{i=0}^{N-1} |g^{(i)}(s)|$$

and the proof is concluded.  $\square$

The next Lemma is crucial for estimating the number of zero points of the switching function  $g$  (corresponding to the number of switching points of the optimal control associated with  $g$ ) and for studying the multiplicity order at zero points of  $g$ .

**Lemma 4.2.2** *Let  $A \in \mathbb{M}_{N \times N}$  and  $b \in \mathbb{R}^N$  be satisfying (4.2.1). Take  $\zeta \in \mathbb{R}^N$ ,  $\|\zeta\| = 1$ , and fix  $T > 0$ . Let  $g(s)$ ,  $s \in [0, T]$ , be defined as in (4.2.34).*

*Then there exist disjoint sets  $I_0, I_1, \dots, I_{N-1}$  and numbers  $\mathcal{N}_i$ , depending only on  $A, b, T$  and  $N$  such that*

$$[0, T] = \bigcup_{i=0}^{N-1} I_i$$

*and, for all  $i = 0, 1, \dots, N-1$ , the set  $I_i$  is the disjoint union of at most  $\mathcal{N}_i$  intervals. Moreover, for each  $i = 0, 1, \dots, N-1$ , for all  $s \in I_i$ , we have*

$$|g^{(i)}(s)| \geq \frac{\mathcal{L}}{N} e^{-\|A\|s}. \quad (4.2.5)$$

**Proof.** We proceed by induction for  $i$  from 0 to  $N-1$ . Set

$$c(s) = \frac{\mathcal{L}e^{-\|A\|s}}{N}, \quad (4.2.6)$$

and

$$J_0 = \{s \in (0, T) \mid |g(s)| < c(s)\}.$$

Since  $J_0$  is open, we can write it as the disjoint union of countably many intervals,

$$J_0 = \bigcup_{k=1}^{\infty} (a_{2k}, a_{2k+1}), \quad (4.2.7)$$

where  $0 < a_2 < a_3 < a_4 < \dots < a_{2k} < a_{2k+1} \dots < T$ .

Observe that, by contradiction, if  $s \in [a_{2k+1}, a_{2k+2}]$ , then  $|g(s)| \geq c(s)$ .

Now, fix  $k$  and consider the intervals  $(a_{2k}, a_{2k+1}), \dots, (a_{2(k+N-1)}, a_{2(k+N)-1})$ . Set, for  $j = 0, 1, \dots, N-1$

$$(a_{2(k+j)}, a_{2(k+j)+1}) := I_j^-,$$



and, for  $j = 0, 1, \dots, N - 2$

$$[a_{2(k+j)+1}, a_{2(k+j+1)}] := I_j^+.$$

Observe that

$$\bigcup_{j=0}^{N-1} I_j^- \cup \bigcup_{j=0}^{N-2} I_j^+ = (a_{2k}, a_{2(k+N)-1}).$$

We are going to give a lower bound on  $|a_{2(k+N)-1} - a_{2k}|$  independent of  $k$ , from which it will follow that the intervals  $(a_{2k}, a_{2k+1})$  are nonempty only for finitely many  $k$ .

Observe that for each  $j = 0, 1, \dots, N - 2$ , there exists at least one point  $c_j^1 \in I_j^+$  such that  $g'(c_j^1) = 0$ . Therefore, there exist at least  $N - 2$  points, say  $c_j^2$  for  $j = 0, 1, \dots, N - 3$ , such that

$$c_j^2 \in (c_j^1, c_{j+1}^1) \text{ and } g''(c_j^2) = 0.$$

Proceeding by induction we see that, for each  $i = 1, \dots, N - 1$ , there exists at least one point  $c_i \in (a_{2k+1}, a_{2(k+N)-1})$  such that  $g^{(i)}(c_i) = 0$ .

Pick any  $s_0 \in (a_{2k}, a_{2k+1})$ . We have

$$|g(s_0)| < c(s_0), \quad (4.2.8)$$

$$\begin{aligned} |g^{(i)}(s_0)| &= |g^{(i)}(s_0) - g^{(i)}(c_i)| = \left| \int_{s_0}^{c_i} g^{(i+1)}(s) ds \right| \\ &\leq \int_{a_{2k}}^{a_{2(k+N)-1}} |g^{(i+1)}(s)| ds \leq (a_{2(k+N)-1} - a_{2k}) e^{\|A\|T} \|A^{i+1}b\|. \end{aligned}$$

Therefore,

$$\sum_{i=1}^{N-1} |g^{(i)}(s_0)| \leq (a_{2(k+N)-1} - a_{2k}) e^{\|A\|T} \sum_{i=1}^{N-1} \|A^{i+1}b\|. \quad (4.2.9)$$

On the other hand, recalling (4.2.3), (4.2.6) and (4.2.8) we have

$$\begin{aligned} \sum_{i=1}^{N-1} |g^{(i)}(s_0)| &\geq \mathcal{L}e^{-\|A\|s_0} - c(s_0) \\ &= \frac{N-1}{N} \mathcal{L}e^{-\|A\|s_0} \geq \frac{N-1}{N} \mathcal{L}e^{-\|A\|T}. \end{aligned} \quad (4.2.10)$$

From (4.2.9) and (4.2.10) we obtain

$$a_{2(k+N)-1} - a_{2k} \geq \frac{(N-1)\mathcal{L}e^{-2\|A\|T}}{N \sum_{i=1}^{N-1} \|A^{i+1}b\|}, \quad (4.2.11)$$

which is the desired estimate. Observe that the right hand side of (4.2.11) depends only on  $A, b, T$  and  $N$ .

We set now  $\mathcal{N}_0$  to be the number of nonempty intervals appearing in (4.2.7), and recall that we just proved that  $\mathcal{N}_0$  depends only on  $A, b, T$  and  $N$ , actually

$$\mathcal{N}_0 \leq \frac{N}{N-1} \frac{T}{\mathcal{L}} e^{2\|A\|T} \sum_{i=1}^{N-1} \|A^{i+1}b\| + N. \quad (4.2.12)$$

Set  $I_0 = [0, T] \setminus J_0$  and observe that we have completed the proof of the lemma for  $i = 0$ .

Fix now a nonempty interval  $(a, b)$  appearing in (4.2.7). Set

$$J_1^{(a,b)} = \{s \in (a, b) \mid |g'(s)| < c(s)\}.$$

Pick any  $s_0 \in J_1^{(a,b)}$  and observe that recalling (4.2.3) and (4.2.6),  $\sum_{i=1}^{N-1} |g^{(i)}(s_0)| \geq \frac{N-1}{N} \mathcal{L} e^{-\|A\|s}$ . By using in  $J_1^{(a,b)}$  the same argument as above, with  $g'$  in place of  $g$ , we see that  $J_1^{(a,b)}$  is the union of finitely many disjoint intervals  $(a_{2k}^1, a_{2k+1}^1)$ ,  $k = 1, 2, \dots, \mathcal{N}_1^{(a,b)}$  where

$$\mathcal{N}_1^{(a,b)} \leq \frac{N}{N-2} \frac{|b-a|}{\mathcal{L}} e^{2\|A\|T} \sum_{j=2}^{N-1} \|A^{j+1}b\| + N - 1. \quad (4.2.13)$$

We define

$$I_1^{(a,b)} = \{s \in (a, b) \mid |g'(s)| \geq c(s)\}$$

and  $I_1$  to be the union of the  $I_1^{(a,b)}$ , over all the at most  $\mathcal{N}_0$  nonempty intervals  $(a, b)$  in  $J_0$ , i.e.

$$I_1 = \bigcup_{l=1}^{\mathcal{N}_0} I_1^{(a_l, b_l)} = \bigcup_{l=1}^{\mathcal{N}_0} \bigcup_{h=1}^{\mathcal{N}_1^{l+1}} (a_l^h, b_l^h).$$

Observe that the number  $\mathcal{N}_1$  of intervals appearing in the above union is bounded from above. More precisely, recalling (4.2.11) we have

$$\mathcal{N}_1 \leq \frac{N}{N-2} \frac{T}{\mathcal{L}} e^{2\|A\|T} \sum_{i=2}^{N-1} \|A^{i+1}b\| + N\mathcal{N}_0. \quad (4.2.14)$$

After this step, we formulate our induction process. We are going to construct, for each  $i = 2, \dots, N-1$ , two disjoint sets  $I_i, J_i$  with the following properties

(ind1) for every  $s \in I_i$ ,  $|g^{(i)}(s)| \geq c(s)$ ,

(ind2) for every  $s \in J_i$ ,  $\sum_{i+1}^{N-1} |g^{(j)}(s)| \geq \frac{N-i-1}{N} c(s)$ ,

(ind3)  $J_i \cup I_i = J_{i-1}$ ,

(ind4)  $J_i$  is a finite union of open intervals, whose number is at most  $\mathcal{N}_i$  and  $\mathcal{N}_i$  depends only on  $T, \mathcal{L}, A, b, N, i$ ,

(ind5)  $I_i$  is the finite union of at most  $\mathcal{N}_i + 1$  intervals.

For  $i = 0, 1$  the above construction was already performed (take  $J_{-1} = (0, T)$ ). Pick any  $i = 2, 3, \dots, N - 3$  and consider the set

$$J_{i+1} := \{s \in J_i \mid |g^{(i+1)}(s)| < c(s)\}.$$

For every connected component  $(a, b)$  of  $J_i$ , we are going to prove that  $J_{i+1}^{(a,b)} := J_{i+1} \cap (a, b)$  is a finite union of intervals, and give a bound on their number  $\mathcal{N}_{i+1}^{(a,b)}$ . Recalling (ind2), for every  $s \in J_{i+1}^{(a,b)}$  we have

$$\sum_{j=i+1}^{N-1} |g^{(j)}(s)| \geq \frac{N-i-1}{N} c(s).$$

By using the same argument developed for  $i = 0$ , with  $g^{(i+1)}$  in place of  $g$ , we see that  $J_{i+1}^{(a,b)}$  is the union of finitely many disjoint open intervals  $(a_{2k}^{i+1}, a_{2k+1}^{i+1})$ ,  $k = 1, 2, \dots, \mathcal{N}_{i+1}^{(a,b)}$ , where

$$\mathcal{N}_{i+1}^{(a,b)} \leq \frac{N}{N-i-2} \frac{|b-a|}{\mathcal{L}} e^{2\|A\|T} \sum_{j=i+2}^{N-1} \|Ab\| + (N-i+1). \quad (4.2.15)$$

We define

$$I_{i+1}^{(a,b)} = \{s \in (a, b) \mid |g^{(i+1)}(s)| \geq c(s)\}$$

and observe that  $I_{i+1}^{(a,b)}$  is the union of at most  $\mathcal{N}_{i+1}^{(a,b)} + 1$  intervals.

We finally set  $I_{i+1}$  to be the union of the  $I_{i+1}^{(a_j, b_j)}$  over all the (at most  $\mathcal{N}_i$ ) connected components  $(a_j, b_j)$  of  $J_i$ . Therefore,  $I_{i+1}$  is the union of at most  $\mathcal{N}_{i+1}$  intervals, where

$$\mathcal{N}_{i+1} = \sum_{j=1}^{\mathcal{N}_i} \mathcal{N}_{i+1}^{(a_j, b_j)} \leq \frac{N}{N-i-2} \frac{T}{\mathcal{L}} e^{2\|A\|T} \sum_{j=i+2}^{N-1} \|A^{j+1}b\| + (N-i)\mathcal{N}_i. \quad (4.2.16)$$

Finally we observe that  $J_{i+1}$  is the union of at most  $\mathcal{N}_{i+1}$  open intervals.

If  $i = N - 2$ , we observe that for each  $s \in J_{N-2}$ , recalling (ind2) we have  $|g^{(N-1)}(s)| \geq c(s)$ . Therefore we set  $J_{N-1} = \emptyset$  and  $I_{N-1} = J_{N-2}$ . The proof is concluded.  $\square$

We are now going to state our main result.

**Theorem 4.2.1** Consider the linear control system

$$\dot{x} = Ax + Bu, \quad (4.2.17)$$

where  $A \in \mathbb{M}_{N \times N}$ ,  $B \in \mathbb{M}_{N \times M}$ ,  $M \leq N$  and  $u = (u_1, u_2, \dots, u_M) \in \mathbb{R}^M$ ,  $|u_j| \leq 1$  for  $j = 1, 2, \dots, M$ .

Assume that (4.2.17) is normal, i.e., for every column  $b_j$ ,  $j = 1, 2, \dots, M$  of  $B$ ,

$$\text{rank} [b_j, Ab_j, \dots, A^{N-1}b_j] = N.$$

Then for all  $T > 0$  there exists a constant  $\gamma > 0$ , depending only on  $N, M, A, B, T$  such that for all  $x, y \in \mathcal{R}^T$ , for all  $\zeta \in N_{\mathcal{R}^T}(x)$ , it holds

$$\langle \zeta, y - x \rangle \leq -\gamma \|\zeta\| \|y - x\|^N. \quad (4.2.18)$$

**Remark 4.2.1** The power  $N$  in (4.2.18) is optimal, as the example,  $\ddot{x} = u$  for  $u \in [-1, 1]$ , shows.

**Proof.** We first consider the case  $M = 1$ , so (4.2.17) reads as

$$\dot{x} = Ax + bu \quad , \quad |u| \leq 1.$$

Fix  $\bar{x} \in \partial \mathcal{R}^T$  together with an optimal control  $\bar{u}(\cdot)$  steering 0 to  $\bar{x}$  in time  $T$ . By Pontryagin's maximum principle, for a.e.  $t \in [0, T]$ ,

$$\bar{u}(t) = \text{sign} \langle \zeta, e^{A(T-t)}b \rangle.$$

Taking  $\bar{y} \in \mathcal{R}^T$  together with a control  $u(\cdot)$  steering 0 to  $\bar{y}$ , we compute:

$$\begin{aligned} \langle \zeta, \bar{y} - \bar{x} \rangle &= \int_0^T \langle \zeta, e^{A(T-t)}b(u(t) - \bar{u}(t)) \rangle dt \\ &= - \int_0^T |\langle \zeta, e^{A(T-t)}b \rangle| |u(t) - \bar{u}(t)| dt. \end{aligned}$$

Set  $K(t) = \frac{1}{2}|u(t) - \bar{u}(t)|$  and observe that  $0 \leq K(t) \leq 1$  a.e.  $t \in [0, T]$ , and

$$\langle \zeta, \bar{y} - \bar{x} \rangle = -2 \int_0^T |\langle \zeta, e^{A(T-t)}b \rangle| K(t) dt = -2 \int_0^T |\langle \zeta, e^{At}b \rangle| K(T-t) dt. \quad (4.2.19)$$

Moreover,

$$\|\bar{y} - \bar{x}\| = \left\| \int_0^T e^{A(T-t)}b(u(t) - \bar{u}(t)) dt \right\| \leq 2e^{T\|A\|} \|b\| \int_0^T K(t) dt. \quad (4.2.20)$$

Set, for  $s \in [0, +\infty)$ ,

$$g(s) = \langle e^{As}b, \zeta \rangle.$$

By Lemma 4.2.2 there exist disjoint sets  $I_0, I_1, \dots, I_{N-1}$  and constants  $N_i$  such that  $[0, T] = \bigcup_{i=0}^{N-1} I_i$ , each  $I_i$  is the disjoint union of at most  $N_i$  intervals and (4.2.5) holds. We rewrite

$$\langle \zeta, \bar{y} - \bar{x} \rangle = -2 \sum_{i=0}^{N-1} \int_{I_i} |g(s)| K_1(s) ds \quad (4.2.21)$$

where  $K_1(s) = K(T - s)$ . We are now going to discuss separately the integrals  $\int_{I_i} |g(s)| K_1(s) ds$ , for all  $i = 0, 1, \dots, N - 1$ .

For  $i = 0$ , we have, recalling (4.2.5) and (4.2.4)

$$\int_{I_0} |g(s)| K_1(s) ds \geq \frac{\mathcal{L}}{N} e^{-\|A\|T} \int_{I_0} K_1(s) ds. \quad (4.2.22)$$

Fix  $i = 1, 2, \dots, N - 1$ , and write, recalling Lemma 4.2.2,

$$\bar{I}_i = \bigcup_{j=1}^{N_i} [a_{i,j}, b_{i,j}]$$

and all *open* intervals  $(a_{i,j}, b_{i,j})$  are disjoint. Recalling (4.2.5), we have, for all  $s \in I_i$ ,  $|g^{(i)}(s)| \geq \frac{\mathcal{L}}{N} e^{-\|A\|T}$ . Fix  $j \in \{1, 2, \dots, N_i\}$ . We now apply inductively Lemma 7.3.2 on  $[a_{i,j}, b_{i,j}]$  with the functions  $g^{(i-k-1)}$  in place of  $f$ , for  $k = 0, \dots, i - 1$ . Let  $k = 0$  and set  $f = g^{(i-1)}$ , the assumption (7.3.1) is satisfied with  $C = \frac{\mathcal{L}}{N} e^{-\|A\|T}$ . Then Lemma 7.3.2 yields that for some point  $c_{i,j}^0 \in [a_{i,j}, b_{i,j}]$  we have both

$$|g^{(i-1)}(s)| \geq C(c_{i,j}^0 - s) \quad \forall s \in [a_{i,j}, c_{i,j}^0],$$

and

$$|g^{(i-1)}(s)| \geq C(s - c_{i,j}^0) \quad \forall s \in [c_{i,j}^0, b_{i,j}].$$

Let  $k = 1$ . By applying Lemma 7.3.2 on each of the two (possibly degenerate) intervals  $[a_{i,j}, c_{i,j}^0]$ ,  $[c_{i,j}^0, b_{i,j}]$  to the function  $f = g^{(i-2)}$  and the constant  $C$ , we find a suitable points  $c_{i,j}^1 \in [a_{i,j}, c_{i,j}^0]$  and  $c_{i,j}^2 \in [c_{i,j}^0, b_{i,j}]$  such that we have both

$$|g^{(i-2)}(s)| \geq \frac{C}{2} (c_{i,j}^1 - s)^2 \quad \forall s \in [a_{i,j}, c_{i,j}^1],$$

$$|g^{(i-2)}(s)| \geq \frac{C}{2} (s - c_{i,j}^1)^2 \quad \forall s \in [c_{i,j}^1, c_{i,j}^0]$$

and

$$|g^{(i-2)}(s)| \geq \frac{C}{2} (c_{i,j}^2 - s)^2 \quad \forall s \in [c_{i,j}^0, c_{i,j}^2],$$

$$|g^{(i-2)}(s)| \geq \frac{C}{2} (s - c_{i,j}^2)^2 \quad \forall s \in [c_{i,j}^2, b_{i,j}].$$

By continuing the induction process until  $k = i - 1$ , we split the interval  $[a_{i,j}, b_{i,j}]$  into at most  $2^i$  intervals  $[a_{i,j} = c_{i,j}^0, c_{i,j}^1], [c_{i,j}^1, c_{i,j}^2], \dots, [c_{i,j}^{2^i-1}, b_{i,j} = c_{i,j}^{2^i}]$  (some of them being possibly degenerate) such that for all  $l = 0, 1, \dots, 2^i - 1$  and  $s \in [c_{i,j}^l, c_{i,j}^{l+1}]$  one has

$$\text{either } |g(s)| \geq \frac{C}{j!} (s - c_{i,j}^l)^i \quad \text{or} \quad |g(s)| \geq \frac{C}{i!} (c_{i,j}^{l+1} - s)^i. \quad (4.2.23)$$

Recalling (4.2.21), and the above discussion we have

$$\begin{aligned} \langle \zeta, \bar{y} - \bar{x} \rangle &= -2 \sum_{i=0}^{N-1} \int_{I_i} |g(s)| K_1(s) ds \\ &= -2 \left[ \int_{I_0} |g(s)| K_1(s) ds + \sum_{i=1}^{N-1} \sum_{j=0}^{\mathcal{N}_i} \sum_{l=0}^{2^i-1} \int_{c_{i,j}^l}^{c_{i,j}^{l+1}} |g(s)| K_1(s) ds \right]. \end{aligned}$$

Recalling (4.2.22) and (4.2.23), we obtain from the above inequality that

$$\langle \zeta, \bar{y} - \bar{x} \rangle \leq -\frac{2\mathcal{L}}{N} e^{-\|A\|T} \left[ \int_{I_0} K_1(s) ds + \sum_{i=1}^{N-1} \sum_{j=0}^{\mathcal{N}_i} \sum_{l=0}^{2^i-1} \int_{c_{i,j}^l}^{c_{i,j}^{l+1}} \frac{|s - \bar{c}_{i,j}^l|^i}{i!} K_1(s) ds \right] \quad (4.2.24)$$

where  $\bar{c}_{i,j}^l$  is either  $c_{i,j}^l$  or  $c_{i,j}^{l+1}$ , according to the two possibilities appearing in (4.2.23). Applying Lemma 7.3.1 to each summand of (4.2.24) we therefore obtain

$$\begin{aligned} \langle \zeta, \bar{y} - \bar{x} \rangle &\leq -2 \frac{\mathcal{L}}{N} e^{-\|A\|T} \left[ \int_{I_0} K_1(s) ds + \sum_{i=1}^{N-1} \sum_{j=0}^{\mathcal{N}_i} \sum_{l=0}^{2^i-1} \frac{\left( \int_{c_{i,j}^l}^{c_{i,j}^{l+1}} K_1(s) ds \right)^{i+1}}{(i+1)!} \right] \\ &\quad \text{(using the convexity of } x \mapsto x^{i+1} \text{ on the positive half line)} \\ &\leq -\frac{2\mathcal{L}}{N} e^{-\|A\|T} \left[ \int_{I_0} K_1(s) ds + \sum_{i=1}^{N-1} \sum_{j=0}^{\mathcal{N}_i} \frac{1}{(i+1)! 2^{i^2}} \left( \int_{a_{i,j}}^{b_{i,j}} K_1(s) ds \right)^{i+1} \right] \\ &\leq -\frac{2\mathcal{L}}{N} e^{-\|A\|T} \left[ \int_{I_0} K_1(s) ds + \sum_{i=1}^{N-1} \frac{1}{(i+1)! 2^{i^2} \mathcal{N}_i^i} \left( \int_{I_i} K_1(s) ds \right)^{i+1} \right]. \end{aligned}$$

Thus,

$$\begin{aligned} \langle \zeta, \bar{y} - \bar{x} \rangle &\leq -\frac{2\mathcal{L}}{N} e^{-\|A\|T} \left[ \frac{1}{|J_0|^{N-1}} \left( \int_{I_0} K_1(s) ds \right)^N \right. \\ &\quad \left. + \sum_{i=1}^{N-1} \frac{1}{(i+1)! 2^{i^2} \mathcal{N}_i^i |I_i|^{N-i-1}} \left( \int_{I_i} K_1(s) ds \right)^N \right]. \quad (4.2.25) \end{aligned}$$

Recalling (4.2.12), (4.2.15) and (4.2.16), we see that  $\mathcal{N}_0 \leq \mathcal{N}_1 \leq \dots \leq \mathcal{N}_N \leq C(A, b, N)(e^{2\|A\|T} + N!)$  where  $C(A, b, N)$  depends only on  $A, b, N$ . Therefore, we obtain finally from (4.2.25) that

$$\langle \zeta, \bar{y} - \bar{x} \rangle \leq -C(A, b, N, T)e^{-\|A\|T} \left( \int_0^T |u(t) - \bar{u}(t)| \right)^N, \quad (4.2.26)$$

where  $C(A, b, N, T)$  is a positive constant, depending only on  $A, b, N, T$  such that

$$\liminf_{T \rightarrow 0} C(A, b, N, T) > 0. \quad (4.2.27)$$

Recalling (4.2.20), we complete the proof for the case  $M = 1$  (i.e., a scalar control) by setting

$$\gamma = 2^N e^{N+1} \|b\|^N C(A, b, N, T).$$

In the case  $M > 1$ , pick  $\bar{x} \in \partial \mathcal{R}^T$  together with an optimal control  $\bar{u}(\cdot) = (\bar{u}_1(\cdot), \bar{u}_2(\cdot), \dots, \bar{u}_M(\cdot))$  steering the origin to  $\bar{x}$  in the optimal time  $T$ , together with  $\bar{y} \in \mathcal{R}^T$  and a control  $u(\cdot) = (u_1(\cdot), u_2(\cdot), \dots, u_M(\cdot))$  steering the origin to  $\bar{y}$  in time  $T$ . Then, for each  $\zeta \in N_{\mathcal{R}^T}(\bar{x})$ ,  $\|\zeta\| = 1$ , we can write

$$\langle \zeta, \bar{y} - \bar{x} \rangle \leq \int_0^T \langle \zeta, e^{A(T-s)} B w(s) \rangle ds = \sum_{i=1}^M \int_0^T \langle \zeta, e^{A(T-s)} b_i w_i(s) \rangle ds, \quad (4.2.28)$$

where  $B = (b_i)_{i=1,2,\dots,M}$  and  $w_i(s) = u_i(s) - \bar{u}_i(s)$ .

Recalling Pontryagin's maximum principle we have also

$$\langle \zeta, \bar{y} - \bar{x} \rangle = - \sum_{i=1}^M \int_0^T |\langle \zeta, e^{A(T-s)} b_i \rangle| |u_i(s) - \bar{u}_i(s)| ds. \quad (4.2.29)$$

Moreover,

$$\|\bar{y} - \bar{x}\| \leq e^{\|A\|T} \sum_{i=1}^M \|b_i\| \int_0^T |u_i(s) - \bar{u}_i(s)| ds. \quad (4.2.30)$$

We now apply  $M$  times (4.2.26) with each summand of the right hand side of (4.2.28) in place of the left hand side of (4.2.26) and obtain, using (4.2.29), that

$$\langle \zeta, \bar{y} - \bar{x} \rangle \leq -C'(A, B, T, N, M) e^{-\|A\|T} \sum_{i=1}^M \left( \int_0^T |u_i(s) - \bar{u}_i(s)| ds \right)^N,$$

where the positive constant  $C'$  depends only on  $A, B, T, N, M$  and

$$\liminf_{T \rightarrow 0} C'(A, B, T, N, M) > 0.$$

We conclude the proof by applying (4.2.30) and setting

$$\gamma = 2^N e^{-(N+1)\|A\|T} \|B\| C''(A, B, T, N, M),$$

where  $C''$  is a constant enjoying the some properties as  $C'$ .  $\square$

#### 4.2.2 Linear nonautonomous control system in the case $N = 2$

The following Lemma is a first step for studying the reachable sets in the case of nonlinear control system by using our linearization approach. It says that under the rank condition (normality) at 0 of the linear nonautonomous control system (4.2.31), the strict convexity of the reachable sets still remain up to sufficiently small time. For future use, we will consider that  $A$  is just measurable instead of continuous.

**Lemma 4.2.3** *Let  $N = 2$ . Consider the linear nonautonomous control system*

$$\dot{x}(t) = A(t)x(t) + B(t)u(t), \quad (4.2.31)$$

where  $A : \mathbb{R}^+ \rightarrow \mathbb{M}_{2 \times 2}$  and  $B : \mathbb{R}^+ \rightarrow \mathbb{M}_{2 \times M}$ ,  $1 \leq M \leq 2$  and  $u = (u_1, u_M)$ ,  $|u_j| \leq 1$  for  $j = 1, M$ , satisfying the following assumptions:

( $C_0$ )  $A(\cdot)$  is measurable and

$$\|A(t) - A(0)\| \leq Lt \text{ for all } t \geq 0,$$

( $C_1$ )  $B(\cdot)$  is of class  $\mathcal{C}^1$  and

$$\left\| \frac{d}{dt} B(t) \right\| \leq 2Lt \text{ for all } t \geq 0,$$

where  $L$  is a positive constant;

( $C_2$ ) Rank  $[b_j(0), A(0)b_j(0)] = 2$  for  $j = 1, M$  where  $B(\cdot) = [b_1(\cdot), b_M(\cdot)]$ .

Then there exist a time  $\mathcal{T} = \mathcal{T}(A, B, L) > 0$  and a constant  $\gamma > 0$  depending only on  $A(0), B(0), L$  such that for every  $0 \leq \tau \leq \mathcal{T}$ , for every  $x, y \in \mathcal{R}^\tau$ , for every  $\zeta \in N_{\mathcal{R}^\tau}(x)$ , we have

$$\langle \zeta, y - x \rangle \leq -\gamma \|\zeta\| \|y - x\|^2.$$

**Remark 4.2.2** *Observe that condition ( $C_0$ ) implies that  $t = 0$  is a continuity point for  $A(\cdot)$ , so that  $A(0)$  in ( $C_0$ ) is meaningful.*

In order to prove Lemma 4.2.3 we need some notation and a preliminary result. We denote by  $M(\cdot, \cdot)$  the fundamental matrix solution of

$$\begin{cases} \frac{\partial}{\partial t} M(t, s) = A(t)M(t, s) \text{ for } t, s \geq 0 \\ M(s, s) = \mathbb{I} \end{cases} \quad (4.2.32)$$

and by  $M_0(\cdot, \cdot)$  the fundamental matrix solution of

$$\begin{cases} \frac{\partial}{\partial t} M_0(t, s) = A(0)M_0(t, s) \text{ for } t, s \geq 0 \\ M_0(s, s) = \mathbb{I} \end{cases} \quad (4.2.33)$$



Let  $b(\cdot)$  be a column of  $B(\cdot)$ , let  $\mathcal{T} > 0$  and define

$$g(t) = \langle M(\mathcal{T}, t)b(t), \zeta \rangle, \quad (4.2.34)$$

and

$$g_0(t) = \langle M_0(\mathcal{T}, t)b(0), \zeta \rangle. \quad (4.2.35)$$

Our preliminary result will permit to transfer to  $g$  the properties of the function  $g_0$  proved in Lemma 4.2.2.

**Lemma 4.2.4** *Let  $g$  be defined according to (4.2.34) and let the assumptions of Lemma 4.2.3 hold. Then for  $\mathcal{T} > 0$  sufficiently small there exist disjoint sets  $I_0, I_1$  and numbers  $\mathcal{N}_0, \mathcal{N}_1$  depending only on  $A(0), B(0), L, \mathcal{T}$  such that*

(a)  $[0, \mathcal{T}] = I_0 \cup I_1$ ;

(b)  $I_i$  is the disjoint of at most  $\mathcal{N}_i$  intervals,  $i = 0, 1$ ;

(c) for each  $x \in I_0$

$$|g(s)| \geq \frac{\mathcal{L}}{4} e^{-\|A(0)\|s};$$

(d)  $g'$  has constant sign in every connected component of  $I_1$  and, for each  $s \in I_1$

$$|g'(s)| \geq \frac{\mathcal{L}}{4} e^{-\|A(0)\|s}.$$

**Remark 4.2.3**  $I_0$  and  $I_1$  are exactly the intervals provided by Lemma 4.2.2 for the case  $N = 2$  with  $g_0$  in place of  $g$ .

**Proof of Lemma 4.2.4.** Let  $g_0$  be defined according with (4.2.35). By using Gronwall's inequality and  $(C_0), (C_1)$  we can find a continuous function  $K(\mathcal{T}) \geq 0$  such that

$$\lim_{\mathcal{T} \rightarrow 0^+} \frac{K(\mathcal{T})}{\mathcal{T}} = K_1 > 0,$$

$$|g(t) - g_0(t)| \leq K(\mathcal{T}) \text{ for all } 0 \leq t \leq \mathcal{T}, \quad (4.2.36)$$

and

$$|g'(t) - g'_0(t)| \leq K(\mathcal{T}) \text{ for all } 0 \leq t \leq \mathcal{T}. \quad (4.2.37)$$

Let  $\mathcal{T} > 0$  be such that

$$K(\mathcal{T}) \leq \frac{\mathcal{L}}{4} e^{-\|A(0)\|\mathcal{T}}.$$

Let  $I_0, I_1$  be given by lemma 4.2.2 with  $g_0$  in place of  $g$ . For each  $s \in I_0$  we have  $|g_0(s)| \geq \frac{\mathcal{L}}{2} e^{-\|A(0)\|s}$ . By (4.2.36), (c) follows.

Fix now a connected component  $J$  of  $I_1$  and observe that, being  $g'_0$  continuous, its sign is constant on  $J$ . Since on  $J$  we have on one hand

that  $|g'_0(s)| \geq \frac{\mathcal{L}}{2}e^{-\|A(0)\|T}$  and on the other hand that  $|g'(s) - g'_0(s)| \leq \frac{\mathcal{L}}{4}e^{-\|A(0)\|T}$ , we obtain both that  $g'$  does not change its sign on  $J$  and satisfies the inequality in (d). The proof is concluded.  $\square$

**Proof of Lemma 4.2.3.** The same argument of the proof of the constant coefficient case can be applied.  $\square$

### 4.3 Nonlinear control system in the case $N = 2$

#### 4.3.1 Strict convexity of the reachable set

In this subsection, we will prove that under the rank condition at 0 together with the Taylor development at 0 of the nonlinear control system, the reachable set is strictly convex up to sufficiently small time.

**Theorem 4.3.1** *Consider the control system*

$$\begin{cases} \dot{x}(t) = F(x(t)) + G(x(t))u(t), \\ x(0) = 0, \end{cases} \quad (4.3.1)$$

where  $u = (u_1, u_M) \in [-1, 1]^2$ ,  $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ ,  $G : \mathbb{R}^2 \rightarrow \mathbb{M}_{2 \times M}$ ,  $1 \leq M \leq 2$ , are of class  $C^{1,1}$  (with Lipschitz constant  $L$ ) and

(i)  $F(0) = 0$ ,

(ii)  $\text{Rank} [G_i(0), DF(0)G_i(0)] = 2$  for  $i = 1, 2$  where  $G = (G_1, G_2)$ ,

(iii)  $DG(0) = 0$ .

Then there exists  $\mathcal{T} > 0$ , depending only on  $L, DF(0), G(0)$  such that for every  $0 < \tau < \mathcal{T}$  the reachable sets  $\mathcal{R}^\tau$  is strictly convex. More precisely, for every  $x_1 \in \partial\mathcal{R}^\tau$  and  $x_2 \in \mathcal{R}^\tau$ , for every  $\zeta \in N_{\mathcal{R}^\tau}^P(x_1)$ , one has

$$\langle \zeta, x_2 - x_1 \rangle \leq -\gamma \|\zeta\| \|x_2 - x_1\|^2. \quad (4.3.2)$$

where  $\gamma$  is a positive constant.

**Proof.** We first prove the Theorem for  $M = 1$  (scalar control).

Fix  $\tau > 0$  and  $x_1 \in \partial\mathcal{R}^\tau$ , together with a optimal control  $u_1(\cdot)$  steering 0 to  $x_1$  and associate trajectory  $x_1(\cdot)$ . Take any  $x_2 \in \mathcal{R}^\tau$  together with  $u_2(\cdot)$  steering 0 to  $x_2$  and associate trajectory  $x_2(\cdot)$ . Set  $x(t) = x_2(t) - x_1(t)$ . Then, for a.e.  $t \in [0, \tau]$ ,

$$\dot{x}(t) = A_1(t)x(t) + G(x_2(t))w(t), \quad (4.3.3)$$

where  $w(t) = u_2(t) - u_1(t)$  and

$$A_1(t) = \int_0^1 DF(x_1(t) + \tau x(t))d\tau + \int_0^1 DG(x_1(t) + \tau x(t))d\tau u_1(t).$$

Let  $z(\cdot)$  be the solution of the linear system which is defined along the optimal trajectory  $x_1(\cdot)$ :

$$\begin{cases} \dot{z}(t) = A(t)z(t) + G(x_1(t))w(t), \\ z(0) = 0, \end{cases} \quad (4.3.4)$$

where  $A(t) = DF(x_1(t)) + DG(x_1(t))u_1(t)$ .

We have

$$\begin{aligned} \frac{d}{dt}\|x(t) - z(t)\| &\leq \|A_1(t)x(t) - A(t)z(t)\| + |G(x_2(t)) - G(x_1(t))||w(t)| \\ &\leq \|A(t)\| \|x(t) - z(t)\| + \|A_1(t) - A(t)\| \|x(t)\| + Lt\|x(t)\| |w(t)| \\ &\leq L_1\|x(t) - z(t)\| + L\|x(t)\|^2 + Lt\|x(t)\| |w(t)|, \end{aligned}$$

where  $L_1 = \|DF(0)\| + 2Le^{2L\tau}$ .

Thus, by Gronwall inequality we get

$$\|x(t) - z(t)\| \leq e^{L_1 t} L \int_0^t (\|x(s)\|^2 + s\|x(s)\||w(s)|) ds. \quad (4.3.5)$$

On the other hand, observing that

$$\frac{d}{dt}\|\dot{x}(t)\| \leq L_2\|x(t)\| + L_3|w(t)|,$$

we also have

$$\|x(t)\| \leq L_3 e^{L_2 t} \int_0^t |w(s)| ds, \quad (4.3.6)$$

where  $L_2 = \|DF(0)\| + 4Le^{2L\tau}$  and  $L_3 = |G(0)| + e^{2L\tau}$ .

From (4.3.5) and (4.3.6), one obtains

$$\|x(t) - z(t)\| \leq L_5 t \left( \int_0^t |w(s)| ds \right)^2, \quad (4.3.7)$$

where  $L_5 = LL_3^2 e^{(L_1+2L_2)\tau} + LL_3 e^{(L_1+L_2)\tau}$ .

Since  $\bar{x}_1 \in \partial\mathcal{R}^\tau$ , by Pontryagin's maximum principle there exists an absolutely continuous function  $\lambda : [0, \tau] \rightarrow \mathbb{R}^2$  with the following properties

$$\dot{\lambda}(t) = -\lambda(t)A^T(t) \quad , \quad \lambda(\tau) = \zeta,$$

$$u_1(t) = \text{sign}\langle \lambda(t), G(x_1(t)) \rangle. \quad (4.3.8)$$

We set now  $b(t) = G(x_1(t))$  and consider the linear nonautonomous control system

$$\begin{cases} \dot{y}(t) = A(t)y(t) + b(t)u(t), \\ y(0) = 0, \end{cases} \quad (4.3.9)$$

together with the trajectory  $y_1(\cdot)$ , corresponding to the control  $u_1(\cdot)$ . Observe that  $A(\cdot)$  is measurable. Moreover, since both  $F$  and  $G$  are Lipschitz with constant  $L$  and  $DG(0) = 0$ , we have

$$\begin{aligned} \|A(t) - A(0)\| &= \|DF(x_1(t)) + DG(x_1(t))u_1(t) - DF(0)\| \\ &\leq 2L\|x_1(t)\| \leq 2L|G(0)|e^{2Lt}. \end{aligned}$$

Moreover,  $b'(t) = DG(x_1(t))\dot{x}_1(t)$  so that

$$|b'(t) - b(0)| = |b'(t)| \leq L\|x_1(t)\|(2L\|x_1(t)\| + |G(0)|) \leq Kt$$

where  $K = Le^{2L\tau}(2Le^{2L\tau} + 1)|G(0)|^2$ .

Therefore, by Lemma 4.2.3, if  $0 \leq \tau < C_1$  where  $C_1 > 0$  depends only on  $L, DF(0), G(0)$  then there exists a constant  $\gamma(\tau) > 0$ , depending only on  $L, DF(0), G(0), \tau$  such that for all  $\langle \zeta, y(\tau) - y_1(\tau) \rangle \leq -\gamma(\tau)\|y(\tau) - y_1(\tau)\|^2$ . In particular, recalling (4.2.26)

$$\langle \zeta, y_2(\tau) - y_1(\tau) \rangle \leq -\gamma_1(\tau) \left( \int_0^\tau w(s) ds \right)^2, \quad (4.3.10)$$

where  $y_2(\cdot)$  is the trajectory of (4.3.9) associated with the control  $u_2(\cdot)$ . Remark that  $\gamma_1(\tau)$  is bounded away from 0 as  $\tau \rightarrow 0^+$ . Moreover, one can see that  $z(t) = y_2(t) - y_1(t)$ . We compute

$$\begin{aligned} \langle \zeta, x_2 - x_1 \rangle &= \langle \zeta, x(\tau) - z(\tau) \rangle + \langle \zeta, z(\tau) \rangle \\ &\leq \|x(\tau) - z(\tau)\| + \langle \zeta, z(\tau) \rangle. \end{aligned}$$

Recalling (4.3.7) and (4.3.10), we obtain

$$\langle \zeta, x_2 - x_1 \rangle \leq (L_5\tau - \gamma_1(\tau)) \left( \int_0^\tau w(s) ds \right)^2. \quad (4.3.11)$$

Thus if  $\tau \leq \frac{\gamma_1(C_1)}{2}$  then

$$\langle \zeta, x_2 - x_1 \rangle \leq -\frac{\gamma_1(\tau)}{2} \left( \int_0^\tau w(s) ds \right)^2.$$

Setting  $\mathcal{T} = \min\{C_1, \frac{\gamma_1(C_1)}{2}\}$  and recalling (4.3.7) we obtain (4.3.2). The proof is completed by applying Proposition 7.2.1.

In the case  $M = 2$ , the proof is done by following entirely the above argument.  $\square$

The following Remark follows immediately from the proof of Theorem 4.3.1.

**Remark 4.3.1** *Let  $x_1(\cdot)$  and  $\lambda(\cdot)$  be in the proof of Theorem 4.3.1. For all  $0 < t \leq \tau = T(x_1)$ , one has  $\lambda(t) \in N_{\mathcal{R}^t}(x_1(T(x_1) - t))$ , more precisely*

$$\langle \lambda(t), y - x_1(T(x_1) - t) \rangle \leq -\gamma\|\lambda(t)\|\|y - x_1(T(x_1) - t)\|^2.$$

for all  $y \in \mathcal{R}^t$ .

### 4.3.2 Optimal points

We will study here the optimality of points near the origin. Before stating our result, we prefer to give the classical definition of optimal point.

**Definition 4.3.1** *Let  $x \in \mathbb{R}^N \setminus \{0\}$ . We say that  $x$  is optimal if and only if there exists a point  $x_1$  such that  $T(x_1) > T(x)$  and  $x = x_1(T(x_1) - T(x))$ , where  $x_1(\cdot)$  is an optimal trajectory steering  $x_1$  to 0 in optimal time  $T(x_1)$ .*

The following result on optimal points is important to prove the positive reach of the epi-graph of the minimum time function, which will be considered in the next subsection.

**Theorem 4.3.2** *Let  $N = 2$  and let the assumption of Theorem 4.3.1 hold. Let  $T > 0$  be such that  $\mathcal{T}$  according with Theorem 4.3.1, for all  $0 \leq \tau < T$ , the reachable set  $\mathcal{R}^\tau$  is strictly convex. Let  $\bar{x}$  be such that  $T(\bar{x}) < T$ . Then  $\bar{x}$  is an optimal point.*

**Proof.** We consider first the case where  $G$  is a vector and the control  $u$  is one-dimensional. Set  $\tau = T(\bar{x})$  and let  $\bar{u}(\cdot)$  be an admissible control steering  $\bar{x}$  to 0 in the optimal time  $\tau$ , together with the associated trajectory  $\bar{x}(\cdot)$ . Set, for all  $t \in [0, \tau]$ ,

$$A(t) = DF(\bar{x}(t)) + DG(\bar{x}(t))\bar{u}(t) \quad , \quad b(t) = G(\bar{x}(t))$$

and let, by Maximum Principle,  $\lambda$  be a solution of

$$\begin{cases} \dot{\lambda}(t) = -\lambda(t)A^T(t), \\ \lambda(\tau) = \zeta, \end{cases} \quad (4.3.12)$$

such that  $\zeta \in N_{\mathcal{R}^\tau}(\bar{x})$  and for a.e.  $t \in [0, \tau]$ ,

$$\bar{u}(t) = \text{sign}\langle \lambda(t), b(t) \rangle.$$

Set, for  $t \in [0, \tau]$ ,

$$g(t) = \langle \lambda(t), b(t) \rangle.$$

We are now going to extend  $\bar{u}(\cdot)$  in an interval  $[\tau, \tau + \delta]$  for a suitable  $\delta > 0$ , with the property that the extended control and its associate trajectory satisfies the Maximum Principle.

Three cases may occur:

- (i)  $g(\tau) := \delta_1 > 0$ ,
- (ii)  $g(\tau) < 0$ ,
- (iii)  $g(\tau) = 0$ .

In the first case, we set  $\bar{u}(t) = 1$  for all  $t > \tau$  and let  $\bar{x}(\cdot)$  the associate trajectory issuing from  $\bar{x}$  at time  $\tau$ . We extend analogously  $A(\cdot)$ ,  $b(\cdot)$ ,  $\lambda(\cdot)$  and  $g(\cdot)$  for  $t > \tau$ . Observe that  $g$  is locally Lipschitz, so that, for  $t > \tau$ ,

$$g(t) = g(\tau) + g(t) - g(\tau) > \delta_1 - L_1(t - \tau)$$

for suitable constant  $L_1$ . Therefore we can find  $\delta > 0$  such that  $0 \leq \tau + \delta < \mathcal{T}$  and  $g(t) > 0$  for all  $t \in [\tau, \tau + \delta]$ , i.e.,

$$\bar{u}(t) = \text{sign } g(t) \quad \forall t \in [\tau, \tau + \delta].$$

The second case is entirely analogous, by substituting 1 with  $-1$ .

We consider now the third case. Let the  $I_0$ ,  $I_1$  be given by Lemma 4.2.4 for the function  $g$  in the interval  $[0, \tau]$ . Observe that necessarily  $\tau \in I_1$ , so that, in particular,  $g'(\tau) \neq 0$ . We set, for  $t > \tau$

$$\bar{u}(t) = 1 \quad \text{if } g'(\tau) > 0$$

or

$$\bar{u}(t) = -1 \quad \text{if } g'(\tau) < 0$$

and let  $\bar{x}(\cdot)$  be the associate trajectory issuing from  $\bar{x}$  at time  $\tau$ . We extend analogous  $A(\cdot)$ ,  $b(\cdot)$ ,  $\lambda(\cdot)$  and  $g(\cdot)$  for  $t > \tau$ . One can compute that

$$g'(t) = \langle \lambda(t), [F, G](\bar{x}(t)) \rangle,$$

where  $[F, G]$  is Lie bracket.

It implies that  $g'$  is continuous. So there exists  $\delta > 0$  such that the sign of  $g'(t)$  equals the sign of  $g'(\tau)$  for all  $t \in [\tau - \delta, \tau + \delta]$ . Therefore our construction of  $\bar{u}(\cdot)$  on  $[0, \tau + \delta]$  is such that for a.e.  $t \in [0, \tau + \delta]$ ,

$$\bar{u}(t) = \text{sign } g(t).$$

Thus, for all  $t \in [0, \tau + \delta]$ ,  $\bar{x}(t) \in \partial\mathcal{R}^t$ . So  $\bar{u}(\cdot)$  steers  $\bar{x}(\tau + \delta)$  to the origin optimally in time  $\tau + \delta$ .

For the case  $G(\cdot) \in \mathbb{M}_{2 \times 2}$ , the proof is entirely done by the above argument.

**Corollary 4.3.1** *Under the same assumptions of Theorem 4.3.2, let  $\tau = T(\bar{x}) < \tau_1 < \mathcal{T}$ . Then there exists  $x_1 \in \partial\mathcal{R}^{\tau_1}$  and a control  $u_1 : [\tau, \tau_1] \rightarrow [-1, 1]^M$  such that the trajectory  $\tilde{x}(\cdot)$  corresponding to the control*

$$\tilde{u}(t) = \begin{cases} \bar{u}(t) & 0 \leq t \leq \tau, \\ u_1(t) & \tau < t \leq \tau_1. \end{cases}$$

*and such that  $\tilde{x}(0) = x_1$  reaches 0 in the optimal time  $\tau_1$  and moreover  $\tilde{x}(\tau_1 - \tau) = \bar{x}$ .*

**Remark 4.3.2** From Corollary 4.3.1,  $\bar{x}(\cdot)$  and  $\lambda(\cdot)$  in the proof of Theorem 4.3.2 can be extended to time  $\mathcal{T}$ . Then the maximized Hamiltonian along  $\bar{x}(\cdot)$  associated with  $\lambda(\cdot)$  is constant in  $[0, \mathcal{T})$ , i.e.,

$$H(\bar{x}(t), \lambda(t)) = C \quad \forall t \in [0, \mathcal{T}).$$

**Proof.** We will now consider here the case  $M = 1$ . From the proof of Theorem 4.3.2, we have that

$$H(\bar{x}(t), \lambda(t)) = \langle \lambda(t), F(\bar{x}(t)) \rangle + |\langle \lambda(t), G(\bar{x}(t)) \rangle|.$$

For a.e.  $t \in [0, \mathcal{T})$ , we compute

$$\begin{aligned} \frac{d}{dt} H(\bar{x}(t), \lambda(t)) \\ = \langle \lambda(t), [G, F](\bar{x}(t)) \rangle \tilde{u}(t) + \langle \lambda(t), [G, F](\bar{x}(t)) \rangle \text{sign} \langle \lambda(t), G(\bar{x}(t)) \rangle. \end{aligned}$$

Since  $\tilde{u}(t) = \text{sign} \langle \lambda(t), G(\bar{x}(t)) \rangle$  for a.e.  $t \in [0, \mathcal{T})$ . We obtain that

$$\frac{d}{dt} H(\bar{x}(t), \lambda(t)) = 0 \text{ for a.e. } t \in [0, \mathcal{T}).$$

Hence, the proof is concluded.

The proof of the case  $M = 2$  is analogous.  $\square$

### 4.3.3 The epi-graph of the minimum time function has positive reach

We are now studying the convexity "type" of the minimum time function  $T(\cdot)$  in which the control dynamics just satisfies a weak controllability condition, i.e., the function  $T(\cdot)$  is just continuous.

**Theorem 4.3.3** Let  $N = 2$  and let the assumptions of Theorem 4.3.1 hold. Let  $\mathcal{T}$  be given by Theorem 4.3.1. Then for every  $0 < \tau < \mathcal{T}$  the epigraph of the minimum time function  $T(\cdot)$  on  $\mathcal{R}^\tau$  has positive reach.

**Corollary 4.3.2** Under the same assumptions of Theorem 4.3.3 the minimum time function satisfies the list of properties in Theorem 2.2.2.

Before beginning the proof of Theorem 4.3.3 we introduce the minimized Hamiltonian and study its sign.

**Definition 4.3.2** Let  $x, \zeta \in \mathbb{R}^N$ . We define the minimized Hamiltonian as

$$h(x, \zeta) = \langle \zeta, F(x) \rangle + \min_{u \in \mathcal{U}} \langle \zeta, G(x)u \rangle.$$

**Proposition 4.3.1** Let the standing assumptions on the dynamics hold. Let  $x$  belong to the boundary of the sublevel set  $\mathcal{R}^\tau$  for some  $\tau > 0$ . Let  $\zeta \in N_{\mathcal{R}^\tau}^F(x)$ . Then  $h(x, \zeta) \leq 0$ .

**Proof.** Let  $\bar{u}(\cdot)$  be an admissible control steering  $x$  to 0 in the optimal time  $\tau$ , together with the associate trajectory  $\bar{x}(\cdot)$ . Then, for all  $0 \leq t \leq \tau$  the point  $\bar{x}(t)$  belongs to  $\mathcal{R}^\tau$ , so that, by definition of Fréchet normal we have

$$\limsup_{t \rightarrow 0^+} \left\langle \zeta, \frac{\bar{x}(t) - x}{\|\bar{x}(t) - x\|} \right\rangle \leq 0.$$

Observing that  $\|x(t) - x\| \leq Kt$  for a suitable constant  $K$ , we have

$$\limsup_{t \rightarrow 0} \left\langle \zeta, \frac{\bar{x}(t) - x}{t} \right\rangle \leq 0.$$

In other words,

$$\begin{aligned} 0 &\geq \limsup_{t \rightarrow 0} \left\langle \zeta, \frac{1}{t} \int_0^t (F(\bar{x}(s)) + G(\bar{x}(s))\bar{u}(s)) ds \right\rangle \\ &= \langle \zeta, F(x) \rangle + \limsup_{t \rightarrow 0} \left\langle \zeta, G(x) \frac{\int_0^t \bar{u}(s) ds}{t} \right\rangle. \end{aligned}$$

Let  $t_n \rightarrow 0$  be sequence such that  $\lim_{n \rightarrow \infty} \frac{1}{t_n} \int_0^{t_n} \bar{u}(s) ds := \tilde{u}$  exists. By the convexity of  $\mathcal{U}$ ,  $\tilde{u} \in \mathcal{U}$ , and so  $h(x, \zeta) = \langle \zeta, F(x) \rangle + \langle \zeta, G(x)\tilde{u} \rangle$ .  $\square$

We are now ready to prove Theorem 4.3.3.

**Proof of Theorem 4.3.3.** Let  $x \neq 0$  such that  $T(x) \leq \mathcal{T}$  and let  $(\bar{u}(\cdot), \bar{x}(\cdot))$  be an optimal pair for  $x$ . By Maximum Principle there exists  $0 \neq \zeta \in N_{\mathcal{R}^{T(x)}}(x)$  such that the adjoint arc  $\lambda$ , with

$$\begin{cases} \dot{\lambda}(t) = \lambda(t)(DF(\bar{x}(t)) + DG(\bar{x}(t))\bar{u}(t))^T, \\ \lambda(T) = \zeta \end{cases}$$

satisfies

$$\langle \lambda(t), F(\bar{x}(t)) + G(\bar{x}(t))\bar{u}(t) \rangle = h(\bar{x}(t), \lambda) \text{ a.e.}$$

We claim that

$$(\zeta, h(x, \zeta)) \in N_{\text{epi}(T)}^P(x, T(x)), \quad (4.3.13)$$

i.e., there exists a constant  $\sigma > 0$  such that, for all  $y \in \mathbb{R}^N$  such that  $0 < T(y) < \mathcal{T}$ , for all  $\beta \geq T(y)$ , we have

$$\langle (\zeta, \theta), (y, \beta) - (x, T(x)) \rangle \leq \sigma \|(\zeta, \theta)\| \left( \|y - x\|^2 + |\beta - T(x)| \right), \quad (4.3.14)$$

where  $\theta = h(x, \zeta)$ , and, moreover

$$\sigma \text{ is independent of } x \text{ and } \zeta. \quad (4.3.15)$$

Indeed, we consider two cases:

- (a)  $T(y) \leq T(x)$ ;



(b)  $T(y) > T(x)$ .

In the first case,  $y \in \mathcal{R}^{T(x)}$ , so that by Theorem 4.3.1

$$\langle \zeta, y - x \rangle \leq 0.$$

If  $\beta \geq T(x)$  then (4.3.14) is automatically satisfied, since  $\theta \leq 0$  by Proposition 4.3.1. If instead  $\beta < T(x)$ , we set  $x_1 = \bar{x}(T(x) - \beta)$ .

We estimate first  $\langle \zeta, y - x_1 \rangle$ . We have since  $y \in \mathcal{R}^\beta$ , recalling Remark 4.3.1, for suitable constants  $K_1, K_2$  given by Gronwall's Lemma,

$$\begin{aligned} \langle \zeta, y - x_1 \rangle &= \langle \lambda(\beta), y - x_1 \rangle + \langle \lambda(T(x)) - \lambda(\beta), y - x_1 \rangle \\ &\leq \langle \lambda(T(x)) - \lambda(\beta), y - x_1 \rangle \leq K_1 \|\lambda(T)\| |T(x) - \beta| \|y - x_1\| \\ &\leq K_1 \|\lambda(T)\| |T(x) - \beta| (\|y - x\| + \|x_1 - x\|) \\ &\leq K_1 \|\zeta\| |T(x) - \beta| (\|y - x\| + K_2 |T(x) - \beta|) \\ &\leq K_3 \|\zeta\| (\|y - x\|^2 + |T(x) - \beta|^2) \end{aligned}$$

for a suitable constant  $K_3$ .

Second, we estimate  $\langle \zeta, x_1 - x \rangle$ . We have

$$\begin{aligned} \langle \zeta, x_1 - x \rangle &= \int_0^{T(x)-\beta} \langle \lambda(T), F(\bar{x}(s)) + G(\bar{x}(s))\bar{u}(s) \rangle ds \\ &= \int_0^{T(x)-\beta} \langle \lambda(s), F(\bar{x}(s)) + G(\bar{x}(s))\bar{u}(s) \rangle ds \\ &+ \int_0^{T(x)-\beta} \langle \lambda(T) - \lambda(s), F(\bar{x}(s)) + G(\bar{x}(s))\bar{u}(s) \rangle ds \\ &\leq (T(x) - \beta) h(x, \zeta) + K_4 \|\zeta\| |T(x) - \beta|^2, \end{aligned}$$

for a suitable constant  $K_4$ , recalling the Maximum Principle and Gronwall's lemma. Therefore,

$$\langle (\zeta, \theta), (y, \beta) - (x, T(x)) \rangle \leq (K_3 + K_4) \|\zeta\| (\|y - x\|^2 + |T(x) - \beta|^2),$$

and the proof for the case (a) is concluded by observing that  $K_3$  and  $K_4$  are independent of  $\zeta$  and  $x$ .

In the second case we need to use the optimality of  $x$ . We observe first that, since  $\theta \leq 0$ , we only need to prove (4.3.14) for  $\beta = T(y)$ . Recalling Corollary 4.3.1, we can extend the control  $\bar{u}$  up to the time  $T(y)$  so that the associated trajectory (still denoted by  $\bar{x}(\cdot)$ ) remains optimal. Let also  $\lambda$  be the extended adjoint vector and denote by  $\tilde{x}(\cdot)$  the trajectory of the reversed dynamics associated with the extended control  $\bar{u}$ , i.e.,

$$\begin{cases} \dot{\tilde{x}}(t) = -F(\tilde{x}(t)) - G(\tilde{x}(t))\bar{u}(t), \\ \tilde{x}(0) = 0 \end{cases}$$

where  $\tilde{u}(t) = \bar{u}(T(y) - t)$ .

Set  $x_1 = \tilde{x}(T(y))$ . We estimate first  $\langle \zeta, y - x_1 \rangle$ . We have, by arguing similarly as before,

$$\begin{aligned} \langle \zeta, y - x_1 \rangle &= \langle \lambda(T(y)), y - x_1 \rangle + \langle \lambda(T(x)) - \lambda(T(y)), y - x_1 \rangle \\ &\quad \text{(the first summand is less than 0} \\ &\quad \text{by the construction in Theorem 4.3.2)} \\ &\leq \langle \lambda(T(x)) - \lambda(T(y)), y - x_1 \rangle \\ &\leq K_5 \|\zeta\| (|T(y) - T(x)|^2 + \|y - x_1\|^2). \end{aligned}$$

On the other hand,

$$\begin{aligned} \langle \zeta, x_1 - x \rangle &= \int_{T(x)}^{T(y)} \langle \zeta, -F(\tilde{x}(s)) - G(\tilde{x}(s))\tilde{u}(s) \rangle ds \\ &= \int_{T(x)}^{T(y)} \langle \lambda(s), -F(\tilde{x}(s)) - G(\tilde{x}(s))\tilde{u}(s) \rangle ds \\ &\quad + \int_{T(x)}^{T(y)} \langle \lambda(T(x)) - \lambda(s), -F(\tilde{x}(s)) - G(\tilde{x}(s))\tilde{u}(s) \rangle ds \\ &\leq \int_{T(x)}^{T(y)} \max_{u \in \mathcal{U}} \langle \lambda(s), -F(\tilde{x}(s)) - G(\tilde{x}(s))u \rangle ds + K_6 \|\zeta\| (T(y) - T(x))^2 \\ &\quad \text{(for a suitable constant } K_6 \text{ given by Gronwall's Lemma).} \end{aligned}$$

Recalling Remark 4.3.2, the maximized Hamiltonian in the integral of the first summand is constant. Therefore we obtain

$$\langle \zeta, x_1 - x \rangle \leq -h(x, \zeta)(T(y) - T(x)) + K_6 \|\zeta\| |T(y) - T(x)|^2.$$

Combining the above estimates we obtain finally

$$\langle (\zeta, \theta), (y, T(y)) - (x, T(x)) \rangle \leq (K_5 + K_6) \|\zeta\| (\|y - x\|^2 + |T(y) - T(x)|^2),$$

and the proof of the claim is concluded, by observing again, that  $K_5, K_6$  are independent of  $x$  and  $\zeta$ .

In order to conclude the proof we observe that  $N_{\text{epi}(T)}^P$  is pointed at every point  $(x, T(x)), x \in \mathcal{R}^\tau$ , since the projection of every  $(\zeta, \theta) \in N_{\text{epi}(T)}^P(x, T(x))$  onto  $\mathbb{R}^N$  is normal to the strictly convex set  $\mathcal{R}^\tau$ . Therefore, we can apply Corollary 3.1 in [47], with  $\Omega_P = \text{int}\mathcal{R}^\tau$ , which shows that  $\text{epi}(T)$  has positive reach.  $\square$

## Part II

# Regularity of a class of continuous functions



## Chapter 5

# External sphere condition and continuous functions

Our aim in this chapter is proving that if the hypograph of a continuous function  $f : \Omega \subseteq \mathbb{R}^N \rightarrow \mathbb{R}$  satisfies an external sphere condition then it has "essentially" positive reach, i.e., the hypograph of the restriction of  $f$  outside a closed set of zero measure has (locally) positive reach. Hence such a function enjoys some properties of a concave function, in particular a.e. twice differentiability.

The result is based on studying the set of *bad points* where the proximal normal cones to such points are not wedged, i.e., the set of horizontal proximal supergradients at "bad point"  $x$ ,  $\partial^\infty f(x)$ , contains a nontrivial subspace. Since the hypograph of  $f$  satisfies an external sphere condition, we can construct a special subspace of  $\partial^\infty f(x)$  which is convex combination of vectors  $v$  such that  $(-v, 0)$  is a proximal normal vector to  $\text{hypo}(f)$  at  $(x, f(x))$  realized by a ball of uniform radius  $\theta > 0$ . Thus, the set of *bad points* is closed in  $\Omega$ . Finally, we prove that the density Lebesgue measure of a bad point is zero by using our special subspace and the inductive method. Therefore, the set of "bad points" has zero Lebesgue measure.

The chapter is organized as follows: Section 5.1 is devoted to definitions and basic facts, while Section 5.2 contains statements of main results. The same section contains also an outline of the proof of Theorem 5.2.1, which is a localized version of the main result and where all the basic arguments appear. Detailed arguments begin in Section 5.3, which contains several lemmas concerning the set of *bad points* (i.e., the normal cone to the hypograph of the function at those points contains at least one line). Section 5.4 is finally devoted to proof of Theorem 5.2.1. On the basis of Theorem 5.2.1, our main theorem will be proved in the same section together with its corollaries.

## 5.1 Notation

We first rewrite quickly some basic notations which concern in the chapter.

Let  $\Omega \subseteq \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be continuous. The hypograph of  $f$  is denoted by

$$\text{hypo}(f) = \{(x, \beta) \mid x \in \Omega, \beta \leq f(x)\}. \quad (5.1.1)$$

The vector  $(-v, \lambda) \in \mathbb{R}^N \times \mathbb{R}$  is a *proximal normal vector* to  $\text{hypo}(f)$  (we will denote this fact that  $(-v, \lambda) \in N_{\text{hypo}(f)}^P(x, f(x))$  at  $(x, f(x))$  iff there exists a constant  $\sigma > 0$  such that for all  $y \in \Omega$  and for all  $\beta \leq f(y)$ , it holds

$$\langle (-v, \lambda), (y, \beta) - (x, f(x)) \rangle \leq \sigma (\|y - x\|^2 + |\beta - f(x)|^2). \quad (5.1.2)$$

Equivalently,  $(-v, \lambda) \in N_{\text{hypo}(f)}^P(x, f(x))$  iff there exists a constant  $\gamma > 0$  such that

$$B_{N+1}((x, f(x)) + \gamma(-v, \lambda), \gamma \|(-v, \lambda)\|) \cap \text{hypo}(f) = \emptyset \quad (5.1.3)$$

where

$$B_k(a, r) = \{z \in \mathbb{R}^k \mid \|z - a\| < r\}$$

is the open ball with center  $a$  and radius  $r$  in  $\mathbb{R}^k$ .

Moreover, the vector  $(-v, \lambda) \in N_{\text{hypo}(f)}^P(x, f(x))$  is *realized by a ball of radius*  $\rho > 0$  if  $(-v, \lambda) \neq 0$  and (5.1.2) is satisfied for  $\sigma = \frac{\|(-v, \lambda)\|}{2\rho}$ .

**Remark 5.1.1** *If  $(-v, \lambda) \in N_{\text{hypo}(f)}^P(x, f(x))$  then  $\lambda \geq 0$ .*

Associated with  $\text{hypo}(f)$ , we define that

1.  $\partial^P f(x) = \{v \mid (-v, 1) \in N_{\text{hypo}(f)}^P(x, f(x))\}$  is set of *proximal supergradients* of  $f$  at  $x$ .
2.  $\partial^\infty f(x) = \{v \mid (-v, 0) \in N_{\text{hypo}(f)}^P(x, f(x))\}$   $v$  is the set a *proximal horizon supergradients* of  $f$  at  $x$ .

We are now giving some new notations. These notations are concerned with the set of *bad points* where the proximal normal cone of  $\text{hypo}(f)$  contains at least one line (i.e., it is not wedged). First we introduce two special types of normal vectors, namely

1. Normal vectors which are limit of unique normals at nearby points

$$N^L(x) = \left\{ \xi \in \mathbb{R}^{N+1} \mid \exists \{x_n\} \rightarrow x \text{ such that} \right. \\ \left. \begin{array}{l} i) f \text{ is Fréchet differentiable at } x_n \text{ and} \\ ii) \xi = \lim_{n \rightarrow \infty} \frac{(-Df(x_n), 1)}{\|(-Df(x_n), 1)\|} \end{array} \right\}.$$

2. Among them we select the horizontal ones

$$N_0^L(x) = N^L(x) \cap (-\partial^\infty f(x), 0).$$

We also denote the subspace which is generated by  $N_0^L(x)$  as

$$H_0(x) = \text{span}\{N_0^L(x)\} = \left\{ \sum_{i=1}^k \alpha_i \xi_i \mid \xi_i \in N_0^L(x) \text{ and } \alpha_i \in \mathbb{R} \right\},$$

and the positive cone which is generated by  $N_0^L(x)$  as

$$H_0^+(x) = \text{span}^+\{N_0^L(x)\} = \left\{ \sum_{i=1}^k \alpha_i \xi_i \mid \xi_i \in N_0^L(x) \text{ and } \alpha_i \geq 0 \right\}.$$

3. The largest vector subspace contained in  $N_{\text{hypo}(f)}^P(x, f(x))$  will be denoted by

$$NL(x) = \left\{ \xi \in N_{\text{hypo}(f)}^P(x, f(x)) \mid -\xi \in N_{\text{hypo}(f)}^P(x, f(x)) \right\}.$$

From Remark 5.1.1, one can see that  $NL(x) \subseteq (-\partial^\infty f(x), 0)$ .

4. We denote the set of *bad points* of  $f$  by

$$BP_f = \{x \in \Omega \mid NL(x) \neq 0\} \quad (5.1.4)$$

At each point  $x \in BP_f$ , we write  $BP_f$  as the union of the two sets

$$BP_f^+(x) = \{y \in BP_f \mid f(y) \geq f(x)\}$$

$$BP_f^-(x) = \{y \in BP_f \mid f(y) \leq f(x)\}.$$

## 5.2 Main results

### 5.2.1 Statement of main results

Our results are the following theorem, together with several corollaries. We recall that the notation  $BP_f$  was defined in (5.1.4).

**Theorem 5.2.1** *Let  $\Omega \subset \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be continuous. Assume that  $\text{hypo}(f)$  satisfies the  $\theta$  – external sphere condition, where  $\theta : \Omega \rightarrow (0, \infty)$  is continuous. Then*

i)  $\Omega_P := \Omega \setminus BP_f$  is open.

ii)  $\mathcal{L}^N(\Omega \setminus \Omega_P) = 0$ .

**Corollary 5.2.1** *Let  $\Omega \subset \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be continuous. Assume that  $\text{hypo}(f)$  satisfies the  $\theta$  – external sphere condition where  $\theta : \Omega \rightarrow [0, \infty)$  is continuous. Then the hypograph of  $f|_{\Omega_P}$  has positive reach.*

**Corollary 5.2.2** *Let  $f : \Omega \rightarrow \mathbb{R}$  be as in the statement of Theorem 5.2.1 then  $f$  satisfies properties (1)-(3) of Theorem 2.2.2.*

In view of Proposition 3.2.1 in Chapter 3, we can apply the previous results to the minimum time function.

**Corollary 5.2.3** *Let  $(f, \mathcal{U})$  be the control system 3.1.1 and  $\mathcal{S}$  be a target in Chapter 3. Under the conditions (H1), (H2), (H3), (H4), there exists an open set  $\mathcal{S}_P^c \subset \mathcal{S}^c$  such that  $\mathcal{L}^N(\mathcal{S}^c \setminus \mathcal{S}_P^c) = 0$  and the restricted continuous function  $T|_{\mathcal{S}_P^c} : \mathcal{S}_P^c \rightarrow [0, +\infty)$  has the hypograph with positive reach.*

**Corollary 5.2.4** *Under the conditions (H1), (H2), (H3), (H4), the minimum time function is twice differentiable a.e. in  $\mathcal{S}^c$ .*

In order to make our proof more clear, we prefer to state our main theorem in a particular case (local case). The arguments are used in the proof of the main part of the proof of Theorem 5.2.1.

**Theorem 5.2.2** *Let  $f : B_N(0, 1) \rightarrow \mathbb{R}$  be continuous and let  $\rho > 0$ . Assume that  $\text{hypo}(f)$  satisfies the  $\rho$ -external sphere condition. Then*

- i)  $BP_f \cup \partial B_N(0, 1)$  is closed.
- ii)  $\mathcal{L}^N(BP_f) = 0$ .

## 5.2.2 Outline of proof of Theorem 5.2.2

The part (i) is precisely Lemma 5.3.4.

To prove the part (ii) we will use induction.

For the case  $N = 1$ . By using Lemma 5.4.1 and Corollary 5.3.5 we obtain that the  $\mathcal{L}^1$ -density of  $BP_f$  at  $x$ ,  $D_{BP_f}^1(x) = \lim_{\sigma \rightarrow \infty} \frac{\mathcal{L}^1(BP_f \cap B_1(x, \sigma))}{\mathcal{L}^1(B_1(x, \sigma))} = 0$  for all  $x \in BP_f$ . Therefore, the proof is completed by the Lebesgue theorem.

In order to get the conclusion for  $N = k + 1$  from the inductive assumption for  $N = k \geq 1$ . We divide the set  $BP_f$  into two parts:

The first part is  $BP_f^{\zeta^+} \cup BP_f^{\zeta^-}$  (see the definition of  $BP_f^{\zeta}$  near Lemma 5.3.7) where  $\zeta^+ = (0, 1)$  and  $\zeta^- = (0, -1)$ . Using Lemma 5.3.7, we get  $\mathcal{L}^N(BP_f^{\zeta^+} \cup BP_f^{\zeta^-}) = 0$ .

To prove  $\mathcal{L}^N[BP_f \setminus (BP_f^{\zeta^+} \cup BP_f^{\zeta^-})] = 0$ , we notice that Lemma 5.3.6 can be used at every point in the open set  $B_N(0, 1) \setminus (BP_f^{\zeta^+} \cup BP_f^{\zeta^-})$ . We need to prove that for all  $B_N(x, r_x) \subset B_N(0, 1) \setminus (BP_f^{\zeta^+} \cup BP_f^{\zeta^-})$ , it holds  $\mathcal{L}^N(BP_f \cap B_N(x, r_x)) = 0$ . Three small steps are considered

Step 1: Let  $\bar{f} = f|_{B_N(x, r_x)}$ . By Lemma 5.3.6, the  $\text{hypo}(\bar{f}_{x_2})$  (See the definition of  $\bar{f}_{x_2}$  near Lemma 5.3.6) satisfies the  $\theta$ -external sphere condition.

Step 2: From Lemma 7.2.3 and the inductive assumption, we obtain that  $\mathcal{L}^{N-1}(BP_{\bar{f}_{x_2}}) = 0$ .

Step 3: We use Fubini's theorem to complete the proof.



### 5.3 Preparatory Lemmas

This section is devoted to several partial results which are needed to prove our main theorem. To simplify our statements, we agree that the continuous function  $f$  in this section is defined on  $B_N(0, 1)$  and  $\text{hypo}(f)$  satisfies the  $\rho$ -external sphere condition for a given constant  $\rho > 0$

#### 5.3.1 Closedness of the set of bad points

The first lemma shows that the proximal normal unit vector to the hypograph of  $f$  at  $(x, f(x))$  where  $f$  is differentiable is unique and is realized by a ball of radius  $\rho$ .

**Lemma 5.3.1** *Let  $x$  be in  $B_N(0, 1)$  such that  $f(\cdot)$  is differentiable at  $x$ . Then  $\frac{(-Df(x), 1)}{\|(-Df(x), 1)\|}$  is the unique proximal normal unit vector to  $\text{hypo}(f)$  at  $(x, f(x))$ . Moreover,  $\frac{(-Df(x), 1)}{\|(-Df(x), 1)\|}$  is realized by a ball of radius  $\rho$ , i.e., for all  $y \in B_N(0, 1)$  and for all  $\beta \leq f(y)$ , it holds:*

$$\left\langle \frac{(-Df(x), 1)}{\|(-Df(x), 1)\|}, (y, \beta) - (x, f(x)) \right\rangle \leq \frac{1}{2\rho} (\|y - x\|^2 + |\beta - f(x)|^2).$$

**Proof.** Since  $f(\cdot)$  is differentiable at  $x$ ,  $\frac{(-Df(x), 1)}{\|(-Df(x), 1)\|}$  is unique Fréchet normal unit vector to the hypograph of  $f(\cdot)$  at  $(x, f(x))$ . Therefore, since  $\text{hypo}(f)$  satisfies the  $\rho$ -external sphere condition,  $\frac{(-Df(x), 1)}{\|(-Df(x), 1)\|}$  is the unique proximal normal unit vector to  $\text{hypo}(f)$  at  $(x, f(x))$ . Thus,  $\frac{(-Df(x), 1)}{\|(-Df(x), 1)\|} \in N_{\text{hypo}(f)}^P(x, f(x))$  is realized by a ball of radius  $\rho$ .  $\square$

From this lemma and the continuity of  $f$ , three corollaries follow.

**Corollary 5.3.1** *Let  $x \in B_N(0, 1)$ . Then*

$$N^L(x) \subseteq N_{\text{hypo}(f)}^P(x, f(x)).$$

*More precisely, for each  $0 \neq \xi \in N^L(x)$  we have that  $\xi$  is a unit proximal normal vector to  $\text{hypo}(f)$  at  $(x, f(x))$  realized by a ball of radius  $\rho$ .*

**Proof.** Let  $\xi \in N^L(x)$ , and take a sequence  $\{x_n\}$  converging to  $x$  such that  $f$  is differentiable at  $x_n$  and  $\left\{ \frac{(-Df(x_n), 1)}{\|(-Df(x_n), 1)\|} \right\}$  converges to  $\xi$ . By Lemma 5.3.1,  $\frac{(-Df(x_n), 1)}{\|(-Df(x_n), 1)\|} \in N_{\text{hypo}(f)}^P(x_n, f(x_n))$  is realized by a ball of radius  $\rho$ , i.e., for all  $y \in B_N(0, 1)$  and for all  $\beta \leq f(y)$ , we have

$$\left\langle \frac{(-Df(x_n), 1)}{\|(-Df(x_n), 1)\|}, (y, \beta) - (x_n, f(x_n)) \right\rangle \leq \frac{1}{2\rho} (\|y - x\|^2 + |\beta - f(x_n)|^2). \quad (5.3.1)$$

By letting  $n$  approach to  $\infty$  in (5.3.1), the inequality

$$\langle \xi, (y, \beta) - (x, f(x)) \rangle \leq \frac{1}{2\rho} (\|y - x\|^2 + |\beta - f(x)|^2)$$

holds for all  $y \in B_N(0, 1)$  and for all  $\beta \leq f(y)$ .

The proof is completed.  $\square$

**Corollary 5.3.2**  $N_0^L(x)$  is closed for all  $x \in B_N(0, 1)$ . Moreover, if  $\xi \in N_0^L(x)$  then  $\xi$  is a proximal normal unit vector to  $\text{hypo}(f)$  at  $(x, f(x))$  realized by a ball of radius  $\rho$ .

**Proof.** Let  $\{\xi_n\} \subseteq N_0^L(x)$  converge to  $\bar{\xi}$ . We need to prove that  $\bar{\xi} \in N_0^L(x)$ . Indeed, for each  $n$ , there exists a sequence  $\{x_n^k\}$  converging to  $x$  such that  $f$  is differentiable at  $x_n^k$  and  $\left\{ \frac{(-Df(x_n^k), 1)}{\|(-Df(x_n^k), 1)\|} \right\}$  converges to a unit vector  $\xi_n \in (-\partial^\infty f(x), 0)$ . For each  $n$  we can take a point  $y_n \in \{x_n^k\}$  such that  $\|y_n - x\| \leq \frac{1}{n}$  and  $\left\| \frac{(-Df(y_n), 1)}{\|(-Df(y_n), 1)\|} - \bar{\xi} \right\| \leq \frac{1}{n}$ . Therefore  $\{y_n\}$  and  $\left\{ \frac{(-Df(y_n), 1)}{\|(-Df(y_n), 1)\|} \right\}$  converge respectively to  $x$  and  $\bar{\xi}$ . This implies that  $\bar{\xi} \in N^L(x)$ . On the other hand, since  $\{\xi_n\} \subseteq N_0^L(x)$  converges to  $\bar{\xi}$  we have  $\bar{\xi} \in (-\partial^\infty f(x), 0)$ . The proof is completed.  $\square$

With a similar proof, we get the third corollary.

**Corollary 5.3.3** Let  $\{x_n\} \in B_N(0, 1)$  converge to  $x \in B_N(0, 1)$  and let  $\xi_n \in N_0^L(x_n)$  converge to  $\bar{\xi}$ , then  $\bar{\xi} \in N_0^L(x)$ .

The next lemma says that if there exists a vector  $0 \neq p_0 \in (-\partial^\infty f(x))$  then we can find a vector in  $N_0^L(x)$ . This vector is found by considering a sequence which converges to  $x$  along the ray  $\{x + tp_0 \mid t > 0\}$  such that  $f$  is differentiable at each point of this sequence. This idea is inspired by the proof of Lemma 4.7 in [19].

**Lemma 5.3.2** Let  $x \in B_N(0, 1)$  such that  $\partial^\infty f(x) \neq \emptyset$ . Then  $N_0^L(x)$  is nonempty.

**Proof.** Let  $0 \neq -p_0 \in \partial^\infty f(x)$ . By the definition of  $\partial^\infty f(x)$ ,  $(p_0, 0) \in N_{\text{hypo}(f)}^P(x, f(x))$ , i.e. there exists a constant  $\sigma_0 > 0$  such that

$$\langle (p_0, 0), (y, \beta) - (x, f(x)) \rangle \leq \sigma_0 (\|y - x\|^2 + |\beta - f(x)|^2) \quad (5.3.2)$$

for all  $y \in B_N(0, 1)$  and for all  $\beta \leq f(y)$ .

Set  $x_n = x + \frac{p_0}{n}$ . By the density theorem (see Theorem 1.3.1 in [25]), for each  $n$  there exists  $z_n$  such that

$$\partial_P f(z_n) \neq \emptyset \quad (5.3.3)$$

$$\|z_n - x_n\| \leq \frac{1}{n^2} \quad (5.3.4)$$

(5.3.3) implies that there exists a vector  $(\zeta_n, -1)$  which is a proximal normal vector to the epigraph of  $f(\cdot)$  at  $(z_n, f(z_n))$ . Therefore, since  $\text{hypo}(f)$  satisfies the  $\rho$ -external sphere condition we obtain that  $f(\cdot)$  is differentiable at  $z_n$ . Recalling Lemma 5.3.1, for all  $z \in B_N(0, 1)$  and for all  $\beta \leq f(z)$ , we have

$$\begin{aligned} & \langle (-Df(z_n), 1), (z, \beta) - (z_n, f(z_n)) \rangle \\ & \leq \frac{1}{2\rho} \|(-Df(z_n), 1)\| (\|z - z_n\|^2 + |\beta - f(z_n)|^2). \end{aligned} \quad (5.3.5)$$

Recalling (5.3.4),  $z_n \in B_N(0, 1)$  for  $n$  large enough. Thus by taking  $y = z_n$  in (5.3.2), we obtain

$$\langle p_0, z_n - x \rangle \leq \sigma_0 (\|z_n - x\|^2 + |\beta - f(x)|^2) \quad (5.3.6)$$

for all  $\beta \leq f(z_n)$ .

We have

$$\begin{aligned} \langle p_0, z_n - x \rangle &= \langle p_0, \frac{p_0}{n} \rangle + \langle p_0, z_n - x_n \rangle \\ &= \frac{\|p_0\|^2}{n} + \langle p_0, z_n - x_n \rangle. \end{aligned}$$

Combining the above inequality with (5.3.4), we get

$$\langle p_0, z_n - x \rangle \geq \frac{\|p_0\|^2}{n} - \frac{\|p_0\|}{n^2}. \quad (5.3.7)$$

Moreover, from (5.3.4) we get

$$\|z_n - x\| = o\left(\frac{1}{n}\right). \quad (5.3.8)$$

Recalling (5.3.6), (5.3.7) and (5.3.8), for  $n$  large enough, the estimate

$$\frac{\|p_0\|^2}{n} \leq o\left(\frac{1}{n^2}\right) + |\beta - f(x)|^2 \quad (5.3.9)$$

holds for all  $\beta \leq f(z_n)$ .

Therefore, there exists a constant  $C > 0$  such that

$$f(x) - f(z_n) \geq \frac{C}{\sqrt{n}}. \quad (5.3.10)$$

for  $n$  large enough.

We are now going to prove that :  $\limsup_{n \rightarrow \infty} \|(-Df(z_n), 1)\| = +\infty$ . Assume by contradiction that there exists a constant  $K > 0$  such that

$$\|(-Df(z_n), 1)\| \leq K \quad \text{for all } n. \quad (5.3.11)$$

By taking  $z = x$  and  $\beta = f(x)$  in (5.3.5) and by recalling (5.3.11) we have

$$(f(x) - f(z_n))\left(1 - \frac{K}{2\rho}(f(x) - f(z_n))\right) \leq K\left(1 + \frac{\|x - z_n\|}{2\rho}\right)\|x - z_n\|. \quad (5.3.12)$$

for  $n$  large enough. Therefore, by (5.3.10) and (5.3.8), we get from the above inequality that there exists a constant  $C_1 > 0$  such that

$$\frac{1}{\sqrt{n}} \leq C_1 \frac{1}{n}.$$

for  $n$  large enough.

This is a contradiction.

We now assume without loss of generality that  $\lim_{n \rightarrow \infty} \frac{(-Df(z_n), 1)}{\|(-Df(z_n), 1)\|} = (-\bar{\zeta}_0, 0)$ . Since  $\{z_n\}$  converges to  $x$ , we have  $(-\bar{\zeta}_0, 0) \in N_0^L(x)$ . The proof is completed.  $\square$

**Corollary 5.3.4** *If  $x \in BP_f$  then  $N_0^L(x)$  is nonempty.*

The following lemma is a crucial observation. At every *bad point*, we can extract a line from  $H_0^+(x) \subseteq NL(x) \subseteq N_{\text{hypo}(f)}^P(x, f(x))$ . It is also pivotal to prove Lemma 5.3.4 and Theorem 5.4.1. The difference between the proof of this lemma and the proof of the previous lemma is the way of choosing a sequence which allows us to get a vector in  $N_0^L(x)$ .

**Lemma 5.3.3** *If  $x \in BP_f$  then  $H_0^+(x)$  contains at least one line.*

**Proof.** We recall that by Corollary 5.3.4,  $N_0^L(x)$  is nonempty. Assume by contradiction that  $H_0^+(x)$  does not contain lines. From Corollary 5.3.2,  $N_0^L(x)$  is compact and does not contain 0. Thus by applying Lemma 7.2.1 for  $C = N_0^L(x)$ , there exists a constant  $\delta_0 > 0$  such that for all  $0 \neq \xi_1, \xi_2 \in H_0^+(x)$ , one has

$$\left\langle \frac{\xi_1}{\|\xi_1\|}, \frac{\xi_2}{\|\xi_2\|} \right\rangle > -1 + \delta_0.$$

Therefore, there exist a vector  $(v_0, 0) \in H_0(x)$  and a constant  $\delta_1 > 0$  such that  $v_0 \in \mathbb{R}^N$ ,  $\|v_0\| = 1$  and

$$\left\langle -(v_0, 0), \frac{\xi}{\|\xi\|} \right\rangle \geq \delta_1 \quad \text{for all } 0 \neq \xi \in H_0^+(x). \quad (5.3.13)$$

Since  $x \in BP_f$  (namely,  $NL(x)$  contains at least one line) there exists a unit vector  $p_0 \in \mathbb{R}^N$  such that  $(p_0, 0) \in NL(x)$  and  $\langle p_0, v_0 \rangle \geq 0$ .

Setting  $v_1 = v_0 + \frac{\delta_1}{2}p_0$ , one can easily get from (5.3.13) that:

$$\left\langle -(v_1, 0), \frac{\xi}{\|\xi\|} \right\rangle \geq \frac{\delta_1}{2} \quad \text{for all } 0 \neq \xi \in H_0^+(x). \quad (5.3.14)$$

Setting  $x_n = x + \frac{v_1}{n}$ . By the density theorem (see Theorem 1.3.1 in [25]), for each  $n$  there exists  $z_n$  such that

$$\partial_P f(z_n) \neq \emptyset \quad (5.3.15)$$

$$\|z_n - x_n\| \leq \frac{1}{n^2} \quad (5.3.16)$$

(5.3.15) implies that there exists a vector  $(\zeta_n, -1)$  which is a proximal normal vector to the epigraph of  $f(\cdot)$  at  $(z_n, f(z_n))$ . Therefore, since  $\text{hypo}(f)$  satisfies the  $\rho$ -external sphere condition we obtain that  $f(\cdot)$  is differentiable at  $z_n$  (see Proposition 3.15, p.51, [21]). Recalling Lemma 5.3.1, for all  $z \in B_N(0, 1)$  and for all  $\beta \leq f(z)$ , we have

$$\begin{aligned} & \langle (-Df(z_n), 1), (z, \beta) - (z_n, f(z_n)) \rangle \\ & \leq \frac{1}{2\rho} \|(-Df(z_n), 1)\| (\|z - z_n\|^2 + |\beta - f(z_n)|^2). \end{aligned} \quad (5.3.17)$$

On the other hand, since  $(p_0, 0) \in NL(x)$ , there exists a constant  $\sigma_0 > 0$  such that

$$\langle (p_0, 0), (y, \beta) - (x, f(x)) \rangle \leq \sigma_0 (\|y - x\|^2 + |\beta - f(x)|^2) \quad (5.3.18)$$

for all  $y \in B_N(0, 1)$  and for all  $\beta \leq f(y)$ .

Recalling (5.3.16),  $z_n \in B_N(0, 1)$  for  $n$  large enough. Thus by taking  $y = z_n$  in (5.3.18), we have

$$\langle p_0, z_n - x \rangle \leq \sigma_0 (\|z_n - x\|^2 + |\beta - f(x)|^2) \quad (5.3.19)$$

for all  $\beta \leq z_n$ .

We have

$$\begin{aligned} \langle p_0, z_n - x \rangle &= \langle p_0, \frac{v_0}{n} \rangle + \langle p_0, \frac{\delta_1}{2n} p_0 \rangle + \langle p_0, z_n - x_n \rangle \\ &\geq \frac{\delta_1}{2n} + \langle p_0, z_n - x_n \rangle. \end{aligned}$$

Combining the above inequality with (5.3.16), we get

$$\langle p_0, z_n - x \rangle \geq \frac{\delta_1}{2n} - \frac{1}{n^2}. \quad (5.3.20)$$

Moreover, from (5.3.16) we get

$$\|z_n - x\| = o\left(\frac{1}{n}\right). \quad (5.3.21)$$

Recalling (5.3.19), (5.3.20) and (5.3.21), for  $n$  large enough, the estimate holds

$$\frac{\delta_1}{2n} \leq o\left(\frac{1}{n^2}\right) + |\beta - f(x)|^2 \quad (5.3.22)$$

for all  $\beta \leq f(z_n)$ .

Therefore, there exists a constant  $C > 0$  such that

$$f(x) - f(z_n) \geq \frac{C}{\sqrt{n}}. \quad (5.3.23)$$

for  $n$  large enough.

We are now going to prove that :  $\limsup_{n \rightarrow \infty} \|(-Df(z_n), 1)\| = +\infty$ .

Assume by contradiction that there exists a constant  $K > 0$  such that

$$\|(-Df(z_n), 1)\| \leq K \quad \text{for all } n. \quad (5.3.24)$$

By taking  $z = x$  and  $\beta = f(x)$  in (5.3.17) and by recalling (5.3.24) we have

$$(f(x) - f(z_n))\left(1 - \frac{K}{2\rho}(f(x) - f(z_n))\right) \leq K\left(1 + \frac{\|x - z_n\|}{2\rho}\right)\|x - z_n\|. \quad (5.3.25)$$

for  $n$  large enough. Therefore, by (5.3.23) and (5.3.21), we get from the above inequality that there exists a constant  $C_1 > 0$  such that

$$\frac{1}{\sqrt{n}} \leq C_1 \frac{1}{n}.$$

for  $n$  large enough.

This is a contradiction.

We now assume without loss of generality that  $\lim_{n \rightarrow \infty} \frac{(-Df(z_n), 1)}{\|(-Df(z_n), 1)\|} = (-\bar{\zeta}_0, 0)$ . Moreover, since  $\{z_n\}$  converges to  $x$ , we have  $(-\bar{\zeta}_0, 0) \in N_0^L(x)$ . On the other hand, by (5.3.23), we can take  $z = x$  and  $\beta = f(z_n)$  in (5.3.17) to get

$$\left\langle \frac{(-Df(z_n), 1)}{\|(-Df(z_n), 1)\|}, \frac{(x - z_n, 0)}{\|x - z_n\|} \right\rangle \leq \frac{\|x - z_n\|}{2\rho}. \quad (5.3.26)$$

Let  $n$  tend to  $+\infty$ . Recalling (5.3.21), (5.3.16) we obtain

$$\langle (-\bar{\zeta}_0, 0), (-v_1, 0) \rangle \leq 0. \quad (5.3.27)$$

Since  $(-\bar{\zeta}_0, 0) \in N_0^L(x)$ , we get a contradiction from (5.3.27) and (5.3.14).  
□

**Lemma 5.3.4**  $BP_f \cup \partial B_N(0, 1)$  is closed.

**Proof.** Letting  $\{x_n\} \subseteq BP_f \cup \partial B_N(0, 1)$  converge to  $x$ , we need to prove that  $x \in BP_f \cup \partial B_N(0, 1) \subseteq \bar{B}_N(0, 1)$ .

If  $x \in \partial B_N(0, 1)$ , there is nothing to prove.

If  $x \in B_N(0, 1)$ , we will prove that  $x \in BP_f$ , namely,  $NL(x)$  contains at least one line.

Assume by contradiction that  $NL(x) = 0$ . In particular,  $H_0^+(x)$  does not

contain lines. Similarly by the previous proof, there exist a vector  $(v_0, 0) \in H_0(x)$  and a constant  $\delta_1 > 0$  such that  $v_0 \in \mathbb{R}^N$ ,  $\|v_0\| = 1$  and

$$\langle -(v_0, 0), \frac{\xi}{\|\xi\|} \rangle \geq \delta_1 \quad \text{for all } 0 \neq \xi \in H_0^+(x). \quad (5.3.28)$$

On the other hand, since  $x \in B_N(0, 1)$  we have  $x_n \in B_N(0, 1)$  for  $n$  large enough. Thus  $x_n \in BP_f$ . From Lemma 5.3.3, for  $n$  large enough,  $H_0^+(x_n)$  contains at least one line. Therefore, for each  $n$  large enough, there exists a vector  $\xi_n \in N_0^L(x_n)$  such that

$$\langle -(v_0, 0), \xi_n \rangle \leq 0. \quad (5.3.29)$$

By Corollary 5.3.2,  $\|\xi_n\| = 1$ . We assume without loss of generality that  $\lim_{n \rightarrow \infty} \xi_n = \bar{\xi}$ . Recalling Corollary 5.3.3, we have that  $\bar{\xi} \in N_0^L(x)$ . Moreover, by taking  $n \rightarrow \infty$  in (5.3.29) we get

$$\langle -(v_0, 0), \bar{\xi} \rangle \leq 0. \quad (5.3.30)$$

Recalling (5.3.28), we get a contradiction.  $\square$

### 5.3.2 Zero Lebesgue measure of some special subsets of the set of bad points

The below lemma is the first step to prove that the  $\mathcal{L}^N$ -density of  $BP_f$  at  $x \in BP_f$  has zero value.

**Lemma 5.3.5** *Define, for  $x \in BP_f$ ,  $F^+(x) = \{y \in B(0, 1) \mid f(y) \geq f(x)\}$ . Then the  $\mathcal{L}^N$ -density of  $F^+(x)$  at  $x$  is zero, i.e.,*

$$D_{F^+(x)}^N(x) := \lim_{\delta \rightarrow 0} \frac{\mathcal{L}^N(B_N(x, \delta) \cap F^+(x))}{\mathcal{L}^N(B_N(x, \delta))} = 0.$$

**Proof.** Since  $x \in BP_f$  (i.e.,  $NL(x)$  contains at least one line), there exists  $(\zeta_0, 0) \in N_{\text{hypo}(f)}^P(x, f(x))$  such that  $(-\zeta_0, 0) \in N_{\text{hypo}(f)}^P(x, f(x))$  and  $\|\zeta_0\| = 1$ . Thus there exists a constant  $\sigma_0 > 0$  such that for all  $y \in B_N(0, 1)$  and for all  $\beta \leq f(y)$ , it holds

$$\begin{cases} \langle (\zeta_0, 0), (y - x, \beta - f(x)) \rangle & \leq \sigma_0 (\|y - x\|^2 + |\beta - f(x)|^2), \\ \langle (-\zeta_0, 0), (y - x, \beta - f(x)) \rangle & \leq \sigma_0 (\|y - x\|^2 + |\beta - f(x)|^2). \end{cases} \quad (5.3.31)$$

Therefore, for all  $y \in F^+(x) \cap B_N(x, \delta)$ , by taking  $\beta = f(x)$  in (5.3.31) we obtain

$$\begin{cases} \langle \zeta_0, y - x \rangle & \leq \sigma_0 \|y - x\|^2 \leq \sigma_0 \delta^2, \\ \langle -\zeta_0, y - x \rangle & \leq \sigma_0 \|y - x\|^2 \leq \sigma_0 \delta^2. \end{cases} \quad (5.3.32)$$

From (5.3.32), the set

$$F^+(x) \cap B_N(x, \delta) \subseteq x + \{t\zeta_0 + v \mid t \in [-\sigma_0 \delta^2, \sigma_0 \delta^2], v \in B_N(0, \delta) \cap \zeta_0^\perp\}$$

where  $\zeta_0^\perp = \{w \in \mathbb{R}^N \mid \langle w, \zeta_0 \rangle = 0\}$ . Therefore,

$$D_{F^+(x)}^N(x) := \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^N(B_N(x, \delta) \cap F^+(x))}{\mathcal{L}^N(B_N(x, \delta))} \leq \lim_{\delta \rightarrow 0^+} \frac{\sigma_0 \delta^{N+1}}{\omega_N \delta^N} = \lim_{\delta \rightarrow 0^+} \frac{\sigma_0 \delta}{\omega_N} = 0$$

where  $\omega_N = \mathcal{L}^N(B_N(0, 1))$ . The proof is completed.  $\square$

Since  $BP_f^+(x) \subseteq F^+(x)$ , the next corollary follows immediately

**Corollary 5.3.5** *If  $x \in BP_f$  then the  $\mathcal{L}^N$ -density of  $BP_f^+(x)$  at  $x$*

$$D_{BP_f^+(x)}^N(x) := \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^N(B_N(x, \delta) \cap BP_f^+(x))}{\mathcal{L}^N(B_N(x, \delta))} = 0.$$

In order to use induction in the proof of Theorem 5.2.2, we need the following two lemmas. In the first lemma, we are working on the cases  $N \geq 2$ . For every vector  $x \in \mathbb{R}^N$  we rewrite  $x = (x_1, x_2)$  where  $x_1 \in \mathbb{R}^{N-1}$  and  $x_2 \in \mathbb{R}$ . For every  $x_2 \in (-1, 1)$ , the function restricted to the first  $n - 1$  variables,  $f_{x_2} : B_{N-1}(0, \sqrt{1 - x_2^2}) \rightarrow \mathbb{R}$ , is denoted by  $f_{x_2}(x_1) = f(x_1, x_2)$  for all  $x_1 \in B_{N-1}(0, \sqrt{1 - x_2^2})$ .

**Lemma 5.3.6** *Let  $(x_1, x_2) \in B_N(0, 1)$  and let  $(\xi_1, \xi_2, \lambda)$  be a proximal normal vector to  $\text{hypo}(f)$  at  $(x_1, x_2, f(x_1, x_2))$  realized by a ball of radius  $\rho$ . If  $(\xi_1, \lambda) \neq 0$  then  $(\xi_1, \lambda)$  is also a proximal vector to  $\text{hypo}(f_{x_2})$  at  $(x_1, f_{x_2}(x_1))$  realized by a ball of radius  $\frac{\|(\xi_1, \lambda)\|}{\|(\xi_1, \xi_2, \lambda)\|} \rho$ .*

**Proof.** The vector  $(\xi_1, \xi_2, \lambda)$  being a proximal normal to the hypograph of  $f$  at  $(x_1, x_2) \in B_N(0, 1)$  realized by a ball of radius  $\rho$  means that for all  $(y_1, y_2) \in \mathbb{R}^N$  and for all  $\beta \leq f(y_1, y_2)$ , we have

$$\begin{aligned} \left\langle \frac{(\xi_1, \xi_2, \lambda)}{\|(\xi_1, \xi_2, \lambda)\|}, (y_1, y_2, \beta) - (x_1, x_2, f(x_1, x_2)) \right\rangle \\ \leq \frac{1}{2\rho} (\|y_1 - x_1\|^2 + |y_2 - x_2|^2 + |\beta - f(x_1, x_2)|^2). \end{aligned} \quad (5.3.33)$$

By taking  $y_2 = x_2$  in (5.3.33), and by replacing  $f(x_1, x_2) = f_{x_2}(x_1)$ ,  $f(y_1, y_2) = f(y_1, x_2) = f_{x_2}(y_1)$  in (5.3.33), we obtain that for all  $y_1 \in B_{N-1}(0, \sqrt{1 - x_2^2})$  and for all  $\beta \leq f_{x_2}(y_1)$ , it holds

$$\left\langle \frac{(\xi_1, \lambda)}{\|(\xi_1, \xi_2, \lambda)\|}, (y_1, \beta) - (x_1, f_{x_2}(x_1)) \right\rangle \leq \frac{1}{2\rho} (\|y_1 - x_1\|^2 + |\beta - f_{x_2}(x_1)|^2). \quad (5.3.34)$$

Since  $(\xi_1, \lambda) \neq 0$ , from (5.3.34) we get that for all  $y_1 \in B_{N-1}(0, \sqrt{1 - x_2^2})$  and for all  $\beta \leq f_{x_2}(y_1)$ , it holds

$$\left\langle \frac{(\xi_1, \lambda)}{\|(\xi_1, \lambda)\|}, (y_1, \beta) - (x_1, f_{x_2}(x_1)) \right\rangle \leq \frac{1}{2\rho \frac{\|(\xi_1, \lambda)\|}{\|(\xi_1, \xi_2, \lambda)\|}} (\|y_1 - x_1\|^2 + |\beta - f_{x_2}(x_1)|^2).$$



The proof is completed.  $\square$

The second lemma is used to treat the case  $(\xi_1, \lambda) = 0$  in Lemma 5.3.6 in the proof of our main theorem. Some notations are needed in this lemma: Let  $\zeta$  be a unit vector in  $\mathbb{R}^N$ , we denote by:

- i)  $N_0^\zeta = \{x \in B_N(0, 1) \mid (\zeta, 0) \in N_{\text{hypo}(f)}^P(x, f(x)) \text{ is realized by a ball of radius } \rho\}$ ,
- ii)  $BP_f^\zeta = BP_f \cap N_0^\zeta$ .

**Lemma 5.3.7**

- i)  $BP_f^\zeta \cup \partial B_N(0, 1)$  is closed.
- ii)  $BP_f^\zeta$  has zero  $N$ -Lebesgue measure.

**Proof of (i).** By Lemma 5.4.17, the set  $BP_f \cup \partial B_N(0, 1)$  is closed. Thus we only need to prove that  $N_0^\zeta \cup \partial B_N(0, 1)$  is closed.

Let  $\{x_n\} \subseteq N_0^\zeta \cup \partial B_N(0, 1)$  converge to  $x$ , we need to show that  $x \in N_0^\zeta \cup \partial B_N(0, 1)$ .

If  $x \in \partial B_N(0, 1)$  there is nothing to prove.

If  $x \in B_N(0, 1)$  then for  $n$  large enough we have  $x_n \in B_N(0, 1)$ . Thus  $x_n \in N_0^\zeta$ , namely,  $(\zeta, 0) \in N_{\text{hypo}(f)}^P(x_n, f(x_n))$  is realized by a ball of radius  $\rho$ , i.e, for all  $z \in B_N(0, 1)$  and for all  $\beta \leq f(z)$ , one has

$$\left\langle \frac{(\zeta, 0)}{\|(\zeta, 0)\|}, (z, \beta) - (x_n, f(x_n)) \right\rangle \leq \frac{1}{2\rho} (\|z - x_n\|^2 + |\beta - f(x_n)|^2). \quad (5.3.35)$$

Since  $\{x_n\}$  converges to  $x$  and  $f(\cdot)$  is continuous, by taking  $n \rightarrow \infty$  we have

$$\left\langle \frac{(\zeta, 0)}{\|(\zeta, 0)\|}, (z, \beta) - (x, f(x)) \right\rangle \leq \frac{1}{2\rho} (\|z - x\|^2 + |\beta - f(x)|^2). \quad (5.3.36)$$

for all  $z \in B_N(0, 1)$  and for all  $\beta \leq f(z)$ .

Thus  $x \in N_0^\zeta$ . The proof is completed.  $\square$

**Proof of (ii).** First, we prove that for all  $x \in BP_f^\zeta$ , it holds

$$D_{BP_f^\zeta}^N(x) = \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^N(B_N(x, \delta) \cap BP_f^\zeta)}{\mathcal{L}^N(B_N(x, \delta))} \leq \frac{1}{2}. \quad (5.3.37)$$

Indeed, since  $BP_f^\zeta \subseteq BP_f$ , recalling Corollary 5.3.5 we obtain

$$D_{BP_f^\zeta \cap BP_f^+(x)}^N(x) = \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^N(B_N(x, \delta) \cap BP_f^\zeta \cap BP_f^+(x))}{\mathcal{L}^N(B_N(x, \delta))} = 0.$$

Thus the inequality (5.3.37) will hold if

$$D_{BP_f^\zeta \cap BP_f^-(x)}^N(x) = \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^N(B_N(x, \delta) \cap BP_f^\zeta \cap BP_f^-(x))}{\mathcal{L}^N(B_N(x, \delta))} \leq \frac{1}{2}. \quad (5.3.38)$$

If  $y \in BP_f^\zeta$ , we have  $(\zeta, 0) \in N_0^\zeta(y)$ . Thus for all  $z \in B_N(0, 1)$  and for all  $\beta \leq f(z)$ , it holds

$$\langle (\zeta, 0), (z - y, \beta - f(y)) \rangle \leq \frac{1}{2\rho} (\|z - y\|^2 + |\beta - f(y)|^2). \quad (5.3.39)$$

Thus, if  $y \in BP_f^\zeta \cap BP_f^-(x)$  we can take  $z = x$  and  $\beta = f(y)$  in 5.3.39 to get

$$\langle \zeta, x - y \rangle \leq \frac{1}{2\rho} \|x - y\|^2. \quad (5.3.40)$$

Therefore, for all  $\delta > 0$  small enough, we have

$$\langle \zeta, x - y \rangle \leq \frac{1}{2\rho} \delta^2 \quad \text{for all } y \in [B_N(x, \delta) \cap BP_f^\zeta \cap BP_f^-(x)]. \quad (5.3.41)$$

(5.3.41) says that

$$[B_N(x, \delta) \cap BP_f^\zeta \cap BP_f^-(x)] \subset x + \{t\zeta + v \mid t \in [-\frac{\delta^2}{2\rho}, \delta], v \in B_N(0, \delta) \cap \zeta^\perp\}$$

where  $\zeta^\perp = \{w \in \mathbb{R}^N \mid \langle w, \zeta \rangle = 0\}$ . Thus, (5.3.38) follows. From (i),  $BP_f^\zeta$  is a Borel set. Moreover, from (5.3.38), the  $\mathcal{L}^N$ -density of  $BP_f^\zeta$  at every point which is in  $BP_f^\zeta$  is less than  $\frac{1}{2}$ . Therefore, by the Lebesgue theorem we have  $\mathcal{L}^N(BP_f^\zeta) = 0$ .  $\square$

## 5.4 Proof of our main results of Chapter 5

### 5.4.1 Proof of Theorem 5.2.2

#### One dimension case

In this part, we are working on  $\mathbb{R}$ . The function  $f(\cdot)$  is defined on  $B_1(0, 1) = \{x \in \mathbb{R} \mid |x| < 1\}$ . Therefore the proximal normal cone  $N_{\text{hypo}(f)}^P(x, f(x)) \subset \mathbb{R}^2$  contains at most one line.

**Lemma 5.4.1** *For all  $x \in BP_f$ , we have  $N_0^L(x) = \{(1, 0), (-1, 0)\}$ .*

**Proof.** Since  $N_0^L(x) \subseteq (-\partial^\infty f(x), 0)$ , we have  $N_0^L(x) \subseteq \{(t, 0) \mid t \in \mathbb{R}\}$ . Therefore, from the fact that  $\|\xi\| = 1$  for all  $\xi \in N_0^L(x)$ , we obtain

$$N_0^L(x) \subseteq \{(1, 0), (-1, 0)\} \quad (5.4.1)$$

Recalling Lemma 5.3.3, the set  $H_0^+(x) = \text{span}^+\{N_0^L(x)\}$  contains at least one line. Thus, the proof is completed by (5.4.1).

The following statement is a one dimensional version of Theorem 5.2.2.

**Theorem 5.4.1** *Let  $f : B_1(0, 1) \rightarrow \mathbb{R}$  be continuous. Assume that  $\text{hypo}(f)$  satisfies the  $\rho$ -external sphere condition. Then*

- i)  $BP_f \cup \partial B_1(0, 1)$  is closed.
- ii)  $\mathcal{L}^1(BP_f) = 0$ .

(i) is the particular case ( $N=1$ ) of Lemma 5.3.4 .

**Proof of (ii).** We prove first that, for all  $x \in BP_f$ , the  $\mathcal{L}^1$ -density of  $BP_f$  at  $x$  is zero, namely,

$$D_{BP_f}^1(x) := \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^1(B_1(x, \delta) \cap BP_f)}{\mathcal{L}^1(B_1(x, \delta))} = 0. \quad (5.4.2)$$

Recalling Corollary 5.3.5 for  $N=1$ , we have

$$D_{BP_f^+(x)}^1(x) = \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^1(B_1(x, \delta) \cap BP_f^+(x))}{\mathcal{L}^1(B_1(x, \delta))} = 0.$$

Therefore, 5.4.2 follows from

$$D_{BP_f^-(x)}^1(x) = \lim_{\delta \rightarrow 0^+} \frac{\mathcal{L}^1(B_1(x, \delta) \cap BP_f^-(x))}{\mathcal{L}^1(B_1(x, \delta))} = 0. \quad (5.4.3)$$

From Lemma 5.4.1, for every  $y \in BP_f$ , we have  $N_0^I(y) = \{(1, 0), (-1, 0)\}$ . Thus, for all  $y \in BP_f$  it holds

$$\begin{cases} \langle (1, 0), (z - y, \beta - f(y)) \rangle & \leq \frac{1}{2\rho} (|z - y|^2 + |\beta - f(y)|^2), \\ \langle (-1, 0), (z - y, \beta - f(y)) \rangle & \leq \frac{1}{2\rho} (|z - y|^2 + |\beta - f(y)|^2). \end{cases} \quad (5.4.4)$$

for all  $z \in \overline{B_1(0, 1)}$  and for all  $\beta \leq f(z)$ .

Since  $f(y) \leq f(x)$  for all  $y \in BP_f^-(x)$ , we can take  $z = x$  and  $\beta = f(y)$  in (5.4.4) to get

$$|x - y| \leq \frac{1}{2\rho} |x - y|^2 \quad \text{for all } y \in BP_f^-(x). \quad (5.4.5)$$

Thus  $B_1(x, \delta) \cap BP_f^-(x) = \{x\}$  for all  $0 < \delta < 2\rho$  and so 5.4.3 follows.

We are now going to complete the proof of (ii).

Since  $BP_f \cup \partial B_1(0, 1)$  is closed,  $BP_f$  is a Borel set. From 5.4.2, the  $\mathcal{L}^1$ -density of  $BP_f$  at  $x$  has zero value for all  $x \in BP_f$ . Therefore, by the Lebesgue theorem, we have  $\mathcal{L}^1(BP_f) = 0$ .  $\square$

### General case

(i) of Theorem 5.2.2 is precisely Lemma 5.3.4.

We are going to prove (ii) of Theorem 5.2.2 by induction.

If  $N = 1$ , (ii) of Theorem 5.2.2 follows from Theorem 5.4.1.

Assume that (ii) of Theorem 5.2.2 holds for  $N = k \geq 1$ . We prove that (ii)

of Theorem 5.2.2 will hold for  $N = k + 1$ .

Let  $\zeta^+ = (0, 1)$  and  $\zeta^- = (0, -1)$  be in  $\mathbb{R}^{k+1}$ . Recalling Lemma 5.3.7, we obtain that  $(BP_f^{\zeta^+} \cup \partial B_{k+1}(0, 1))$  and  $(BP_f^{\zeta^-} \cup \partial B_{k+1}(0, 1))$  are closed. Moreover,

$$\mathcal{L}^{k+1}(BP_f^{\zeta^+}) = \mathcal{L}^{k+1}(BP_f^{\zeta^-}) = 0. \quad (5.4.6)$$

Set  $E = B_{k+1}(0, 1) \setminus [N_0^{\zeta^+} \cup N_0^{\zeta^-} \cup \partial B_{k+1}(0, 1)]$ . One can easily see that  $E$  is an open set in  $\mathbb{R}^{k+1}$ . From (5.4.6), the conclusion of (ii) of Theorem 5.2.2 follows from the equality

$$\mathcal{L}^{k+1}(E \cap BP_f) = 0. \quad (5.4.7)$$

Recalling Lemma 5.3.4,  $BP_f \cap \partial B_{k+1}(0, 1)$  is closed. Thus  $E \cap BP_f$  is a Borel set. Therefore, by the Lebesgue theorem, (5.4.7) will follow if for every  $x \in E \cap BP_f$ , the  $\mathcal{L}^{k+1}$ -density  $D_{E \cap BP_f}^{k+1}(x)$  at  $x$  has zero value.

We divide the proof into several steps:

The first step is pivotal (see the below inequality (5.4.8)) to show that the restricted functions (defined before Lemma 5.3.6) which are restricted from the function  $f|_{B_{k+1}(x, r_x)}$  where  $x \in E$ , have the hypograph satisfying the  $\rho_x$ -external sphere condition.

*Step1:* Let  $x \in E$ . Since  $E$  is open, there exists  $r_x > 0$  such that  $\overline{B_{k+1}(x, r_x)} \subset E$ . By the external sphere assumption on  $f$ , for each  $y \in B_{k+1}(x, r_x)$ , there exists  $0 \neq (\xi_1^y, \xi_2^y, \lambda^y) \in N_{\text{hypo}(f)}^P(y, f(y))$  realized by a ball of radius  $\rho$  where  $\xi_1^y \in \mathbb{R}^k$  and  $\xi_2^y, \lambda^y \in \mathbb{R}$ . We claim that there exists a constant  $\alpha_x > 0$  such that

$$\frac{\|(\xi_1^y, \lambda^y)\|}{\|(\xi_1^y, \xi_2^y, \lambda^y)\|} \geq \alpha_x > 0 \quad \text{for all } y \in B_{k+1}(x, r_x). \quad (5.4.8)$$

Assume by contradiction that there exists a sequence  $\{y_n\} \subseteq B_{k+1}(x, r_x)$  such that

$$\lim_{n \rightarrow \infty} \frac{\|(\xi_1^{y_n}, \lambda^{y_n})\|}{\|(\xi_1^{y_n}, \xi_2^{y_n}, \lambda^{y_n})\|} = 0. \quad (5.4.9)$$

Assume without loss of generality that  $\lim_{n \rightarrow \infty} y_n = \bar{y} \in \overline{B_{k+1}(x, r_x)}$  and  $\lim_{n \rightarrow \infty} \frac{(\xi_1^{y_n}, \xi_2^{y_n}, \lambda^{y_n})}{\|(\xi_1^{y_n}, \xi_2^{y_n}, \lambda^{y_n})\|} = (\bar{\xi}_1, \bar{\xi}_2, \bar{\lambda})$ . From (5.4.9), one can see that

$$(\bar{\xi}_1, \bar{\xi}_2, \bar{\lambda}) \in \{(0, 1, 0), (0, -1, 0)\} = \{(\zeta^+, 0), (\zeta^-, 0)\}. \quad (5.4.10)$$

Moreover,  $(\bar{\xi}_1, \bar{\xi}_2, \bar{\lambda})$  is a proximal normal vector to  $\text{hypo}(f)$  at  $(\bar{y}, f(\bar{y}))$  realized by a ball of radius  $\rho$ . Indeed, since  $0 \neq (\xi_1^{y_n}, \xi_2^{y_n}, \lambda^{y_n})$  and  $(\xi_1^{y_n}, \xi_2^{y_n}, \lambda^{y_n}) \in N_{\text{hypo}(f)}^P(y_n, f(y_n))$  is realized by a ball of radius  $\rho$ , we have

$$\left\langle \frac{(\xi_1^n, \xi_2^n, \lambda^n)}{\|(\xi_1^n, \xi_2^n, \lambda^n)\|}, (z, \beta) - (y_n, f(y_n)) \right\rangle \leq \frac{1}{2\rho} (\|z - y_n\|^2 + |\beta - f(y_n)|^2)$$

for all  $z \in B_{k+1}(0, 1)$  and for all  $\beta \leq f(z)$ .

By taking  $n \rightarrow \infty$ , we obtain that

$$\langle (\bar{\xi}_1, \bar{\xi}_2, \bar{\lambda}), (z, \beta) - (\bar{y}, f(\bar{y})) \rangle \leq \frac{1}{2\rho} (\|z - \bar{y}\|^2 + |\beta - f(\bar{y})|^2)$$

for all  $z \in B_{k+1}(0, 1)$  and for all  $\beta \leq f(z)$ .

Therefore, by (5.4.10), we get  $\bar{y} \in N_0^{\zeta^+} \cup N_0^{\zeta^-}$ . This is a contradiction because  $\bar{y} \in \overline{B_{k+1}(x, r_x)} \subset E = B_{k+1}(0, 1) \setminus [N_0^{\zeta^+} \cup N_0^{\zeta^-} \cup \partial B_{k+1}(0, 1)]$ .

The second step allows us to make a connection between the set of *bad points* of  $f$  and the set of *bad points* of restricted functions of  $f$ .

*Step 2:* Let  $x \in E \cap BP_f$ . We claim that there exists a line  $\{t\xi_x \mid t \in \mathbb{R}\} \subseteq N_{\text{hypo}(f)}^P(x)$  such that  $\{t\xi_x \mid t \in \mathbb{R}\} \neq \{t(\zeta^+, 0) \mid t \in \mathbb{R}\}$ .

Assume by contraction, since  $x \in BP_f$ , i.e.  $NL(x) \neq 0$ , we have  $NL(x) = \{t(\zeta^+, 0) \mid t \in \mathbb{R}\}$ . Recalling Lemma 5.3.3, the set  $H_0^+(x) \subseteq NL(x)$  contains at least one line. Therefore  $H_0^+(x) = \{t(\zeta^+, 0) \mid t \in \mathbb{R}\}$  which implies that  $(\zeta^+, 0) \in N_0^L(x)$ . Recalling Corollary 5.3.2,  $(\zeta^+, 0) \in N_{\text{hypo}(f)}^P(x, f(x))$  is realized by a ball of radius  $\rho$ . Thus  $x \in N_0^{\zeta^+}$  and this is a contradiction because  $x \in E$ .

In the next step, we are going to prove that  $\mathcal{L}^{k+1}(B_{k+1}(x, r_x) \cap BP_f) = 0$  by our inductive assumption .

*Step 3:* Let  $\bar{f} = f|_{B_{k+1}(x, r_x)} : B_{k+1}(x, r_x) \rightarrow \mathbb{R}$  be the restricted function of  $f$  on  $B_{k+1}(x, r_x)$ . From Lemma 7.2.2, the continuous function  $\bar{f}$  has  $\text{hypo}(\bar{f})$  satisfying the  $\rho$ -external sphere condition, and

$$BP_f \cap B_{k+1}(x, r_x) = BP_{\bar{f}}. \quad (5.4.11)$$

Moreover, two properties which we claimed in Step 1 and Step 2 still hold for the function  $\bar{f}$ .

Since (5.4.11) holds, we only need to prove  $\mathcal{L}^{k+1}(BP_{\bar{f}}) = 0$ .

In order to make the proof more clear, we restate our above problem by replacing  $x = 0$ ,  $r_x = 1$  and  $\bar{f} = f$ . The statement is that

Let  $f : B_{k+1}(0, 1) \rightarrow \mathbb{R}$  be continuous. Assume that  $\text{hypo}(f)$  satisfies  $\rho$ -external

*sphere condition*. Moreover,

i) For all  $y \in B_{k+1}(0, 1)$ , there exists a non zero vector  $(\xi_1^y, \xi_2^y, \lambda^y) \in N_{\text{hypo}(f)}^P(y, f(y))$  realized by a ball of radius  $\rho$  such that

$$\frac{\|(\xi_1^y, \lambda^y)\|}{\|(\xi_1^y, \xi_2^y, \lambda^y)\|} \geq \alpha_0 > 0. \quad (5.4.12)$$

ii) For all  $x \in BP_f$ , there exists a line  $\{t\xi_x \mid t \in \mathbb{R}\} \subseteq NL(x)$  such that  $\{t\xi_x \mid t \in \mathbb{R}\} \neq \{t(\zeta^+, 0) \mid t \in \mathbb{R}\}$ .

Then  $\mathcal{L}^{k+1}(BP_f) = 0$ .

*Proof.* Since  $k \geq 1$ , for every  $x \in \mathbb{R}^{k+1}$ , we write  $x = (x_1, x_2)$  where

$x_1 \in \mathbb{R}^k$  and  $x_2 \in \mathbb{R}$ . For each  $x_2 \in (-1, 1)$ , the restricted function  $f_{x_2} : B_k(0, \sqrt{1-x_2^2}) \rightarrow \mathbb{R}$  is denoted by  $f_{x_2}(x_1) = f(x_1, x_2)$  for all  $x_1 \in B_k(0, \sqrt{1-x_2^2})$ .

First, we claim that  $\text{hypo}(f_{x_2})$  satisfies  $\rho\alpha_0$ -external sphere condition. Indeed by assumption (i) of the above statement we have that, for each  $x_1 \in B_k(0, \sqrt{1-x_2^2})$ , or  $(x_1, x_2) \in B_{k+1}(0, 1)$ , there exists a vector

$$0 \neq (\xi_1^{(x_1, x_2)}, \xi_2^{(x_1, x_2)}, \lambda^{(x_1, x_2)}) \in N_{\text{hypo}(f)}^P((x_1, x_2), f(x_1, x_2))$$

realized by a ball of radius  $\rho$  such that

$$\frac{\|(\xi_1^{(x_1, x_2)}, \lambda^{(x_1, x_2)})\|}{\|(\xi_1^{(x_1, x_2)}, \xi_2^{(x_1, x_2)}, \lambda^{(x_1, x_2)})\|} \geq \alpha_0 > 0. \quad (5.4.13)$$

Recalling Lemma 5.3.6 for  $N = k + 1 \geq 2$  and observing that  $(\xi_1, \xi_2, \lambda) = (\xi_1^{(x_1, x_2)}, \xi_2^{(x_1, x_2)}, \lambda^{(x_1, x_2)})$ , and by (5.4.13) we obtain that  $(\xi_1^{(x_1, x_2)}, \lambda^{(x_1, x_2)})$  is also a proximal normal vector to  $\text{hypo}(f_{x_2})$  at  $(x_1, f_{x_2}(x_1))$  realized by a ball of radius  $\rho\alpha_0$ .

Second, we claim that

$$\mathcal{L}^k(BP_{f_{x_2}}) = 0 \quad \text{for all } x_2 \in (-1, 1). \quad (5.4.14)$$

Indeed, set  $\gamma(x_2) = \frac{1}{\sqrt{1-x_2^2}}$  and let  $h_{x_2} = f_{x_2}^{\gamma(x_2)}$  be the  $\gamma(x_2)$ -stretched function of  $f_{x_2}$  (see Lemma 7.2.3). By Lemma 7.2.3 and by the first step, the continuous function  $h_{x_2} : B_k(0, 1) \rightarrow \mathbb{R}$  has hypograph satisfying the  $\rho_1$ -external sphere condition where  $\rho_1 = \rho\alpha_0 \frac{(1-x_2^2)^{\frac{1}{2}}}{(2-x_2^2)^{\frac{3}{2}}}$ . Therefore, by the inductive assumption, we have

$$\mathcal{L}^k(BP_{h_{x_2}}) = 0. \quad (5.4.15)$$

Moreover, recalling Corollary 7.2.1 for  $g = f_{x_2}$  and  $\gamma = \gamma(x_2)$  we get

$$BP_{h_{x_2}} = (1-x_2^2)^{-\frac{1}{2}} BP_{f_{x_2}}. \quad (5.4.16)$$

Combining (5.4.15) and (5.4.16), we get (5.4.14).

Thirdly, we claim that

$$BP_f \subseteq \bigcup_{x_2 \in (-1, 1)} BP_{f_{x_2}} \times \{x_2\}. \quad (5.4.17)$$

Assume  $x = (x_1, x_2) \in BP_f$ . By (ii) there exists a line  $\{t\xi_x \mid t \in \mathbb{R}\} \subseteq NL(x) \subseteq (-\partial^\infty f(x), 0)$  such that  $\{t\xi_x \mid t \in \mathbb{R}\} \neq \{t(\zeta^+, 0) \mid t \in \mathbb{R}\}$  and  $\|\xi_x\| = 1$ . Therefore,  $\xi_x = (\xi_1, \xi_2, 0)$  and  $-\xi_x = (-\xi_1, -\xi_2, 0)$  are proximal

normal vectors to  $\text{hypo}(f)$  at  $(x, f(x))$  realized by a ball of radius  $\sigma$  where  $\sigma > 0$ ,  $0 \neq \xi_1 \in \mathbb{R}^k$ ,  $x_2 \in \mathbb{R}$  and  $\|(\xi_1, \xi_2)\| = 1$ . Recalling Lemma 5.3.6, we obtain that  $(\xi_1, 0)$  and  $(-\xi_1, 0)$  are proximal normal vectors to the hypograph of  $f_{x_2}$  at  $(x_1, f_{x_2}(x_1))$ . This implies that  $N_{\text{hypo}f_{x_2}}^P(x_1, f_{x_2}(x_1))$  contains the line  $\{t(\xi_1, 0) \mid t \in \mathbb{R}\}$ . Thus,  $x_1 \in BP_{f_{x_2}}$  or  $(x_1, x_2) \in (BP_{f_{x_1}}, x_2)$ .

Finally, since  $BP_f$  is a Borel set contained in  $B_{k+1}(0, 1)$ , the indicator function  $\mathbf{1}_{BP_f}$  is in  $\mathbb{L}^{k+1}(B_{k+1}(0, 1))$ . From Fubini's theorem, we have

$$\mathcal{L}^{k+1}(BP_f) = \int_{B_{k+1}(0,1)} \mathbf{1}_{BP_f} dx = \int_{-1}^1 \int_{B_k(0, \sqrt{1-x_2^2})} \mathbf{1}_{BP_f} dx_1 dx_2. \quad (5.4.18)$$

Combining the above equality and (5.4.17), we get

$$\mathcal{L}^{k+1}(BP_f) \leq \int_{-1}^1 \int_{B_k(0, \sqrt{1-x_2^2})} \mathbf{1}_{BP_{f_{x_2}}} dx_1 dx_2 = \int_{-1}^1 \mathcal{L}^k(BP_{f_{x_2}}) dx_1. \quad (5.4.19)$$

The proof is completed using (5.4.19) and (5.4.14).  $\square$

### 5.4.2 Proof of Theorem 5.2.1

*Proof of (i).* It is equivalent to prove that  $BP_f \cup \partial\Omega \subset \overline{\Omega}$  is closed. Let  $\{x_n\} \subseteq BP_f \cup \partial\Omega$  converge to  $x$ . We need to show that  $x \in BP_f \cup \partial\Omega$ .

If  $x \in \partial\Omega$ , there is nothing to prove.

If  $x \in \Omega$ , we will prove  $x \in BP_f$ . Indeed, there exist  $r_x > 0$  and  $M > 0$  such that  $x_n \in B_N(x, r_x) \subset \overline{B}_N(x, r_x) \subset \Omega$  for all  $n > M$ . From Lemma 7.2.2, we have  $x_n \in BP_{f|_{B_N(x, r_x)}}$  for all  $n > M$ . On the other hand, from Corollary 7.2.1, and (i) of Theorem 5.2.2, one can easily see that the set  $BP_{f|_{B_N(x, r_x)}} \cup \partial B_N(x, r_x)$  is closed. Therefore, the sequence  $\{x_n\}$  converge to  $x \in BP_{f|_{B_N(x, r_x)}} \cup \partial B_N(x, r_x)$ . Recalling again Lemma 7.2.2, we obtain  $x \in BP_f$ .

*Proof of (ii).* Since  $BP_f \cup \partial\Omega$  is closed,  $BP_f$  is a Borel set. Therefore, it is sufficient to prove that for all  $x \in BP_f$ , the  $\mathcal{L}^N$ -density of  $BP_f$  at  $x$  has zero value, i.e, for all  $x \in BP_f$

$$D_{BP_f}^N(x) = \lim_{\delta \rightarrow 0} \frac{\mathcal{L}^N(BP_f \cap B_N(x, \delta))}{\mathcal{L}^N(B_N(x, \delta))} = 0. \quad (5.4.20)$$

Indeed, for all  $x \in BP_f \subseteq \Omega$ , there exists  $r_x > 0$  such that  $\overline{B}_N(x, r_x) \subset \Omega$ . From Lemma 7.2.2, Lemma 7.2.3, Corollary 7.2.1 and Theorem 5.2.2, one can easily get

$$\mathcal{L}^N(BP_f \cap B(x, r_x)) = \mathcal{L}^N(BP_{f|_{B_N(x, r_x)}}) = 0, \quad (5.4.21)$$

and (5.4.20) follows.  $\square$

### 5.4.3 Proof of Colloraries

**Proof of Corollary 5.2.1.** From Theorem 5.2.1 we have

The set  $\Omega_P$  is open. The function  $f|_{\Omega_P} : \Omega_P \rightarrow \mathbb{R}$  is a continuous function and

i) The set  $\text{hypo}(f|_{\Omega_P})$  satisfies the  $\theta$  – *external sphere condition*.

ii) For every  $x \in \Omega_P$ , the set  $N_{\text{hypo}(f|_{\Omega_P})}^P(x, f|_{\Omega_P}(x))$  is pointed.

The remainder of the proof is done by the argument in [19]. More precisely, one can prove that  $N_{\text{hypo}(f|_{\Omega_P})}^P(x, f|_{\Omega_P}(x)) = \text{Co}\{tN^L(x) \mid t \geq 0\}$  (see Lemma 4.7, Theorem 4.1, Theorem 3.1, Theorem 3.2 in [19]). From Corollary 5.3.1 in this paper, if  $\xi \in N^L(x)$  then  $\xi \in N_{\text{hypo}(f|_{\Omega_P})}^P(x, f|_{\Omega_P}(x))$  is realized by a ball of radius  $\theta(x)$ , the proof is completed by following the proof of Theorem 3.3 in [19].  $\square$

**Proof of Corollary 5.2.3.** Using the Proposition (3.1) in [19], the  $\text{hypo}(T)$  satisfies the  $\theta$  – *external sphere condition*. Applying Corollary 5.2.1 for  $f = T(\cdot)$ , we get the conclusion.  $\square$



## Chapter 6

# Rectifiability of the set of bad points

We prove here some rectifiability properties of the set of bad points  $BP_f$  of  $f$  where the hypograph of  $f$  doesn't require an external sphere condition. The set  $BP_f$  will be also defined more generally (see Section 6.1 for the definition). We partition the set  $BP_f$  (see (6.1.6)) into sets  $BP_{f,k}, k = 1, \dots, N$ , where, roughly speaking, the suffix  $k$  corresponds to the dimension of the largest vector space contained in the set  $\partial^{F,\infty} f$  of Fréchet horizon supergradients of  $f$  (see Section 6.1 for the definition). We are able to prove that  $BP_{f,k}$  is countably  $(N - k)$ -rectifiable.

**Theorem 6.0.2** *Let  $\Omega \subseteq \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be upper semi-continuous. Then the set  $BP_{f,k}$  is countably  $(N - k)$ -rectifiable.*

Moreover, under an external sphere condition on the hypograph of  $f$ , the definition of  $BP_f$  here will coincide with the one in the previous chapter. Therefore, we also refine Theorem 5.2.2 as follows:

**Theorem 6.0.3** *Let  $\Omega \subseteq \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be continuous. If the hypograph of  $f$  satisfies the  $\theta$ -exterior sphere condition for some  $\theta > 0$ , then the set of bad points  $BP_f$  is locally  $(N - 1)$ -rectifiable. In particular,  $\mathcal{H}^{N-1}(BP_f \cap K)$  is finite for any compact set  $K \subset \mathbb{R}^N$ .*

Finally, in Section 6.3 we provide an example showing that, in general, the set  $BP_{f,k}, k \geq 2$  may not have finite  $(N - k)$ -Hausdorff measure even under the exterior sphere condition.

### 6.1 Notations

To make the reader easy to follow, we prefer to rewrite shortly some basic notations.

Let  $\Omega \subseteq \mathbb{R}^N$  be open and let  $f : \Omega \rightarrow \mathbb{R}$  be upper semi-continuous. The hypograph of  $f$  is denoted by

$$\text{hypo}(f) = \{(x, \beta) \mid x \in \Omega, \beta \leq f(x)\}. \quad (6.1.1)$$

The vector  $(-v, \lambda) \in \mathbb{R}^N \times \mathbb{R}$  is a Fréchet normal vector to  $\text{hypo}(f)$  at  $(x, f(x))$  iff

$$\limsup_{\text{hypo}(f) \ni (y, \beta) \rightarrow (x, f(x))} \left\langle (-v, \lambda), \frac{(y, \beta) - (x, f(x))}{|y - x| + |\beta - f(x)|} \right\rangle \leq 0. \quad (6.1.2)$$

We denote by  $N_{\text{hypo}(f)}^F(x, f(x))$  the set of Fréchet normal vectors to  $\text{hypo}(f)$  at  $(x, f(x))$ .

**Remark 6.1.1** If  $(-v, \lambda) \in N_{\text{hypo}(f)}^F(x, f(x))$  then  $\lambda \geq 0$ .

Recalling that  $N_{\text{hypo}(f)}^P(x, f(x))$  is the set of proximal normal vectors to  $\text{hypo}(f)$  at  $(x, f(x))$ , we have:

**Remark 6.1.2**  $N_{\text{hypo}(f)}^P(x, f(x)) \subseteq N_{\text{hypo}(f)}^F(x, f(x))$  for all  $x \in \Omega$ .

Associated with  $\text{hypo}(f)$ , we define that

1.  $\partial^F f(x) = \{v \mid (-v, 1) \in N_{\text{hypo}(f)}^F(x, f(x))\}$  is set of Fréchet supergradients of  $f$  at  $x$ .
2.  $\partial^{F, \infty} f(x) = \{v \mid (-v, 0) \in N_{\text{hypo}(f)}^P(x, f(x))\}$   $v$  is the set a Fréchet horizon supergradients of  $f$  at  $x$ .

The largest vector subspace contained in  $N_{\text{hypo}(f)}^F(x, f(x))$  will be denoted by

$$NL(x) = \{ \xi \in N_{\text{hypo}(f)}^F(x, f(x)) \mid -\xi \in N_{\text{hypo}(f)}^F(x, f(x)) \}. \quad (6.1.3)$$

From Remark 6.1.1, one can see that  $NL(x) \subseteq \{(v, 0) \mid -v \in \partial^\infty f(x)\}$ . Let us define

$$V_x := \{v \in \mathbb{R}^N \mid (v, 0) \in NL(x)\}; \quad (6.1.4)$$

clearly,  $V_x$  is the largest vector space contained in  $\partial^\infty f(x)$  and  $\dim V_x = \dim NL(x)$ . We say that  $v \in V_x$  is realized by a ball of radius  $\theta$  if  $(v, 0) \in N_{\text{hypo}(f)}^P(x, f(x))$  is realized by a ball of radius  $\theta$ .

The set of *bad points*  $BP_f$  of  $f$  is defined by

$$BP_f = \{x \in \Omega \mid NL(x) \neq \{0\}\}. \quad (6.1.5)$$

According to the dimension of  $NL(x)$ , for  $k = 1, \dots, N$  we introduce

$$BP_{f,k} = \{x \in BP_f \mid \dim NL(x) = k\} = \{x \in BP_f \mid \dim V_x = k\}. \quad (6.1.6)$$

It is clear that  $BP_f = \bigcup_{k=1}^N BP_{f,k}$ .

Now, let  $k \geq 0$  and  $A, B \subset \mathbb{R}^N$  be fixed. We recall that:

- (i)  $\mathcal{H}_k(A)$  is the  $k$ -dimensional Hausdorff measure of  $A$ ;
- (ii)  $d_{\mathcal{H}}(A, B)$  is the Hausdorff distance between  $A$  and  $B$ .

Finally, we will denote by  $G(N, k)$  the Grassmann manifold of all  $k$  – dimensional vector subspaces of  $\mathbb{R}^N$ ; we endow  $G(N, k)$  with the distance

$$d_G(V_1, V_2) := d_H(V_1 \cap S^{N-1}, V_2 \cap S^{N-1}).$$

The metric space  $(G(N, k), d_G)$  is separable and, in particular, the following property holds:

$$\forall R > 0 \exists V_1, \dots, V_m \in G(N, k) \text{ s.t. } G(N, k) \subset \bigcup_{i=1}^m B_G(V_i, R) \quad (6.1.7)$$

where  $B_G(V_i, R)$  denote the open ball (with respect to  $d_G$ ) with center  $V_i$  and radius  $R$ .

## 6.2 Rectifiability results for the set of bad points

### 6.2.1 Preparatory Lemmas

Let  $V \in G(N, k)$  be fixed; each  $z \in \mathbb{R}^N$  can be written in a unique way as  $z = z_V + z_{V^\perp}$  where  $z_V \in V$  and  $z_{V^\perp} \in V^\perp$ . For  $\alpha \in (0, 1)$  we denote by  $C_\alpha(V)$  the open cone along  $V$  of aperture  $1/\alpha$  defined by

$$C_\alpha(V) := \{z \in \mathbb{R}^N \mid \|z_V\| > \alpha \|z\|\}.$$

If  $x \in \mathbb{R}^N$  we set

$$C_\alpha(x, V) := x + C_\alpha(V) = \{z \in \mathbb{R}^N \mid \|(z - x)_V\| > \alpha \|z - x\|\};$$

It is easily seen that

$$z \in C_\alpha(x, V) \iff \exists v \in V \cap S^{N-1} \text{ such that } \langle v, z - x \rangle > \alpha \|z - x\|. \quad (6.2.1)$$

We also point out the following implication:

$$d_G(V_1, V_2) < R \implies C_{\alpha+R}(x, V_1) \subset C_\alpha(x, V_2) \quad (6.2.2)$$

which holds provided  $\alpha + R < 1$ . To prove (6.2.2) it is enough to notice that for any  $z \in C_{\alpha+R}(x, V_1)$

$$\begin{aligned} & \text{there exists } v_1 \in V_1 \cap S^{N-1} \text{ such that } \langle v_1, z - x \rangle > (\alpha + R) \|z - x\| \\ & \text{there exists } v_2 \in V_2 \cap S^{N-1} \text{ such that } \|v_1 - v_2\| \leq R \end{aligned}$$

whence

$$\langle v_2, z - x \rangle = \langle v_1, z - x \rangle - \langle v_1 - v_2, z - x \rangle > \alpha \|z - x\|,$$

i.e.,  $z \in C_\alpha(x, V_2)$ .

For any fixed  $\rho > 0$ , let us introduce the sets

$$BP_{f,k}^\rho = \left\{ x \in BP_{f,k} \mid \left\langle v_x, \frac{y-x}{|y-x|+|\beta-f(x)|} \right\rangle \leq \frac{\|v_x\|}{8} \right. \\ \left. \forall v_x \in V_x, y \in B(x, \rho), \beta < f(y) \right\}. \quad (6.2.3)$$

**Remark 6.2.1** If  $\rho_1 > \rho_2 > 0$  then  $BP_{f,k}^{\rho_1} \subseteq BP_{f,k}^{\rho_2}$ .

As the following Lemma shows, the sets  $BP_{f,k}^\rho$  give a partition of  $BP_{f,k}$ .

**Lemma 6.2.1** *We have*

$$BP_{f,k} = \cup_{\rho>0} BP_{f,k}^\rho. \quad (6.2.4)$$

*In particular, from Remark 6.2.1 it holds*

$$BP_{f,k} = \cup_{i \in \mathbb{N} \setminus \{0\}} BP_{f,k}^{1/i}. \quad (6.2.5)$$

**Proof.** Fix  $x \in BP_{f,k}$  and let  $v_1, v_2, \dots, v_k$  be an orthonormal basis for  $V_x$ . By the definition of  $V_x$  we have  $-v_i \in V_x$  for all  $i \in \{1, 2, \dots, k\}$ . Recalling (6.1.4), (6.1.3) and (6.1.2), there exists a constant  $\rho_x > 0$  such that  $B(x, \rho_x) \subset \Omega$  and for all  $i \in \{1, 2, \dots, k\}$  one has

$$\left\langle v_i, \frac{y-x}{|y-x|+|\beta-f(x)|} \right\rangle \leq \frac{1}{8\sqrt{k}} \text{ and } \left\langle -v_i, \frac{y-x}{|y-x|+|\beta-f(x)|} \right\rangle \leq \frac{1}{8\sqrt{k}}$$

for all  $y \in B(x, \rho_x)$  and  $\beta \leq f(y)$ . Thus

$$\left| \left\langle v_i, \frac{y-x}{|y-x|+|\beta-f(x)|} \right\rangle \right| \leq \frac{1}{8\sqrt{k}} \quad (6.2.6)$$

for all  $y \in B(x, \rho_x)$  and  $\beta \leq f(y)$ .

Fix  $v_x \in V_x$ ; we have  $v_x = \sum_{i=1}^k \alpha_i v_i$  for suitable  $\alpha_i \in \mathbb{R}$ . From (6.2.6), we get

$$\left\langle v_x, \frac{y-x}{|y-x|+|\beta-f(x)|} \right\rangle \leq \frac{\sum_{i=1}^k |\alpha_i|}{8\sqrt{k}}$$

for all  $y \in B(x, \rho_x)$  and  $\beta \leq f(y)$ . On the other hand,

$$\|v_x\| = \left( \sum_{i=1}^k \alpha_i^2 \right)^{1/2} \geq \frac{\sum_{i=1}^k |\alpha_i|}{\sqrt{k}}.$$

Therefore

$$\left\langle v_x, \frac{y-x}{|y-x|+|\beta-f(x)|} \right\rangle \leq \frac{\|v_x\|}{8}$$

for all  $y \in B(x, \rho_x)$  and  $\beta \leq f(y)$ . Thus  $x \in BP_{f,k}^{\rho_x}$  and the proof is accomplished.  $\square$

In view of a rectifiability result for the sets  $BP_{f,k}$ , we begin with a technical result.

**Lemma 6.2.2** *Let  $a \in \mathbb{R}^N$ ,  $\rho > 0$  and  $x, y \in BP_{f,k}^\rho \cap B(a, \frac{\rho}{2})$  be such that  $d_G(V_x, V_y) < \frac{1}{8}$ ; then*

$$y \in \mathbb{R}^N \setminus C_{\frac{1}{4}}(x, V_x).$$

**Proof.** Since  $x, y \in B(a, \frac{\rho}{2})$ , we have  $x \in B(y, \rho)$  and  $y \in B(x, \rho)$ . Therefore, from (6.2.3) if  $v_x \in V_x \cap S^{N-1}$  we have

$$\langle v_x, y - x \rangle \leq \frac{1}{8}(\|y - x\| + |\beta - f(x)|) \quad \text{for all } \beta \leq f(y). \quad (6.2.7)$$

Similarly, for any  $v_y \in V_y \cap S^{N-1}$  we obtain

$$\langle v_y, y - x \rangle \leq \frac{1}{8}(\|y - x\| + |\beta - f(y)|) \quad \text{for all } \beta \leq f(x). \quad (6.2.8)$$

We have to distinguish two cases: if  $f(y) \geq f(x)$ , we choose  $\beta = f(x)$  in (6.2.7) to get

$$\langle v_x, y - x \rangle \leq \frac{1}{8}\|y - x\| \quad \forall v_x \in V_x \cap S^{N-1}.$$

Recalling (6.2.1), this implies that  $y \notin C_{\frac{1}{4}}(x, V_x)$ , as desired.

If  $f(y) \leq f(x)$ , we choose  $\beta = f(y)$  in (6.2.8) to get

$$\langle v_y, y - x \rangle \leq \frac{1}{8}\|y - x\| \quad \forall v_y \in V_y \cap S^{N-1}.$$

Since  $d_G(V_x, V_y) < \frac{1}{8}$ , for any  $v_x \in V_x \cap S^{N-1}$  there exists  $v_y = v_y(v_x) \in V_y \cap S^{N-1}$  such that  $\|v_x - v_y\| < \frac{1}{8}$ . Therefore, for any  $v_x \in V_x \cap S^{N-1}$  it holds

$$\langle v_x, y - x \rangle \leq \langle v_y, y - x \rangle + |\langle v_x - v_y, y - x \rangle| \leq \frac{1}{4}\|y - x\| \quad (6.2.9)$$

i.e.  $y \notin C_{\frac{1}{4}}(x, V_x)$ , as desired.  $\square$

We now fix  $R := 1/16$  and let  $V_1, \dots, V_m \in G(N, k)$  be given by (6.1.7). We thus divide  $BP_{f,k}^\rho$  into  $m$  sets

$$BP_{f,k}^\rho = \bigcup_{j=1}^m BP_{f,k}^{\rho,j} \quad (6.2.10)$$

where

$$BP_{f,k}^{\rho,j} = \{x \in BP_{f,k}^\rho \mid d_G(V_x, V_j) < 1/16\}.$$

For  $j = 1, \dots, m$  we denote by  $\pi_j$  the orthogonal projection  $\mathbb{R}^n \rightarrow V_j^\perp$ ; clearly,  $\pi_j(z) = z_{V_j^\perp} = z - z_{V_j}$ .

**Lemma 6.2.3** *The projection  $\pi_j : BP_{f,k}^{\rho,j} \cap B(a, \rho/2) \rightarrow \pi_j(BP_{f,k}^{\rho,j} \cap B(a, \rho/2))$  is invertible and its inverse map is Lipschitz continuous with Lipschitz constant at most 2.*

**Proof.** Let  $x, y \in BP_{f,k}^{\rho,j} \cap B(a, \rho/2)$  be fixed. We have  $d_G(V_x, V_y) < 1/8$  and Lemma 6.2.2 ensures that  $y \notin C_{1/4}(x, V_x)$ . Since  $d_G(V_x, V_j) < 1/16$ , by (6.2.2) we deduce that  $C_{1/2}(x, V_j) \subseteq C_{5/16}(x, V_j) \subseteq C_{1/4}(x, V_x)$  and, in particular, that  $y \notin C_{1/2}(x, V_j)$ . This implies that  $\|(y-x)_{V_j}\| \leq \frac{1}{2}\|y-x\|$ , whence

$$\|\pi_j(y) - \pi_j(x)\| = \|\pi_j(y-x)\| = \|(y-x) - (y-x)_{V_j}\| \geq \frac{1}{2}\|y-x\|.$$

This is enough to conclude.  $\square$

The rectifiability of the sets  $BP_{f,k}^\rho$  is now a consequence of Lemma 6.2.3.

## 6.2.2 Proof of main results

**Theorem 6.2.1** *The set  $BP_{f,k}^\rho \cap K$  is  $(N-k)$ -rectifiable for any  $\rho > 0$  and any compact set  $K \subset \mathbb{R}^N$ ; in particular*

$$\mathcal{H}^{N-k}(BP_{f,k}^\rho \cap K) < +\infty. \quad (6.2.11)$$

**Proof.** It will be sufficient to show that for any  $j = 1, \dots, m$  the set  $BP_{f,k}^{\rho,j} \cap K$  is  $k$ -rectifiable. Since  $K$  is compact, there exist  $a_1, \dots, a_h \in \mathbb{R}^N$  such that

$$BP_{f,k}^{\rho,j} \cap K \subset \bigcup_{i=1}^h (BP_{f,k}^{\rho,j} \cap B(a_i, \rho/2)).$$

By Lemma 6.2.3, for any  $i = 1, \dots, h$  the set  $BP_{f,k}^{\rho,j} \cap B(a_i, \rho/2)$  is the image of

$$\pi_j^{-1} : \pi_j(BP_{f,k}^{\rho,j} \cap B(a_i, \rho/2)) \rightarrow \mathbb{R}^N,$$

i.e. of a Lipschitz map defined on a bounded subset of  $V_j^\perp \equiv \mathbb{R}^{N-k}$  with Lipschitz constant at most 2. In particular,  $BP_{f,k}^{\rho,j} \cap B(a_i, \rho/2)$  is  $(N-k)$ -rectifiable and this allows to conclude.  $\square$

We can finally pass to the proof of our main results.

**Proof of Theorem 6.0.2.** It is an easy consequence of Lemma 6.2.1 and Theorem 6.2.1.  $\square$

Before passing to the proof of Theorem 6.0.3, we would like to discuss the relation between  $BP_f$  and the set of bad points  $BP_f^P$  considered in [47], namely,

$$BP_f^P := \{x \in \Omega \mid NL^P(x) \neq \{0\}\},$$

where  $NL^P(x) = \{ \xi \in N_{\text{hypo}(f)}^P(x, f(x)) \mid -\xi \in N_{\text{hypo}(f)}^P(x, f(x)) \}$ . From Remark 6.1.2 it is clear that  $BP_f^P \subseteq BP_f$ , but in general the two sets do not coincide.

However, the equality  $BP_f = BP_f^P$  holds under the assumptions of Theorem 6.0.3. Indeed, from Corollary 3.1 in [47] it follows that the hypograph of  $f|_{\Omega_P}$  has positive reach, where  $\Omega_P$  is the open set defined by  $\Omega_P := \Omega \setminus BP_f^P$ . Therefore (see [31, Proposition 6.2 and 4.2] and [40, Theorem 4.8 (12)]) one has

$$N_{\text{hypo}(f|_{\Omega_P})}^P(x, f|_{\Omega_P}(x)) = N_{\text{hypo}(f|_{\Omega_P})}^F(x, f|_{\Omega_P}(x)) \quad \text{for all } x \in \Omega_P.$$

and thus

$$N_{\text{hypo}(f)}^P(x, f(x)) = N_{\text{hypo}(f)}^F(x, f(x)) \quad \text{for all } x \in \Omega_P.$$

Consequently,  $NL(x) = NL^P(x)$  for all  $x \in \Omega_P$ . By the definition of  $BP_f^P$ , we have  $NL^P(x) = \{0\}$  for all  $x \in \Omega_P$ . This implies that  $NL(x) = \{0\}$  for all  $x \in \Omega_P$ , i.e.  $BP_f \cap \Omega_P = \emptyset$ . Thus,  $BP_f \subseteq BP_f^P$ , as claimed.  $\square$

**Proof of Theorem 6.0.3.** Recalling 6.0.2, we have  $\mathcal{H}^{N-1}(BP_{f,k}) = 0$  for all  $k \in \{2, 3, \dots, N\}$ . Since

$$BP_f = BP_{f,1} \cup \bigcup_{k=2}^N BP_{f,k},$$

the proof will be accomplished after proving that the set  $BP_{f,1}$  is locally  $(N-1)$ -rectifiable. From the definition (6.1.6), for every  $x \in BP_{f,1}$  the set

$$V_x = \{tv_x \mid v_x \in \mathbb{R}^N, \|v_x\| = 1 \text{ and } t \in \mathbb{R}\}$$

is a line along  $v_x$ . Therefore by [47, Lemma 4.3],  $(\pm v_x, 0) \in N_{\text{hypo}(f)}^P(x, f(x))$  is realized by a ball of radius  $\theta$ , i.e.

$$\langle \pm v_x, y - x \rangle \leq \frac{1}{2\theta} (\|y - x\|^2 + |\beta - f(x)|^2) \quad \forall y \in \Omega, \beta \leq f(x).$$

From the above inequality, reasoning as in the proof of Lemma 6.2.2 one can obtain that the following holds. If  $a \in \mathbb{R}^N$ ,  $\rho \in (0, \theta/8]$ ,  $x, y \in BP_{f,1} \cap B(a, \frac{\rho}{2})$  are such that  $d_G(V_x, V_y) < \frac{1}{8}$ , then

$$y \in \mathbb{R}^N \setminus C_{\frac{1}{4}}(x, V_x).$$

From this fact, the local  $(N-1)$ -rectifiability of  $BP_{f,1}$  follows (up to considering  $BP_{f,1}$  instead of  $BP_{f,k}^\rho$ ) as in the proof of Theorem 6.2.1.  $\square$

### 6.3 A counterexample

By virtue of Theorem 6.0.3, the set of bad points  $BP_f$  is locally  $(N - 1)$ -rectifiable provided the  $\theta$ -exterior sphere condition holds. On the contrary, an analogous  $(N - k)$ -rectifiability result does not hold for  $BP_{f,k}$ ; in other words, Theorem 6.0.2 cannot be refined to show that  $\mathcal{H}^{N-k}(BP_{f,k} \cap K) < \infty$  for any compact set  $K \subset \mathbb{R}^N$ . We are going to provide an example of a continuous function  $f : (-1, 1) \times (-1, 1) \rightarrow \mathbb{R}$  satisfying the  $\theta$ -exterior sphere condition with  $\theta = 1$  and such that  $\mathcal{H}^0(BP_{f,2} \cap K) = +\infty$  for any neighborhood  $K$  of the origin. It will be clear from the construction that what is missing is a uniform control on the radii of exterior balls (recall that, by Theorem 6.2.1,  $BP_{f,k}^\rho$  is locally  $(N - k)$ -rectifiable for any  $\rho > 0$ ).

Let  $\Omega := (-1, 1) \times (-1, 1)$ ; for  $n \in \mathbb{N}$ ,  $n \geq 0$  let us define  $x_n^+, x_n^- \in \overline{\Omega}$  by

$$x_n^+ := (2^{-n}, 0), \quad x_n^- := (-2^{-n}, 0).$$

We also set

$$c_n^+ := \frac{x_n^+ + x_{n+1}^+}{2} = (3 \cdot 2^{-n-2}, 0) \in \Omega, \quad c_n^- := \frac{x_n^- + x_{n+1}^-}{2} = (-3 \cdot 2^{-n-2}, 0) \in \Omega$$

and

$$r_n := \frac{\|x_n^+ - x_{n+1}^+\|}{2} = \frac{\|x_n^- - x_{n+1}^-\|}{2} = 2^{-n-2}.$$

Notice that the closed balls  $\overline{B(c_n^\pm, r_n)}$  are pairwise disjoint except for the case of consecutive balls, which instead are tangent, i.e., for any  $n \geq 1$  one has

$$\overline{B(c_n^+, r_n)} \cap \overline{B(c_{n-1}^+, r_{n-1})} = \{x_n^+\}, \quad \overline{B(c_n^-, r_n)} \cap \overline{B(c_{n-1}^-, r_{n-1})} = \{x_n^-\}.$$

Define  $f_1 : \Omega \rightarrow \mathbb{R}$  by

$$f_1(x) = \begin{cases} -\sqrt{r_n^2 - \|x - c_n^+\|^2} & \text{if } x \in B(c_n^+, r_n) \\ -\sqrt{r_n^2 - \|x - c_n^-\|^2} & \text{if } x \in B(c_n^-, r_n) \\ 0 & \text{if } x \in \Omega \setminus (\bigcup_n B(c_n^+, r_n) \cup \bigcup_n B(c_n^-, r_n)). \end{cases}$$

It is easily seen that  $f_1$  is continuous and that  $\{x_n^+, x_n^- : n \geq 1\} \subset BP_{f_1}$ ; more precisely

$$\begin{aligned} (1, 0) \in \partial^\infty f_1(x_n^+) & \text{ is realized by a ball of radius } r_{n-1} \\ (-1, 0) \in \partial^\infty f_1(x_n^+) & \text{ is realized by a ball of radius } r_n \\ (1, 0) \in \partial^\infty f_1(x_n^-) & \text{ is realized by a ball of radius } r_n \\ (-1, 0) \in \partial^\infty f_1(x_n^-) & \text{ is realized by a ball of radius } r_{n-1}. \end{aligned} \tag{6.3.1}$$

For any  $x = (\xi, \eta) \in \Omega$  we also define

$$f_2(x) = -\sqrt{\eta^2 - |\eta|} = -\sqrt{1 - (1 - |\eta|)^2}.$$



One can easily check that  $f_2$  is continuous on  $\Omega$  and that  $BP_{f_2} = \{(\xi, 0) : \xi \in (-1, 1)\}$ ; more precisely, for any  $\xi \in (-1, 1)$

$$(0, 1), (-1, 0) \in \partial^\infty f_2(\xi, 0) \text{ are realized by balls of radius 1.} \quad (6.3.2)$$

Notice also that  $f_1(x_n^\pm) = f_2(x_n^\pm) = 0$  for any  $n \geq 1$ . Therefore, the function  $f := \inf\{f_1, f_2\}$  is continuous on  $\Omega$  and  $f(x_n^\pm) = f_1(x_n^\pm) = f_2(x_n^\pm) = 0$ . Taking (6.3.1) and (6.3.2) into account we obtain that

$$(1, 0), (-1, 0), (0, 1), (0, -1) \in \partial^\infty f(x_n^\pm) \quad \text{for any } n \geq 1$$

whence

$$\{x_n^+, x_n^- : n \geq 1\} \subset BP_{f,2}$$

which in turn implies  $\mathcal{H}^0(BP_{f,2}) = \infty$ , as desired.  $\square$



# Chapter 7

## Appendix

### 7.1 Appendix A

In this section, under the assumptions  $(H_1)$  and  $(H_2)$  on (3.1.1) in Chapter 3, we prove some elementary estimates which are needed in Lemma 3.3.1, Lemma 3.3.2 and Lemma 3.3.3. For future use, we set

$$\begin{aligned} K_1 &= \max_{u \in \mathcal{U}} \|f(0, u)\|, \\ K_2 &= \max_{u \in \mathcal{U}} \|D_x f(0, u)\|, \\ L_2(s, \delta) &= L_1 e^{Ls} \delta + \frac{L_1(e^{Ls} - 1)K_1}{L} + K_2 \quad \text{for all } s, \delta \geq 0. \end{aligned} \quad (7.1.1)$$

**Lemma 7.1.1** *Let  $\alpha(\cdot) := y^{x,u}(\cdot)$  be the solution of (3.1.1). The following estimates hold true for all  $t > 0$ :*

- (i)  $\|\alpha(t) - x\| \leq \frac{(L\|x\| + K_1)(e^{Lt} - 1)}{L}$ .
- (ii)  $\|\alpha(t)\| \leq e^{Lt} \|x\| + \frac{(e^{Lt} - 1)K_1}{L}$ .
- (iii)  $\|f(\alpha(t), u(t))\| \leq L e^{Lt} \|x\| + e^{Lt} K_1$ .
- (iv)  $\|D_x f(\alpha(t), u(t))\| \leq L_2(t, \|x\|)$ .

**Proof.** Since  $\alpha(\cdot)$  is the solution of (3.1.1), for all  $t > 0$  we have

$$\begin{aligned} \|\alpha(t) - x\| &= \left\| \int_0^t f(\alpha(s), u(s)) ds \right\| \leq \int_0^t \|f(\alpha(s), u(s))\| ds \\ &\leq \int_0^t \|f(\alpha(s), u(s)) - f(x, u(s))\| ds \\ &\quad + \int_0^t \|f(x, u(s)) - f(0, u(s))\| ds + \int_0^t \|f(0, u(s))\| ds \\ &\leq L \int_0^t \|\alpha(s) - x\| ds + L \|x\| t + K_1 t. \end{aligned}$$

Applying Gronwall's inequality we obtain

$$\|\alpha(t) - x\| \leq \frac{(L\|x\| + K_1)(e^{Lt} - 1)}{L}, \quad (7.1.2)$$

whence

$$\|\alpha(t)\| \leq e^{Lt}\|x\| + \frac{(e^{Lt} - 1)K_1}{L}. \quad (7.1.3)$$

Recalling the condition  $(H_2)$ , we obtain

$$\|f(\alpha(t), u(t))\| \leq Le^{Lt}\|x\| + e^{Lt}K_1 \quad (7.1.4)$$

and also

$$\|D_x f(\alpha(t), u(t))\| \leq L_1 e^{Lt}\|x\| + \frac{L_1(e^{Lt} - 1)K_1}{L} + K_2. \quad (7.1.5)$$

The proof is concluded.  $\square$

In the next Lemma, we will give some estimates related to the limiting adjoint trajectories  $M^T(\cdot)$  in Chapter 3.

**Lemma 7.1.2** *Let  $x \in \mathcal{S}^c$ , set  $r = T(x) > 0$ , and take  $\bar{x} \in \mathcal{S}_x$  and  $M(\cdot) \in \mathcal{M}_{\bar{x}}$ . Then*

$$(i) \quad \|M(t)\| \leq e^{L_2(t, \|x\|)t} \text{ for all } t \in [0, r],$$

$$(ii) \quad \|M(t)^{-1}\| \leq e^{L_2(t, \|x\|)t} \text{ for all } t \in [0, r].$$

**Proof.** Let  $x_n \rightarrow x$ ,  $\{\bar{u}_n\} \subset \mathcal{U}_{\text{ad}}$  be such that  $\{y^{x_n, \bar{u}_n}(\cdot)\} \subset \mathcal{T}_{\bar{x}}$  and  $M(\cdot, x_n, \bar{u}_n)$  converges to  $M(\cdot)$  uniformly on  $[0, T(x)]$ . By (iv) in Lemma 7.1.1 and Theorem 2.2.1, p. 23, in [13], we obtain that for all  $w \in \mathbb{R}^N$

$$\|M(t, x_n, \bar{u}_n)w\| \leq e^{[L_1 e^{Lt}\|x\| + \frac{L_1(e^{Lt} - 1)K_1}{L} + K_2]t} \|w\|.$$

Taking  $n \rightarrow \infty$  we conclude the proof of (i).

The proof of (ii) proceeds exactly as the proof of (i), by replacing  $M(\cdot, x_n, \bar{u}_n)$  with  $M(\cdot, x_n, \bar{u}_n)^{-1}$ .  $\square$

The following result is essentially Theorem 2.2.4, pp. 25, 26 in [13].

**Lemma 7.1.3** *Let  $A_1, A_2 : [0, T] \rightarrow \mathcal{M}^{N \times N}$  be matrices with  $L^\infty$ -entries, and set  $\|A_i\| = L_i$ ,  $i = 1, 2$ . Let  $M_1, M_2$  be the fundamental solution of, respectively,*

$$\begin{aligned} \dot{p}(t) &= A_1(t)p(t), & p(0) &= \mathbb{I}^{N \times N} \\ \dot{p}(t) &= A_2(t)p(t), & p(0) &= \mathbb{I}^{N \times N}. \end{aligned}$$

Then, for every  $t \in [0, T]$  and every unit vector  $v \in \mathbb{R}^N$  we have

$$\|(M_2(t) - M_1(t))v\| \leq e^{(L_1 + L_2)t} \int_0^t \|A_2(s) - A_1(s)\| ds.$$

## 7.2 Appendix B

The first Lemma is to prove that the angle of a wedged cone  $H_C^+$  which is generated by a compact set  $C \subset \mathbb{R}^N \setminus \{0\}$  is far from  $\pi$ .

**Lemma 7.2.1** *Let  $C \in \mathbb{R}^N$  be a compact set which does not contain 0. We denote the positive cone generated by  $C$  as*

$$H_C^+ = \text{span}^+(C) = \left\{ \sum_{i=1}^k \alpha_i c_i \mid c_i \in C \text{ and } \alpha_i \geq 0 \right\}.$$

Assume that  $H_C^+$  is wedged. Then:

i)  $H_C^+$  is closed.

ii) There exists a constant  $\delta_0 > 0$  such that for all  $0 \neq x_1, x_2 \in H_C^+$ , it holds

$$\left\langle \frac{x_1}{\|x_1\|}, \frac{x_2}{\|x_2\|} \right\rangle > -1 + \delta_0. \quad (7.2.1)$$

**Proof of (i).** Let a sequence  $\{x_n\} \subset H_C^+$  converge to  $x$ . We need to prove that  $x \in H_C^+$ . By Caratheodory theorem, we can write

$$x_n = \sum_{i=1}^{N+1} \alpha_n^i c_n^i, \text{ where } \alpha_n^i \geq 0, c_n^i \in C. \quad (7.2.2)$$

Assume without loss of generality that  $\lim_{n \rightarrow \infty} c_n^i = \bar{c}^i \in C$  for all  $i \in \{1, 2, \dots, N+1\}$ .

If  $\sum_{i=1}^{N+1} \alpha_n^i$  is unbounded, we extract subsequences  $\{\alpha_{n_k}^i\} \subseteq \{\alpha_n^i\}$  such that

$$\frac{\alpha_{n_k}^i}{\sum_{i=1}^{N+1} \alpha_{n_k}^i} = \bar{\alpha}^i \geq 0 \text{ and } \lim_{n_k \rightarrow \infty} \sum_{i=1}^{N+1} \alpha_{n_k}^i = +\infty.$$

Therefore, from (7.2.2) and  $\lim_{n \rightarrow \infty} x^n = x$  we get

$$\sum_{i=1}^{N+1} \bar{\alpha}^i \bar{c}^i = \lim_{n_k \rightarrow \infty} \frac{x_{n_k}}{\sum_{i=1}^{N+1} \alpha_{n_k}^i} = 0. \quad (7.2.3)$$

Note that  $\bar{\alpha}^i \geq 0$ ,  $\sum_{i=1}^{N+1} \bar{\alpha}^i = 1$  and  $\bar{c}^i \neq 0$ . We recall (7.2.3) to obtain that the cone  $H_C^+$  contains at least one line. This is a contradiction.

Thus  $\sum_{i=1}^{N+1} \alpha_n^i$  is bounded. It implies that the sequences  $\{\alpha_n^i\}$  are bounded for all  $i \in \{1, 2, \dots, N+1\}$  since  $\alpha_n^i \geq 0$ . We extract subsequences  $\{\alpha_{n_k}^i\} \subseteq \{\alpha_n^i\}$  such that

$$\lim_{n_k \rightarrow \infty} \alpha_{n_k}^i = \bar{\alpha}^i \geq 0 \text{ for all } i \in \{1, 2, \dots, N+1\}.$$

From the above equality and (7.2.2), we have

$$x = \lim_{n \rightarrow \infty} x_n = \lim_{n_k \rightarrow \infty} x_{n_k} = \lim_{n_k \rightarrow \infty} \sum_{i=1}^{N+1} \alpha_{n_k}^i c_{n_k}^i = \sum_{i=1}^{N+1} \overline{\alpha^i} \overline{c^i}.$$

This implies  $x \in H_C^+$ .

**Proof of (ii).** Assume by contradiction that there exist two sequences  $\{x_1^n\}$ ,  $\{x_2^n\}$  contained in  $H_C^+$  such that  $\|x_1^n\| = \|x_2^n\| = 1$  and

$$\lim_{n \rightarrow \infty} \langle x_1^n, x_2^n \rangle = -1. \quad (7.2.4)$$

Assume without loss of generality that  $\lim_{n \rightarrow \infty} x_1^n = \overline{x_1}$  and  $\lim_{n \rightarrow \infty} x_2^n = \overline{x_2}$ . Recalling (7.2.4), we obtain that  $-\overline{x_1} = \overline{x_2}$ . Moreover, since  $H_C^+$  is closed, we have  $x_1, x_2 \in H_C^+$ . Therefore  $H_C^+$  contains at least one line. This is a contradiction.  $\square$

The following Proposition provides a sufficient condition for the strict convexity of a set.

**Proposition 7.2.1** *Let  $K \subset \mathbb{R}^N$  be compact and assume that there exist  $\gamma > 0$  and  $p > 1$  with the following property: for every  $x \in \partial K$ , there exists  $\zeta \neq 0$  such that for every  $y \in K$  one has*

$$\langle \zeta, y - x \rangle \leq -\gamma \|\zeta\| \|y - x\|^p. \quad (7.2.5)$$

*Then  $K$  is convex (with nonempty interior) and (7.2.5) is satisfied by all  $\zeta \in N_K(x)$  for each  $x \in \partial K$ .*

**Proof.** We show first that  $K$  is convex. To this aim, assume by contradiction that there exist three points  $x_1 \neq x_2 \in K$  such that the segment  $[x_1, x_2]$  is not contained in  $K$ . Let  $0 < t < 1$  be such that  $x_t = (1-t)x_1 + tx_2 \in \partial K$  and let  $\zeta \neq 0$  be such that (7.2.5) holds with  $x_t$  in place of  $x$ . Obviously,  $\zeta \perp x_2 - x_1$  and this is a contradiction. It is also easy to see that  $K$  must have nonempty interior. Since  $K$  is convex with nonempty interior, for each  $x \in \partial K$  the normal cone  $N_K(x)$  is pointed and so it is the convex hull of its exposed rays (see [54]).

We now see Theorem 4.6 in [27] and see that for every unit vector  $w$  belonging to our exposed ray of  $N_K(x)$ , there exists a sequence  $x_n \rightarrow x$  such that

$$N_K(x_n) = \mathbb{R}^+ w_n, \|w_n\| = 1.$$

Of course (7.2.5) holds with  $x_n$  (resp,  $w_n$ ) in place of  $x$  (resp,  $\zeta$ ), so that by passing to the limit,  $w$  also satisfies (7.2.5). By taking convex combinations, we conclude the proof.  $\square$

The second lemma is necessary to use Theorem 5.2.2 in the proof of the main theorem in Chapter 4.

**Lemma 7.2.2** *Let  $\Omega \subseteq \mathbb{R}^N$  be open and let  $g : \Omega \rightarrow \mathbb{R}$  be continuous. Assume that  $\text{hypo}(g)$  satisfies the  $\theta$ -external sphere condition where  $\theta : \Omega \rightarrow [0, +\infty)$  is continuous. Let, for all  $x \in \Omega$ ,  $r_x > 0$  be such that  $\overline{B}_N(x, r_x) \subset \Omega$ . Then*

- i) The hypograph of the restricted function  $g|_{B_N(x, r_x)} : B_N(x, r_x) \rightarrow \mathbb{R}$  satisfies the  $\theta_x$ -external sphere condition with  $\theta_x = \max\{\theta(y) \mid y \in \overline{B}_N(x, r_x)\}$ .*
- ii)  $BP_g \cap B_N(x, r_x) = BP_{g|_{B_N(x, r_x)}}$ .*

**Proof of (i).** Let  $z \in B_N(x, r_x)$ , there exists a vector  $0 \neq \xi \in N_{\text{hypo}(g)}^P(z, g(z))$  realized by a ball of radius  $\theta(z)$ , i.e, for all  $y \in \Omega$  and for  $\beta \leq g(y)$ , it holds

$$\left\langle \frac{\xi}{\|\xi\|}, (y, \beta) - (z, g(z)) \right\rangle \leq \theta(z) (\|y - z\|^2 + |\beta - g(z)|^2). \quad (7.2.6)$$

Thus, for all  $y \in B_N(x, r_x)$  and for all  $\beta \leq g|_{B_N(x, r_x)}(y)$ , we have

$$\left\langle \frac{\xi}{\|\xi\|}, (y, \beta) - (z, g|_{B_N(x, r_x)}(z)) \right\rangle \leq \theta_x (\|y - z\|^2 + |\beta - g|_{B_N(x, r_x)}(z)|^2). \quad (7.2.7)$$

The proof is completed.  $\square$

*Proof of (ii).* It is similar to the previous proof. Indeed, if  $0 \neq \xi \in N_{\text{hypo}(g)}^P(z, g(z))$  then  $0 \neq \xi \in N_{\text{hypo}(g|_{B_N(x, r_x)})}^P(z, g|_{B_N(x, r_x)}(z))$ . Therefore,  $BP_g \cap B_N(x, r_x) \subseteq BP_{g|_{B_N(x, r_x)}}$ .

We are going now to prove  $BP_{g|_{B_N(x, r_x)}} \subseteq BP_g$ . It is sufficient to prove that if  $0 \neq \xi \in N_{\text{hypo}(g|_{B_N(x, r_x)})}^P(z, g|_{B_N(x, r_x)}(z))$  then  $0 \neq \xi \in N_{\text{hypo}(g)}^P(z, g(z))$ . Indeed,  $0 \neq \xi \in N_{\text{hypo}(g|_{B_N(x, r_x)})}^P(z, g|_{B_N(x, r_x)}(z))$ , i.e, there exists a constant  $\sigma > 0$  such that for all  $y \in B_N(x, r_x)$  and for all  $\beta \leq g|_{B_N(x, r_x)}(y)$ , it holds

$$\left\langle \frac{\xi}{\|\xi\|}, (y, \beta) - (z, g|_{B_N(x, r_x)}(z)) \right\rangle \leq \sigma (\|y - z\|^2 + |\beta - g|_{B_N(x, r_x)}(z)|^2). \quad (7.2.8)$$

Therefore, for all  $y \in B_N(x, r_x)$  and for all  $\beta \leq g(y)$ , one has

$$\left\langle \frac{\xi}{\|\xi\|}, (y, \beta) - (z, g(z)) \right\rangle \leq \sigma (\|y - z\|^2 + |\beta - g(z)|^2). \quad (7.2.9)$$

Since  $z \in B_N(x, r_x)$ , one can easily get from (7.2.9) that there exists a constant  $\sigma_1 > 0$  such that the inequality

$$\left\langle \frac{\xi}{\|\xi\|}, (y, \beta) - (z, g(z)) \right\rangle \leq \sigma_1 (\|y - z\|^2 + |\beta - g(z)|^2)$$

holds for all  $y \in \Omega$  and for all  $\beta \leq g(y)$ .

It means that  $\xi \in N_{\text{hypo}(g)}^P(z, g(z))$ . The proof is completed.  $\square$

The last one is a technical lemma which is used in Chapter 4.

**Lemma 7.2.3** *Let  $g : \Omega \rightarrow \mathbb{R}$  be continuous and let  $\gamma > 0$ . We denote by  $g^\gamma : \gamma\Omega \rightarrow \mathbb{R}$ , the  $\gamma$ -stretched function of  $g$ , as follows:*

$$g^\gamma(y) = g\left(\frac{y}{\gamma}\right) \quad \text{for all } y \in \gamma\Omega.$$

*Assume that  $(\xi, \lambda)$  is a proximal normal vector to  $\text{hypo}(g)$  at  $(x, g(x))$  realized by a ball of radius  $\rho$ . Then  $(\frac{\xi}{\gamma}, \lambda)$  is a proximal normal vector to  $\text{hypo}(g^\gamma)$  at  $(\gamma x, g^\gamma(\gamma x))$  realized by a ball of radius  $\rho \frac{\gamma^2}{(1+\gamma^2)^{3/2}}$ .*

**Proof.** For all  $z \in \Omega$  and for all  $\beta \leq g(z)$ , it holds

$$\left\langle \frac{(\xi, \lambda)}{\|(\xi, \lambda)\|}, (z, \beta) - (x, g(x)) \right\rangle \leq \frac{1}{2\rho} (\|z - x\|^2 + |\beta - g(x)|^2).$$

Equivalently, for all  $\gamma z \in \gamma\Omega$  and for all  $\beta \leq g^\gamma(\gamma z)$ , it holds

$$\left\langle \frac{(\frac{\xi}{\gamma}, \lambda)}{\|(\frac{\xi}{\gamma}, \lambda)\|}, (\gamma z, \beta) - (\gamma x, g^\gamma(\gamma x)) \right\rangle \leq \frac{1}{2\rho} \left( \frac{1}{\gamma^2} \|\gamma z - \gamma x\|^2 + |\beta - g^\gamma(\gamma x)|^2 \right). \quad (7.2.10)$$

Since  $\|(\xi, \lambda)\| \leq \sqrt{\gamma^2 + 1} \|(\frac{\xi}{\gamma}, \lambda)\|$ , one can easily get from (7.2.10) that for all  $\bar{z} = \gamma z \in \gamma\Omega$  and for all  $\beta \leq g^\gamma(\bar{z})$ , it holds

$$\left\langle \frac{(\frac{\xi}{\gamma}, \lambda)}{\|(\frac{\xi}{\gamma}, \lambda)\|}, (\bar{z}, \beta) - (\gamma x, g^\gamma(\gamma x)) \right\rangle \leq \frac{1}{2\rho \frac{\gamma^2}{(1+\gamma^2)^{3/2}}} (\|\bar{z} - \gamma x\|^2 + |\beta - g^\gamma(\gamma x)|^2) \quad . \quad (7.2.11)$$

The proof is completed.  $\square$

The following result is an immediate consequence of the previous lemma.

**Corollary 7.2.1** *for every  $\gamma > 0$ , it holds*

$$BP_{g^\gamma} = \gamma BP_g.$$

### 7.3 Appendix C

We will give here some inequalities on one variable functions which are used in Chapter 4.

**Lemma 7.3.1** *Let  $K : (a, b) \rightarrow [0, 1]$  be measurable and let  $k \in \mathbb{N}$ . Then*

$$\int_a^b (t - a)^k K(t) dt \geq \frac{1}{k+1} \left( \int_a^b K(t) dt \right)^{k+1},$$

and

$$\int_a^b (b - t)^k K(t) dt \geq \frac{1}{k+1} \left( \int_a^b K(t) dt \right)^{k+1}.$$



**Proof.** Indeed,

$$\int_a^b K(t)dt = k! \int_a^b \int_{t_k}^b \dots \int_{t_1}^b K(t_0)dt_0 \dots dt_k$$

Since  $K(t) \in [0, 1]$  for a.e.  $t \in [0, 1]$ , we obtain that

$$\int_a^b K(t)dt = k! \int_a^b K(t_k) \int_{t_k}^b K(t_{k-1}) \dots \int_{t_1}^b K(t_0)dt_0 \dots dt_k$$

By using induction, one can easily prove that

$$\int_a^b K(t_k) \int_{t_k}^b K(t_{k-1}) \dots \int_{t_1}^b K(t_0)dt_0 \dots dt_k = \frac{1}{k+1} \left( \int_a^b K(t)dt \right)^{k+1}$$

The proof is completed.  $\square$

**Lemma 7.3.2** *Let  $f : [a, b] \rightarrow \mathbb{R}$  be of class  $C^1$  and fix  $k \geq 0$ . Assume that there exists  $C \in \mathbb{R}$  such that*

$$|f'(s)| \geq C(s-a)^k \quad \forall s \in [a, b]. \quad (7.3.1)$$

*Then, either  $f$  has no zeros in  $(a, b)$  and then, for all  $s \in (a, b)$  either*

$$|f(s)| \geq \frac{C}{k+1}(s-a)^{k+1} \quad \text{or} \quad |f(s)| \geq \frac{C}{k+1}(b-s)^{k+1}$$

*or there exists  $c \in (a, b)$  such that  $g(c) = 0$  and then, for all  $s \in [a, c]$ ,*

$$|f(s)| \geq \frac{C}{k+1}(c-s)^{k+1}$$

*and for all  $s \in [c, b]$*

$$|f(s)| \geq \frac{C}{k+1}(s-c)^{k+1}.$$

*The same conclusion hold if (7.3.1) is substituted by*

$$|f'(s)| \geq C(b-s)^k \quad \forall s \in [a, b].$$

**Proof.** Observe that by our assumptions  $f'$  has constant sign on  $[a, b]$ . We treat the case  $f' \geq 0$ , while the other one can be obtained by taking  $-f$ . So (7.3.1) now reads as

$$f'(s) \geq C(s-a)^k \quad \forall s \in [a, b].$$

If  $f$  has no zeros, we have two cases, namely  $f(s) > 0$  for all  $s \in (a, b)$  or  $f(s) < 0$  for all  $s \in (a, b)$ . For the first case

$$f(s) - f(a) = \int_a^s f'(t)dt \geq C \int_a^s (t-a)dt = \frac{C}{k+1}(s-a)^{k+1},$$

therefore,

$$f(s) \geq \frac{C}{k+1}(s-a)^{k+1}.$$

In the second case,

$$\begin{aligned} f(b) - f(s) &= \int_s^b f'(t)dt \geq C \int_s^b (t-a)^k dt \\ &= \frac{C}{k+1} [(b-a)^{k+1} - (s-a)^{k+1}] \geq \frac{C}{k+1} (b-s)^{k+1}. \end{aligned}$$

Therefore,

$$f(s) \leq f(b) - \frac{C}{k+1}(b-s)^{k+1} \leq -\frac{C}{k+1}(b-s)^{k+1}.$$

Assume now that there exists  $c \in (a, b)$  such that  $f(c) = 0$ . Then, for all  $c \in [a, c]$  we have

$$-f(s) = f(c) - f(s) = \int_s^c f'(t)dt \geq \frac{C}{k+1}(c-s)^{k+1},$$

while for all  $s \in [c, b]$  we have

$$\begin{aligned} f(s) &= f(s) - f(c) = \int_c^s f'(t)dt \geq \frac{C}{k+1} [(s-a)^{k+1} - (c-a)^{k+1}] \\ &\geq \frac{C}{k+1}(s-c)^{k+1}. \end{aligned}$$

and the proof is completed. □

# Bibliography

- [1] G. Alberti, L. Ambrosio, P. Cannarsa, On the singularities of convex functions, *Manuscr. Math.* **76** (1992), 421-435.
- [2] G. Alberti, *On the structure of singular sets of convex functions*, Calc. Var. Partial Differential Equations 2 (1994), 17-27.
- [3] P. Albano & P. Cannarsa, *Singularities of semiconcave functions in Banach spaces*, in: Stochastic Analysis, Control, Optimization and Applications, W. M. McEneaney, G. G. Yin & Q. Zhang (eds), Birkhauser, Boston, 1999, pp. 171-190.
- [4] P. Albano & P. Cannarsa, *Structural properties of singularities of semiconcave functions*, Annali Scuola Norm. Sup. Pisa Sci. Fis. Mat. 28 (1999), 719-740.
- [5] L. Ambrosio, *Corso introduttivo alla Teoria Geometrica della Misura ed alle Superfici Minime*, Scuola Normale Superiore, Pisa (1997).
- [6] L. Ambrosio & P. Tilli, *Topics on analysis in metric spaces*, Oxford Lecture Series in Mathematics and its Applications, 25. Oxford University Press, Oxford, 2004.
- [7] L. Ambrosio, P. Cannarsa & H. M. Soner, *On the propagation of singularities of semi-convex functions*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 20 (1993), 597-616.
- [8] L. Ambrosio, N. Fusco, D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford Science Publications, Clarendon Press, Oxford (2000).
- [9] J.P. Aubin, A. Cellina, *Differential inclusions*, Springer-Verlag, Berlin, 1984.
- [10] J. P. Aubin, H. Frankowska, *Set-Valued Analysis*, Birkhäuser, Boston (1990).

- 
- [11] M. Bardi, I. Capuzzo-Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton–Jacobi–Bellman Equations*, Birkhäuser, Boston (1997).
- [12] U. Boscain, B. Piccoli, *Optimal syntheses for control systems on 2-D manifolds*, Springer, Berlin (2004).
- [13] A. Bressan, B. Piccoli, *Introduction to the mathematical theory of control*, AIMS, Springfield (2007).
- [14] A. Canino, On  $p$ -convex sets and geodesics, *J. Differential Equations* **75** (1988), 118–157.
- [15] A. Canino, Local properties of geodesics on  $p$ -convex sets, *Ann. Mat. Pura Appl.* (4) **159** (1991), 17–44.
- [16] P. Cannarsa, P. Cardaliaguet, Perimeter estimates for reachable sets of control systems, *J. Convex Anal.* **13** (2006), pp. 253–267.
- [17] P. Cannarsa, H. Frankowska, Interior sphere property of attainable sets and time optimal control problems, *COCV.12* (2006), pp. 350–370.
- [18] P. Cannarsa, M Castelpietra, Lipschitz continuity and local semiconcavity for exit time problems with state constraints, *J. Differential equation* **245** (2008), 616–636.
- [19] P. Cannarsa, Khai T. Nguyen, External sphere condition and time optimal control for differential inclusions, in preparation.
- [20] P. Cannarsa, C. Sinestrari, Convexity properties of the minimum time function, *Calc. Var.* **3** (1995), 273–298.
- [21] P. Cannarsa, C. Sinestrari, *Semiconcave functions, Hamilton–Jacobi Equations, and Optimal Control*, Birkhäuser, Boston (2004).
- [22] P. Cannarsa, P. R. Wolenski, Semiconcavity of the value function for a class of differential inclusion, *Discrete and Continuous Dynamical Systems-Series*, to appear.
- [23] P. Cardaliaguet, On the regularity of semipermeable surfaces in control theory with application to the optimal exit-time problem, I, II, *SIAM J. Control Optim.* **35** (1997), 1638–1671.
- [24] F. H. CLARKE, *Optimization and Nonsmooth Analysis*, In *Applied Mathematics*.
- [25] F. H. Clarke, Yu. S. Ledyaev, R. J. Stern, P. R. Wolenski, *Nonsmooth Analysis and Control Theory*, Springer, New York (1998).

- 
- [26] F. H. Clarke, R. J. Stern, P. R. Wolenski, Proximal smoothness and the lower- $\mathcal{C}^2$  property, *J. Convex Anal.* **2** (1995), 117-144.
- [27] G. Colombo, A. Marigonda, Differentiability properties for a class of non-convex functions, *Calc. Var.* **25** (2006), 1-31.
- [28] G. Colombo, A. Marigonda, Singularities for a class of non-convex sets and functions, and viscosity solutions of some Hamilton-Jacobi equations, *J. Convex Anal.* **15** (2008), 105-129.
- [29] G. Colombo, A. Marigonda, P. R. Wolenski, Some new regularity properties for the minimal time function, *Siam J. Control Optim.* **44** (2006), 2285-2299.
- [30] G. Colombo, A. Marigonda, P. R. Wolenski, A representation formula for the generalized gradient for a class of non-Lipschitz functions, in preparation
- [31] G. Colombo, V. V. Goncharov, Variational Inequalities and Regularity Properties of Closed Sets in Hilbert Spaces, *J. Convex Anal.* **8** (2001), 197-221.
- [32] G. Colombo, Khai T. Nguyen, Quantitative isoperimetric inequalities for a class of nonconvex sets, *Cal. Var.* **37** (2010), 141-166.
- [33] G. Colombo, Khai T. Nguyen, On the structure of the minimum time function, *Siam J. Control Optim.* **48** (2010), 4776-4814.
- [34] G. Colombo, Khai T. Nguyen, The minimum time function around the origin, in preparation.
- [35] G. Colombo, L. Thibault, Prox-regular sets and applications, in *Handbook of non-convex analysis*, D.Y.Gao and D. Motreanu editors, Wiley, to appear.
- [36] P. Erdős, *Some remarks on the measurability of certain sets*, Bull. Amer. Math. Soc. 51 (1945), 728-731.
- [37] L. C. Evans, R. F. Gariepy, *Measure Theory and Fine Properties of Functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton (1992).
- [38] K. Falconer, *Fractal geometry. Mathematical foundations and applications*, Wiley, Chichester (1990).
- [39] H. Federer, Curvature Measures, *Trans. Amer. Math. Soc.* **93** (1959), 418-491.
- [40] H. Federer, *Geometric Measure Theory*, Springer, 1969.

- [41] H. Hermes, J. P. LaSalle, *Functional analysis and time optimal control*, Academic Press, New York-London (1969).
- [42] M. Krastanov, M. Quincampoix, Local small time controllability and attainability of a set for nonlinear control system, *ESAIM Control Optim. Calc. Var.* **6** (2001), 499-516.
- [43] A. Marino, M. Tosques, Some variational problems with lack of convexity, in *Methods of Nonconvex Analysis*, Proceedings from CIME, Varenna 1989 (A. Cellina, ed.), Springer, Berlin (1990), 58-83.
- [44] B. S. Mordukhovich, *Variational Analysis and Generalized Differentiation I*, Springer, Berlin (2006).
- [45] A. Marigonda, Second order conditions for the controllability of nonlinear systems with drift, *Comm. in Pure and Applied Analysis* **5** (2006), 861-885 .
- [46] P. Mattila, *Geometry of sets and measures in Euclidean spaces. Fractals and rectifiability*, Cambridge Studies in Advanced Mathematics, 44, Cambridge University Press, Cambridge, 1995.
- [47] Khai T. Nguyen, Hypographs satisfying an external sphere condition and the regularity of the minimum time function, *Journal of Mathematical Analysis and Applications (JMAA)* **372** (2010), 611–628.
- [48] Khai T. Nguyen, D. Vittone, Rectifiability for special singularities of non-lipschitz functions, submitted.
- [49] C. Nour, R. J. Stern and J. Takche, Proximal smoothness and the exterior sphere condition, *J. Convex Anal.* **16** (2009), 501-514.
- [50] C. Nour, R. J. Stern and J. Takche, The  $\theta$ -exterior sphere condition,  $\varphi$ -convexity, and local semiconcavity, *Nonlinear Anal.* **73** (2010) 573-589.
- [51] P. Nistri, M. Quincampoix, On open-loop and feedback attainability of a closed set for nonlinear control systems, *J. Math. Anal. Appl.* **270** (2002), 474-487.
- [52] R. A. Poliquin, R. T. Rockafellar, Prox-regular functions in variational analysis, *Trans. Am. Math. Soc.* **348** (1996), 1805-1838.
- [53] R. A. Poliquin, R. T. Rockafellar, L. Thibault, Local differentiability of distance functions, *Trans. Amer. Math. Soc.* **352** (2000), 5231-5249.

- [54] R. T. Rockafellar, Clarke's tangent cones and the boundaries of closed sets in  $\mathbb{R}^n$ , *Nonlinear Analysis, Theory, Methods and Applications*. 3 (1979), 145-154.
- [55] R. T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton (1972).
- [56] P. Soravia, Optimality principles and representation formulas for viscosity solutions of Hamilton-Jacobi equations. II. Equations of control problems with state constraints, *Differential and Integral Equations* 12 (1999), 275-293.
- [57] P. Soravia, On Aronsson equation and deterministic optimal control, *Appl Math Optim* 59 (2009), 175-201.
- [58] R. T. Rockafellar, R. J.-B. Wets, *Variational Analysis*, Springer, Berlin (1998).
- [59] M. Tosques, Equivalence between generalized gradients and subdifferentials for a suitable class of lower semicontinuous functions, in *Generalized Convexity*, Proceedings, Pécs 1992 (S. Komlósi, T. Rapcsák, S. Schaible eds.), Springer, Berlin (1994), 116-133.
- [60] V. Veliov, Lipschitz continuity of the value function in optimal control, *J. Optim. Theory Appl.* 94 (1997), 335-363.
- [61] L. Veselý, *On the multiplicity points of monotone operators on separable Banach spaces*, Comment. Math. Univ. Carolin. 27 (1986), 551-570.
- [62] L. Veselý, *On the multiplicity points of monotone operators on separable Banach spaces*, Comment. Math. Univ. Carolin. 28 (1987), 295-299.
- [63] L. Zajíček, *On the points of multiplicity of monotone operators*, Comment. Math. Univ. Carolin. 19 (1978), 179-189.
- [64] L. Zajíček, *On the differentiation of convex functions in finite and infinite dimensional spaces*, Czechoslovak Math. J. 29 (1979), 340-348.
- [65] P. R. Wolenski, Z. Yu, Proximal analysis and the minimal time function, *SIAM J. of Control and Opt.* 36 (1998), 1048-1072.